



HAL
open science

A Short Note On The Solutions Of Fermat's Diohantine Equation In Some Neutrosophic Rings

Katy D Ahmad, Mikail Bal, Malath Aswad

► **To cite this version:**

Katy D Ahmad, Mikail Bal, Malath Aswad. A Short Note On The Solutions Of Fermat's Diohantine Equation In Some Neutrosophic Rings. *journal of neutrosophic and fuzzy systems*, 2022, pp.55-60. 10.54216/JNFS.010201 . hal-03524528

HAL Id: hal-03524528

<https://hal.science/hal-03524528v1>

Submitted on 28 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A Short Note On The Solutions Of Fermat's Diophantine Equation In Some Neutrosophic Rings

Katy D. Ahmad^{1,*}, Mikail Bal², Malath Aswad³

¹Islamic University Of Gaza, Palestine

²Gaziantep University, Turkey

³Malath Aswad, Albaath university, Syria

Emails : katyon765@gmail.com^{1,*}, mikailbal46@hotmail.com², malaz.aswad@yahoo.com³

Abstract: This paper is dedicated to study the concept of Fermat's triples in rings. Also, it determines the possible Fermat's triples in the neutrosophic ring of integers $Z(I)$. Also, it discussed these triples in several finite commutative rings such as Z_n .

Keywords: Neutrosophic Ring, Fermat's Diophantine equation, Neutrosophic Integer

1.Introduction

Number Theory is always a rich material for pure mathematical ideas. Where concepts such as ideals, and cyclic groups were derived from number theory [16].

The equation $X^n + Y^n = Z^n; n = 2$, appeared firstly from Pythagoras famous theorem, and then it was generalized to the integers.

In the literature, Fermat presented his famous conjecture that the Diophantine equation $X^n + Y^n = Z^n; n \geq 3$ has only trivial solutions in the ring of integers. This conjecture was solved completely in 1993 by Andrew Wiles [16].

Neutrosophy is a new kind of generalized logic founded by Smarandache [1]. It represents a useful tool in the study of spaces [2], rings [6,7,9], number theory [15], and geometry [10].

In this work, we extend the Fermat's equation $X^n + Y^n = Z^n; n \geq 3$ to Neutrosophic algebraic rings, where we define the concept of Fermat's triples in any ring as the solutions of the previous equation, and

we handle some special cases in several finite rings such as Z_n (integers modulo n) and neutrosophic rings of integers.

This work is motivated by finding novel applications and connections between number theory and algebraic structures. Also, it will provide some new open questions, which will represent the future of this theory.

Main discussion

Definition: [5]

Let $(R, +, \times)$ be a ring, $R(I) = \{a + bI \mid a, b \in R\}$ is called the neutrosophic ring where I is a neutrosophic element with condition $I^2 = I$.

Definition 2:

Let R be a ring, $F = (X, Y, Z)$ be a triple, where $X, Y, Z \in R$. F is called a general Fermat's triple if and only if $X^n + Y^n = Z^n$; for all integers $n \geq 3$.

Definition 3:

Let R be a ring, $F = (X, Y, Z)$ be a triple, where $X, Y, Z \in R$. F is called an n -Fermat's triple if and only if $X^n + Y^n = Z^n$; for a fixed integer $n \geq 3$.

Example 4:

Let Z_3 be the ring of integers modulo 3, then $(0, 1, 1)$ is a general Fermat's triple, that is because for all integers $n \geq 3$, $0^n + 1^n = 1^n$

Example 5:

Let Z_5 be the ring of integers modulo 5, then $(0, 2, 1)$ is a 4-Fermat's triple, that is because:

$$0^4 + 2^4 = 1^4.$$

Theorem 6: [16]

Let Z be the ring of integers, then:

- (a) The set of general Fermat's triples is $\{(0, 1, 1), (1, 0, 1), (0, 0, 0)\}$.
- (b) For every fixed integer $n \geq 3$, the set of n -Fermat's triples is $\{(0, 1, 1), (1, 0, 1), (0, 0, 0)\}$.

Theorem 7:

Let R be any ring, F be the set of all general Fermat's triples in R , F_n be the set of all n -Fermat's triples in R , then $F = \bigcap_{n=1}^{\infty} F_n$.

Proof:

The proof holds directly from the definition.

Theorem 8:

Let $Z(I) = \{a + bI; a, b \in Z \text{ and } I^2 = I\}$ be the ring of neutrosophic integers. The equation $X^n + Y^n = Z^n; n \geq 3$, has nine solutions where $X, Y, Z \in Z(I)$.

Proof.

$$\begin{aligned} X^n + Y^n = Z^n &\Leftrightarrow x_0^n + I[(x_0 + x_1)^n - x_0^n] + y_0^n + I[(y_0 + y_1)^n - y_0^n] \\ &= z_0^n + I[(z_0 + z_1)^n - z_0^n] \end{aligned}$$

$$X^n + Y^n = Z^n \Leftrightarrow \begin{cases} x_0^n + y_0^n = z_0^n \dots (1) \\ (x_0 + x_1)^n + (y_0 + y_1)^n = (z_0 + z_1)^n \dots (2) \end{cases}$$

Now, solutions of (1) is.

$$\begin{cases} x_0 = y_0 = z_0 = 0 \dots (a) \\ x_0 = z_0 = 1, y_0 = 0 \dots (b) \\ y_0 = z_0 = 1, x_0 = 0 \dots (c) \end{cases}$$

And solutions of (2) is.

$$\begin{cases} x_0 + x_1 = y_0 + y_1 = z_0 + z_1 = 0 \dots (d) \\ x_0 + x_1 = z_0 + z_1 = 1, y_0 + y_1 = 0 \dots (e) \\ y_0 + y_1 = z_0 + z_1 = 1, x_0 + x_1 = 0 \dots (f) \end{cases}$$

We discuss possible cases.

Case1. If $(x_0 = y_0 = z_0 = 0)$ and $(x_0 + x_1 = y_0 + y_1 = z_0 + z_1 = 0)$, then $X = Y = Z = 0$.

Case2. If $(x_0 = y_0 = z_0 = 0)$ and $(x_0 + x_1 = z_0 + z_1 = 1, y_0 + y_1 = 0)$, then $X = Z = 1, Y = 0$.

Case3. If $(x_0 = y_0 = z_0 = 0)$ and $(y_0 + y_1 = z_0 + z_1 = 1, x_0 + x_1 = 0)$, then $Y = Z = 1, X = 0$.

Case4. If $(x_0 = z_0 = 1, y_0 = 0)$ and $(x_0 + x_1 = z_0 + z_1 = y_0 + y_1 = 0)$, then $X = 1 - I, Y = 0, Z = 1 - I$.

Case5. If $(x_0 = z_0 = 1, y_0 = 0)$ and $(x_0 + x_1 = z_0 + z_1 = 1, y_0 + y_1 = 0)$, then $X = 1, Y = 0, Z = 1$.

DOI: <https://doi.org/10.54216/JNFS.010201>

Case6. If $(x_0 = z_0 = 1, y_0 = 0)$ and $(z_0 + z_1 = y_0 + y_1 = 1, x_0 + x_1 = 0)$, then $X = 1 - I, Y = I, Z = 1$.

Case7. If $(z_0 = y_0 = 1, x_0 = 0)$ and $(z_0 + z_1 = y_0 + y_1 = x_0 + x_1 = 0)$, then $X = 0, Y = 1 - I, Z = 1 - I$.

Case8. If $(z_0 = y_0 = 1, x_0 = 0)$ and $(x_0 + x_1 = z_0 + z_1 = 1, y_0 + y_1 = 0)$, then $X = 0, Y = 1 - I, Z = 1 - I$.

Case9. If $(x_0 = z_0 = 1, y_0 = 0)$ and $(y_0 + y_1 = z_0 + z_1 = 1, x_0 + x_1 = 0)$, then $X = 0, Y = 1, Z = 1$.

Theorem 9:

Let Z_2 be the ring of integers modulo 2, hence it has exactly 4 general Fermat's triples.

Proof:

Consider the equation $X^n + Y^n = Z^n; n \geq 3$, where $X, Y, Z \in Z_2$. We have the following solutions:

$(1,1,0), (0,1,1), (1,0,1), (0,0,0)$.

Theorem 10:

Let Z_3 be the ring of integers modulo 3. If n is odd then it has exactly 9 n -Fermat's triples.

Proof:

Consider the equation $X^n + Y^n = Z^n; n \geq 3$

Where n is odd. It has the following solutions:

$(0,0,0), (1,0,1), (0,1,1), (2,0,2), (0,2,2), (2,2,1), (2,1,0), (1,2,0), (1,1,2)$.

Theorem 11:

Let Z_3 be the ring of integers modulo 3. If n is even, then it has exactly 5 n -Fermat's triples.

Proof:

Consider the equation $X^n + Y^n = Z^n; n \geq 3$

Where n is even. It has the following solutions:

$(0,0,0), (1,0,1), (0,1,1), (0,2,1), (2,0,1)$.

Remark 12:

According to the previous two theorems, the set of general Fermat's triples in Z_3 is

$\{(0,0,0), (1,0,1), (0,1,1)\}$.

Conclusion

In this paper, we have presented an algorithm to solve the Fermat's Diophantine equation in neutrosophic rings. Also, we solved this Diophantine equation in some numerical rings modulo n.

As a future research direction, we aim to solve the Fermat's Diophantine equation in refined neutrosophic rings and n-refined neutrosophic rings respectively.

References

- [1] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set, Neutrosophic Probability", American Research Press. Rehoboth, 2003.
- [2] Abobala, M., "Neutrosophic Real Inner Product Spaces", Neutrosophic Sets and Systems, Vol. 43, 2021.
- [3] Das, R., Smarandache, F., and Tripathy, B., "Neutrosophic Fuzzy Matrices and Some Algebraic Operations", Neutrosophic Sets and Systems, Vol. 32, pp. 401-409, 2020
- [4] Kandasamy, V.W.B., and Smarandache, F., "Some Neutrosophic Algebraic Structures and Neutrosophic N-Algebraic Structures", Hexis, Phonex, Arizona, 2006.
- [5] Agboola, A.A.A., Akinola, A.D., and Oyebola, O.Y., " Neutrosophic Rings I" , International J.Mathcombin, Vol 4,pp 1-14. 2011

- [6] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [7] Chalapathi, T., and Madhavi, L., "Neutrosophic Boolean Rings", *Neutrosophic Sets and Systems*, Vol. 33, pp. 57-66. 2020
- [8] F. Smarandache, *Neutrosophic Theory and Applications*, Le Quy Don Technical University, Faculty of Information technology, Hanoi, Vietnam, 17th May 2016.
- [9] Abobala, M., "A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", *International Journal of Mathematics and Mathematical Sciences*, hindawi, 2021
- [10] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", *Neutrosophic Sets and Systems*, Vol. 43, 2021.
- [11] Abobala, M., Bal, M., and Hatip, A., "A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 17, 2021.
- [12] Hatip, A., and Olgun, N., "On Refined Neutrosophic R-Module", *International Journal of Neutrosophic Science*, Vol. 7, pp.87-96. 2020.
- [13] Hajjari, A., and Ali, R., "A Contribution To Kothe's Conjecture In Refined Neutrosophic Rings", *International Journal of Neutrosophic Science*", Vol. 16, 2021.
- [14] Olgun, N., Hatip, A., Bal, M., and Abobala, M., "A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices", *International Journal of Neutrosophic Science*, Vol. 16, pp. 72-79, 2021.
- [15] Ibrahim, M., and Abobala, M., "An Introduction To Refined Neutrosophic Number Theory", *Neutrosophic Sets and Systems*, Vol. 45, 2021.
- [16] Wiles, A., "Modular Elliptic Curves and Fermat's Last Theorem", *Annals Of Mathematics*, 1995.