



HAL
open science

Ecological aspects of the dispersal of a melastomataceae by marmosets and howler monkeys (Primates : Platyrrhini) in a semideciduous forest of southeastern Brazil

Rodolfo Antônio de Figueiredo, Claudia Aparecida Longatti

► **To cite this version:**

Rodolfo Antônio de Figueiredo, Claudia Aparecida Longatti. Ecological aspects of the dispersal of a melastomataceae by marmosets and howler monkeys (Primates : Platyrrhini) in a semideciduous forest of southeastern Brazil. *Revue d'Écologie (La Terre et La Vie)*, 1997, 52 (1), pp.3-8. <hal-03529121>

HAL Id: hal-03529121

<https://hal.science/hal-03529121v1>

Submitted on 17 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

ECOLOGICAL ASPECTS OF THE DISPERSAL
OF A MELASTOMATACEAE
BY MARMOSETS AND HOWLER MONKEYS
(PRIMATES : PLATYRRHINI)
IN A SEMIDECIDUOUS FOREST OF SOUTHEASTERN BRAZIL

Rodolfo Antônio de FIGUEIREDO* & Claudia Aparecida LONGATTI**

Forest fragments generally show lower biodiversity than preserved areas, but some primate species seem to be apt to live in such small remnants (Lovejoy *et al.*, 1986 ; Schwarzkopf & Rylands, 1989). Diet and feeding behavior of neotropical primates have been studied (Guillotin *et al.*, 1994 ; Julliot, 1994), but their role in tropical ecosystems is not completely understood (Heymann, 1993). In the Old World, monkeys consume fruits brightly colored, with succulent pulp and are potential dispersers of the seeds (Hladik & Hladik, 1967 ; Gautier-Hion *et al.*, 1985). Neotropical primate species are also capable to disseminate several plant species in the rainforest since seeds pass undamaged through their digestive tracts (Hladik & Hladik, 1969 ; Chapman, 1989 ; Julliot, 1994).

The faunae surviving in semideciduous forest fragments of southeastern Brazil, as well as the ecological relationships between primates and plant species in these areas, are poorly known (see Morellato & Leitão-Filho, 1995). Figueiredo (1993, 1994) found Howlers as inadequate dispersers of *Ficus* spp. seeds, and some studies proposed that small seeded plants are dispersed mainly by small monkeys, like Marmosets, while large seeded plants by Howler monkeys (Galetti & Pedroni, 1994 ; Galetti *et al.*, 1994).

This study verifies the dispersal quality of Howlers and Marmosets on the small seeds of a Melastomataceae.

STUDY AREA

Fieldwork was conducted in a forest fragment of 30 ha at the Estação Experimental de Jundiá (Instituto Agrônômico de Campinas), southeastern Brazil (23° 06' S, 46° 55' W, 715 m alt.). The area is characterized by a cold and dry winter (May to August) and a warm and rainy summer (November to February). Figure 1 shows the mean temperature and the monthly precipitation during the study period in the area.

* Departamento de Botânica, Instituto de Biologia, C.P. 6109, Universidade Estadual de Campinas, 13083.970 Campinas, SP, Brasil.

** Faculdade de Ciências, Associação « Padre Anchieta » de Ensino, Jundiá, SP, Brasil.




Figure 1. — Average temperature and precipitation in a semideciduous forest of SE Brazil.

The vegetation consists of secondary growth semideciduous forest. This type of vegetation is typical of southeastern Brazil, and descriptions can be found in Morellato (1992) and Morellato & Leitão-Filho (1995). The fauna and flora of the study area are little known, but semideciduous forests generally show high flora diversity (Morellato, 1992). Three monkey species were found living in the forest: *Alouatta fusca fusca* (Ihering, 1914) (the Northern brown howler monkey), *Callithrix penicillata* (É. Geoffroy, 1812) (the Black-tufted-ear marmoset) and *Callicebus personatus nigrifrons* (Spix, 1823) (the Black-fronted titi monkey) (primate names follow Rylands *et al.*, 1995). Only the first two monkey species were found ingesting seeds of *Miconia cinnamomifolia* (DC.) Naud. (Melastomataceae), a common tree in gaps and borders of the area. This plant species reaches heights up to 22 m, with trunks up to 50 cm dbh, and produced thousands of purple fruits between April and July 1995. The fruits have a diameter of 3.9 ± 0.4 mm, and 33.1 ± 8.7 seeds within ($N = 10$).

METHODS

Primates were observed feeding on ten *M. cinnamomifolia* individuals. Faeces of monkey species containing melastome seeds were searched along established trails in forest edge and towards the interior of the forest ($n = 10$; a total of 2 500 m). The faeces of Howlers and Marmosets were easy to be

distinguished from each other because the former monkey species consumes a great quantity of leaves in addition to fruits. These surveys, and the germination experiments, were done in the last week of July 1995.

One thousand seeds of *Miconia cinnamomifolia* were chosen randomly from faecal droplets of each monkey species (total 2 000 ingested seeds). As a control, it was chosen 1 000 seeds from randomly collected mature fruits underneath the fruiting trees (1 000 non-ingested seeds). Germination sets of 100 seeds in 10 replicates were made for each of the three treatments. Half of the replicates of each treatment were sown on filter paper in covered Petri plates, kept moistened with distilled water and placed in an uncontrolled light and temperature regime inside the laboratory. The other half of the replicates was sown on Petri plates filled with soil, because differences in germination rates observed in laboratory may not be confirmed when seeds experience a more real situation (Figueiredo & Perin, 1995). The soil samples were collected near the site of natural occurrence of *M. cinnamomifolia*. These plates were covered to maintain moisture and exclude seed predators and placed in a forest gap.

The number of germinated seeds in each replicate was verified after two months, and the differences observed were tested using analysis of variance (one-way ANOVA).

RESULTS

Howler monkeys were observed feeding on *M. cinnamomifolia* in two occasions. Their feeding bouts lasted ca. 20 minutes. They eat whole mature and immature fruits. Eighteen Howlers' faeces were found in gaps and forest edges. Their contents revealed hundreds of *M. cinnamomifolia* seeds, as well as pieces of mature and immature *Miconia* fruits, and some seeds of *Syagrus oleraceae* (Arecaceae), *Myrcianthes pungens* (Myrtaceae), and *Copaifera langsdorffii* (Caesalpiniaceae).

Marmosets were observed on *M. cinnamomifolia* on nine occasions. They fed only on mature fruits, mashing and swallowing the whole fruit. Twenty-five faecal droplets, with *M. cinnamomifolia* and *Ficus* sp. seeds, were also found in forest edges and gaps.

The number of *Miconia* seeds germinated in each treatment are shown in Table I (for laboratory trials) and in Table II (for field ones). Laboratory experiments indicated that the germination rates of Marmoset-ingested seeds were similar to control ones ($F = 2.96$, $p = 0.13$). Howler-ingested seeds, however, showed a smaller germination rate when compared both with control ($F = 184.07$, $p < 0.01$) and Marmoset lots ($F = 149.33$, $p < 0.01$). In the field experiments, germination rates were lower than those observed in the laboratory. The results, however, were similar: Howler-ingested seeds germinated less than control ($F = 95.34$, $p < 0.01$) and than Marmoset-ingested seeds ($F = 138.03$, $p < 0.01$), and Marmoset-ingested and control seeds did not differ significantly ($F = 1.35$, $p = 0.29$).

DISCUSSION

Primate ingestion did not inhibit germination of *Miconia cinnamomifolia* seeds. The reduction of germination rates observed for Howler-ingested seeds is

TABLE I

Number of vertebrate-ingested and non-ingested Miconia cinnamomifolia seeds germinated in laboratory trials.

Treatment	Replicates					$\bar{x} \pm SD$
	1	2	3	4	5	
Control	69	73	74	68	71	71.0 ± 2.6
Howler-ingested	39	43	36	47	42	41.4 ± 4.2
Marmoset-ingested	69	73	80	79	74	75.0 ± 4.5

TABLE II

Number of vertebrate-ingested and non-ingested Miconia cinnamomifolia seeds germinated in field germination trials.

Treatment	Replicates					$\bar{x} \pm SD$
	1	2	3	4	5	
Control	46	49	47	36	41	43.8 ± 5.3
Howler-ingested	10	06	18	15	15	12.8 ± 4.8
Marmoset-ingested	39	44	38	41	42	40.8 ± 2.4

probably due to the behaviour of feeding on immature fruits showed by this monkey. Howlers, therefore, can be considered a non-specialized disperser of *M. cinnamomifolia*. Almost half of the Howler-ingested seeds germinated, hence Howlers can contribute to increase the recruitment of the plant species in gaps and forest edges. Marmosets, contrariwise, are better seed dispersers. Marmoset activity, mainly in successional, disturbed and edge habitats (Stevenson & Rylands, 1988), may increase seedling recruitment of *M. cinnamomifolia* in the forest fragment.

Fragmentation of native forests and hunting can provoke extinction or extreme population declines of several primate species (Lovejoy *et al.*, 1986), and this can lead to long-term effects on the recruitment of several plant species (Howe, 1984). Some studies showed the importance of monkeys on seed dispersal and germination of several plant species (Izawa, 1975 ; Garber, 1986 ; Galetti *et al.*, 1994 ; Zang & Wang, 1995). In this study, both primate species may contribute to regeneration of forest edges and gaps by dispersing the seeds of a pioneer tree. The maintenance and regeneration of forest remnants are very important in southeastern Brazil, because the Northern howler monkey and the Black-fronted titi monkey are on the verge of extinction (Mittermeier, 1986 ; Fonseca *et al.*, 1994), and the Black-tufted-ear marmoset is threatened by deforestation of its small geographic range (Emmons, 1990).

The present study strengthened the hypothesis that smaller monkey species are better dispersers of small seeds than larger monkey species like Howlers.

ACKNOWLEDGEMENTS

The authors are indebted to Fernando P. Martins for permission to work in the Estação Experimental de Jundiaí, and José Luiz Hernandes for discussions. Rodolfo A. de Figueiredo acknowledges a doctoral fellowship from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, process number 140055/94-0). Two anonymous reviewers greatly improved an early version of the manuscript.

SUMMARY

Feeding behavior and digestion effect on fruits and seeds of a pioneer tree (*Miconia cinnamomifolia*) by primates (*Alouatta fusca fusca* and *Callithrix penicillata*) were investigated in a forest fragment of southeastern Brazil. Howler monkeys consumed mature as well as immature fruits, and the germination rates of ingested seeds were reduced when compared with non-ingested ones. Marmosets defecated unharmed seeds and were considered better seed dispersers than Howlers. These results strengthened the hypothesis that plant species with small seeds are dispersed mainly by small monkey species.

RÉSUMÉ

Nous avons étudié le comportement alimentaire et l'effet de la digestion sur les graines d'un arbre néotropical (*Miconia cinnamomifolia*) par les primates (*Alouatta fusca fusca* et *Callithrix penicillata*) dans une forêt du sud-est du Brésil. *Alouatta fusca* consomme en partie des fruits immatures et les taux de germination des graines ingérées sont réduits. *Callithrix penicillata* est un meilleur disperseur de graines de *M. cinnamomifolia*. Ces résultats confortent l'hypothèse selon laquelle les petites graines sont disséminées par des petits animaux.

REFERENCES

- CHAPMAN, C.A. (1989). — Primate seed dispersal: the fate of dispersed seeds. *Biotropica*, 21 : 148-154.
- EMMONS, L.H. (1990). — *Neotropical rainforest mammals*. The University of Chicago Press, Chicago.
- FIGUEIREDO, R.A. de (1993). — Ingestion of *Ficus enormis* seeds by howler monkeys (*Alouatta fusca*) in Brazil: effects on seed germination. *J. Trop. Ecol.*, 9 : 541-543.
- FIGUEIREDO, R.A. de (1994). — *Fenologia e polinização de três espécies de figueiras em mata semidecídua de planalto na região de Campinas*, SP. Master Thesis. Universidade Estadual de Campinas, Campinas, SP.
- FIGUEIREDO, R.A. de & PERIN, E. (1995). — Germination ecology of *Ficus luschnathiana* drupelets after bird and bat ingestion. *Acta Oecol.*, 16 : 71-75.

- FONSECA, G.A.B., RYLANDS, A.B., COSTA, C.M.R., MACHADO, R.B. & LEITE, Y.L.R. (1994). — *Livro vermelho dos mamíferos brasileiros ameaçados de extinção*. Fundação Biodiversitas, Belo Horizonte.
- GALETTI, M. & PEDRONI, F. (1994). — Seasonal diet of capuchin monkeys (*Cebus apella*) in a semideciduous forest in south-east Brazil. *J. Trop. Ecol.*, 10 : 27-39.
- GALETTI, M., PEDRONI, F. & MORELLATO, L.P.C. (1994). — Diet of the brown howler monkey *Alouatta fusca* in a forest fragment in southeastern Brazil. *Mammalia*, 58 : 111-118.
- GARBER, P.A. (1986). — The ecology of seed dispersal in two species of callitrichid primates (*Saguinus mystax* and *Saguinus fuscicollis*). *Am. J. Primatol.*, 10 : 155-170.
- GAUTIER-HION, A., DUPLANTIER, J.M., QURIS, R., FEER, F., SOURD, C., DECOUX, J.P., DUBOST, G., EMMONS, L., ERARD, C., HECKESTWEILER, P., MOUNGAZI, A., ROUSSILHON, C. & THIOLLAY, J.M. (1985). — Fruit characters as a basis of fruit choice and seed dispersal in a tropical forest vertebrate community. *Oecologia*, 65 : 324-337.
- GUILLOTIN, M., DUBOST, G. & SABATIER, D. (1994). — Food choice and food competition among the three major primate species of French Guiana. *J. Zool. Lond.*, 233 : 551-579.
- HEYMANN, E.W. (1993). — The role of primates in tropical ecosystems. P. 103-108 in : W. Barthlott, C.M. Naumann, K. Schmidt-Loske & K.L. Schuchmann (eds) *Animal-plant interactions in tropical environments*. Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn.
- HLADIK, A. & HLADIK, C.M. (1969). — Rapports trophiques entre végétation et primates dans la forêt de Barro Colorado. *Terre et Vie*, 1 : 25-117.
- HLADIK, C.M. & HLADIK, A. (1967). — Observations sur le rôle des primates dans la dissémination des végétaux de la forêt gabonaise. *Biol. Gabon*, 3 : 43-58.
- HOWE, H.F. (1984). — Implications for seed dispersal by animals for tropical reserve management. *Biol. Conserv.*, 30 : 261-281.
- HOWE, H.F. (1993). — Specialized and generalized dispersal systems : where does 'the paradigm' stand ? *Vegetatio*, 107/108 : 3-13.
- IZAWA, K. (1975). — Foods and feeding behavior of monkeys in the upper Amazon Basin. *Primates*, 16 : 295-316.
- JULLIOT, C. (1994). — Frugivory and seed dispersal by red howler monkeys : evolutionary aspects. *Rev. Ecol. (Terre Vie)*, 49 : 331-341.
- LOVEJOY, T.E., BIERREGAARD, R.O., RYLANDS, A.B., MALCOLM, J.R., QUINTELA, C.E., HARPER, L.H., BROWN, K.S., POWELL, A.H., POWELL, G.V., SCHUVART, H.O. & HAYS, M.B. (1986). — Edge and other effects of isolation on Amazon forest fragments. P. 257-285 in M.E. Soulé (ed) *Conservation biology : the science of scarcity and diversity*. Sinauer Associates Inc., Sunderland, Mass.
- MCKEY, D. (1975). — The ecology of coevolved seed dispersal systems. P. 159-191 in L.E. Gilbert & P. Raven (eds) *Coevolution of animals and plants*. Univ. Texas Press, Austin.
- MORELLATO, L.P.C. (1992). — *História natural da Serra do Japi : ecologia e preservação de uma área florestal no sudeste do Brasil*. Editora da Unicamp, Campinas.
- MORELLATO, L.P.C. & LEITÃO-FILHO, H.F. (1995). — *Ecologia e preservação de uma floresta tropical urbana : Reserva de Santa Genebra*. Editora da Unicamp, Campinas.
- MITTERMEIER, R.A. (1986). — A global overview of primate conservation. P. 325-340 in J.G. Else & P.C. Lee (eds) *Primate ecology and conservation*. Cambridge University Press, Cambridge.
- NUNES, A. (1995). Um teste de germinação em sementes dispersas por macacos-aranha em Maracá, Roraima ; Brasil. *Studies on Neotropical Fauna and Environment*, 30 : 31-36.
- RYLANDS, A.B., MITTERMEIER, R.A. & LUNA, E.R. (1995). — A species list for the New World Primates (Platyrrhini) : distribution by country, endemism, and conservation status according to the mace-land system. *Neotropical Primates*, 3 : 113-160.
- SCHWARZKOPF, L. & RYLANDS, A.B. (1989). — Primate species richness in relation to habitat structure in Amazonian rainforest fragments. *Biological Conservation*, 48 : 1-12.
- SNOW, D.W. (1971). — Evolutionary aspects of fruit-eating in birds. *Ibis*, 113 : 194-202.
- STEVENSON, M.F. & RYLANDS, A.B. (1988). — The marmosets, genus *Callithrix*. P. 131-222 in R.A. Mittermeier, A.F. Coimbra-Filho, A.B. Rylands & G.A.B. Fonseca (eds) *Ecology and behaviour of neotropical primates*. World Wildlife Fund, Washington D.C.
- STILES, F.G. & ROSSELLI, L. (1993). — Consumption of fruits of the Melastomataceae by birds : how diffuse is coevolution ? *Vegetatio*, 107-108 : 57-73.
- ZHANG, S.-Y. & WANG, L.-X. (1995). — Fruit consumption and seed dispersal of *Ziziphus cinnamomum* (Rhamnaceae) by two sympatric primates (*Cebus apella* and *Atelles paniscus*) in French Guiana. *Biotropica*, 27 : 397-401.