



Dengue viremia kinetics in asymptomatic and symptomatic infection

Ponpan Matangkasombut, Kajohnpong Manopwisedjaroen, Nada Pitabut, Sasikanya Thaloengsok, Swangjit Suraamornkul, Tawatchai Yingtaweesak, Veasna Duong, Anavaj Sakuntabhai, Richard E. Paul, Pratap Singhasivanon

► To cite this version:

Ponpan Matangkasombut, Kajohnpong Manopwisedjaroen, Nada Pitabut, Sasikanya Thaloengsok, Swangjit Suraamornkul, et al.. Dengue viremia kinetics in asymptomatic and symptomatic infection. International Journal of Infectious Diseases, 2020, 101, pp.90 - 97. <10.1016/j.ijid.2020.09.1446>. <hal-03533008>

HAL Id: hal-03533008

<https://hal.science/hal-03533008v1>

Submitted on 18 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License



Dengue viremia kinetics in asymptomatic and symptomatic infection



Ponpan Matangkasombut^{a,b,*}, Kajohnpong Manopwisedjaroen^{a,1}, Nada Pitabut^{c,d}, Sasikanya Thaloengsok^{a,2}, Swangjit Suraamornkul^e, Tawatchai Yingtaeasak^f, Veasna Duong^g, Anavaj Sakuntabhai^h, Richard Paulⁱ, Pratap Singhasivanon^{i,**}

^a Department of Microbiology, Faculty of Science, Mahidol University, Bangkok, Thailand

^b Systems Biology of Diseases Research Unit, Faculty of Science, Mahidol University, Bangkok, Thailand

^c Office of Research Services, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

^d Faculty of Medicine, King Mongkut's Institute of Technology Ladkrabang, Bangkok, Thailand

^e Faculty of Medicine, Vajira Hospital, Bangkok, Thailand

^f Thasongyang Hospital, Tak, Thailand

^g Virology Unit, Institut Pasteur du Cambodge, Institut Pasteur International Network, PO Box 983, Phnom Penh, Cambodia

^h Institut Pasteur, Functional Genetics of Infectious Diseases Unit, UMR 2000 (CNRS), Paris 75015, France

ⁱ Department of Tropical Hygiene, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

ARTICLE INFO

Article history:

Received 19 August 2020

Received in revised form 22 September 2020

Accepted 22 September 2020

Keywords:

Dengue
Asymptomatic
Inapparent
Virus kinetics
Viral decay
Viremia duration

ABSTRACT

Background: Dengue infection is a global health threat. While symptomatic cases contribute to morbidity and mortality, the majority of infected people are asymptomatic but serve as an important reservoir. However, the kinetics of viremia in asymptomatic infections remains unknown.

Methods: We enrolled 279 hospital-based symptomatic index cases and quantified dengue virus (DENV) RNA at enrollment and at the day of defervescence. To identify asymptomatic cases, 175 household members of index cases were monitored for clinical symptoms during follow-up, and blood was taken twice weekly to test for and quantify DENV RNA until cleared.

Results: We detected DENV in thirteen asymptomatic household members (7.43%). Their DENV serotypes were primarily the same as those of their family index cases. The median peak DENV viremia in asymptomatic subjects was lower than that of symptomatic individuals during the febrile phase, and the viral decay rate was slower in asymptomatic infections.

Conclusions: DENV level and kinetics in asymptomatic individuals differed significantly from those of symptomatic cases. Despite the lower viremia, the slower decay rate in asymptomatic infections could lead to their prolonging the infectious reservoir. The improvement of transmission control to prevent such long-lived asymptomatic infections from transmitting the DENV is needed.

© 2020 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Dengue is the most important arthropod-borne viral infection worldwide, infecting an estimated 390 million people annually (Bhatt et al., 2013). The incidence of dengue virus (DENV) infection has been rising over the last five decades (Gubler, 2020). The outcome of DENV infection ranges from asymptomatic/inapparent infection, mild self-limited dengue fever (DF) to severe dengue hemorrhagic fever (DHF), with the potential development of life-threatening dengue shock syndrome (DSS). However, the majority of infected individuals have no or insufficient symptoms to result in a clinical presentation; nevertheless, they could serve as a

* Corresponding author at: Department of Microbiology, Faculty of Science, Mahidol University, 272 Rama VI Road, Rajatewee, Bangkok, 10400 Thailand.

** Corresponding author: Department of Tropical Hygiene, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand.

E-mail addresses: ponpan.mat@mahidol.edu (P. Matangkasombut), pratap.sin@mahidol.ac.th (P. Singhasivanon).

¹ Kajohnpong Manopwisedjaroen: Current affiliation and address: Mahidol vivax research unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand 10400.

² Sasikanya Thaloengsok: Current affiliation and address: Department of Bacterial and Parasitic Diseases, Armed Forces Research Institute of Medical Sciences (AFRIMS), Bangkok, Thailand 10400.

significant reservoir for DENV transmission (Endy, 2002; Endy et al., 2011; Grange et al., 2014; ten Bosch et al., 2018).

The level of DENV viremia is one of the most important determinants of human infectiousness to mosquitoes (Nguyen et al., 2013; Duong et al., 2015), and the duration of infection is a crucial parameter determining the epidemiological dynamics of any pathogen and the subsequent R_0 (Anderson and May, 1991). In symptomatic DENV infections, classical experimental infection studies showed that DENV was infectious from two days before and five days after illness onset (Siler et al., 1926; Simmons et al., 1931; Nishiura and Halstead, 2007). The duration of infectiousness, estimated via the success of transmission to mosquitoes or viral isolation, was found to range from between 1–7 days with a mean of 4–5 days and with longer durations in primary than secondary infections (Siler et al., 1926; Duyen et al., 2020; Kuberski et al., 1977). More recent studies using molecular detection of the virus have revealed a similar duration of infection, with a range lasting up to six days post-onset of fever and with some variation according to serotype, disease severity, and 1° vs. 2° infections (Duyen et al., 2020). In experimental non-human primate sylvatic DENV infections,

viremia duration ranged from three to five days, depending on the DENV serotype (Althouse et al., 2014). To date, the kinetics of DENV in individuals with asymptomatic DENV infections remains unknown.

We conducted a cohort study to identify and follow asymptomatic DENV infected individuals, DF, and DHF patients prospectively to characterize their viremia kinetics. Asymptomatic DENV infected individuals were identified by investigating the presence of DENV by Reverse transcription PCR (RT-PCR) in household members (HHM) of symptomatic DENV infected patients (index cases) (Dussart et al., 2012). Once identified, we followed these individuals prospectively, monitored their symptoms, and measured the level of DENV in blood samples twice a week until DENV was cleared from the circulation. This knowledge of DENV kinetics in asymptomatic individuals is crucial for predicting the level and duration of infectiousness in this important DENV reservoir.




Figure 1. Study design for index cases and household members investigation. D1 (day of enrollment); Ddef (Day of defervescence); Wk2 (two weeks after enrollment); Mo2 (two months after enrollment); DENV PCR (RT-PCR of DENV result).

Materials and methods

Ethics statement

The study was approved by the Institutional Review Board of Faculty of Medicine Vajira Hospital (No.015/12) and the Faculty of Tropical Medicine Mahidol University (TMEC 13–041). All subjects or their legal guardians signed written informed consent before study participation.

Study population

Subjects were recruited from two study sites in Bangkok (Vajira hospital and the Faculty of Tropical Medicine, Mahidol University) and one study site in Tak province (Tasongyang hospital). Subjects were recruited from Vajira hospital in 2012–2015, and the Faculty of Tropical Medicine and Tak sites were added in 2015. A total of 279 index cases were enrolled

Index case sample and data collection

Demographic and clinical data of index cases were collected with standardized case report forms. Blood samples were collected for RT-PCR twice: on the day of enrollment (D1) and on the day fever subsided (Day of defervescence, Ddef). Additional blood samples were collected once a day during the febrile phase between D1 and Ddef and at 2-week and 2-month follow-up time points for a heme-agglutination inhibition (HI) test and other immunological studies. Index cases were classified into DF and DHF according to WHO criteria (World Health Organization, 1997) (Figure 1).

Household member samples and data collection

Once the index cases were confirmed for DENV infection, their HHM were enrolled within 1–2 days. Their body temperature and clinical symptoms were recorded on standardized case report forms throughout the follow-up period. Blood samples were taken for DENV RT-PCR on the first day of enrollment (D1) and then between 24–72 hours later. The result of DENV RT-PCR became available within 24 h after blood collection and dictated subsequent investigation and follow-up. If DENV was detectable by RT-PCR, two drops of blood were taken onto filter paper (Aubry et al., 2012) every 2–4 days until DENV become undetectable twice in a row (or the subject was lost to follow-up). The clinical follow-up also ended when DENV was undetectable twice consecutively by RT-PCR (Figure 1).

DENV detection, quantification and serotype determination by PCR

Serotype-specific nested RT-PCR was performed on all samples to detect the presence of DENV RNA and determine the DENV serotype. Quantitative RT-PCR (qRT-PCR) was used to quantify the DENV viral load when possible. First, viral RNA was extracted from blood samples using the QIAamp viral RNA mini kit (QIAGEN, Germany) according to the manufacturer's instructions and stored at –80 °C until used. Serotype-specific nested RT-PCR was performed according to the published protocol (Lanciotti et al., 1992). The serotype and quantity of DENV RNA were determined by the qRT-PCR assay, as previously described (Duong et al., 2015) (supplementary Table 1).

Primary/secondary DENV infection determination

In-house IgM capture ELISA (Duong et al., 2015) and HI assays were performed as previously described (Clarke, 1958). Paired plasma during acute infection and two weeks or two months follow-up time points (when available) were used for HI. Primary and secondary infections were determined using WHO/TDR (1997) criteria (World Health Organization, 1997).

Statistical analysis

DENV decay rate in index cases was calculated based on the DENV viral load on D1 subtracted by that on Ddef and divided by the number of days between D1 and Ddef. Decay rates in asymptomatic HHM were also calculated for every time point when RT-PCR was performed on an individual.

For risk factor analyses of viral load, decay rate, day to defervescence, and day to viral clearance, a Generalized Linear Model with Poisson error structure was fitted to test the association with the following explanatory variables: age (continuous), gender (M/F), year, site, infection severity (DHF vs. DF) and serotype and where indicated, log10 transformed viral load at D1. The number of individuals included in each analysis varied: all viral load samples were analyzed irrespective of whether they were subsequently lost to follow-up; decay rate included only individuals who were not lost to follow-up and who showed a decrease in viremia from D1 to Ddef (only eight individuals had an increase in viremia from D1 to Ddef); days to defervescence included all individuals not lost to follow-up irrespective of whether viremia increased from D1 to Ddef; day to viral clearance included only those with zero viremia at Ddef.

To assess whether there was a difference in the decay rate between asymptomatic household members and their index cases, a Generalized linear mixed model with Poisson error structure was fitted to decay rate with age, gender, serotype (DENV-3 vs. DENV-4), infection type (Index case vs. Asymptomatic HHM) as explanatory variables and household ID as the random factor. The viral load at D1 and Ddef of the index cases and the peak viral load of the asymptomatic HHM were compared by the Mann-Whitney *U* test. For all analyses, a dispersion parameter was estimated and used to account for overdispersion in the data. Analyses were performed in Genstat version 15 (VSN International, 2017).

Table 1
Index cases characteristics.

		Number of cases (percentage)
Severity	DF	177 (63.44%)
	DHF	102 (36.55%)
Year	2012	31 (11.11%)
	2013	118 (42.29%)
	2014	16 (5.73%)
	2015	114 (40.86%)
Serotype	1	39 (13.98%)
	2	38 (13.62%)
	3	89 (31.90%)
	4	96 (34.40%)
	indeterminate ^a	17 (6.09%)
Primary/secondary infection	Primary	22 (8.24%)
	Secondary	208 (74.55%)
	Indeterminate	49 (17.5%)
Gender	Male	136 (48.75%)
	Female	143 (51.25%)
Age	Children(≤15)	88 (31.54%)
	Adults (>15)	191 (68.46%)

^a These subjects had a positive result for NS1 or IgM, but DENV RNA undetectable by PCR and, therefore, unable to determine serotype.

Results

Index cases characteristics

290 index cases were initially enrolled in the study, of which 279 cases had confirmed DENV infection from 138 households. DENV was detected by RT-PCR in 262 of these 279 cases (94.27%). Of those 17 with negative RT-PCR, eleven had anti-DENV IgM, four had detectable NS1, and two had both anti-DENV IgM and NS1. The majority of index cases had DF (63.44%), All four DENV serotypes circulated during the study years, with DENV-4 (34.40%) and DENV-3 (31.90%) being more prevalent than the other serotypes. An almost equal number of male (48.75%) and female (51.25%) subjects were enrolled, and the majority were adults (68.46%) (Table 1).

DENV kinetics in index cases

Overall, the mean viral load in DENV positive individuals on the day of enrollment (D1) was 7.24×10^9 viral copies/mL (SEM: 3.45×10^9). The time to defervescence ranged from 1 to 7 days post day of recruitment (the average time was 2.28 days, SEM 0.08). Of those individuals that cleared their viral load completely by the day of defervescence, the mean time to clearance was 2.77 days (SEM 0.15).

DENV viral load on D1 was higher in DHF than DF (Relative Risk (RR) = 5.88, $P < 0.001$), decreased with age (RR = 0.92, $P = 0.003$), was lower in male than female gender (RR = 0.31, $P < 0.001$) and DENV-1 (RR = 0.25, $P = 0.001$), DENV-2 (RR = 0.08, $P = 0.002$), and DENV-4 (RR = 0.06, $P = 0.006$) as compared to DENV-3 (Table 2, top). The full minimum adequate model explained 29.2% of the variation in viral load.

DENV decay rate (from D1 to Ddef) was faster in DHF than DF (RR = 1.44, $P = 0.004$) and DENV-1 (RR = 1.43, $P = 0.014$) as compared to DENV-3 (Table 2, middle) and with an increased viral load on D1 (RR = 6.34, $P < 0.001$). The full minimum adequate model explained 94.7% of the variation in the decay rate.

Time taken to defervescence decreased with age (RR = 0.99, $P = 0.003$) and increased with viral load at D1 (RR = 1.12, $P < 0.001$) and for DENV-4 (RR = 1.53, $P < 0.001$) as compared to DENV-3 (Table 2, bottom). The full minimum adequate model explained 27.4% of the variation in days to defervescence. A similar result was found when using only those individuals who had a completely cleared viremia on the day of defervescence.

Rate of asymptomatic DENV infections in household members of dengue index cases

Overall, thirteen subjects of 175 HHM (7.43%) from the 138 households investigated had asymptomatic DENV infections (hereon called "asymptomatic HHM") as determined by the presence of DENV RNA by RT-PCR and absence of symptoms during the follow-up period. These thirteen asymptomatic HHM were from eleven households out of the 138 households investigated (7.97%). There was an additional one HHM with DENV viremia that subsequently developed symptoms (pre-symptomatic) in Tak. A further two HHM had had a recent clinical dengue infection within two weeks before their family index cases were diagnosed. Overall, the attack rate with more than one infection in a household (including asymptomatic, pre-symptomatic, and recent DENV infection) was 14/138 households (10.14%). The proportion of households with asymptomatic HHM was 6/53 (11.32%) in Bangkok and 5/85 (5.88%) in Tak (Figure 2).

Table 2

Factors associated with index case DENV kinetics.

A. Day 1 Viral load		N	Mean Log ₁₀ Viremia (SEM)	RR (95% CI)	P-value
Age (years)		222	7.31 (0.14)	0.92 (0.87–0.97)	0.003
Gender	F	113	7.12 (0.21)	Ref	
	M	109	7.51 (0.18)	0.31 (0.16–0.62)	<0.001
Severity	DF	142	7.34 (0.17)	Ref	
	DHF	80	7.27 (0.23)	5.88 (2.92–11.82)	<0.001
Serotype	1	38	7.11 (0.40)	0.25 (0.11–0.57)	0.001
	2	31	6.82 (0.40)	0.08 (0.02–0.40)	0.002
	3	80	7.54 (0.24)	Ref	
	4	73	7.39 (0.17)	0.06 (0.01–0.43)	0.006
B. Decay Rate		N	Mean (SEM) decay rate/day RR		P-value
Severity	DF	114	7.47 (6.83)	Ref	
	DHF	69	7.63 (7.03)	1.44 (1.12–1.84)	0.004
Serotype	1	30	7.71 (7.13)	1.43 (1.08–1.90)	0.014
	2	25	7.66 (7.13)	0.32 (0.79–2.08)	0.319
	3	69	7.55 (6.97)	Ref	
	4	59	7.30 (6.92)	0.18 (0.24–1.32)	0.182
Log10 Viral load D 1		183	7.54 (6.90)	6.34 (5.49–7.32)	<0.001
C. Days to Defervescence		N	Mean (SEM) days	RR	P-value
Age (years)		225	2.38 (0.08)	0.990 (0.986–0.997)	0.003
Serotype	1	32	2.24 (0.18)	1.13 (0.93–1.37)	0.21
	2	35	2.07 (0.19)	1.04 (0.84–1.29)	0.69
	3	79	1.98 (0.11)	Ref	
	4	79	3.03 (0.16)	1.53 (1.31–1.78)	<.001
Log10 Viral load D 1		225	2.38 (0.08)	1.12 (1.08–1.16)	<.001

Shown in this table are the number of samples analyzed for each of the significant risk factors, the Relative Risk (RR) and associated P-value and the dependent variable estimate with standard error from the final fit in the multivariate log-linear regression (see Methods).




Figure 2. Numbers of index cases and household members investigated and numbers of asymptomatic dengue viremia in Bangkok and Tak study sites.

Characteristics of subjects with asymptomatic dengue infection and their family index cases

The characteristics of each of the thirteen asymptomatic HHM and their family index cases are shown in Supplementary Table 2. There were two households from the Bangkok site that had two asymptomatic HHM in the same house. Most but not all of the asymptomatic HHM 9/13 (69.23%) harbored the same DENV serotypes as their family index cases. Interestingly, two asymptomatic HHM (H1 and H2) had co-infection with two DENV serotypes (DENV-3 and -4), while their family index case was infected with only DENV-4. An additional asymptomatic HHM (H9) had a different DENV serotype from the family index case; there was one index case (family of H10) for whom we do not know the DENV serotype. Most family index cases were male 9/11 (81.82%), while 5/13 (38.46%) of asymptomatic HHM were male. Most, 10/11 (90.90%), family index cases had DF, and only one had DHF.

The kinetics of DENV viremia in asymptomatic infections

Overall, the DENV kinetics varied widely among individuals (Supplementary Fig. 1). While some asymptomatic HHM had high viremia but rapidly cleared (H1, H2, H6, H11), others had lower viremia that lasted longer (H3, H4). H9 likely had a very low viral load below the detection limit of qRT-PCR, but detectable by nested PCR. Unfortunately, several subjects were lost to follow-up before the virus was cleared from the circulation. Taken together, the DENV kinetics in asymptomatic HHM are variable, but DENV viremia could persist up to 1–2 weeks after the detection of index cases in the household.

Factors affecting DENV kinetics in asymptomatic infections

Excluding the single instance of a DENV-1 infection, we assessed risk factors associated with viral load, decay, and clearance rate. The mean maximum viral load was 3.89×10^6 viral copies/mL (SEM 1.35×10^6); there was no association with any explanatory variables (age, gender, serotype, mixed or single serotype infection: $P > 0.05$). The time needed for DENV clearance




Figure 3. Dengue viral load in asymptomatic dengue infected HHM compared to index cases. The peak viral load of asymptomatic dengue infected HHM (asymptomatic (peak)) was compared to the viral load of index cases at the day of enrollment (D1) and day defervescence (Ddef). Mann-Whitney was used to compare two groups indicated, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

from the maximum (measured) viral load was found to decrease with increasing maximum viral load ($\chi^2_1 = 5.54$, $P = 0.019$) and to be also faster in mixed serotype infections (Single serotype infections: Mean 6.4 days SEM 1.5; Mixed serotype infections: 3.0 days SEM 0.7. $\chi^2_1 = 5.02$, $P = 0.025$).

Comparing asymptomatic HHM and index case viral kinetics

The peak viremias of asymptomatic DENV infections were lower than those of index cases at the D1 when patients were still febrile. At Ddef, index cases' viral loads dropped markedly from D1




Figure 4. Dengue viral load decay rate by day for asymptomatic HHM infections and index cases from the day of recruitment (D1) to the day of defervescence (Ddef). The decay rate is log (10) transformed for visual clarity.

and were even lower than those of asymptomatic HHM peak viral load (Figure 3).

Asymptomatic infections were associated with a slower decay rate than index cases (Index case: mean 4.31×10^8 / day SEM: 2.91×10^8 ; Asymptomatic HHM: 5.32×10^5 / day, SEM 2.89×10^5 . $\chi^2_{21} = 72.0$, $P = 0.003$) (Figure 4).

Discussion

A DENFREE cohort of DENV infected patients with their household members in Thailand was established to identify asymptomatic DENV infections to provide the first description of viral kinetics within asymptomatic infections and compare them with that in symptomatic clinical presentations from the vicinity and/or the same household. In the symptomatic index cases with a measurable virus at enrollment (D1), viremias were generally higher than those previously reported (Duyen et al., 2020; Yeh et al., 2020), but time to defervescence and/or viral clearance significantly faster (Duyen et al., 2020; Yeh et al., 2020; Gubler et al., 1978; Gubler et al., 1979; Vaughn et al., 1997; Libraty et al., 2002). Significant variation within viral load and time to both defervescence and viral clearance was observed. Notably, patients with DHF and DENV-3 had a higher viral load, and those who were male and older had lower viral loads. A trend for increased viremia with disease severity has been observed previously (Perdomo-Celis et al., 2017), and variation amongst serotypes has also been noted (Duyen et al., 2020). Previous studies have repeatedly found shorter durations of infection in secondary infections as compared to primary infections but which also had higher viremias (Kuberski et al., 1977; Duyen et al., 2020; Yeh et al., 2020; Libraty et al., 2002). In our study, we had too few primary infections to make a reasonable comparison. The viral decay rate from enrollment to defervescence was faster in patients with DHF than those with a higher viral load at enrollment. Despite this, time to defervescence was still longer with higher enrollment viral load.

From index case houses, 7.43% of HHM of index cases had DENV viremia without any symptoms. Most of these asymptomatic HHM had the same DENV serotypes as their family index cases. The kinetics of DENV of these asymptomatic HHM were highly variable among individuals. There were no factors associated with the maximum measured viral load. A higher decay rate was found in

DENV-4 infections, and time to viral clearance was faster with increasing viral load and mixed serotype infections. Thus, in both index cases and asymptomatic infections, high viral load was associated with more rapid viral decay, although this did not lead to clearance at defervescence in symptomatic cases. Significant among serotype differences in viral kinetics were observed here as often before (Duyen et al., 2020), and both among and within-serotype differences in the immune response have been recently highlighted (Katzelnick et al., 2015), albeit not in the context of asymptomatic infections.

Although the exact duration of viremia is difficult to estimate, the viremia lasted up to two weeks in some of the asymptomatic infections, and the decay rate was slower than that of index cases. Similarly, compared with their index cases, the maximum level of viremia in asymptomatic HHM was lower than that of index cases at enrollment but higher than that on the day of defervescence, consistent with the observed slower overall decay rate in asymptomatic infections. This observation supports previous studies that found lower viremia in asymptomatic infections than those of symptomatic infections (Duong et al., 2015; Dussart et al., 2012; Sowath et al., 2019).

Overall, the kinetics of asymptomatic infections differ from symptomatic infections in the magnitude of the viremia and the rate of clearance, suggesting these infections last longer but with a lower viremia. We and others have previously described fundamental immunological differences in the immune responses associated with symptomatic and asymptomatic infections (García et al., 2010; Simon-Lorière et al., 2017; Halstead and O'Rourke, 1977). A polymorphism in the FcγRIIA was found to be associated with inapparent infection vs. DF or DHF in the Cuban population (García et al., 2010). Asymptomatic DENV infected individuals have been found to have increased T cell responses with feedback regulation when compared to symptomatic counterparts (Simon-Lorière et al., 2017). Classically secondary infections are associated with more severe disease due to the phenomenon of ADE and/or cross-reactive T cells (Halstead and O'Rourke, 1977), but whether or not this leads to a decreased risk of an infection being inapparent remains moot (Grange et al., 2014; Clapham and Cummings, 2020). Post-secondary infections will likely induce different immune responsiveness and have also been found to impact upon inapparent rates (Olkowski et al., 2013), which could introduce additional variability in the viral kinetics in both asymptomatic and symptomatic infections. In our study, however, the vast majority of infections were secondary, although it is notable that increasing age had a significant impact on viral clearance rates, suggesting a potential impact of post-secondary infections. More recently, the importance of the interplay between viral genotype within serotype and the interaction with the immune response has been found to be significant (Katzelnick et al., 2015; OhAinle et al., 2020), thereby introducing additional variability that we can not capture in our study.

We have previously shown that asymptomatic infections are as, if not more, infectious to mosquitoes than symptomatic infections (Duong et al., 2015). The likelihood that such asymptomatic infections also last longer does suggest that their epidemiological contribution is even more important. In symptomatic dengue infections, the level of dengue viremia was shown to be the most critical factor for transmission to mosquitoes (Duong et al., 2015). Thus, both those asymptomatic cases with higher but shorter-lived viremia and those with lower but longer-lasting viremia could contribute to transmitting the disease.

A wide range of different epidemiological studies has attempted to ascertain the extent of asymptomatic infections and associated risk factors (Endy, 2002; Endy et al., 2011; Grange et al., 2014; Gordon et al., 2013; Balmaseda et al., 2010; Montoya

et al., 2013; Morrison et al., 2010). It is widely agreed that inapparent, sub-clinical infections are prevalent, and thus this silent infectious reservoir will be a significant contributor to transmission. That we observed asymptomatic infections lasting for two weeks will prove particularly problematic in preventing their role as a reservoir through traditional fumigating approaches around index cases: mosquito re-invasion of a fumigated neighborhood occurs far quicker than such slow-decaying infections. A more pro-active program encouraging individual level protection of household members of index cases from being bitten by mosquitoes is likely the only reasonable approach to reduce any epidemiological contribution of potentially infected asymptomatic HHM. The relatively low percentage of infected HHM found in this study is likely underestimated, further underscoring the futility of single time point surveys to see whether HHM of index cases are infected: the variable intrinsic incubation period undermines the utility of active "case" detection.

In conclusion, his work not only the significant role that asymptomatic infections can play in DENV epidemiology but also emphasizes the need for alternative strategies to prevent infected individuals from spreading the virus in the course of their daily mobility.

Funding

This study was supported by the European Union Seventh Framework Program (FP7/2007–2013) [under Grant Agreement #282,378 (DENFREE)] (to AS, RP, PS, PM) and a National Research University grant (Mahidol, to AS, PM). The funders had no role in study design, data collection, analysis, decision to publish, or manuscript preparation.

Authors contribution

RP, AS, PS, PM designed the study; KM, ST performed experiments; NP clinical data collection and cohort management; PM, VD supervised experiments; PM, RP analyzed data; SS, TY oversaw patient recruitment and clinical data collection; PM oversaw cohort sample collection; RP, AS coordinated the multinational DENFREE project; PS, PM managed DENFREE: Thailand cohort; PM, RP wrote the manuscript; PM, VD, RP edited the manuscript, and all authors read and approved the final manuscript.

Conflict of interest statement

All authors declared no conflict of interest.

Acknowledgments

We thank Wilawan Chan-in for assistance with figure preparation. We thank all subjects and their families for their participation in this study. We are grateful to the nursing team for assistance in subject recruitment and follow-up. We thank the Faculty of Science, Mahidol University, Central Instrument Facility, Thailand Research Fund (TRG5880121), and Anandamahidol foundation for PM support.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ijid.2020.09.1446>.

References

- Bhatt S, Gething PW, Brady OJ, et al. The global distribution and burden of dengue. *Nature* 2013;496(7446 April 25):504–7.
- Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol* 2002;10:100–3.
- Endy TP. Epidemiology of Inapparent and Symptomatic Acute Dengue Virus Infection: A Prospective Study of Primary School Children in Kamphaeng Phet, Thailand. *Am J Epidemiol* 2002;156(1 July 1):40–51.
- Endy TP, Anderson KB, Nisalak A, et al. Determinants of Inapparent and Symptomatic Dengue Infection in a Prospective Study of Primary School Children in Kamphaeng Phet, Thailand. Harris E, editor. *PLoS Negl Trop Dis* 2011;5(3 March 1):e975.
- Grange L, Simon-Loriere E, Sakuntabhai A. Epidemiological Risk Factors Associated with High Global Frequency of Inapparent Dengue Virus Infections. *Front Immunol* 2014;11(June):5.
- ten Bosch QA, Clapham HE, Lambrechts L, et al. Contributions from the silent majority dominate dengue virus transmission. Ferguson NM, editor. *PLOS Pathog* 2018;14(5 May 3):e1006965.
- Nguyen NM, Kien DTH, Tuan TV, et al. Host and viral features of human dengue cases shape the population of infected and infectious *Aedes aegypti* mosquitoes. *Proc Natl Acad Sci* 2013;110(22 May 28):9072–7.
- Duong V, Lambrechts L, Paul RE, et al. Asymptomatic humans transmit dengue virus to mosquitoes. *Proc Natl Acad Sci* 2015;112(47 November 24):14688–93.
- Anderson RM, May RM. Infectious diseases of humans: dynamics and control. Oxford: Oxford University Press; 1991. p. 757 ISBN: 0-19-854599-16.
- Siler JF, Hall MW, Hitchens AP. Dengue. Its history, epidemiology, mechanism of transmission, etiology, clinical manifestations, immunity, and prevention. *Philippine J Sci* 1926;29:1–304.
- Simmons JS, St. John JH, Reynolds FHK. Experimental studies of dengue. *Philippine J Sci* 1931;44:1–247.
- Nishiura H, Halstead SB. Natural history of dengue virus (DENV)-1 and DENV-4 infections: reanalysis of classic studies. *J Infect Dis* 2007;195(7 April 1):1007–13.
- Kuberski T, Rosen L, Reed D, Mataika J. Clinical and laboratory observations on patients with primary and secondary dengue type 1 infections with hemorrhagic manifestations in Fiji. *Am J Trop Med Hyg* 1977;26:775–83.
- Duyen HT, Ngoc TV, Ha do T. Kinetics of plasma viremia and soluble nonstructural protein 1 concentrations in dengue: differential effects according to serotype and immune status. *J Infect Dis* 2020;203(9):1292–300, doi:<http://dx.doi.org/10.1093/infdis/jir014>.
- Althouse BM, Durbin AP, Hanley KA. Viral kinetics of primary dengue virus infection in non-human primates: a systematic review and individual pooled analysis. *Virology* 2014;. doi:<http://dx.doi.org/10.1016/j.virol.2014.01.015> Mar;452:453:237–246.
- Dussart P, Baril L, Petit L, et al. Clinical and Virological Study of Dengue Cases and the Members of Their Households: The Multinational DENFRAME Project. *PLoS Negl Trop Dis* 2012;6(1 January 24):e1482.
- World Health Organization. Dengue haemorrhagic fever: diagnosis, treatment, prevention and control. 2nd edition 1997 Geneva.
- Aubry M, Roche C, Dupont-Rouzeyrol M. Use of serum and blood samples on filter paper to improve the surveillance of Dengue in Pacific Island Countries. *J Clin Virol* 2012;55(1 September):23–9, doi:<http://dx.doi.org/10.1016/j.jcv.2012.05.010>.
- Lanciotti RS, Calisher CH, Gubler DJ, Chang GJ, Vorndam AV. Rapid detection and typing of dengue viruses from clinical samples by using reverse transcriptase-polymerase chain reaction. *J Clin Microbiol* 1992;30(3 March):545–51.
- Clarke DH, Casals J. Techniques for hemagglutination and hemagglutination-inhibition with arthropod-borne viruses. *Am J Trop Med Hyg* 1958;7(5 September):561–73.
- VSN International. Genstat for Windows. VSN International, Hemel Hempstead, UK. 19th edition Genstat.co.uk: Web page; 2017.
- Yeh WT, Chen RF, Wang L. Implications of previous subclinical dengue infection but not virus load in dengue hemorrhagic fever. *FEMS Immunol Med Microbiol* 2020;48(1):84–90.
- Gubler DJ, Reed D, Rosen L, Hitchcock Jr JR. Epidemiologic, clinical, and virologic observations on dengue in the Kingdom of Tonga. *Am J Trop Med Hyg* 1978;27:581–9.
- Gubler DJ, Suharyono W, Lubis I, Eram S, Sulianti Saroso J. Epidemic dengue hemorrhagic fever in rural Indonesia. I. Virological and epidemiological studies. *Am J Trop Med Hyg* 1979;28:701–10.
- Vaughn DW, Green S, Kalayanaroj S, et al. Dengue in the early febrile phase: viremia and antibody responses. *J Infect Dis* 1997;176:322–30.
- Libraty DH, Endy TP, Hough HS, et al. Differing influences of virus burden and immune activation on disease severity in secondary dengue-3 virus infections. *J Infect Dis* 2002;185:1213–21.
- Perdomo-Celis F, Salgado DM, Narváez CF. Magnitude of viremia, antigenemia and infection of circulating monocytes in children with mild and severe dengue. *Acta Trop* 2017;167:1–8, doi:<http://dx.doi.org/10.1016/j.actatropica.2016.12.011>.
- Katzelnick LC, Fonville JM, Gromowski GD. Dengue viruses cluster antigenically but not as discrete serotypes. *Science* 2015;349(6254 September 18):1338–43, doi:<http://dx.doi.org/10.1126/science.aac5017>.
- Sowath Ly, Camille F, Veasna D, et al. Asymptomatic Dengue Virus Infections Asymptomatic Dengue Virus Infections, Cambodia, 2012–2013. *Emerg Infect Dis* 2019;.

- García G, Sierra B, Pérez AB. Asymptomatic Dengue Infection in a Cuban Population Confirms the Protective Role of the RR Variant of the FcγRIIIa Polymorphism. *Am J Trop Med Hyg* 2010;82:1153–6, doi:http://dx.doi.org/10.4269/ajtmh.2010.09-0353.
- Simon-Lorière E, Duong V, Tawfik A, et al. Increased adaptive immune responses and proper feedback regulation protect against clinical dengue. *Sci Transl Med* 2017;9(405 August 30):eaal5088.
- Halstead SB, O'Rourke EJ. Dengue viruses and mononuclear phagocytes. I. Infection enhancement by non-neutralizing antibody. *J Exp Med* 1977;146:201–17.
- Clapham HE, Cummings DAT, Johansson MA. Immune status alters the probability of apparent illness due to dengue virus infection: Evidence from a pooled analysis across multiple cohort and cluster studies. *PLoS Negl Trop Dis* 2020;11(9): e0005926, doi:http://dx.doi.org/10.1371/journal.pntd.0005926.
- Olkowski S, Forshey BM, Morrison AC. Reduced risk of disease during post-secondary dengue virus infections. *J Infect Dis* 2013;208:1026–33.
- OhAinle M, Balmaseda A, Macalalad AR, et al. Dynamics of dengue disease severity determined by the interplay between viral genetics and serotype-specific immunity. *Sci Transl Med* 2020;3:114–28, doi:http://dx.doi.org/10.1126/scitranslmed.3003084.
- Gordon A, Kuan G, Mercado JC, et al. The Nicaraguan Pediatric Dengue Cohort Study: Incidence of Inapparent and Symptomatic Dengue Virus Infections, 2004–2010. Pettigrew MM, editor. *PLoS Negl Trop Dis* 2013;7(September (9)):e2462.
- Balmaseda A, Standish K, Mercado JC, et al. Trends in Patterns of Dengue Transmission over 4 Years in a Pediatric Cohort Study in Nicaragua. *J Infect Dis* 2010;201(1 January 1):5–14.
- Montoya M, Gresh L, Mercado JC. Symptomatic versus inapparent outcome in repeat dengue virus infections is influenced by the time interval between infections and study year. *PLoS Negl Trop Dis* 2013;7:e2357, doi:http://dx.doi.org/10.1371/journal.pntd.0002357.
- Morrison AC, Minnick SL, Rocha C. Epidemiology of dengue virus in Iquitos, Peru 1999 to 2005: interepidemic and epidemic patterns of transmission. *PLoS Negl Trop Dis* 2010;4:e670, doi:http://dx.doi.org/10.1371/journal.pntd.0000670.