



HAL
open science

Bilaplacian on a Riemannian manifold and Levi-Civita connection

Rémi Léandre

► **To cite this version:**

Rémi Léandre. Bilaplacian on a Riemannian manifold and Levi-Civita connection. ASMDA 2017, 2017, London, United Kingdom. hal-03534041

HAL Id: hal-03534041

<https://hal.science/hal-03534041>

Submitted on 19 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bilaplacian on a Riemannian manifold and Levi-Civita connection

Rémi Léandre

Laboratoire de Mathématiques, Université de Franche-Comté,
25030, Besançon, FRANCE.
email: remi.leandre@univ-fcomte.fr

July 3, 2017

1 Introduction

It is classical in stochastic analysis that the horizontal lift of a diffusion is useful in order to construct canonically the Brownian motion on a Riemannian manifold ([1], [2]). We extend in this note this classical relation on the Brownian motion and the (degenerated) process associated to the horizontal Laplacian to the case of a Bilaplacian. See [7] in the subelliptic case.

We consider a compact Riemannian oriented manifold M endowed with its normalized Riemannian measure dx . x is the generic element of M which is of dimension m . We consider the special orthonormal frame bundle $SO(M)$ endowed with the Levi-Civita connection with canonical projection π on M . u is the generic element of $SO(M)$. We consider the canonical vector fields X_i on $SO(M)$ and the associated horizontal Laplacian

$$L = \sum_{i=1}^m X_i^2 \tag{1}$$

Δ is the Laplace-Beltrami operator on M and Δ^2 the associated Bilaplacian. The Bilaplacian is elliptic, symmetric on $L^2(dx)$ and by elliptic theory ([5], [6]) generates a unique contraction semi group P_t^Δ on $L^2(dx)$ with generic element f .

We can glue on the fiber the normalized "Haar" measures (We omitt the details) such that we get a probability measure du on $SO(M)$. L^2 is symmetric, densely defined on $L^2(du)$ and therefore admits a self-adjoint extension which generates a contraction semi-group P_t^L on $L^2(du)$.

Theorem 1 *If f is a smooth function on M , we have if $\pi u = x$*

$$P_t^\Delta[f](x) = P_t^L[f \circ \pi](u) \tag{2}$$

This theorem enters in our general program to extend stochastic analysis tools to the general theory of linear semi-group (See [8] and [9] for reviews).

2 Proof of the theorem

If u_0 belongs to $SO(\mathbb{R}^m)$ we get clearly

$$P_t^L[f \circ \pi](uu_0) = P_t^L[f \circ \pi(.u_0)](u) = P_t^L[f \circ \pi](u) \quad (3)$$

Therefore, $u \rightarrow P_t^L[f \circ \pi](u)$ defines a function on M . This function when t is moving defines a semi-group. Namely

$$P_{t+s}^L[f \circ \pi](u) = P_t^L[P_s^L[f \circ \pi]](u) = P_t^L[P_s^L[f \circ \pi] \circ \pi](u) \quad (4)$$

where is the right-hand side $P_s^L[f \circ \pi]$ is seen as a function on M .

We are therefore in presence of two semi-groups on $L^2(dx)$. But

$$L(f \circ \pi) = (\Delta f) \circ \pi \quad (5)$$

such that

$$L^2(f \circ \pi) = (\Delta^2 f) \circ \pi \quad (6)$$

Since there is only one semi-group generated by Δ^2 , the result holds.

◇

References

- [1] K.D. Elworthy, *Stochastic differential equations on manifolds*. LMS Lectures Notes 70. Cambridge, England: Cambridge-University Press (1982).
- [2] N. Ikeda and S.Watanabe, *Stochastic differential equations and diffusion processes* Sec. Edi, Amsterdam, Netherland: North-Holland (1989).
- [3] S. Kobayashi and N. Nomizu, *Foundations of differential geometry I* New-York, U.S.A.: Interscience 1963.
- [4] S.Kobayashi and N. Nomizu, *Foundations of differential geometry II* New-York, U.S.A.: Interscience, 1969.
- [5] L. Hoermander, *The analysis of linear partial operators III*. Berlin, Germany: Springer, 1984.
- [6] L. Hoermander, *The analysis of linear partial operators IV*. Berlin, Germany: Springer, 1984.
- [7] R. Léandre, "A geometrical hypoelliptic diffusion", In *Differential geometry and its applications*.J. Bures and al eds.Prague, Cech republic: matfyzpress, 2004, pp 517-523.
- [8] R. Léandre, "Malliavin Calculus of Bismut type in semi-group theory", *Far-East. J. Math. Sci*, 30, pp 1-26 (2008).
- [9] R. Léandre, "Stochastic analysis for a non-markovian generator: an introduction", *Russian Journal of Mathematical Physics*, 22, pp 39–52, 2015.