



## Searching for the Mesozoic pelagic carbonate revolution

Baptiste Suchéras-Marx, Emanuela Mattioli, Fabienne Giraud, Julien Plancq,

Jorijntje Henderiks

### ► To cite this version:

Baptiste Suchéras-Marx, Emanuela Mattioli, Fabienne Giraud, Julien Plancq, Jorijntje Henderiks.  
Searching for the Mesozoic pelagic carbonate revolution. Paleocean - Journées de paléocéanographie,  
2015, Brest, France. hal-03538142

HAL Id: hal-03538142

<https://hal.science/hal-03538142>

Submitted on 24 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License



# Searching for the Mesozoic pelagic carbonate revolution

Baptiste Suchéras-Marx<sup>1</sup>

Emanuela Mattioli<sup>2</sup>, Fabienne Giraud<sup>3</sup>, Julien Plancq<sup>4</sup>, Jorijntje Henderiks<sup>5</sup>

1 CEREGE UM34, Aix-Marseille Université, CNRS, IRD, 13454 Aix-en-Provence, France.



3 ISTerre, UMR 5275, Université de Grenoble 1, 1381 rue de la piscine, BP 53, 38041 Grenoble Cedex 9, France.

4 Laboratoire de Morphodynamique Continentale et Côtière (M2C), UMR CNRS 6143, Université Caen-Basse Normandie, Campus 1, 24 rue des tilleuls, Caen, France.

5 Paleobiology Program, Department of Earth Sciences, Uppsala University, Villavägen 16, SE-75 236 Uppsala, Sweden.

## Introduction :

Carbonate accumulation is largely sustained by **pelagic carbonate producers**, in the form of calcite platelets (calcareous nannofossils including coccoliths) produced by calcifying haptophyte algae and the calcite tests of planktic foraminifera. Although deep-sea carbonates (calcareous oozes or chalks) have production and accumulation rates that are orders of magnitude lower than shallow water carbonates, they cover a basin area that is orders of magnitude larger. Pelagic carbonate producers were not present all along the Phanerozoic. The first part of the Phanerozoic is considered as a time interval where the **carbonate production was supported by benthic producers** in carbonate platforms and is referred as *Neritan Ocean* (Zeebe and Westbroek, 2003). The second part of the Phanerozoic is referred as *Cretan Ocean* where the **carbonate production was supported by pelagic producers** still relevant in modern oceans. The transition from Neritan to Cretan oceans is considered to be an important **revolution in oceans** history, deeply changing the organization of the carbonate system and, among other, leading to the formation of the carbonate compensation depth (CCD). This pelagic carbonate revolution, named by Roth (1986) the *Kuenen Event* is supposed to occur during the Mesozoic in relation with the emergence of calcareous nannofossil. In this study, we explore the **timing** of the *Kuenen Event* based on **quantification of calcareous nannofossils accumulation** and **mass accumulation**, calcareous nannofossil diversity and assemblage mean size and qualitative observations of pelagic carbonate deposits.



## Nannofossil flux evolution :

We have compiled **quantitative data** of absolute abundances and burial **fluxes** of calcareous nannofossils over the past 200 Myr. The nannofossil burial flux **increases** from the Early Jurassic up to a **maximum during the Early Cretaceous**. After, the burial fluxes fluctuate but maximum values are stable. Meanwhile, the **nannofossil diversity increases** during the Mesozoic up to a maximum at the K/Pg boundary. After high diversity during the Paleocene/Eocene interval, nannofossil species abundance decreases reaching minimum and relative low diversity in the Neogene. The coccolith assemblage size mean has the same evolution as diversity suggesting **dependance** of size from diversity. Comparing those three records, we propose a model of nannofossil evolution for the Mesozoic in two steps: 1) **Invasion** of environments from the Early Jurassic to the Early Cretaceous; 2) **Specialization** of nannofossils leading to compartmentalization of ecological niches until the K/Pg boundary.

## Carbonate production revolution :

Calcareous nannofossils first occurrence is observed during the **Late Triassic** but with low diversity and abundance. According to qualitative observations, **nannofossil-dominated deposits** only occurred during the **Late Jurassic** or the **Early Cretaceous**. But both observations are challenged by quantitative studies. Common occurrence of chalk or carbonate oozes produced by planktic producers started during the **Late Cretaceous**. The occurrence of pelagic sediments resulting from planktic producers is **not synchronous** with the Early Cretaceous nannofossil flux maximum. The **planktic foraminifera** diversified and increased in abundance at the **end of the Early Cretaceous**. Hence, the emergence of pelagic carbonate accumulation in the Late Cretaceous is related to the **concurrent nannofossil and planktic foraminifera productions**. We consider the *Kuenen Event* to correspond to a **long term event** started with the appearance of calcareous nannofossils and ended with planktic foraminifera diversification.

## Perspectives :

There is **no clear abiotic forcing** identified that has lead to the increase in nannofossil and the emergence of pelagic carbonate accumulation. Despite relative similarities between nannofossil flux and pCO<sub>2</sub>, the **mechanistic link** between both is not yet identified. The carbonate pelagic revolution may be the results of **concurrent biotic evolution** and **abiotic changes** that need further explorations.