



# Lagrangian approach to one-dimensional constrained systems

Charlotte Perrin

## ► To cite this version:

Charlotte Perrin. Lagrangian approach to one-dimensional constrained systems. Dispersive Hydrodynamics and oceanography: from experiments to theory, Aug 2017, Les Houches, France. hal-03538831

HAL Id: hal-03538831

<https://hal.science/hal-03538831>

Submitted on 21 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Lagrangian approach to one-dimensional constrained systems

Charlotte Perrin  
joint work with Michael Westdickenberg

Institut für Mathematik  
RWTH Aachen University

Dispersive Hydrodynamics and oceanography  
August 2017

# Constrained systems in fluid mechanics

- partially free surface flows: flows in closed channels, floating structures, etc.
- mixtures

→ existence of a max. constraint (height of the flow, volume fraction, etc)  $\rho \leq \rho^*$

## typical system

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla p = 0 \\ 0 \leq \rho \leq \rho^*, \ (\rho^* - \rho)p = 0, \ p \geq 0 \end{cases}$$

in the following:  $d = 1, \rho^* = 1$


## two-phase flow

- pressureless equations in the free domain
- $\operatorname{div} u \geq 0$  on  $\rho = \rho^*$
- activation of the pressure  $p$  on  $\rho = \rho^*$   
Lagrange multiplier associated to the constraint

# Modelling of immersed granular flows

- mixture: solids particles / liquid

- ▶ volume fraction of grains



- constitutive laws in the congested zones ?

- ▶ 1d model with memory effects proposed by Lefebvre-Lepot & Maury (2012)



$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_x p = \rho f \\ \textcolor{blue}{\partial_t \gamma + u \partial_x \gamma = -p} \\ 0 \leq \rho \leq 1, \quad (1 - \rho) \gamma = 0, \quad \gamma \leq 0 \end{cases}$$

# Memory effects, case of a single particle (Maury, 2007)

vanishing liquid viscosity  $\nu = \varepsilon$

$$\ddot{q}_\varepsilon + \varepsilon \frac{\dot{q}_\varepsilon}{q_\varepsilon} = f$$

$$\varepsilon \rightarrow 0$$



$$\begin{cases} \dot{q} + \gamma = u^0 + \int_0^t f(s) ds \\ \gamma \leq 0, \quad q \geq 0, \quad q\gamma = 0 \end{cases}$$



$$\begin{cases} \ddot{q} - \mu = f \\ \dot{\gamma} = -\mu \\ \text{spt}(\mu) \in \{t, q(t) = 0\} \\ \dot{q}(t^+) = P_{C_{q,\gamma}} \dot{q}(t^-) \\ \gamma \leq 0, \quad q \geq 0 \end{cases}$$

# Macroscopic models

- Lefebvre-Lepot, Maury (2012): Micro. variable  $q \rightarrow$  Macro. variable  $1 - \rho$

## suspension

compressible with  
singular viscosity

$$\begin{cases} \partial_t \rho_\varepsilon + \partial_x (\rho_\varepsilon u_\varepsilon) = 0 \\ \partial_t (\rho_\varepsilon u_\varepsilon) + \partial_x (\rho_\varepsilon u_\varepsilon^2) \\ \quad - \partial_x (\lambda_\varepsilon(\rho_\varepsilon) \partial_x u_\varepsilon) = \rho f \\ \lambda_\varepsilon(\rho_\varepsilon) = \frac{\varepsilon}{1 - \rho_\varepsilon} \end{cases}$$

## granular

two-phase flow with  
adhesion potential

$$\xrightarrow[\varepsilon \rightarrow 0]{\text{FORMAL}} \begin{cases} \partial_t \rho + \partial_x (\rho u) = 0 \\ \partial_t (\rho u) + \partial_x (\rho u^2) + \partial_x \mathbf{p} = \rho f \\ \partial_t \gamma + u \partial_x \gamma = -p \\ 0 \leq \rho \leq 1, (1 - \rho)\gamma = 0, \gamma \leq 0 \end{cases}$$

**Issue:** existence of solutions to the granular model

## Plan of the talk:

- ① Lagrangian approach to pressureless gas equations
- ② addition of the maximal density constraint
- ③ granular system with memory effects

# Lagrangian approach to pressureless gas eq.

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) = 0 \end{cases} \quad \begin{array}{l} \text{Bouchut (1994)} \\ \bullet \text{ concentration phenomena on the density} \end{array}$$

- some references:
  - ▶ approx. by sticky particles: Brenier, Grenier (1998), Natile, Savaré (2009)
  - ▶ viscous regularization: Sobolevskii (1997), Boudin (2000)
  - ▶ hydrodynamical limit: Jabin, Rey (2016)

# Lagrangian approach to pressureless gas eq.

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) = 0 \end{cases}$$

Bouchut (1994)

- concentration phenomena on the density

- some references:

- approx. by sticky particles: Brenier, Grenier (1998), Natile, Savaré (2009)  
→ Lagrangian point of view via the monotone rearrangement
- viscous regularization: Sobolevskii (1997), Boudin (2000)
- hydrodynamical limit: Jabin, Rey (2016)

# Dynamics of sticky particles

- $N$  particles  $P_i = (m_i, x_i, u_i)$  such that  $\sum_{i=1}^N m_i = 1$ ,  $x_1 < x_2 < \dots < x_N$
- free motion ...

$$x_i(t) = x_i(0) + t u_i(0), \quad u_i(t) = u_i(0)$$

... until the first time of collision  $t^1$  when  $x_j(t^1) = x_{j+1}(t^1) = \dots = x_k(t^1)$   
particles  $P_j, P_{j+1}, \dots, P_k$  form a new particle  $P$  with

- ▶ mass  $m = m_j + \dots + m_k$
- ▶ velocity  $u = \frac{m_j u_j(t^1) + m_{j+1} u_{j+1}(t^1) + \dots + m_k u_k(t^1)}{m}$

then  $P$  evolves freely with velocity  $u$  until the next collision

- measure description:  $\rho_t = \sum_{i=1}^{N(t)} m_i \delta_{x_i(t)}$ ,  $(\rho u)_t = \sum_{i=1}^{N(t)} m_i u_i \delta_{x_i(t)}$  satisfy
- $$\begin{cases} \partial_t \rho + \partial_x (\rho u) = 0 \\ \partial_t (\rho u) + \partial_x (\rho u^2) = 0 \end{cases} \quad \text{in } \mathcal{D}'$$

# Lagrangian representation of the sticky particles dynamics

- use of the **monotone rearrangement**  $X_\rho : (0, 1) \rightarrow \mathbb{R}$   
(pseudo inverse of the cumulative distributive function)

$$X_\rho(w) := \inf \{x \in \mathbb{R}, \rho((-\infty, x]) > w\}$$

→  $X_\rho$  is nondecreasing and right-continuous

- it pushes the Lebesgue measure on  $(0, 1)$  onto  $\rho$



$$\rho = (X_\rho)_\# \mathcal{L}_{(0,1)}^1 \quad \text{i.e.}$$

$$\int_{\mathbb{R}} \varphi(x) d\rho(x) = \int_0^1 \varphi(X_\rho(w)) dw$$

$$\rho_t \in \mathcal{P}_2(\mathbb{R}) \longleftrightarrow X_t \in K$$

$K = \text{cone of non-decreasing functions of } L^2(0, 1)$

$$\rho = 0.5 \delta_{\{-0.5\}} + 0.2 \delta_{\{0.5\}} + 0.3 \delta_{\{1\}}$$





- velocity  $U_t = u_t \circ X_t \in \mathcal{H}_{X_t}$ ,  $\mathcal{H}_{X_t} = \left\{ U = u \circ X_t \text{ for some } u \in L^2_{\rho_t}(\mathbb{R}) \right\}$   
 $\rightarrow U_t$  is constant where  $X_t$  is constant

**Property of sticky particles**  $\mathcal{H}_{X_t} \subset \mathcal{H}_{X_s}$ ,  $U_t = P_{\mathcal{H}_{X_t}}(U_s) \quad \forall s \leq t$

## Lagrangian representation


The sticky particles system satisfies  $X_t = P_K(X_0 + tU_0)$

- $X_0 + tU_0$  corresponds to the "collision free" motion



ref: Natile, Savaré (2009)

two other Lagr. characterizations  
(differential inclusions) are derived



## Direct construction of weak sol. to the pressureless eq.

- definition of the transport  $X_t := P_K(X_0 + tU_0)$

$\implies t \mapsto X_t$  is Lipschitz continuous

$\implies$  definition of the Lagrangian velocity  $U_t := \frac{d}{dt}X_t$  a.e.  $t$

- proof of

$$U_t = P_{\mathcal{H}_{X_t}}(U_0) \quad \text{a.e. } t$$

$\rightarrow U_t$  is constant where  $X_t$  is constant

$\rightarrow \exists u_t \in L^2_{\rho_t}$  such that  $U_t = u_t \circ X_t$

- the Eulerian variables satisfy the weak formulation of the pressureless system

**Remark:** one does not use the sticky properties

$$(\mathcal{H}_{X_t} \subset \mathcal{H}_{X_s}, U_t = P_{\mathcal{H}_{X_t}}(U_s) \ \forall s \leq t)$$

ref: Cavalletti, Sedjro, Westdickenberg (2015)

# Constrained pressureless system

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_x p = 0 \\ 0 \leq \rho \leq 1, \quad (1 - \rho)p = 0, \quad p \geq 0 \end{cases}$$

Bouchut, Brenier, Cortes, Ripoll (2000)

- no more concentration phenomena

- some references

- ▶ theoretical existence results: Berthelin in 1d (2002), multi-d (2016)
- ▶ numerical studies: Degond, Hua, Navoret (2011), Preux (2016)

# Max. density constraint from the transport point of view

$$\rho_t := (X_t)_\# \mathcal{L}^1_{|(0,1)}$$

**Proposition**  $\rho_t \ll \mathcal{L}^1$  if and only if  $\partial_w X_t > 0$  a.e. on  $(0, 1)$  and in this case

$$\rho_t(X_t(w)) = \frac{1}{\partial_w X_t(w)} \quad \text{a.e. on } (0, 1)$$

**Consequence**

$$\rho_t \leq 1 \iff X_t \in \tilde{K} = K + \text{id}$$

$\tilde{K}$  is a convex subset of  $L^2(0, 1)$

in the following  $X_t = S_t + \text{id}$ ,  $S_t \in K$

**Congested domain**




$$\Omega_t = \left\{ w \in (0, 1), S_t \text{ is constant on a neighborhood of } w \right\}$$

# Sticky blocks

- particles  $P_i = (m_i, x_i, u_i) \rightarrow$  blocks  $B_i = (\ell_i, x_i, u_i)$
- sticky dynamics  $\rightarrow \mathcal{H}_{S_t} \subset \mathcal{H}_{S_\tau} \quad \forall \tau \leq t$   
property of the Lagrangian velocity  $U_t = P_{\mathcal{H}_{S_t}}(U_\tau) \quad \forall \tau \leq t$

## Lagrangian representation

The sticky blocks system satisfies  $X_t = P_{\tilde{\mathcal{K}}}(X_0 + tU_0)$



# Resulting pressure

- define  $\Gamma_t$  such that  $\partial_w \Gamma_t := U_t - U_0 = P_{\mathcal{H}_{S_t}}(U_0) - U_0$

**properties:**  $\text{spt } \Gamma_t \subset \Omega_t, \quad \Gamma_t \leq 0$

- define  $P_t := -\frac{d}{dt} \Gamma_t$  in  $\mathcal{D}' \rightarrow \text{spt } P_t \subset \Omega_t$

**sticky property**  $U_t = P_{\mathcal{H}_{S_t}}(U_\tau) \quad \forall \tau \leq t \rightarrow P_t \geq 0$

- the associated Eulerian variables:  $\gamma_t = \Gamma_t \circ X_t^{-1}, p_t = P_t \circ X_t^{-1}$  satisfy in  $\mathcal{D}'$

$$\partial_t \gamma + u \partial_x \gamma = -p \quad \text{on } \{\rho = 1\}$$

## Theorem (Perrin, Westdickenberg)

Let  $\rho_0 \in \mathcal{P}_2(\mathbb{R})$  with  $\rho_0 \leq 1$  a.e.,  $u_0 \in L^2_{\rho_0}(\mathbb{R})$ , there exists a global weak sol. to

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_x p = 0 \\ 0 \leq \rho \leq 1, \quad p \geq 0, \quad \text{spt } p \subset \{\rho = 1\} \end{cases}$$

**remark:** Berthelin's hypothesis  $\rho_0 \in L^1(\mathbb{R})$ ,  $0 \leq \rho_0 \leq 1$ ,  $u_0 \in L^\infty(\mathbb{R})$ .

# Granular system with memory effects

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x (\rho u) = 0 \\ \partial_t (\rho u) + \partial_x (\rho u^2) + \partial_x p = pf \\ \partial_t \gamma + u \partial_x \gamma = -p \\ 0 \leq \rho \leq 1 \\ (1 - \rho)\gamma = 0, \quad \gamma \leq 0 \end{array} \right.$$

- no sticky dynamics due to the external force  $f$
- no sign on the pressure  $p$

- introduction of system by Lefebvre, Maury (2012) but no existence result
- similar system with additional viscosity: C.P. (2016)
  - ▶ weak solutions obtained by a singular limit from compressible NS eq.

**Strategy:** construct directly weak sol. without any discrete approximation

# Definition of the transport

- set of admissible transports

maximal density constraint  $\rho_t \leq 1 \implies X_t = S_t + \text{id} \in \tilde{K}, S_t \in K$

- free trajectory and effective transport

▶ free velocity  $U_t^{\text{free}} = U_0 + \int_0^t f(s, X_s) ds$

▶ associated free trajectory  $X_0 + \int_0^t U_s^{\text{free}} ds$

we would like to define

$$X_t = P_{\tilde{K}} \left( X_0 + \int_0^t U_s^{\text{free}} ds \right)$$

→ conditions on  $f$  to ensure the existence and uniqueness of  $X_t \forall t$

# Velocity and adhesion potential in Lagrangian coordinates

- $t \mapsto X_t$  is Lipschitz continuous
- definition of the Lagrangian velocity  $U_t := \frac{d}{dt} X_t$  a.a.  $t$

$$\implies U_t = P_{\mathcal{H}_{S_t}}(U_t^{\text{free}}) \quad \text{where } \mathcal{H}_{S_t} := \{u \circ S_t, u \in L^2_{\rho_t}\}$$

- adhesion potential

$$\partial_w \Gamma_t := U_t - U_t^{\text{free}} + \partial_w \Gamma_0$$

$$\begin{cases} \text{spt } \Gamma_0 \in \Omega_0 \\ \Gamma_0 \leq 0 \end{cases} \implies \begin{cases} \text{spt } \Gamma_t \in \Omega_t \\ \Gamma_t \leq 0 \end{cases}$$

# Reformulation of the granular system

we remove the pressure from the eq. by use of the **effective velocity**  $v = u - \partial_x \gamma$

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2) + \partial_x p = \rho f \\ \partial_t \gamma + u \partial_x \gamma = -p \\ 0 \leq \rho \leq 1, \quad (1 - \rho)\gamma = 0, \quad \gamma \leq 0 \end{cases}$$

→

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho v) + \partial_x(\rho u v) = \rho f \\ v = u - \partial_x \gamma \\ 0 \leq \rho \leq 1, \quad (1 - \rho)\gamma = 0, \quad \gamma \leq 0 \end{cases}$$

## Theorem (Perrin, Westdickenberg (ArXiv))

Assume that

- $f \in L^\infty(0, T; \text{Lip} \cap L^\infty(\mathbb{R}))$
- $\rho_0 \in \mathcal{P}_2(\mathbb{R})$  with  $\rho_0 \leq 1$  a.e.
- $(u_0, \gamma_0) \in L^2_{\rho_0}(\mathbb{R}) \times W^{1,2}(\mathbb{R})$  with  $(1 - \rho_0)\gamma_0 = 0$ ,  $\gamma_0 \leq 0$  a.e.

Let  $X_0$  be such that  $\rho_0 = (X_0)_\# \mathcal{L}_{(0,1)}^1$  and  $U_0 := u_0 \circ X_0$ ,  $\Gamma_0 := \gamma_0 \circ X_0$ .

One can then define

$$U_\tau^{free} := U_0 + \int_0^\tau f(s, X_s) ds, \quad X_t := P_{\tilde{K}} \left( X_0 + \int_0^t U_\tau^{free} d\tau \right) \quad \forall t$$
$$U_t := \frac{d}{dt} X_t, \quad \partial_w \Gamma_t := U_t - U_t^{free} + \partial_w \Gamma_0 \quad \text{a.e. } t$$

and the associated family of Eulerian variables  $(\rho_t, u_t, \gamma_t)$  is a weak solution of


$$\begin{cases} \partial_t \rho + \partial_x (\rho u) = 0 \\ \partial_t (\rho(u - \partial_x \gamma)) + \partial_x (\rho u (u - \partial_x \gamma)) = \rho f \\ 0 \leq \rho \leq 1, \quad (1 - \rho)\gamma = 0, \quad \gamma \leq 0 \end{cases}$$

# Numerical simulation

→ minimization at each time step of  $\|X_0 + \int_0^{t_h} U_s^{\text{free}} \, ds - X\|_2^2$

under the constraint  $X - \text{id} \in K$

→ Python software for convex optimization CVXOPT



# Conclusion

- Lagrangian approach
  - ▶ Euler under max. constraint → improvement of the existing theoretical result
  - ▶ system with memory effects → first existence result
  - ▶ easy implementation
- direct extension to the heterogeneous case  $\rho \leq \rho^*(t, x)$  with

$$\partial_t \rho^* + u \partial_x \rho^* = 0$$