



Identification and partial characterization of antifungal and antibacterial activities of two *Bacillus* sp. strains isolated from salt soil in Tunisia

Ben Maachia Sihem, Errakhi Rafik, Florence Mathieu, Chérif Mohamed, Ahmed Lebrihi

► To cite this version:

Ben Maachia Sihem, Errakhi Rafik, Florence Mathieu, Chérif Mohamed, Ahmed Lebrihi. Identification and partial characterization of antifungal and antibacterial activities of two *Bacillus* sp. strains isolated from salt soil in Tunisia. *African Journal of Microbiology Research*, 2011, vol. 5 (n° 13), pp. 1599-1608. 10.5897/AJMR11.073 . hal-03539451

HAL Id: hal-03539451

<https://hal.science/hal-03539451>

Submitted on 21 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 5980

To link to this article: DOI:10.5897/AJMR11.073
URL: <http://dx.doi.org/10.5897/AJMR11.073>

To cite this version: Sihem, Ben Maachia and Rafik, Errakhi and Mathieu, Florence and Mohamed, Chérif and Lebrihi, Ahmed (2011) Identification and partial characterization of antifungal and antibacterial activities of two *Bacillus* sp. strains isolated from salt soil in Tunisia. *African Journal of Microbiology Research*, vol. 5 (n°13). pp. 1599-1608. ISSN 1996-0808

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes.diff.inp-toulouse.fr

Identification and partial characterization of antifungal and antibacterial activities of two *Bacillus* sp. strains isolated from salt soil in Tunisia

Ben Maachia Sihem^{1,2,4*}, Errakhi Rafik³, Mathieu Florence¹, Chérif Mohamed⁴ and Lebrihi Ahmed¹

¹Laboratoire de Génie chimique UMR 5503 (CNRS/INPT/UPS), Institut National Polytechnique de Toulouse, 1, av. de l'Agrobiopole, 31326 Castanet-Tolosan, France.

²Centre Régionale de Recherches en Agriculture Oasienne, Dégache, Tozeur, Tunisia.

³Plateforme de biotechnologie AGRONUTRITION SAS 3, avenue de l'Orchidée Parc Activestre 31390 CARBONNE – France.

⁴Laboratoire de phytopathologie, Institut National Agronomique de Tunisie, Tunis, Tunisie.

Two *Bacillus* sp. strains (B29 and B27) isolated from soil in the South of Tunisia were tested for their abilities to produce antimicrobial compounds. Both strains showed antimicrobial activity against Gram-positive and Gram-negative bacteria, yeasts and fungi. The produced compounds were extracted by using four different solvents. The hexane solvent allowed to obtain maximum of activity of the strain B29. The activity of the strain B27 was not elucidated by the four solvents used. Bio-autography results of B29 hexane extract revealed presence of different antibiotics and antifungal compounds with different R_f values of 0.3 and 0.76 for antifungal compounds and of 0.12, 0.14, 0.19 and 0.3 for antibacterial ones. Two active fractions were isolated from the culture broth of the strain B29 by semi-preparative high performance liquid chromatography (HPLC). The partial sequencing of the 16S rDNA gene was used to identify the two *Bacillus* strains. They may be assigned to new *Bacillus* species.

Key words: Antimicrobial activity, *Bacillus* sp., characterization, salt soil, Tunisia.

INTRODUCTION

An analysis of the number of pathogenic bacteria and fungi indicated that they become resistant to antibiotics in common use. These antibacterial and antifungal resistances are presently an urgent focus of research and new antibiotics are necessary to combat these pathogens. An investigation of the potential sources of this molecules indicated that over 60% of approved drugs are derived from natural compounds (Cragg et al., 1997; Newman et al., 2003). Many organisms, including microorganisms, plants and some animals, have the capability to synthesize bioactive natural products. These natural products constitute a library of compounds with a large and privileged structural diversity, showing a variety of biological activities. In fact, many of them have been

used for agricultural or pharmaceutical applications (Salas and Mendez, 2007). The search for new, safer, broad-spectrum antifungal antibiotic with greater potency has been progressing slowly (Gupte et al., 2002). In the course of screening for new antibiotics, several research studies are currently oriented towards isolation of new microorganism's species from different soils and ecosystems (Mellouli et al., 2003; Errakhi et al., 2007). Within microorganisms, the *Bacillus* species are one of the largest source of bioactive natural products. Various studies have confirmed that *Bacillus* species have a wide range of antimicrobial activities since they are used as antifungal (Milner et al., 1995), antibacterial (Yilmaz et al., 2006), antiviral (Steller et al., 1999), antiameobocytic (Galvez et al., 1994) and antimycoplasma agents (Peypoux et al., 1999). These are often complex secondary metabolites with great potential for biotechnological and biopharmaceutical applications. A well-known class

*Corresponding author. E-mail: maachiasihem@yahoo.fr.

of such compounds includes the lipopeptides surfactin (Cho et al., 2003), fengycin (Vanittanakom et al., 1986), and the iturin compounds (Cho et al., 2009), iturin A, B and C (Peypoux et al., 1978), mycosubtilins (peypoux et al., 1976), bacillomycins (Peypoux et al., 1984), plipastatin (Tsuge et al., 1996) and di- and tripeptides such as bacilysin (Walker and Abraham, 1970) are peptide antibiotics with potent antimicrobial activities.

In fact, the list of novel microorganisms, especially *Bacillus*, and products found in microbiologically unexplored ecosystems around the world suggest that a careful exploration of other habitats might continue to be useful. South of Tunisia is an explored ecosystem and may be a source of new microorganisms producing new compounds. In the present work, we describe a preliminary characterization of two bacterial strains isolated from south of Tunisia and to assess their antimicrobial activity based on molecular methods as well as the production, the isolation and the partial characterization of their antibiotics.

MATERIALS AND METHODS

Microorganism's strains

Bacillus strains, denoted B27 and B29, were previously isolated from salt soil in the south of Tunisia and identified as *Bacillus* sp. These bacteria were routinely grown on nutrient agar (NA) medium at 30°C for further treatments. *Escherichia coli* CIP 102 400, *Staphylococcus aureus* CIP658, *Saccharomyces cerevisiae* ATCC 9226, *Mycobacterium phlei* CIP105 389, *Micrococcus luteus* CIPA270, *Pseudomonas aeruginosa* CIP 100720, *Mucor ramannianus* NRLL 1829 (strain collection of Laboratoire de génie chimique, Toulouse) and *Botrytis cinerea* (strain isolated from leaves of vine plants) as indicator microorganisms for the biological assays.

Antimicrobial activity

The antimicrobial and antifungal activity of the isolates was determined by the plate diffusion method (Barakate et al., 2002; Errakhi et al., 2007) against Gram-positive bacteria (*M. luteus*, *S. aureus*, and *M. phlei*), Gram-negative bacteria (*E. coli* and *P. aeruginosa*), yeasts (*S. cerevisiae*), and filamentous fungi (*M. ramannianus* and *B. cinerea*). Isolates were grown on NA medium and two discs (6 mm in diameter) were cut and placed on Bennett medium (beef extract (Merck, Germany) 1 g/l; glucose (Merck) 10 g/l; peptone (Merck) 2 g/l; yeast extract (Merck) 1 g/l and agar (Difco) 15 g/l; for bacteria) and potato dextrose agar (PDA) medium (for fungi and yeasts) which were seeded with appropriate test organism. Plates were first kept in a refrigerator (4°C) for at least 2 h to allow the diffusion of any antibiotics produced, then incubated at 30°C. Inhibition zones were determined after 24 h for bacteria and yeast and after 48 h for fungi.

Production of antimicrobial activity in various media

Antimicrobial activity of the two *Bacillus* strains was evaluated on 4 media that is YMA (peptone 0.3%, yeast extract 0.5%, mannitol

2.5% and agar 15 g/l), NA, PDA and Bennett (BT). The two bacteria were inoculated on plates of 90 mm diameter on each of this media and incubated at 30°C for 24 h. The activity against *M. luteus*, *E. coli*, *P. aeruginosa*, *S. cerevisiae* and *M. ramannianus* was recorded by the plate diffusion method. Three discs (10 mm in diameter), for each bacteria and each medium, were cut and placed on BT (for bacteria) and PDA (for fungi and yeast) which were seeded with appropriate test organisms. Plates were first kept in a refrigerator (4°C) for at least 2 h to allow the diffusion of antibiotics produced, then incubated at 30°C. Inhibition zones were determined after 24 h for bacteria and yeast and after 48 h for fungi.

Kinetics of antimicrobial products of the *Bacillus* sp. strains in potato dextrose medium

The two *Bacillus* strains were selected for their interesting antifungal properties. Kinetics of antimicrobial products was assessed on potato dextrose broth (PDB) medium. The pH media was adjusted to 7.2 and 100 ml aliquots were dispensed into 1 l Erlenmeyer flasks. The inoculated flasks were put on a shaker (250 rpm; 30°C). 2 ml of the liquid culture were centrifuged to remove most of the bacterial cells, the supernatant was passed through a sterile filter 0.20 µm membranes, and the resulting broth was used for antifungal activity determination. The activity against *M. luteus*, *S. cerevisiae*, *M. ramannianus*, and *B. cinerea* was regularly recorded each 2 h by the agar diffusion method (well technique; each well of 10 mm in diameter was filled with 50 µl of supernatant). Fungi, bacteria, and yeast spore suspensions were prepared by suspending loopfuls of hyphae and colonies from PDA slant incubated at 22°C (*B. cinerea*) and 30°C for the other organisms. The suspension was filtered with sterile cotton (for *B. cinerea*). A 10⁸ CFU/ml spore suspension was obtained through dilutions and counting with a hemacytometer and adjusted to 10⁵ CFU/ml by serial dilution. 4 ml of the spore suspension was suspended under the PDA plates, containing the wells with the supernatant. The Petri dishes were incubated for 4 h at 4°C to allow the diffusion of produced antimicrobial metabolites and then incubated at 30°C for 24 to 48 h. The relative inhibition data is presented as the diameter of development inhibition around wells.

Production, extraction and detection of antibiotics

Each *Bacillus* strain was cultivated in PDB medium in four 500 ml Erlenmeyer flasks each containing 100 ml of medium. The cultures were incubated at 30°C for 3 days under constant agitation of 250 rpm. The liquid culture was centrifuged for 20 min at 8000 g to remove the bacteria. The supernatant was shared in 4 equal volumes of 100 ml, and then each was extracted with 200 ml of organic solvent. A range of extraction solvents was screened for effectiveness, including n-hexane, dichloromethane, n-butanol and ethyl acetate. The organic extracts were evaporated to dryness using a Rotavapor (Laborota 4000). The resulting dry extracts were recuperated in 2 ml of methanol and subjected to biological assay (25 µl) (disk of 6 mm in diameter, Institut Pasteur) against *M. luteus*, *S. cerevisiae*, *M. ramannianus*, *P. aeruginosa*, and *B. cinerea*. The solvents which gave the highest inhibition diameter were then retained for the detection of antibiotics by bioautography.

Purification and partial characterization of antifungal products of *Bacillus* sp. strains

Preparative chromatography with silica gel plates (Merck Art. 5735, Kiesselgel 60F254) was used for the partial purification of antimicrobial products. For this, 20 µl of n-hexane extract were

spotted onto 20 × 20 cm silica gel plates, and then developed with butanol-acetic acid-water (3:1:1). The developed thin layer chromatography (TLC) plates were air-dried overnight at 37°C to remove all traces of solvents. They were then placed in a plastic bioassay dish (23 × 23 × 2.2 cm³, Fisher Scientific Labosi) and overlaid with 100 ml of PDB medium (containing 7 g/l of agar) seeded with *M. luteus* or *M. ramannianus*. After the agar had set, the plate was incubated at 30°C. After 2 days, reddish brown-colored fungal growth or opaque bacterial growth was visible. Clear areas due to inhibition of microorganism growth indicated the location of antibiotic compounds on the TLC plates. The retention factor (RF) of each spot was recorded.

The liquid cultures of B27 and B29 strains at 30°C for three days were harvested to remove the biomass. The cell-free supernatant fluid was extracted with hexane for B29 and with ethyl acetate for B27. Antibiotics were purified by HPLC (Waters: controller 600, pump 600, dual λ absorption detector 2487, Linear Recorder); Column C18 (250 × 7.8 mm UP ODS); mobile phase: Linear gradient of methanol-H₂O from 0 to 100% for 40 min, Flow rate: 1.5 ml/min, detection: UV at 220 nm. Different peaks were collected separately, concentrated and then tested against *M. ramannianus* and *M. luteus*. After purification, the antibiotics were subjected to spectroscopic studies. The absorption spectrum of active extracts in methanol were recorded in the UV region (180 to 500 nm) by using a UV-visible Prekin-Elmer (Lambda 20) spectrophotometer and compared with those of known antifungal antibiotics.

Identification of bacteria

DNA extraction

Single isolated colonies of the two bacterial cultures were taken from agar plates and suspended in 500 μ l of lysis buffer (400 mM Tris-HCl (pH 8.0), 60 mM EDTA (pH 8.0), 150 mM NaCl, 1% sodium dodecyl sulfate). The tube is then left at room temperature for 10 min. After adding 150 μ l of potassium acetate (pH 4.8), the tube is vortexed briefly and spun at >10 000 g for 1 min. After transferring the supernatant to a new Eppendorf tube, an equal volume of isopropyl alcohol is added. The tube is mixed by inversion briefly, spun at >10 000 g for 2 min and the supernatant is discarded. The resultant DNA pellet is washed in 300 μ l of 70% ethanol. After the pellet is spun at 10 000 rpm for 1 min, the supernatant is discarded. The DNA pellet is air dried and dissolved in 50 μ l of sterile ultrapure water (Liu et al., 2000).

PCR amplification

PCR amplification of almost full length 16SrRNA gene was carried out with specific primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492r (5'-GGTTACCTTGTTACGACTT-3'). A 50 μ l reaction volume PCR was performed using about 100 ng of genomic DNA, 10X Reaction buffer, 10 mM (each) deoxynucleoside triphosphates, 25 mM MgCl₂ and 0.5 U of DNA polymerase. The PCR was performed in an automated Gene Amp PCR System thermal cycler under the following conditions. The amplification conditions were as follows 98°C for 4 min (denaturation), 55°C for 1 min (annealing), 72°C for 3 min (elongation) and 72°C for 10 min final elongation. Expected PCR product of around 1.5 Kb was checked by electrophoresis of 5 μ l of the PCR product on 1% agarose gel in 1X TBE buffer and stained with ethium bromide 0.5 μ g / ml.

The PCR products obtained were submitted to GenomExpress for sequence determination. The same primers as above and an automated sequencer were used for this purpose. The analysis of sequences was done at NCBI server

(<http://www.ncbi.nlm.nih.gov/BLAST>). All the analyzed sequences have been deposited in genbank.

Phylogenetic analysis

Phylogenetic and molecular evolutionary analyses were conducted using softwares included in MEGA version 3.0 (Kumar et al., 2004) package. The 16S rDNA sequence of B27 and B29 strains was aligned using the CLUSTAL W program (Thompson et al., 1994) against corresponding nucleotide sequences of representatives of the genus *Bacillus* retrieved from GenBank. Evolutionary distance matrices were generated as described by Jukes and Cantor (1969) and a phylogenetic tree was inferred by the Neighbor joining method (Saitou and Nei, 1987). Tree topologies were evaluated by bootstrap analysis (Felsenstein, 1985) based on 1000 resembling of the neighbor joining data set.

RESULTS

Antimicrobial activity

In order to assess whether the two *Bacillus* strains isolated from the salt soil originating from south of Tunisia habitats may be a useful source for natural bioactive compounds, the antimicrobial activity of the isolates was determined (Table 1). The two isolates exhibited antibiotic activity towards the Gram-negative bacteria *E. coli* and *P. aeruginosa*, the Gram-positive bacteria *M. luteus*, the yeast *S. cerevisiae* and the filamentous fungi *M. ramannianus* and *B. cinerea*. The activity against the Gram-positive bacteria, *S. aureus* and *M. phlei*, was not recorded.

Choice of media for production of antibiotic

The antifungal and antimicrobial productions in various media are illustrated in Table 2. PDA was found to be the better medium in promoting the production of antifungal and antimicrobial compounds active against micro-organisms tested and was there for retained for later studies.

Kinetics of antimicrobial products of the *Bacillus* sp. strains in potato dextrose medium

Bacillus sp. strains produce metabolites that display antifungal activity. The growth inhibition of the two fungi *B. cinerea* and *M. ramannianus*, the yeast *S. cerevisiae* and the bacteria *M. luteus* by the broth of steady-statecell-free supernatants at different time of growth of the bacteria show the presence of halo of inhibition that began at the stationary phase of the growth of the two bacteria (Figures 1 and 2). The results of antagonism tests have revealed an important antimicrobial activity of the two *Bacillus* strains, especially against filamentous fungi. Indeed, their activity is very strong against the two

Table 1. Antimicrobial activity of B27 and B29 *Bacillus* sp. strains against microorganisms tested.

Microorganisms	Diameter of inhibition (mm)	
	B27	B29
Gram-positive bacteria		
<i>M. luteus</i> CIPA270	24.25 (\pm 0.85)	25 (\pm 0)
<i>S. aureus</i> CIP658	-	-
<i>M. phlei</i> CIP105 389	-	-
Gram-negative bacteria		
<i>E. coli</i> CIP102 400	20.5 (\pm 0.5)	21.75 (\pm 0.75)
<i>P. aeruginosa</i> CIP 100720	20.5 (\pm 0.5)	21.25 (\pm 0.47)
Yeasts		
<i>S. cerevisiae</i> ATCC 9226	22.5 (\pm 0.28)	24 (\pm 0.40)
Filamentous fungi		
<i>M. ramannianus</i> NRLL 1829	17.25 (\pm 0.28)	19.25 (\pm 0.47)
<i>B. cinerea</i>	21.75 (\pm 0.62)	24 (\pm 0.40)

Values given are the mean of four samples (\pm standard error of the arithmetic mean); (-) were not inhibited by the strain.

Table 2. Antimicrobial activity of B27 and B29 *Bacillus* sp. strains in various culture media.

	YMA		NA		PDA		BT	
	B27	B29	B27	B29	B27	B29	B27	B29
<i>M. luteus</i> CIPA270	32 \pm 0.4	35 \pm 0.3	30 \pm 0	35 \pm 0.3	45 \pm 0.22	55 \pm 0.13	35 \pm 0.31	45 \pm 0.4
<i>E. coli</i> CIP102 400	12 \pm 1.1	19 \pm 0.7	12 \pm 0.65	18 \pm 0.8	13 \pm 0.13	20 \pm 0.4	12 \pm 0.51	16 \pm 0.12
<i>P. aeruginosa</i> CIP 100720	15 \pm 0.75	21 \pm 0.1	15 \pm 0.01	24 \pm 0.21	30 \pm 0.11	25 \pm 0	22 \pm 0	23 \pm 0
<i>S. cerevisiae</i> ATCC 9226	12 \pm 0.4	16 \pm 0	14 \pm 0.43	16 \pm 0	17 \pm 0.21	17 \pm 0.22	15 \pm 0	16 \pm 0.11
<i>M. ramannianus</i> NRLL 1829	12 \pm 1.1	17 \pm 0.2	12 \pm 1.0	12 \pm 0.12	17 \pm 0.22	23 \pm 0.11	12 \pm 0.13	22 \pm 0.21

Values given are the mean of four samples \pm standard error of the arithmetic mean.

species *M. ramannianus* and *B. cinerea*. Concerning the antibacterial activity, the two *Bacillus* are strongly active against *M. luteus* and *S. cerevisiae*. The kinetic of production of the antibiotics was realized on PDB medium. Antifungal and antibacterial activities appear after 14 h against *M. ramannianus*, 18 h against *M. luteus*, 20 h against *B. cinerea* and 28 h against *S. cerevisiae* for the B29 strain (Figure 1). While for B27, the antimicrobial activity has been recorded after 18 h of culture against *M. ramannianus*, *M. luteus* and *B. cinerea* and 24 h against *S. cerevisiae*. These activities increase immediately and remain more or less stable thereafter (Figure 2).

Extraction of antimicrobial activities

The results of the antimicrobial assays of the isolate B29

are shown in Figure 3. The bacterium B29 exhibited antibiotic activity towards *M. luteus*, *S. cerevisiae*, *M. ramannianus* and *B. cinerea*. The activity against *P. aeruginosa* was absent. The different fractions obtained from B27 isolate culture did not show any antibiotic activity against all microorganisms tested.

The highest activity was exhibited by the strain B29, with an inhibition zone of 25 mm against *S. cerevisiae*, 20 mm against *M. ramannianus* and 15 mm against *B. cinerea* in the hexane fraction (Figure 3). This bacterium showed important inhibitory activities against bacteria. In fact, the ethyl acetate fraction induced an inhibition zone of 25 mm towards *M. luteus* colony compared to 30 mm noted with the butanol fraction. The hexane has been therefore chosen as extraction solvent for the *Bacillus* strain B29. Therefore, this fraction was retained for later studies. The B27 strain exhibited strong antifungal and antimicrobial activity against the microorganisms tested.




Figure 1. Time course of growth and antibiotics production of *Bacillus* sp. strain B29 on PDB medium at 30°C. The measure of the diameter inhibition against microorganisms was undertaken by the discs method (25 μ L extract by disc). Values represent diameter inhibition in mm, including the discs diameter (6 mm).




Figure 2. Time course of growth and antibiotics production of *Bacillus* sp. strain B27 on PDB medium at 30°C. The measure of the diameter inhibition against microorganisms was undertaken by the discs method (25 μ L extract by disc). Values represent diameter inhibition in mm, including the discs diameter (6 mm).




Figure 3. Choice of extraction solvent of antibiotic activities produced on PDB media by the *Bacillus* sp. strain B29. Values represent diameters of inhibition in mm (including the disc diameter that is 6 mm) against *M. r.* *M. ramannianus*; *B. c.* *B. cinerea*; *S. c.* *S. cerevisiae*; *P. a.* *P. aeruginosa* and *M. l.* *M. luteus*.

This activity was not extracted by n-butanol or by the other organic solvents such as ethyl acetate, dichloromethane and n-hexane.

Partial characterization of antifungal products of *Bacillus* B29 strain

The cell-free supernatant of 1.5 L of liquid culture of B29 strain was extracted by hexane. The organic phase was concentrated to dryness, recuperated in methanol and chromatographed by TLC. On analytic TLC plates developed in BAW solvent system, many active spots were detected by bioautography. Antifungal spots at 0.30 and 0.76 Rf values and antibacterial spots at 0.12, 0.14, 0.19 and 0.30 Rf values. The antibiotic localized at 0.30 Rf value showed strong antibacterial and antifungal activities.

Antimicrobial activity guided fractionation of the broth of B29 culture showed that the hexane extract exhibited inhibitory activity against the filamentous fungi tested, and the butanol extract showed activity against the bacterial strains. Further fractionation of the hexane extract was undertaken by reverse phase HPLC. Figure 4 shows a representative HPLC trace of hexane extract when separated on a C18 semi-preparative HPLC column and detected at 220 nm. Seventeen fractions could be clearly resolved under the conditions employed. Of the 17 fractions that were tested for antimicrobial activity, fractions 15 and 16 were found to be active against *M. ramannianus* and *M. luteus*. The remaining fractions did not exhibit antimicrobial activity. Antibiotic 15

shown maxima, UV-VIS spectrum absorbance in methanol, at 210, 224 and 272 nm. Antibiotic 16 exhibited the most interesting activity. Its UV-VIS spectrum in methanol showed maxima at 203, 224 and 270 nm.

Molecular identification and phylogenetic analysis of bacteria

Data from the 16S rDNA gene provide a limited number of characters that are informative for the phylogeny of the *Bacillus* strain. Aspect of tree presented in Figures 5 and 6 appears to be informative. Based on the sequence homology and phylogenetic analysis, organism was found to be *Bacillus* (1104513.1). It is more closely related to *Bacillus* sp. Z19 (EU236746.1). For the *Bacillus* B29 strain (1104513.2), it was found to be more closely related to *Bacillus amyloliquefaciens* strain EXWB3-03 (EU334107.1). The partial sequence analysis of the 16S rDNA gene and BLAST sequence comparison in the GenBank database showed an alignment of 1414/1415 bases (99% similarity) of the strain B27 (accession number 1104513.1) with the sequence of *B. subtilis* strain PCL 1605 (accession number DQ779882.1), similar results have been also obtained for *B. mojavensis* strain BCRC 17531 (accession number DQ993678.1). For the strain B29 (accession number 1104513.2), the 16S rDNA gene partial sequence aligned with the sequence of *B. amyloliquefaciens* strain TB2 (accession number EU359773.1), though at the lower similarity level of 99% with an alignment of 1414/1415 bases and with the *B. subtilis* strain Pab02 (accession number EU346662.1).




Figure 4. HPLC trace of hexane extract of the culture broth of *Bacillus* sp. strain B29. Seventeen fractions are of relatively high intensity, are easily identified. In addition several peaks of lower intensity are also observed. Majority of the observed antimicrobial activity appears to arise from compounds in peaks 15 and 16.




Figure 5. Phylogenetic tree derived from parsimony analysis of 16S rDNA gene from *Bacillus* sp. strain B27 (accession number 1104513.1). The 1415 bases determined for B27 were aligned to the corresponding regions from bacteria. The number at the nodes indicate the levels of bootstrap support based on neighbor-joining analyses of 1000 resampled data sets; only values over 50% are given. Bar, 0.0002 nt substitution per nt position.




Figure 6. Phylogenetic tree derived from parsimony analysis of 16S rDNA gene from *Bacillus* sp. strain B29 (accession number 1104513.2). The 1415 bases determined for B29 were aligned to the corresponding regions from bacteria. The number at the nodes indicate the levels of bootstrap support based on neighbor-joining analyses of 1000 resampled data sets; only values over 50% are given. Bar, 0.0005 nt substitution per nt position.

DISCUSSION

Microorganisms produce some of the most important compounds with pesticidal activity. They are the source of lifesaving treatments for bacterial and fungal infections. In spite of the tremendous success of the past secondary metabolite research, the number of terrestrial antibiotics seems currently to approach a saturation curve with an apparent limit in the near future. The increasing number of duplications and the urgent demand for few leading structures in pharmacology and agriculture, have enforced the search for metabolites in so far untouched habitats. Our interest focused on microorganisms belonging to the *Bacillus* genus, the members of which have demonstrated interesting antimicrobial activity. In this article, we have described the use of *Bacillus* strains B27 and B29 isolated from salt soil as well as their secreted secondary metabolites to act as antimicrobial product against a large range of microorganisms tested (bacteria, yeasts and filamentous fungi). Cho et al. (2009) have recently described the use of many strains of *Bacillus* sp. as antifungal and antibiotic agents.

The extent of the antibiosis of the two *Bacillus* isolates against the test organisms, evaluated in terms of reduced radial growth, is almost similar. The two isolates exhibited a strong activity (inhibition zone > 20 mm) against Gram-positive and Gram-negative bacteria, yeast and filamentous fungi. The strong activity expressed by a large zone of inhibition on agar plates indicated, as mentioned by Barakate et al. (2002), that those two isolates produce water soluble antimicrobial metabolites which may play an important role in the biocontrol of plant diseases.

Results of antibiotics activity expressed in terms of the diameter of the inhibition zone showed differences in the percentage of antibiosis and specificity of efficacy; this may imply that the two investigated *Bacillus* isolates belong to different species or to the same one but they produced different bioactive compounds exhibiting inhibitory activity against a large number of microorganisms.

These results were confirmed by the time course of growth and antibiotics production of the two strains. In fact, we noted many differences that concerned the specificity of the antibiotics and their time of production.

The chemical nature of the compounds responsible for the observed activity in the organic solvent extracts, respectively for B29 and B27 strains, are being elucidated. The *Bacillus* species are widely recognized as a rich source of antimicrobial agents (Gebhardt et al., 2002). Many antibiotics including cyclic peptides, cyclic lipopeptides and novel thiopeptides have been reported from *Bacillus* sp. (Nagai et al., 2003).

The bioautography of the hexane extract of the B29 strain has revealed several products active against bacteria and fungus suggesting the presence of antibiotics having probably different chemical structures. The B29 strain active fractions 15 and 16 have a time of retention of 47.5 and 48.35 min respectively. The UV-

visible spectra in methanol of these two fractions exhibited maxima at 270 nm suggesting the presence of an aromatic ring compound (Badji et al., 2005; Zitouni et al., 2005). The 16S rDNA gene sequence analysis has been described to be necessary mainly to detect some misidentification of *Bacillus* and related strains (Guinebreiere et al., 2001; Wu et al., 2006). Sequencing the nucleotide positions 70 to 344 of the 50 end region of the 16S rDNA gene may represent a very efficient index for the identification of *Bacillus* and *Paenibacillus* species being this region the most informative one. In fact, about 275 pb of the 50 end region of the 16S rDNA gene were the hypervariant region in the gene and it was highly specific for each *Bacillus* type strain, but also highly conserved within the species (Goto et al., 2000). The 16S rDNA sequence of B27 and B29 strains were compared with those of other *Bacillus* species. The similarity level was about 99% with *B. subtilis* and *B. mojavensis* for the B27 strain and was about 99% with *B. amyloliquefaciens* and *B. subtilis* for the B29 strain. However, it is clear from phylogenetic analysis that these two strains did not cluster with neither *B. subtilis*, nor *B. amyloliquefaciens* nor *B. mojavensis* and represented distinct phyletic lines suggesting new genomic species. All results suggest that both strains may be new species of *Bacillus*. The antifungal activities exhibited especially against *B. cinerea* showed that both strains may play an important role in biological control. Further studies are needed to determine the ability of these strains to control disease caused by *B. cinerea* on *Vitis vinifera*.

ACKNOWLEDGEMENTS

The authors would like to thank Prof. Zouba A., the head of the Centre Régional des Recherches en Agriculture Oasienne Tunisia, for financial contribution.

REFERENCES

- Badji B, Riba A, Mathieu F, Lebrihi A, Sabaou N (2005). Activité antifongique d'une souche d'*Actinomadura* d'origine saharienne sur divers champignons pathogènes et toxigènes. J. de de Mycol. Méd., 15: 211–219.
- Barakate M, Ouhdouch Y, Oufdou KH, Beaulieu C (2002). Characterization of rhizospheric soil *Streptomyces* from Moroccan habits and their antimicrobial activities. World J. Microbiol. Biotechnol., 18: 49–54.
- Cho MK, Math RK, Hong SY, Islam S Md A, Mandanna DK, Cho JJ, Yun MG, Kim JM, Yun HD (2009). Iturin produced by *Bacillus pumilus* HY1 from Korean soybean sauce (Kanjang) inhibits growth of aflatoxin producing fungi. Food Control, 20: 402–406.
- Cho SJ, Hong SY, Kim JY, Park SR, Kim MK, Lim WJ (2003). Endophytic *Bacillus* sp. CY22 from a balloon flower (*Platycodon grandiflorum*) produces surfactin isoforms. J. Microbiol. Biotechnol., 13: 859–865.
- Cragg GM, Newman DJ, Snader KM (1997). Natural products in drug discovery and development. J. Nat. Prod., 60: 52–60.
- Errakhi R, Bouteau F, Lebrihi A, Barakate M (2007). Evidences of biological control capacities of *Streptomyces* spp. against *Sclerotium rolfsii* responsible for damping-off disease in sugar beet (*Beta*

- vulgaris* L.). Wold J. Microbiol. Biotechnol., 23: 1503–1509.
- Felsenstein J (1985). Confidence limits on phylogenies: an approach using the bootstrap. Evolution, 39: 783–791.
- Galvez A, Maqueda M, Cordovilla P, Martinez-Bueno M, Lebbadi M, Valdivia E (1994). Characterization and biological activity against *Naegleria fowleri* of amonebicins produced by *Bacillus licheniformis* D-13. Antimicrob. Agents Chemother., 38: 1314–1319.
- Gebhardt K, Schimana J, Muller J, Fielder HP, Kallenborn HG, holzenkampfer M, Krastel P, Zeeck A, Vater J, HOLTzel A, Schmid DG, Rheinheimer J, Dettner K (2002). Screening for biologically active metabolites with endosymbiotic bacilli isolated from arthropods. FEMS Microbiol. Lett., 217: 199–205.
- Goto K, Omura T, Hara Y, Sadaie Y (2000). Application of the partial 16S rDNA sequence as an index for rapid identification of species in the genus *Bacillus*. J. Gen. Appl. Microbiol., 46: 1–8.
- Guinebretiere MH, Berge O, Normand P, Morris C, Carlin F, Nguyen-The C (2001). Identification of bacteria in pasteurized zucchini purées stored at different temperatures and comparison with those found in other pasteurized vegetable purées. Appl. Environ. Microbiol., 67: 4520–4530.
- Gupte M, Kulkarni P, Ganguli BN (2002). Antifungal antibiotics. Appl. Microbiol. Biotechnol., 58:46–57.
- Jukes TH, Cantor CR (1969). Evolution of protein molecules. In: Munro, H.N. (Ed.), Mammalian Protein Metabolism, vol. 3. Academic Press, New York, pp. 21–132.
- Liu D, Coloe S, Baird R, Pedersen J (2000). Rapid mini-preparation of fungal DNA for PCR. J. Clin. Microbiol., 38: 471.
- Mellouli L, Mehdi RB, Sioud S, Salem M, Bejar S (2003). Isolation, purification and partial characterization of antibacterial activities produced by a newly isolated *Streptomyces* sp. US24 strain. Res. Microbiol., 154: 345–352.
- Milner JL, Raffel SJ, Lethbridge BJ, Handesman J (1995). Culture conditions that influence accumulation of zwittermicin a by *Bacillus cereus* UW85. Appl. Microbiol. Biotechnol., 43: 685–691.
- Nagai K, Kamigiri K, Arao N, Suzumura K, Kawano y, Yamaoka M, Zhang H, Watanabe M, Suzuki K (2003). YM-266183 and YM-266184, novel thiopeptide antibiotics produced by *Bacillus cereus* isolated from marine sponge. J. Antibiot., 56: 123–128.
- Newman DJ, Gragg GM, Snader KM (2003). Natural products as source of new drugs over the period 1981-2002. J. Nat. Prod., 66: 1022–1037.
- Peypoux F, Michel G, Delcambe L (1976). Structure de la mycosubtine, antibiotique isolé de *Bacillus subtilis*. Eur. J. Biochem., 63: 391–398.
- Peypoux F, Pommier MT, Das BC, Besson F, Delcambe L, Michel G (1984). Structures of bacillomycin D and bacillomycin L peptidolipid antibiotics from *Bacillus subtilis*. J. Antibiot., 77: 1600–1604.
- Peypoux F, Bonmatin JM, Wallach J (1999). Recent trends in the biochemistry of surfactin. Appl. Microbiol. Biotechnol., 51: 553–563.
- Peypoux F, Guinand M, Michel G, Delcambe L, Das BC, Lederer E (1978). Structure of iturin A, a peptidolipid antibiotic from *Bacillus subtilis*. Biochemistry, 17: 3992–3996.
- Saitou N, Nei M (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol. Biol. Evol., 4: 406–425.
- Salas JA, Mendez C (2007). Engineering the glycosylation of natural products in actinomycetes. Trends Microbiol., 15: 219–232.
- Steller S, Vollenbroich D, Leenders F, Stein T, Conrad B, Hofemeisterr J, Jaques P, Thonart P, Vater J (1999). Structural and functional organization of the fengycin synthase multienzyme system from *Bacillus subtilis* b213 and A1/ 3. Chem. Biol., 6: 31–41.
- Thompson JD, Higgins DG, Gibson TJ (1994). CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighing, position specific gap penalties and weight matrix choice. Nucl. Acids Res., 22: 4673–4680.
- Tsuge K, Ano T, Shoda M (1996). Isolation of a gene essential for biosynthesis of the lipopeptide antibiotics plipastatin B1 and surfactin in *Bacillus subtilis* YB8. Arch. Microbiol., 165: 243–251.
- Vanittanakom N, Loeffle W, Koch U, Jung G (1986). Fengycine-a novel antifungal lipopeptide antibiotic produced by *Bacillus subtilis*: F-29-3. J. Antibiot., (Tokyo) 39: 888–901.
- Walker JE, Abraham EP (1970). The structure of bacilysin and other products of *Bacillus subtilis*. Biochem. J., 118: 563–570.
- Wu XY, Walker MJ, Hornitzky M, Chin J (2006). Development of a group-specific PCR combined with ARDRA for the identification of *Bacillus* species of environmental significance. J. Microbiol. Meth., 64:107–119.
- Yılma M, Soran H, Beytli Y (2006). Antimicrobial activities of some *Bacillus* spp. strains isolated from the soil. Microbiol. Res., 161: 127–131.
- Zitouni A, Boudjella H, Lamari L, Badji B, Mathieu F, Lebrihi A, Sabaou N (2005). *Nocardioopsis* and *Saccharothrix* genera in Saharan soils in Algeria: Isolation, biological activities and partial characterization of antibiotics. Res. Microbiol., 156: 984–993.