



**HAL**  
open science

## How Music Can Literally Heal the Heart

Elaine Chew, Psyche Loui, Grace Leslie, Caroline Palmer, Jonathan Berger,  
Edward Large, Nicolò F. Bernardi, Suzanne B. Hanser, Julian F. Thayer,  
Michael A. Casey, et al.

► **To cite this version:**

Elaine Chew, Psyche Loui, Grace Leslie, Caroline Palmer, Jonathan Berger, et al.. How Music Can Literally Heal the Heart. 2021. hal-03541629

**HAL Id: hal-03541629**

**<https://hal.science/hal-03541629v1>**

Submitted on 24 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## M E D I C I N E

# How Music Can Literally Heal the Heart

Its structural attributes and physiological effects make it an ideal tool for learning cardiology, studying heart-brain interactions and dispensing neurocardiac therapy

By

Elaine Chew, Psyche Loui, Grace Leslie, Caroline Palmer, Jonathan Berger, Edward W. Large, Nicolò F. Bernardi, Suzanne Hanser, Julian F. Thayer, Michael A. Casey, Pier D. Lambiase  
on September 18, 2021



Credit: Dragan Todorovic *Getty Images*

In a maverick method, nephrologist Michael Field taught medical students to decipher different heart murmurs through their stethoscopes, trills, grace notes, and decrescendos to describe the distinctive sounds of heart valves snapping closed, and blood ebbing through leaky valves in plumbing disorders of the heart.

Separately, in music based on electrocardiographic (ECG) traces of heart rhythm disorders, one of us—musician-mathematician Elaine Chew—used music notation to

capture the signature rhythms of electrical anomalies of the heart. Collaged from extant music fragments matching the heartbeats, Brubeck's *Blue Rondo à la Turk* provided the 2:4:3 rhythmic tattoo of ventricular early beats, Piazzolla's *Le Grand Tango* remixed produced the irregular rhythms of atrial fibrillation. *Little Etudes for piano*, with pedagogical descriptions by cardiologist Pier Lambiase, provided a layperson's introduction to electrical aberrations of the heart.

The reason these heart-music mappings work is because abnormal heart rhythms tend to form simple inter-beat-interval ratios. In fact, the distinctive rhythms in Beethoven's music so closely resemble those of heart rhythm disorders that cardiologists have speculated that they may be transcriptions of Beethoven's possible arrhythmia, his interoceptive awareness of his own heartbeat enhanced by his deafness.

**Holter Highlights - Excerpts from a patient with AF and other rhyth...**



This is but one of multiple reasons music should be part of every heart physician's toolkit. Music and the heart have been romantically linked in popular consciousness due to their shared connections to human emotions and the brain. History is replete with examples of emotionally charged events followed almost immediately by the death of the person. The surgeon John Hunter famously pronounced, "My life is at the mercy of any scoundrel who should put me in a passion," before collapsing and dying after a heated board room meeting.

Cardiologists Peter Taggart and Pier Lambiase have been studying how emotions alter the conductive properties of individual heart cells. Mental stress changes the recovery period of heart cells after each heartbeat, called the action potential duration. Taggart co-authored a study in which patients whose hearts were paced at a steady rate watch the harrowing “cut the rope” scene from *Vertical Limit* (2000). The patients’ action potential duration shortened under the stress. This may explain how more extreme stress coupled with underlying cardiac disease could precipitate life-threatening arrhythmias.

Acute stress produces dramatic effects in the heart, but slow-burning chronic stress due to protracted insecurity also predispose sufferers to disease and mortality. The sympathetic nervous system’s default state of high alertness is suppressed when safety is perceived; these safety brakes are lifted under duress. The Generalized Unsafty Theory of Stress co-written by psychophysiological Julian Thayer links the unconsciously perceived unsafety of prolonged stressors like low social status, early life adversity or loneliness to hypervigilance that increases the odds of developing heart disease.

### The Making of Arrhythmia Suite, III. Ventricular Ectopics with Short ...



Music moves us in part because it draws on our primal intuitions about the heartbeat. Until the mid-19th century when it was replaced by the mechanical metronome, the human heartbeat provided the standard unit of measure for musical time. In his 1496 treatise, the *Practica Musicae*, the composer-theorist Franchinus Gaffurius wrote that

the proper measure of the musical beat should be the pulse of a healthy human, noting that the pulses of “fevered persons” undergo an increase or become unequal in ways that worry physicians.

When we connect to the pulse of the music, we sense another’s physiological states. The steady pulse at the beginning of Schubert’s *Trio, Op. 100*, sets a strong but serene pace for its haunting melody. The breathless octaves in the opening of *Der Erlkönig* evokes the rapid heart palpitations of the fevered boy in his father’s arms, galloping through the stormy, windswept night. Hearing just heartbeats, pulse-only music, has been found to increase listeners’ ability to sense what others are feeling in a study co-authored by musician-scientist Grace Leslie.

Music changes our heartrates, breathing, and blood pressure, and alters our heart rate variability, indicators of cardiac and mental health. Neuroscientist Psyche Loui and colleagues have traced music-induced physiological changes to a central node in the brain’s networks, called the anterior insular, with dense connections to the vagus nerve, responsible for unconscious regulation of body functions.



Sign up for *Scientific American's* free newsletters.

[Sign Up](#)

---

The anterior insula is associated with empathetic mirroring of external and internal experiences. It is also connected to parts of the brain responsible for hearing (the auditory cortices) and for pleasure (the dopaminergic reward system). These auditory and reward network pathways likely subserve the mind’s ability to form predictions and expectations during music listening. The systematic fulfilment and violation of expectations are thought to underlie emotion and meaning in music.

Music is an ideal catalyst for inducing physiological changes in heart-brain studies because it can be dissected systematically into features based on note content and the way this content is communicated in performance. Evidence suggests that these musical attributes trigger brain responses at a basic level. Analyzing listeners’ brain imaging data in the OpenfMRI Study Forrest dataset, composer-neuroscientist Michael Casey found

that specific music features induced predictable activation patterns in regions of listeners' brains. The activation patterns were consistent enough for machines to infer the music the listener heard or its genre simply from their fMRI scans.

Music features have also been linked to physiological responses. In a study co-authored by physicians Luciano Bernardi and Peter Sleight, loudness increases in vocal and orchestral music produced vascular constriction and blood pressure increases proportionate to these crescendos. Verdi arias with ten-second-long phrases—the period of Mayer waves, the body's natural blood pressure oscillation—caused listeners' heart and respiratory signals to sync with the music envelop. Such unconscious physiological responses are thought to be the progenitors of music-induced emotions.

### Arrhythmia Suite, III. Ventricular Ectopics with Short Ventricular Tac...



Music also has a communal impact on human physiology. People listening to the same music tend to synchronize not only their movements, but also their breathing and heart rhythms. Some of this heartbeat coherence is due to breathing together, but partial coherence (linear relationships) remained higher between the heartbeats of people vocalizing long notes together, over the baseline or breathing together, even after removing the effect of respiration.

The cognitive and physical demands of playing music also have measurable effects on musicians' heart rhythms and breathing patterns. Psychologists Caroline Palmer and

Shannon Wright showed that repetitiveness of musicians' heart rhythms show greater rigidity (predictability) when playing unfamiliar musical melodies, and also when playing first thing after waking in the morning rather than in the evening.

For cardiac patients, music-based interventions can also modulate cerebral blood flow, reduce pre-operative anxiety and post-operative stress, improve surgery outcomes, and lower cortisol levels. Music interventions are found to significantly affect heartrate and blood pressure in coronary heart disease patients. Listening to relaxing music not only reduced heart and respiration rates but also oxygen demand of the heart in patients who have had a heart attack.

### Music: An Underutilised Tool in Neurocardiology?



Technological advances in biofeedback sensors means that physiological parameters like heartbeats and heart rate variability can be harnessed to guide music interventions in cardiac therapy. Physiological feedback can be used to select or shape music to influence listeners' heart rates and breathing, for example, to increase heart rate variability. With widespread adoption of biofeedback devices, the tailoring of music interventions to individual cognitive or neural-cardiac states is now well within reach enabling a “musical prescription” for improved mental and physical wellbeing.

*This is an opinion and analysis article; the views expressed by the author or authors are not necessarily those of Scientific American.*

*This article has been developed from an exploratory multidisciplinary seminar on Music and the Heart at the Radcliffe Institute for Advanced Study at Harvard University involving physicians, neuroscientists and musicians with a specific interest in music and its effects on human physiology.*

*EC's and PDL's work is supported in part by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement Nos. 788960 and 957532).*

*EL's work is supported in part by the National Science Foundation (NSF) under an NSF STTR Phase I Grant No. 2014870. Any opinions, findings and conclusions, or recommendations expressed in this material are those of the author(s), and do not necessarily reflect those of the funding agencies.*

ADVERTISEMENT

**ABOUT THE AUTHOR(S)**

**Elaine Chew**, Ph.D., F.T.C.L., is a pianist-mathematician and principal investigator of the European Research Council-funded project COSMOS at the French National Center for Scientific Research (CNRS), where she is based at the Sciences and Technologies of Music and Sound Laboratory located at the Institute for Research and Coordination of Acoustics/Music (IRCAM).

**Psyche Loui**, Ph.D., is a neuroscientist-violinist and an associate professor and director of the Music, Imaging, and Neural Dynamics (MIND) Laboratory at Northeastern University.

**Grace Leslie**, Ph.D., a flutist, electronic musician and scientist, is an assistant professor of music technology at the Georgia Institute of Technology, where she directs the Brain Music Laboratory.

**Caroline Palmer**, F.R.S.C., is the Canada Research Chair in the Cognitive Neuroscience of Performance and a professor of psychology at McGill University, where she directs the Sequence Production Laboratory.

**Jonathan Berger** is the Denning Family Provostial Professor of Music and a Bass University Fellow in Undergraduate Education at Stanford University, and a composer-researcher at the Center for Computer Research in Music and Acoustics (CCRMA).

**Edward W. Large**, Ph.D., is director of the Music Dynamics Laboratory and the Theoretical Neuroscience Laboratory at the University of Connecticut, where he is a professor in the Department of Psychological Sciences and the Department of Physics.

**Nicolò F. Bernardi**, Ph.D., A.C.C., is a personal and professional coach and consultant who is trained as a cognitive neuroscientist, music therapist and yoga teacher.



**Suzanne Hanser**, Ed.D., is chair emerita and professor of music therapy at Berklee College of Music and president of the International Association for Music & Medicine.

**Julian F. Thayer**, Ph.D., a bassist and composer is the Distinguished University Professor in the Department of Psychological Science at the University of California, Irvine and Ohio Eminent Scholar Professor in Health Psychology Emeritus at the Ohio State University.

**Michael A. Casey** is a professor in the Department of Computer Science and the Department of Music at Dartmouth College.

**Pier D. Lambiase** is professor of cardiology at University College London Institute of Cardiovascular Science and Barts Heart Center and co-director of cardiovascular research at Barts Health NHS Trust.

---

Scientific American is part of Springer Nature, which owns or has commercial relations with thousands of scientific publications (many of them can be found at [www.springernature.com/us](http://www.springernature.com/us)). Scientific American maintains a strict policy of editorial independence in reporting developments in science to our readers.

© 2022 SCIENTIFIC AMERICAN, A DIVISION OF SPRINGER NATURE AMERICA, INC.

ALL RIGHTS RESERVED.