



HAL
open science

Neurophysiology of gaze orientation: Core neuronal networks

Laurent Goffart, Julie Quinet, Clara Bourrelly

► **To cite this version:**

Laurent Goffart, Julie Quinet, Clara Bourrelly. Neurophysiology of gaze orientation: Core neuronal networks. 2022. hal-03541980v2

HAL Id: hal-03541980

<https://hal.science/hal-03541980v2>

Preprint submitted on 6 Mar 2023 (v2), last revised 18 Dec 2023 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 For a publication in Encyclopedia of the Human Brain, edited by Jordan Grafman,
2 Angela Sirigu & Jean-René Duhamel (section editors)

3
4

5 Neurophysiology of gaze orientation: 6 Core neuronal networks

7 Laurent Goffart^{1,2}, Julie Quinet³ and Clara Bourrelly⁴

8 1 : Aix Marseille Université, CNRS, Institut de Neurosciences de la Timone, Marseille, France

9 2 : Aix Marseille Université, CNRS, Centre Gilles Gaston Granger, Aix-en-Provence, France

10 3 : Department of Optometry and Vision Science, The University of Alabama at Birmingham, Birmingham, USA

11 4 : University of Pittsburgh, Center for the Neural Basis of Cognition, Pittsburgh, USA

12
13

14 Abstract

15 The appearance of an object triggers a shift of gaze toward its location. This orienting response
16 consists of a rapid rotation of the eyes, the saccade, sometimes accompanied by a head
17 rotation. In this chapter, instead of describing the path leading from the target-evoked retinal
18 activity to the changes in muscle tension, we shall take the reverse path. Starting from the
19 muscle contractions, we shall proceed upstream and describe the core neuronal networks
20 that, in the brainstem and cerebellum, enables us to rapidly and accurately orient the foveae
21 towards visual targets located at different eccentricities and depths.

22
23



24

25 **Keywords**

26 Brain stem

27 Cerebellum

28 Eye movement

29 Fixation

30 Gaze

31 Head movement

32 Monkey

33 Neurophysiology

34 Non-human primate

35 Orienting

36 Pursuit

37 Reticular formation

38 Saccade

39 Tracking

40 Vergence

41

42

43

44

45

46

47 Key points/objectives

48 Complementing several past reviews, this chapter brings a synthesis of the considerable
49 knowledge that neurophysiologists and neuroanatomists gathered in the monkey during the
50 last six decades, with the aim to explain the neuronal underpinnings of one of the most basic
51 spatial abilities, i.e., directing gaze toward the location where something just happened.

52

53

84 compartmentalization of extraocular muscles may lead to revise this view (Demer, 2015).
85 Moreover, precise measurements made in human subjects revealed a transient torsion during
86 horizontal saccades (Straumann et al. 1995), suggesting brief changes in the activation of the
87 other extraocular muscles.

88 The neural control is more complicated when gaze moves vertically. Vertical
89 movements involve not only steady tension of LR and MR muscles, but also a complex
90 cooperation between the four other muscles. The contraction of superior and inferior rectus
91 muscles supraduct (or elevate) and depress (or infraduct) each eye, respectively. These actions
92 are stronger when the eye is in abduction. The contraction of oblique muscles causes torsional
93 movements but also supraduct or infraduct the eye, more strongly when the eye is adducted.
94 If we consider the movements made when the eyes are centered in the orbit, in addition to
95 elevating the optic axis (turquoise arrows in Fig. 1B), the contraction of the superior rectus
96 (SR) muscle fibers causes adduction (black arrows) and incyclotorsion (or incycloduction; blue
97 arrows) of the eyeball. Likewise, in addition to lowering the optic axis (yellow arrows in Fig.
98 1B), the contraction of the inferior rectus (IR) muscle fibers causes adduction and
99 excyclotorsion (or excycloduction; green arrows). These secondary actions can be
100 counteracted by the contraction of oblique muscles, for the contraction of inferior oblique
101 (IO) muscle fibers causes elevation, abduction and excycloduction, whereas the contraction of
102 superior oblique (SO) muscle fibers causes depression, abduction and incycloduction (Demer
103 2019).

104 Binocular conjugate movements are made when the head rotates while a visual target
105 is being fixated (vestibulo-ocular reflex), when the whole visual scene is moving (optokinetic
106 reflex) and whenever a target appears or is imagined in the peripheral visual field (saccades).
107 We shall first examine the neurophysiology of saccade generation when the head does not
108 move. Then, we shall study the case when a rotation of the head accompanies the saccade.
109 During these combined eye-head movements, two couplings can happen: the eyes and the
110 head move either in the same direction or in opposite directions. Next, we shall see the
111 neurophysiology of disconjugate movements of the eyes, which occur when gaze is shifted
112 between targets located at different depths (vergence eye movements). Finally, we shall have
113 an overview of the complex network that enables one to fixate and track a continuously
114 moving target (smooth pursuit eye movements).

115 Binocular conjugate movements

116 Horizontal saccades

117 Figure 2 schematizes the current view of the premotor neurophysiology of horizontal
118 saccades in the pontine and mesencephalic reticular formation. During leftward saccades (Fig.
119 2A), the fast-twitch muscle fibers of the right eye's MR and of the left eye's LR contract while
120 those of antagonist muscles (right eye's LR and left eye's MR) relax. Motor neurons in the
121 abducens nucleus (ABD, located in the pontine reticular formation) and in the oculomotor
122 nucleus (OMN in the midbrain tegmentum) emit bursts of action potentials that cause the
123 phasic contraction of muscle fibers (Fuchs and Luschei 1970; Fuchs et al. 1988). The burst
124 discharge of cells in the OMN is triggered and supported by spikes emitted by internuclear
125 neurons (AIN) located in the contralateral abducens nucleus. The bursts of abducens motor
126 (MN) and internuclear (AIN) neurons are driven by afferent input from excitatory burst
127 neurons (EBN) located in the ipsilateral paramedian pontine reticular formation (PPRF,
128 synapses a) (Keller, 1974; Luschei and Fuchs, 1972; Strassman et al., 1986a). The abducens
129 neurons also receive excitatory input from internuclear burst-tonic neurons in the
130 contralateral oculomotor nucleus (not shown). Discharging during saccades and increasing
131 their firing rate during convergence, these cells are possibly involved in the motor drive for
132 conjugate saccades because in addition to cause exophoria, lidocaine injection in the medial
133 rectus subdivision of the oculomotor nucleus causes hypometria and slowing of abducting
134 saccades of the contralateral eye (Clendaniel and Mays, 1994).

135 Otherwise silent, the EBN fire trains of action potentials whose interspike interval is
136 smaller for larger saccades. Contrary to statements made in numerous articles, the
137 instantaneous firing rate of EBN is not a replica of instantaneous eye velocity. For each
138 saccade, the interspike interval is much more constant than expected if this claim were true
139 (i.e, if their discharge determined instantaneous eye velocity) (Hu et al., 2007). The suggestion
140 that the firing rate of motoneurons would "encode" the instantaneous eye velocity is also
141 brought into question by a study in the cat (Davis-López de Carrizosa et al., 2011), which
142 showed that a model factoring in muscle tension and its first derivative accounted for the
143 firing rate of motor neurons better than a model factoring in eye position and its first and
144 second derivatives (velocity and acceleration).

174 Vertical cardinal saccades

175 The neuronal network underlying the generation of vertical saccades is located in the
176 midbrain tegmentum (Büttner and Büttner-Ennever, 2006). For upward saccades (Fig. 3A), the
177 agonist muscles are the IO and the SR and the antagonist muscles the IR and the SO. As
178 reported above, in addition to elevating the eye, the contraction of SR muscle fibers causes
179 incycloduction, whereas contraction of IO muscle fibers causes excycloduction. Thus, the
180 incycloduction caused by the motor commands to the SR combines with the excycloduction
181 caused by the commands to the IO. The observation of negligible torsional component during
182 upward saccades suggests that the two torsions cancel each other out. Such a balance of force
183 likely involves bilateral activity in the midbrain because for each eye, the motor neurons
184 innervating the SR muscle are located in the contralateral OMN whereas the motor neurons
185 innervating the IO muscle are located in the ipsilateral OMN (Porter et al., 1983). The two sets
186 of commands during upward saccades must be adjusted in a timely manner, and any
187 asymmetry in the bilateral motor drive would engender a torsional component.

188 **< Figure 3 near here >**

189 The excitatory commands responsible for generating upward saccades are issued by
190 burst neurons located in the left and right rostral interstitial nucleus of the medial longitudinal
191 fasciculus (RIMLF) (Büttner et al., 1977). These commands are doubly bilateral. Indeed, on the
192 one hand, the burst emitted by premotor neurons (uEBN) in the left RIMLF excite the
193 motoneurons innervating the SR muscles of *both* eyes (black synapses a) and motor neurons
194 innervating the IO muscles of *both* eyes as well (black synapses b in Fig. 3) (Moschovakis et al.,
195 1991a). On the other hand, the burst emitted by uEBNs in the opposite RIMLF excite the same
196 motoneuronal group (blue synapses a and b). Moreover, in addition to recruiting the motor
197 neurons, the uEBN also excite neurons that inhibit the motor neurons that innervate the
198 antagonist muscles (IR and SO, synapses c). Located in the interstitial nucleus of Cajal, these
199 cells (uIBN) inhibit the motor neurons in the contralateral oculomotor nucleus, i.e., cells
200 responsible for the contraction of the IR of the contralateral eye (synapses d) and motor
201 neurons responsible for the contraction of the SO of the ipsilateral eye (synapses e). However,
202 contrary to uEBN, uIBN do not send bilateral projections to the motor neurons innervating the

203 SO muscle fibers. Their projections make synaptic contacts only with motor neurons on the
204 contralateral side (Horn et al., 2003).

205 As for the upward saccades, the discharge of neurons located in each side of the brain
206 is capable of causing a downward movement of *both* eyes (Fig. 3B). The premotor excitatory
207 burst neurons leading to downward saccades (dEBN) are also located in the RIMLF
208 (Moschovakis et al., 1991b). However, unlike the uEBN, the dEBN do not send bilateral
209 projections to the motoneurons; their projection is ipsilateral only. During downward
210 saccades, the SO and IR muscle fibers of *both* eyes contract because the dEBN excite 1) the
211 motor neurons in the ipsilateral OMN, which cause the contraction of IR muscle fibers of the
212 *ipsilateral* eye (synapses a), and 2) the motor neurons which, in the ipsilateral trochlear
213 nucleus, cause the contraction of SO muscle fibers of the *contralateral* eye (synapses b).
214 Moreover, in addition to lowering the eyes, the contraction of SO muscle fibers causes an
215 incyclotorsion of the eye while the contraction of IR muscle fibers causes their excyclotorsion.
216 The commands from the ipsilateral OMN for excyclotorsion combine with commands from the
217 contralateral trochlear nucleus for incyclotorsion. Precisely how these opposite torsional
218 components cancel each other out during downward saccades remains unexplained. A
219 bilateral adjustment at the level of premotor burst neurons seems inevitable. Moreover, the
220 neural processes in charge of the relaxation of antagonist muscles during downward saccades
221 remain to be characterized.

222 Even though the number of cells and the number of synaptic boutons likely differ
223 between the left and right sides of the reticular formation, the connectivity patterns involved
224 in generating leftward and rightward eye movements are symmetrical with respect to the
225 brainstem midsagittal plane (see Fig. 2 and 5). By contrast, the networks involved in the
226 generation of downward and upward eye movements are not symmetrical with respect to the
227 midsagittal plane (Fig 3).

228 During vertical saccades, the EBN involved in the generation of horizontal saccades do
229 not fire. Some IBN in the dPGRF emit action potentials whose number increases with the
230 saccade size (Scudder et al., 1988). For vertical saccades to remain straight while their size
231 increases, one possibility is that the motor and internuclear cells in the abducens nucleus emit
232 additional spikes, thus reducing the contrapulsive effect of IBN spikes. This does not seem to

233 be the case because MN and AIN in the abducens nucleus do not emit action potentials during
234 vertical saccades. A more likely possibility is that the contrapulsion exerted by IBN in the left
235 and right dPGRF cancel each other out. If no perfect symmetry exists between the set of active
236 neural elements (number of neurons and of synaptic boutons) located on each part of the
237 reticular formation, then the observation of strictly vertical saccades leads us to infer that a
238 process adjusts the bilateral activity of IBN during vertical saccades.

239 The bilateral control is not restricted to the group of IBNs, which are primarily involved
240 in the generation of horizontal saccades. In addition to elevating both eyes, the discharge of
241 motor neurons in the right OMN causes, as a secondary action and via the contraction of the
242 right eye's IO and the left eye's SR, a rightward deflection of both eyes during vertical saccades
243 while the discharge of motor neurons in the opposite side causes a leftward deflection (Fig.
244 3A, top row). Likewise, in addition to lowering both eyes, the dEBN in the right RIMLF lead to
245 the contraction of the left eye's SO and the right eye's IR, causing a leftward deflection of their
246 trajectory while dEBN in the left RIMLF cause a rightward deflection during vertical saccades
247 (Fig. 3B, top row). The observation of straight vertical saccades (with no horizontal deviation)
248 implies then that the horizontal deflections caused by the secondary action of the muscles
249 elevating or depressing the eyes are also canceled in the motor drives.

250 Therefore, the generation of straight vertical saccades rests upon a balance of bilateral
251 motor activation, which itself depends upon an adjustment of premotor commands that takes
252 into account the structural (neurons and their connectivity) and functional (firing properties,
253 secondary and tertiary consequence of muscle contractions) asymmetries between the
254 oculomotor territories that are distributed on either side of the brain stem. The saccade-
255 related neurons in the left and right fastigial nuclei and in the oculomotor vermis regulate this
256 bilateral balance. Indeed, most of them burst during vertical saccades and their asymmetrical
257 perturbation horizontally deflects the trajectory of vertical saccades (Goffart et al., 2003;
258 Goffart et al., 2004; Bourrelly et al. 2021). Bilateral activity in the two deep superior colliculi is
259 also involved with their medial halves for driving upward saccades and their lateral halves
260 driving downward saccades. The contribution of the deep superior colliculi and of the caudal
261 fastigial nuclei differ insofar as horizontal saccades involve bursting activity of cells in only one
262 superior colliculus, on the contralateral side (Sparks et al., 1976), whereas saccade-related
263 neurons burst in the caudal fastigial nuclei on both sides (Fuchs et al., 1993).

264 Combined horizontal and vertical saccades

265 When a target appears off the cardinal visual axes, the orienting response does not
266 consist of horizontal and vertical saccades made in sequential order. Both horizontal and
267 vertical components are generated in such a manner that their onset and offset are relatively
268 simultaneous. These combined horizontal-vertical saccades are conventionally called
269 “oblique” saccades. Considering the muscle diversity (lengths, contractile properties and
270 orbital attachments) involved in the horizontal and vertical components and the fact that their
271 motor innervations are not located in nearby regions of the brainstem, their simultaneous
272 onsets and the limited curvature of their spatial trajectory are empirical observations
273 suggesting coupling mechanisms.

274 In the medial pontine reticular formation, the nucleus raphe interpositus (RIP) hosts a
275 population of cells that exhibit a sustained firing rate during intersaccadic intervals but pauses
276 during all saccades, regardless of their direction and amplitude (Keller 1974; Phillips et al.,
277 1999; Strassman et al., 1987). These cells are called omnidirectional pause (or omnipause)
278 neurons (OPN) and are thought to prevent the eyes from moving during visual fixation through
279 their inhibitory post-synaptic influence upon the premotor burst neurons (Fig. 2). Moreover,
280 with their pause, OPN could also be responsible for the synchronous onsets of horizontal and
281 vertical saccades, as they project to burst neurons in both the horizontal and vertical gaze
282 centers (Ohgaki et al., 1989). So far, however, experimental studies did not document onset
283 asynchronies during the inactivation or the lesion of RIP. No fixation instability was reported
284 either (Kaneko, 1996; Soetedjo et al., 2000). This absence of deficit in visual fixation may have
285 resulted from the particular testing conditions because the consequence during delayed or
286 memory-guided saccade tasks was not tested. The observation of irrepressible saccades to
287 the peripheral target during the delay interval would have unambiguously confirmed the role
288 of OPN in maintaining steady the gaze direction during visual fixation, as electrical
289 microstimulation studies suggested (Gandhi and Sparks 2007; Keller 1974). However, these
290 studies are not sufficiently conclusive because the interruption and delaying of saccades
291 caused by electrical stimulation might result from the (undesired) retrograde recruitment of
292 an inhibitory input to pause neurons. A projection from inhibitory burst neurons to the pause
293 neurons has been reported in the cat (Takahashi et al. 2022). However, it remains to be

294 confirmed in the monkey because so far, the available data do not support it (Strassman et al.
295 1986b).

296 Turning back to the generation of combined horizontal and vertical saccades, one
297 study reported crucial observations signaling a process that couples their time course during
298 oblique saccades. When the horizontal component is experimentally slowed by the local
299 injection of a pharmacological agent in the pontine reticular formation, the vertical
300 component is also slowed (Barton et al., 2003; Sparks et al., 2002). In some of these
301 experiments, the duration of both components was prolonged and the accuracy of saccades
302 was preserved, supporting further the persistence and preservation of the target-related
303 commands. The neural connectivity underlying this “component stretching mechanism” in
304 primates have not yet been elucidated.

305 Combined eye and head movements

306 Saccades are frequently associated with a head movement when the target
307 eccentricity approaches and exceeds the limits of the oculomotor range. Moving the head
308 enables to look at targets located more eccentrically in the visual field, and to re-center the
309 eyes in their orbit. The deviation of the eyes in the orbit and of the head relative to the trunk
310 add together to determine the direction of gaze.

311 Two functional modes characterize orienting gaze movements toward a static target:
312 a mode during which both eyes and the head rotate in the same direction and a mode during
313 which they move in opposite directions. The second mode happens sometimes when the head
314 movement precedes the eye saccade, mostly when the target location is known or foreseeable
315 (Bizzi et al. 1972). It happens more frequently later because the eyes move much faster than
316 the head and because the restoration of the equilibrium of bilateral activity that determines
317 gaze direction has been completed before the equilibrium that determines head direction.
318 Before describing how gaze is being stabilized and ends its orienting movement, we shall
319 examine the first mode, i.e., how despite the vestibulo-ocular reflex, the eyes and the head
320 can move in the same direction as during horizontal gaze shifts (Figure 4).

321 Combining a head movement with a saccade

322 In this section, we limit our description to the horizontal eye movements that are made
323 while the head also moves horizontally. Whether the head is moving or not, behavioral studies
324 show that the gaze movements are equally accurate. In the monkey, around twenty muscles
325 are involved to change the orientation of the head (Richmond et al. 2001). Among them are
326 the muscles whose contraction changes the orientation of the head relative to the trunk, the
327 muscles that stabilize the head during its rotation and the muscles that do not exhibit any
328 change (Lestienne et al. 1995). The co-contraction of the left and right *rectus capitis posterior*
329 *minor* (RCPm) stabilize the atlas while the contraction of the *rectus capitis posterior major*
330 (RCPM) and *obliquus capitis inferior* (OCI) produce an ipsilateral horizontal rotation of the
331 head. The observation that the contralateral muscles exhibit no electromyographic change
332 indicates that stopping the head rotation is not caused by muscle co-contraction (Lestienne et
333 al. 1995). For movement amplitudes greater than 20 degrees, the contraction of the *splenius*
334 *capitis* muscle adds to that of the RCPM and OCI muscles. At the end of the orienting
335 movement, the co-contraction of the RCPm, RCPM, OCI and *obliquus capitis superior* (OCS)
336 muscles altogether stabilizes the head (Corneil et al., 2001; Lestienne et al. 1995).

337 Thus, orienting gaze shifts mobilize several muscles in parallel when a movement of
338 the head accompanies the saccade. The synergy results from activities distributed within a
339 complex set of motor and premotor neurons located in the reticular and vestibular nuclei
340 (Cullen 2009; Sparks et al., 2001). Contrary those involved in the cat (Takahashi and Shinoda
341 2018), the premotor processes remain incompletely characterized in the monkey.
342 Contemporary knowledge is not sufficient to reach a synthetic understanding that is
343 comparable to the neurophysiology of saccades.

344 The recruitment of EBN and reticulo-spinal neurons (RSN) by descending commands
345 from bursting cells in the deep superior colliculus (grey-colored synapses a in Fig. 4) leads to
346 the excitation of the pools of extraocular motoneurons (ABD) in the abducens nucleus and
347 neck motoneurons (NMN) in the cervical spinal cord (green and blue synapses b), and thus, to
348 the contraction of the agonist extraocular and neck muscle fibers, respectively. Contrary to
349 the cat, collicular connections with EBN do not seem to be direct in the monkey (Keller et al.,
350 2000). The EBN and RSN also excite the IBN (synapse a'), thereby suppressing the activity of

351 cells that otherwise act to facilitate the contraction of antagonist extraocular (synapses c) and
352 neck muscles (synapse g), respectively. Finally, they also excite secondary vestibular neurons
353 which are inhibitory (IVN-2; blue and green colored synapses d) and located in the ipsilateral
354 medial vestibular nucleus. Within the vestibular nuclei, the discharge of IVN-2 inhibits the
355 secondary vestibular neurons, IVN-1 and EVN-1 (purple-colored synapse e). By inhibiting IVN-
356 1, the motor neurons innervating the agonist extraocular and neck muscles fibers (red
357 synapses f and f') are disinhibited, and the contraction of these muscles is facilitated. By
358 inhibiting the EVN-1, the IVN-2 prevent the primary vestibular neurons (PVN), whose firing is
359 enhanced by the head movement, from causing a counter-rotation of the eyes in the orbits
360 (synapse h). The contraction of antagonist neck muscles through vestibulospinal pathways is
361 also prevented (synapse h').

362 **< Figure 4 near here >**

363 **Stopping gaze while the head terminates its motion**

364 The second mode of combined eye head movements is similar to the vestibulo-ocular
365 reflex recorded when the head is passively rotated while subjects maintain their gaze directed
366 toward a visual target. During active eye and head movements, the premotor input to motor
367 neurons makes a transition from commands fed by target-related (visual or auditory) signals
368 to commands fed by head movement-related signals (Cullen 2009; Cullen and Roy, 2004). The
369 latter consist of trains of action potentials that originate in the three semicircular canals buried
370 within the temporal bone. Each semicircular canal consists of a ring within which a fluid
371 (endolymph) flows and mechanically deflects the stereocilia of multiple cells whenever the
372 head rotation accelerates or decelerates. This deflection either increases or decreases the
373 discharge of primary vestibular neurons (PVN) with a rate proportional to head velocity.

374 The two lateral canals are located in the left and right inner ears and lie roughly in a
375 "plane" which is tilted not only relative to the Reid plane, but also to the "plane" passing
376 through the medial and lateral rectus muscles (Cox and Jeffery, 2008; Ostriker et al., 1985).
377 These canals are often called "horizontal" because they are primarily activated during head
378 rotations about the Earth gravity axis. During such rotations, PVN on the side toward which
379 the nose is rotating (ipsilateral side) increase their firing rate while those on the contralateral
380 side decrease their firing rate. These bilateral changes modify the firing rate of post-synaptic

381 neurons (secondary vestibular neurons) located in the medial and superior vestibular nuclei
382 (McCrea et al., 1987b). On either side of the brain's midsagittal plane, these nuclei are
383 bilaterally organized and house two categories of neurons: type 1 and type 2 neurons. Type 1
384 neurons increase their firing rate during head rotations toward the side (left or right)
385 containing the nucleus in which their discharge is recorded. During contraversive head
386 rotations, they decrease their discharge. In addition, they pause during saccades (McCrea et
387 al. 1987). In the monkey, these neurons have been called Position Vestibular Pause (PVP)
388 neurons. Type 2 neurons exhibit the opposite pattern, increasing their firing rate during
389 contraversive head rotations and decreasing it during ipsiversive head rotations (Fuchs et al.
390 2005; Roy and Cullen, 2002; 2003).

391 **< Figure 5 near here >**

392 Figure 5 schematizes the mechanism by which vestibular signals contribute to ending
393 a leftward movement of gaze toward a visual target. The end of the orienting movement is
394 characterized by a time interval during which the gaze direction remains stationary relative to
395 the trunk while the head slowly terminates its motion. The saccades by both eyes give way to
396 their rotation in the opposite direction. As the head rotates, the primary vestibular neurons
397 (PVN) excite the type I vestibular neurons (EVN-1) in the ipsilateral medial and ventrolateral
398 vestibular nuclei (Broussard et al. 1995). In turn, the EVN-1 initiate and drive the counter-
399 rotation of both eyes in the orbit by exciting the motor (LRMN) and internuclear (AIN) neurons
400 in the contralateral abducens nucleus (red-colored synapses a). Thus, the abducens neurons
401 can resume their activity from the pause imposed by the inhibitory input from IBN in the left
402 Med RF (see blue-colored synapses b in Fig. 2). The LR muscle fibers of the right eye can
403 contract, as well as the MR muscle fibers of the left eye, which are activated via motoneurons
404 located in the OMN. The latter cells (MRMN) are also excited by neurons in the ipsilateral
405 nucleus of the ascending tract of Deiters (ATDN; green-colored synapse b). In parallel, the
406 spikes emitted by other type 1 vestibular neurons (IVN-1) choke the activity the LRMN and
407 AIN in the abducens nucleus (red-colored synapses c), preventing them from responding to
408 any residual descending (target-related) commands. The recruitment of type 2 neurons (IVN-
409 2) in the contralateral vestibular nuclei (red-colored synapse d) complements the cessation of
410 agonist commands by inhibiting the activity of EVN-1, thereby reducing their excitatory
411 influence upon the motor and internuclear neurons (synapses e). IVN-2 also inhibit the activity

412 of IVN-1, which in turn favors the disinhibition of cells (synapses e) in charge of the counter-
413 rotation of the eyes under the influence of EVN-1 (synapses a) (Broussard et al., 1995; Scudder
414 and Fuchs, 1992).

415 The functional anatomy schematized in Fig. 5 enables us to see how the disinhibition
416 of type 1 vestibular neurons can quickly stop the eye saccade during ipsilateral orienting
417 movements. The spikes of IVN-1 (inhibitory synapses c) choke the burst commands while the
418 spikes of EVN-1 order the eyes to counter-rotate as the head terminates its motion (excitatory
419 synapses a). This mechanism also helps us to understand the hypometria of contralateral eye
420 and head movements after unilateral lesion of the caudal fastigial nucleus (for instance the
421 right nucleus) (Quinet and Goffart, 2007). Contralateral (leftward) saccades are hypometric
422 because the agonist drive from EBN lacks of the excitatory input from inactivated (right)
423 fastigial cells, and because the saccade-related burst of neurons in the opposite (unaffected)
424 fastigial nucleus excite IBN in the contralateral (right) reticular formation. These IBN inhibit
425 IVN-2 in the opposite (left) vestibular nuclei, disinhibiting the excitatory influence of EVN-1
426 (initiating the counter-rotation of the eyes) and the inhibitory influence of IVN-1 upon
427 ipsilateral motor neurons (inhibiting the saccade-related drive). The latter mechanism also
428 explains the hypometria of head movements. Their amplitude is truncated because the IVN-1
429 inhibits the ipsilateral neck motor neurons (NMN; synapse e) while the EVN-1 facilitates the
430 contraction of contralateral neck muscles (synapse f).

431 **< Figure 6 near here >**

432 As for the generation of upward and downward saccades (Fig. 3), the neural
433 connectivity involved in the generation of downward and upward slow eye movements are
434 not symmetrical (McCrea et al. 1987a). Figure 6A schematizes the network that stabilizes gaze
435 during downward gaze movements. While the head pitches down and slowly terminates its
436 motion, both eyes rotate upward in the orbits. The upward eye movement is the outcome of
437 contracting SR and IO muscles fibers. Driven by the increased firing of PVN connected with the
438 left anterior canal, excitatory vestibular neurons (EVN-U) in the left medial vestibular nucleus
439 activate the motor neurons innervating the left eye's SR and the right eye's IO muscle fibers
440 (synapses a).

441 In addition to elevating both eyes, the contraction of these fibers causes an
442 incyclotorsion of the left eye and an excyclotorsion of the right eye, as if both eyes were
443 counteracting a tilt of the head toward the left shoulder. This clockwise torsion of both eyes
444 can be attenuated by the torsion caused by the excitatory input from EVN-U in the right MVN.
445 Because of their excitation by the right anterior canal, PVN drive the firing rate of EVN-U,
446 which in turn excite the motor neurons that innervate the right eye's SR and the left eye's IO
447 muscle fibers (synapses b), causing the right eye's incyclotorsion and left eye's excyclotorsion.
448 The absence of torsion during strictly upward slow eye movements suggests that the output
449 of the left and right vestibulo-oculomotor channels have been adjusted so that the torsional
450 signals balance out.

451 Regarding the antagonist muscles, their inhibition is promoted by IVN-U located in the
452 superior vestibular nuclei. These cells inhibit the motor neurons innervating the IR of the
453 ipsilateral eye (synapses c) and those innervating the SO muscle of the contralateral eye
454 (synapse d). Contrary to the network involved in horizontal eye and head movements (Fig. 5),
455 the neural mechanisms triggering the counter-rotation of the eyes when the head pitches
456 down are not yet elucidated. However, the inputs are necessarily bilateral and balanced since
457 the eyes move by the same amount and their torsion canceled.

458 Finally, figure 6B schematizes the neuronal network involved in stopping an upward
459 gaze shift while the head slowly terminates its upward motion. Both eyes rotate downward in
460 the orbits because of the contraction of IR and SO muscle fibers. This is driven by the increased
461 firing of primary vestibular neurons (PVN) connected with the left posterior canal. Excitatory
462 vestibular neurons (EVN-D) in the left medial vestibular and ventral lateral vestibular nuclei
463 activate the motor neurons innervating the fibers of the right eye's IR and the left eye's SO
464 muscles (synapses a).

465 As above, in addition to depressing both eyes, the contraction of these fibers cause an
466 excyclotorsion of the right eye and an incyclotorsion of the left eye, as if the eyes are
467 counteracting a tilt of the head toward the right shoulder. Once again, the binocular torsion
468 is attenuated by a torsion made in the opposite direction and caused by excitatory input from
469 the right posterior canal to the motor neurons innervating the left eye's IR and the right eye's
470 SO muscle fibers (synapses b). Inhibitory vestibular neurons (IVN-D) located in the superior

471 vestibular nuclei prevent the antagonist muscles from contracting by inhibiting the motor
472 neurons innervating the SR of the contralateral eye (synapses c) and possibly also the motor
473 neurons innervating the IO muscle of the ipsilateral eye (synapse d). As for the neural
474 mechanisms ending downward gaze shifts, the neural mechanisms triggering the counter-
475 rotation of the eyes when the head pitches up are not completely identified. Nevertheless, if
476 the eyes do not exhibit any torsional component, the motor commands are necessarily
477 bilateral and balanced.

478 In summary, by studying the networks underlying the generation of slow eye
479 movements, we discover a complexity that radically differs from the straightforward solutions
480 that human engineers would have designed. One puzzling question concerns the constraints
481 that led the torsional components caused by the contraction of elevator or depressor muscles
482 to cancel each other out. Moreover, the connectivity that we have described is not complete,
483 because new eye movements can be produced in response to “unusual” combinations of head
484 rotations and visual stimulation. Experiments in the cat have shown that the visual stimulation
485 exerts a significant influence on the functional connectivity yielding the vestibulo-oculomotor
486 responses. Horizontal compensatory eye movements can be generated during vertical head
487 rotations in the dark after training sessions during which the visual field moved horizontally
488 whenever the head rotated vertically. The horizontal nystagmus persisted in the absence of
489 any new visuo-vestibular experience (Schultheis and Robinson, 1981).

490 Fixating a static visual target

491 Considering the line of sight as the direction of gaze measured when a subject fixates
492 a visual target, the binocular fixation point designates the intersection between the two lines
493 of sight (Collewijn et al., 1997). Of course, the quantification of movements must be
494 distinguished from their neurophysiological correspondence. Looking at a visual target
495 obviously does not reduce to the excitation of one single photoreceptor on the fovea of each
496 eye. We will therefore rather speak of a binocular fixation zone. On the motor side, the
497 orientation of the eyes is maintained steady by the sustained discharge of a tonic neurons
498 distributed in the nucleus prepositus hypoglossi (NPH), the vestibular nuclei (in the medullary
499 reticular formation) and in the interstitial nucleus of Cajal (Fukushima et al. 1992; McFarland
500 and Fuchs, 1992; Moschovakis, 1997). The more deviated the eyes in the orbit, the higher their

501 firing rate. In addition to connective tissues that wrap each eyeball within the orbit, two
502 neuronal networks are involved in maintaining the rotation axes of the eyes within Listing's
503 plane and deviating them horizontally and vertically. These networks are relatively
504 independent from each other insofar as a lesion in the NPH primarily affects the ability to
505 maintain gaze direction within the horizontal plane, with marginal effects on vertical gaze
506 direction (Kaneko, 1997; 1999). After the saccades, whose amplitude is unaffected, the eyes
507 exhibit an exponential drift toward their primary orientation (roughly straight ahead). By
508 contrast, lesion in the NIC causes a failure to hold vertical deviations of the eyes (i.e. both
509 upward and downward) but not horizontal deviations. A torsional drift of the eyes is also
510 observed when the lesion is unilateral (Helmchen et al., 1998).

511 Considering now the case when the foveation changes between targets or locations
512 situated at different depths in the three-dimensional physical space, three visuomotor
513 processes occur: 1) vergence, 2) accommodation and 3) variation of pupil diameter.
514 Convergence designates the symmetrical eye movements made when the MR muscle fibers
515 of both eyes contract, diminishing the distance of the binocular fixation zone relative to the
516 head. Divergence designates those movements made when the distance increases because of
517 a binocular relaxation of MR muscle fibers. The contraction level of intraocular muscles also
518 changes (McDougal and Gamlin, 2015). During convergence, the contraction of ciliary muscles
519 (CM) reduces the tension in zonular fibers, changing the refractive power of the lens as its
520 anterior surface bulges forward (increase in convexity). Thus, the target images are brought
521 into focus on the foveae. This accommodation is associated with a myosis, i.e., the constriction
522 of the iris and the reduction of the pupil diameter, thereby increasing the depth of field. The
523 myosis itself results from the contraction of the sphincter pupillae (SP) combined with the
524 relaxation of the dilator pupillae (DP). Altogether, these three visuomotor processes
525 constitute the so-called the near response (or near triad).

526 Although these processes are generated jointly, clinical studies report pathological
527 cases in which they are dissociated and the neuroanatomy helps us to understand why. Figure
528 7 shows that the contraction of MR muscle fibers is controlled by the firing rate of motor
529 neurons (MRMN) located in the oculomotor nucleus whereas the intrinsic musculature of each
530 eye is controlled by motor neurons located in the ciliary ganglions (CG) and premotor neurons
531 in the Edinger-Westphal preganglionic nucleus (EWpg). In both cases, the innervation is

532 ipsilateral (Gamlin, 1999). The binocular control and the near triad starts with their afferent
533 input from two major regions of the midbrain tegmentum: the supraoculomotor area (SOA)
534 and the central mesencephalic reticular formation (cMRF) (May et al., 2018; 2019).

535 **< Figure 7 near here >**

536 The SOA is primarily involved in the generation of the slow near response and its
537 sustained maintenance. Therein, we find burst neurons and tonic neurons, with separate
538 subgroups characterized by different discharge patterns during the generation of diverging
539 and converging eye movements. More common just lateral to the MLF, four categories of
540 neurons are found: convergence burst neurons, divergence burst neurons, convergence tonic
541 neurons and divergence tonic neurons (Mays, 1984; Mays et al., 1986). Whether divergence
542 neurons exert an inhibitory influence upon the firing rate of convergence neurons is not
543 known yet. Neurons in SOA send projections to the medial rectus MN that are ipsilateral,
544 whereas the projections to the EWpg are bilateral (May et al., 2018).

545 In the cMRF, we find neurons that emit a burst discharge whenever the near response
546 accompanies a saccadic eye movement: saccade convergence burst neurons and saccade
547 divergence burst neurons. The former fire during asymmetrical saccades associated with an
548 increase of the vergence angle and the latter fire during asymmetrical saccades associated
549 with a divergence (Quinet et al., 2020). Each cMRF innervates the SOA and the EWpg
550 bilaterally. It also sends ipsilateral projections to the sites in the OMN in which the motor
551 neurons that innervate the singly-innervated muscle fibers are located (May et al., 2018).

552 The bilateral projections from SOA to the EWpg, and from the cMRF to both EWpg and
553 SOA are completed by bilateral afferents from caudal fastigial nuclei (CFN) (Bohlen et al. 2021;
554 May et al. 1992) whose contribution in the near-triad has been reported (Bohlen et al., 2021;
555 Gamlin and Zhang, 1996). In addition to send bilateral projections to the midbrain tegmentum
556 and contralateral projections to the pontomedullary reticular formation (where EBN and IBN
557 are located, Fig. 2), the CFN also project bilaterally to the rostral part of the deep superior
558 colliculi, with a contralateral predominance (May et al. 1990). Therein, some cells either
559 increase (convergence neurons) or decrease (divergence neurons) their firing rate during
560 vergence eye movements (Bohlen et al., 2021; Gamlin 2002; Gamlin et al., 1996; Zhang and
561 Gamlin, 1996).

562 In addition to modulations during vergence (van Horn et al., 2013), recording studies
563 in the rostral superior colliculi also revealed the presence of cells that burst during miniature
564 saccades (also called microsaccades), like more caudal cells discharge during larger saccades
565 (Hafed and Krauzlis 2012). Local pharmacological inactivation confirmed their causal role in
566 the generation of microsaccades (Hafed et al. 2009). However, it also unraveled an implication
567 of rostral SC in specifying the direction of gaze when a target is presented (or even imagined)
568 in the central visual field, static or moving (Hafed et al. 2008). Indeed, unbalanced activity
569 between the two rostral SC causes a fixation offset to the extent that gaze direction is not
570 directed anymore to the target center but toward an offset location. The same deficit happens
571 after unilateral pharmacological perturbation of CFN, suggesting that the sustained firing rate
572 of its neurons adjusts the balance of activity between the two rostral SC during visual fixation
573 (Bourrelly et al., 2018b; Goffart et al., 2012; Guerrasio et al., 2010). Thus, the direction of gaze
574 would correspond to an equilibrium of commands that counterbalance with each other,
575 involving fastigial projections to the left and right rostral SC but also to the left and right nuclei
576 *reticularis tegmenti pontis* (NRTP). Through their projections to NRTP, the two rostral SC and
577 CFN have access to neurons which, in the flocculus and paraflocculus, influence the firing rate
578 of tonic neurons that innervate the motor and internuclear neurons in the abducens nuclei
579 (Belton and McCrea 2000; Lisberger et al. 1994; Noda and Mikami, 1986; Noda and Suzuki,
580 1979; Noda et al., 1990).

581 In summary, the knowledge of the detailed neuronal circuitry underlying the
582 generation of disjunctive eye movements is not advanced as much as the circuitry for
583 conjugate saccades (Mays and Gamlin, 1995a, b) but a lot of progress has been made during
584 the last decade (e.g., May et al., 2022). Soon it will be possible to understand the complex
585 neurophysiology that underlies the generation of vergence eye movements and their
586 dysfunction (Das 2016; Lennerstrand 2007).

587 Tracking a moving visual target

588 A target drifting in the visual field often elicits a primary saccade (called interceptive
589 saccade) at the end of which gaze does *not* land ahead of the moving target. Interceptive
590 saccades either accurately capture (foveate) or fall behind the target. Then, a slow eye
591 movement follows with a velocity that rarely matches the target's. A textured background

592 (yielding visual motion in the direction opposite to the eye movement) does not seem to be
593 responsible for the pursuit slower than target velocity because the consequences of its
594 suppression are very weak (Keller and Khan, 1986). A subsequent correction saccade (called
595 catch-up saccade) restores the foveation. Thus, unless the target trajectory is already known
596 and foreseeable, gaze lags behind the target most of the time (Goffart et al., 2017a), especially
597 when the target is small: the visual tracking is mostly composed of catch-up saccades
598 interspersed by intervals during which the eyes drift roughly in the same direction as the target
599 and with a lesser speed. In well-trained subjects, the catch-up saccades can be eliminated
600 (Botschko et al. 2018; Bourrelly et al., 2016). Moreover, increasing the target size enhances
601 the velocity of slow eye movements and reduces the size and number of catch-up saccades
602 (Pola and Wyatt, 1985). This point is important because most experimental studies use a small
603 spot of light as a target.

604 Contrary to a static target which yields a bounded activity, a moving spot evokes on
605 the retina a streak of activity the tail of which corresponds to past locations of the target and
606 the leading edge the most recent location. In the deep superior colliculus, as for saccades to
607 a static target, the population of saccade-related burst neurons is expanded (Anderson et al.,
608 1998; Goossens and van Opstal 2006; Sparks et al., 1976). However, it does not recruit cells
609 that fire during saccades to future target locations. The population consists of a continuum of
610 cells ranging from neurons issuing commands related to past locations of the target to neurons
611 issuing commands related to its current location (Goffart et al., 2017). At the level of motor
612 neurons, other input must therefore complement the collicular signals. This input stems, at
613 least in part, from motion signals processed in area MT because its lesion leads to saccades
614 landing at locations that the target crossed earlier. Thus, the ability to foveate the visual target
615 may consist of adding a “predictive” component based on the integration of putative target-
616 related “velocity” signals to collicular signals (Keller et al., 1996; Optican, 2009). However, an
617 alternative explanation is possible. While considering that any target recruits an extended
618 assembly of neurons, the generation of accurate interceptive saccades may result from the
619 attenuation of the synaptic weight of spikes emitted by the cells that were the earliest excited
620 by the passage of the target, combined with the promotion of the synaptic weight of spikes
621 emitted by the latest and most recently excited cells.

652 organization, such that a reduced firing rate on one side enhances the firing rate on the
653 opposite side. Thus, by suppressing the activity of the neurons that innervate the antagonist
654 muscle fibers (red synapses a), the inhibitory tonic neurons (ITN) in the NPH/MVN contribute
655 to the relaxation of these muscles, while disinhibiting the ipsilateral abducens and
656 internuclear neurons (blue synapses a) and promoting an ipsiversive movement of both eyes.

657 Neurons in NPH/MVN are monosynaptically inhibited (synapse b) by Purkinje cells in
658 the flocculus and the ventral paraflocculus (Lisberger et al. 1994), which receive excitatory
659 input from the three major pontine nuclei (synapses c) involved in relaying visual motion
660 signals to the cerebellum: the rostral part of nucleus reticularis tegmenti pontis (rNRTP), the
661 dorsolateral pontine nucleus (DLPN) and the dorsomedian pontine nucleus (DMPN). These
662 pontine nuclei contribute to the inhibition that Purkinje cells exert upon NPH/MVN neurons.
663 After a chemical lesion in the dorsolateral pontine nucleus (DLPN) or in the rostral nucleus
664 reticularis tegmenti pontis (rNRTP), ipsilesional pursuit eye movements are severely impaired
665 (May et al., 1988; Ono et al. 2003; Suzuki et al., 1999). The suppression of pontine input likely
666 reduces the floccular inhibition of ITN neurons, which in turn impede the recruitment of
667 agonist motor and internuclear neurons. The consequences of suppressing the input from
668 DMPN have not been tested yet. However, an impairment of ipsilesional pursuit is expected
669 because the DMPN and the contralateral caudal fastigial nucleus exhibit strong reciprocal
670 connections, and inactivation of the latter severely impairs the generation of contralesional
671 pursuit eye movements (Bourrelly et al., 2018b).

672 The NPH/MVN are also the target of signals from neurons in the pretectal nucleus of
673 the optic tract (NOT; synapse d) (Mustari et al. 1994), some of which are sensitive to the
674 motion of a small visual target (Fuchs et al., 1992; Mustari and Fuchs, 1990). These neurons
675 fire during ipsilateral target motions (Mustari et al., 1990) and their loss severely impairs the
676 ability to generate ipsilesional slow pursuit eye movements (Yakushin et al., 2000). NOT
677 receives visual signals from the retina and from the middle temporal (MT) and medial superior
678 temporal (MST) areas of cerebral cortex (synapse f) wherein numerous cells are sensitive to
679 target motion as well (Distler et al., 2002). Projections of the frontal eye field (FEF) to NOT and
680 to NPH have been documented (synapses e and e'). This access of frontal cortex to premotor
681 neurons does not indicate a voluntary control of pursuit eye movements since the presence
682 of a target is required for initiating a smooth pursuit. The NOT contributes to the maintenance

683 of ipsilateral pursuit eye movements by its direct projections to tonic neurons in the ipsilateral
684 NPH/MVN (synapse d), but also to the ipsilateral pontine nuclei (synapses g).

685 Through their projections to the ipsilateral rostral SC (synapse j), Gaba-ergic cells in
686 NOT may contribute to suppression of the generation of saccades made in the direction
687 opposite to the target motion. Simultaneously, they may depress the excitatory drive that
688 NRTTP exerts upon the ipsilateral paraflocculus (synapse k).

689 Finally, the secondary excitatory vestibular neurons (EVN-1 or PVP cells) that are
690 involved in moving the eyes in the direction opposite to the head movement (Fig. 5) increase
691 their firing rate during contraversive pursuit eye movements. However, as they do not seem
692 to be inhibited by Purkinje cells located in the floccular region, the origin of the discharge
693 modulation during contraversive pursuit remains to be identified. Direct projections from the
694 frontal eye fields to the NPH/MVN might support this influence.

695 Conclusion

696 Studying orienting movements of gaze is a convenient means for investigating how
697 intertwined networks of neurons in the central nervous system transforms sensory signals into
698 motor commands, but also, how it enables an entire animal to visually locate and capture an
699 object in the physical environment and to adapt when intrinsic damage or contextual changes
700 alter their execution.

701 During the last decades, neurophysiologists and neuroanatomists gathered in the
702 monkey considerable knowledge that enabled identifying the core networks involved in the
703 generation of eye movements (Büttner-Ennever, 2006; Fuchs et al., 1985; Goffart et al., 2017a;
704 Grandin et al., 2002; Henn et al., 1984; Horn and Leigh, 2011; Krauzlis, 2004; Krauzlis et al.
705 2017; Moschovakis et al., 1996; Raphan and Cohen, 1978; Scudder et al., 2002; Soetedjo et
706 al., 2019; Sparks, 2002; Sparks and Mays, 1990; Ugolini et al. 2006). Technical developments
707 offered the possibility to measure precisely the time course of these movements and to study
708 correlations between the firing rate of neurons and kinematic parameters such as movement
709 amplitude, velocity, acceleration and various differences (e.g., gaze error, velocity error, etc.).
710 However, these correlation studies should not lead us to believe that a one-to-one
711 correspondence exists between the multiple networks within which neuronal activities

712 propagate, and the extrinsic and homogeneous medium (mathematical space) with which we
713 quantify the motion of rigid objects such as the eyeballs, the head and targets. Such a mapping
714 is not at all obvious; it is even questionable. Contrary to the physical space, the medium of
715 neuronal activity is neither homogeneous nor passive. Unlike most objects at which we look
716 and that we manipulate, the corresponding brain activity is not rigid. Even the mere spot of
717 activity that a small static object evokes on the retina yields multiple parallel flows of activity
718 that make its correspondence in the brain spatially distributed, temporally extended and
719 context-dependent (Goffart et al. 2018). Moreover, reducing neurons or neuronal chains to
720 units encoding geometric or kinematic relations between gaze and target conceals the muscle
721 forces and the multiple antagonisms that we have described between muscles (Miller et al.
722 2002). Furthermore, the velocity profile of the movement and the time course of force
723 development do not match (Miller and Robins, 1992). During horizontal saccades for example,
724 the duration of the interval during which the muscle force increases is on average 60% shorter
725 than the duration of the total change in eye orientation. A force decay happens well before
726 the change in eye position ends and the larger the saccade, the longer its lead-time relative to
727 saccade arrest (Lennerstrand et al., 1993). Finally, the temporal series of numerical values
728 quantifying the changes during eye reorientation hides the antagonisms between those
729 multiple channels that generate each movement. To our knowledge, no study has reported
730 identical time courses between the kinematic and neurophysiological recordings that would
731 support the assumption of an isomorphism.

732 In the majority of models proposed during the last decades, the movements of the
733 eyes and the head were considered as driven by (intrinsic) error signals encoding displacement
734 vectors in physical space (e.g., Lisberger et al. 1987). These models have definitely been useful
735 to communicate concepts and to bring a comprehensive picture of the complexity underlying
736 the generation of movements. However, embedding geometric and kinematic notions within
737 the inner functioning of the brain (i.e., mapping intrinsic neuronal signals with extrinsic
738 behavioral measurements) may be neurophysiologically misleading because different
739 constraints characterize the neurophysiological and kinematic descriptions. In fact, rather
740 than outcomes of processes reducing geometric or kinematic errors, the saccade and pursuit
741 deceleration may resume merely to transitions from imbalance to equilibria opposing

742 populations of neurons whose activity leads to mutually antagonist movement tendencies
743 (Goffart, 2019).

744 Further empirical investigation is still required to determine and explain several other
745 issues such as: whether and how the networks underlying orienting movements of the eyes
746 and head interact with those generating other types of goal-directed action like hand reaching
747 movements or whole-body locomotion; whether and how they interact with the networks
748 involved in the navigation and the memory of locations; whether and how they support the
749 learning of new skills and possibly the acquisition of more abstract knowledge such as
750 geometry or counting; etc. With the recent multiplication of cognitive studies that use eye-
751 tracking techniques to explore with quantitative methods the so-called “inner space”
752 (Johansson et al., 2013; Strohmaier et al. 2020), more effort is required to characterize what
753 exactly are those covert processes that eye movements would express. Neurophysiological
754 and neuroanatomical works must also continue to reveal the innervation of extraocular
755 muscles compartments and unravel the multiplicity of channels and neuronal networks that
756 enable a living organism to interact visually with its natural and social environment.

757 By studying the simple and seemingly straightforward problem of orienting the eyes
758 toward the location of a physical object, we see how the neurophysiological and
759 neuroanatomical studies performed with animals enables us to understand the symptoms
760 exhibited by patients suffering from eye movement disorders (Horn and Leigh, 2011; Rucker
761 and Lavin, 2021; Sharpe, 2008; Sharpe and Wong, 2005). They also show that evolutionary
762 and embryological developments led to patterns that are much more complex and redundant
763 than the straightforward strategies of human engineers. Contrary to the case where a design
764 precedes the realization of a machine for solving a specific problem, the Physiology teaches
765 us that any behavioral observation is the outcome of a multiple networks and parallel chains
766 of neurons. It also teaches us how the multiplicity and the diversity of elements enable an
767 organism to solve several problems in parallel and even take up new challenges.

768

769 Bibliography

770

771 Anderson, R. W., Keller, E. L., Gandhi, N. J., Das, S., 1998. Two-dimensional saccade-related
772 population activity in superior colliculus in monkey. *J. Neurophysiol.* 80, 798-817.

773 Balliet, R., Nakayama, K., 1978. Training of voluntary torsion. *Invest. Ophthalm. Visual Sci.* 17,
774 303-314.

775 Barton EJ, Nelson JS, Gandhi, N.J., Sparks, D.L., 2003 Effects of partial lidocaine inactivation of
776 the paramedian pontine reticular formation on saccades of macaques. *J. Neurophysiol.* 90,
777 372–386.

778 Basso, M.A., Krauzlis, R.J., Wurtz, R.H., 2000. Activation and inactivation of rostral superior
779 colliculus neurons during smooth-pursuit eye movements in monkeys. *J. Neurophysiol.* 84,
780 892-908.

781 Belton, T., McCrea, R.A., 2000 Role of the cerebellar flocculus region in the coordination of
782 eye and head movements during gaze pursuit. *J. Neurophysiol.* 84, 1614-1626.

783 Bizzi, E., Kalil, R.E., Morasso, P., 1972. Two modes of active eye-head coordination in monkeys.
784 *Brain Res.*, 40, 45-48.

785 Bohlen, M., Gamlin, P.D., Warren, S., May, P.J., 2021. Cerebellar projections to the macaque
786 midbrain tegmentum: possible near response connections. *Visual Neurosci.* 38, E007.

787 Botschko Y., Yarkoni M., Joshua M., 2018. Smooth pursuit eye movement of monkeys naive to
788 laboratory setups with pictures and artificial stimuli. *Front. Syst. Neurosci.* 12, 15.

789 Burrelly, C., Quinet, J., Goffart, L., 2018a. The caudal fastigial nucleus and the steering of
790 saccades toward a moving visual target. *J. Neurophysiol.* 120, 421-438.

791 Burrelly, C., Quinet, J., Goffart, L., 2018b. Pursuit disorder and saccade dysmetria after caudal
792 fastigial inactivation in the monkey. *J. Neurophysiol.* 120, 1640-1654.

793 Burrelly, C., Quinet, J., Goffart, L., 2021. Bilateral control of interceptive saccades: evidence
794 from the ipsipulsion of vertical saccades after caudal fastigial inactivation. *J. Neurophysiol.*
795 125, 2068-2083.

796 Burrelly, C., Quinet, J., Cavanagh, P., Goffart, L., 2016. Learning the trajectory of a moving
797 visual target and evolution of its tracking in the monkey. *J. Neurophysiol.* 116, 2739-2751.

798 Broussard, D.M., DeCharms, R.C., Lisberger, S.G., 1995. Inputs from the ipsilateral and
799 contralateral vestibular apparatus to behaviorally characterized abducens neurons in rhesus
800 monkeys. *J. Neurophysiol.* 74, 2445-2459.

801 Büttner-Ennever, J.A., 2006. Neuroanatomy of the oculomotor system. *Prog. Brain Res.* Vol.
802 151.

803 Büttner, U., Büttner-Ennever, J.A., 2006. Present concepts of oculomotor organization. *Prog.*
804 *Brain Res.* 151, 1-42.

805 Büttner, U., Buttner-Ennever J., Henn V., 1977. Vertical eye movement related unit activity in
806 the rostral mesencephalic reticular formation of the alert monkey. *Brain Res.* 130, 239–252.

807 Clendaniel, R.A., Mays, L.E., 1994. Characteristics of antidromically identified oculomotor
808 internuclear neurons during vergence and versional eye movements. *J. Neurophysiol.* 71,
809 1111–1127.

810 Collewijn, H., Erkelens, C.J., Steinman, R.M., 1997. Trajectories of the human binocular fixation
811 point during conjugate and non-conjugate gaze-shifts. *Vision Res.* 37, 1049-1069.

812 Corneil, B.D., Olivier, E., Richmond, F.J., Loeb, G.E., Munoz, D.P., 2001. Neck muscles in the
813 rhesus monkey. II. Electromyographic patterns of activation underlying postures and
814 movements. *J. Neurophysiol.* 86, 1729-1749.

815 Cox, P.G., Jeffery, N., 2008. Geometry of the semicircular canals and extraocular muscles in
816 rodents, lagomorphs, felids and modern humans. *J. Anat.* 213, 583-596.

817 Crawford, J.D., Vilis, T., 1992. Symmetry of oculomotor burst neuron coordinates about
818 Listing's plane. *J. Neurophysiol.* 68, 432-448.

819 Cullen, K.E., 2009. Eye and head movements. *Encyclopedia of neuroscience*, 157-167.

820 Cullen, K.E., Roy, J.E., 2004. Signal processing in the vestibular system during active versus
821 passive head movements. *J. Neurophysiol.* 91, 1919-1933.

822 Demer, J.L., 2015. Compartmentalization of extraocular muscle function. *Eye* 29, 157-162.

823 Demer, J.L., 2019. *Anatomy and physiology of the extraocular muscles and surrounding
824 tissues.* Ophthalmology. 5th ed. Philadelphia, PA: Elsevier.

825 Das, V.E., 2016. Strabismus and the oculomotor system: insights from macaque models. *Annu.
826 Rev. Vis. Sci.* 2, 37-59.

827 Davis-López de Carrizosa, M.A., Morado-Díaz, C.J., Miller, J.M., de la Cruz, R.R., Pastor, A.M.,
828 2011. Dual encoding of muscle tension and eye position by abducens motoneurons. *J.
829 Neurosci.* 31, 2271–2279.

830 Distler, C., Mustari, M., Hoffmann, K.P., 2002. Cortical projections to the nucleus of the optic
831 tract and dorsal terminal nucleus and to the dorsolateral pontine nucleus in macaques, a dual
832 retrograde tracing study. *J. Comp. Neurol.* 444, 144-158.

833 Fuchs, A.F., Kaneko, C.R., Scudder, C.A., 1985. Brainstem control of saccadic eye movements.
834 *Annu. Rev. Neurosci.* 8, 307–337.

835 Fuchs, A.F., Ling, L., Phillips, J.O., 2005. Behavior of the position vestibular pause (PVP)
836 interneurons of the vestibuloocular reflex during head-free gaze shifts in the monkey. *J.
837 Neurophysiol.* 94, 4481-4490.

838 Fuchs, A.F., Luschei, E.S., 1970. Firing patterns of abducens neurons of alert monkeys in
839 relationship to horizontal eye movement. *J. Neurophysiol.* 33, 382–392.

840 Fuchs, A.F., Mustari, M.J., Robinson, F.R., Kaneko, C.R., 1992. Visual signals in the nucleus of
841 the optic tract and their brain stem destinations. *Ann. N.Y. Acad. Sci.* 656, 266-276.

842 Fuchs, A.F., Scudder, C. A., Kaneko, C.R., 1988. Discharge patterns and recruitment order of
843 identified motoneurons and internuclear neurons in the monkey abducens nucleus. *J.*
844 *Neurophysiol.*, 60, 1874-1895.

845 Fuchs, A.F., Robinson, F.R., Straube, A., 1993. Role of the caudal fastigial nucleus in saccade
846 generation. I. Neuronal discharge pattern. *J. Neurophysiol.*, 70, 1723-1740.

847 Fukushima, K., Kaneko, C.R.S., Fuchs, A.F., 1992. The neuronal substrate of integration in the
848 oculomotor system. *Progr. Neurobiol.* 39, 609-639.

849 Gamlin, P.D., 1999. Subcortical neural circuits for ocular accommodation and vergence in
850 primates. *Ophthalm. Physiol. Opt.* 19, 81-89.

851 Gamlin, P.D., 2002. Neural mechanisms for the control of vergence eye movements. *Ann. N.Y.*
852 *Acad. Sci.* 956, 264-272.

853 Gamlin, P.D., 2006. The pretectum: connections and oculomotor-related roles. *Prog. Brain*
854 *Res.* 151, 379-405.

855 Gamlin, P.D., Yoon, K., Zhang, H., 1996. The role of cerebro-ponto-cerebellar pathways in the
856 control of vergence eye movements. *Eye (London)* 10, 167-171.

857 Gamlin, P.D.R. & Zhang, H.Y. (1996). Effects of muscimol blockade of the posterior fastigial
858 nucleus on vergence and ocular accommodation in the primate. *Soc. Neurosci. Abstr.* 22,
859 2034.

860 Gandhi, N.J., Sparks, D.L., 2007. Dissociation of eye and head components of gaze shifts by
861 stimulation of the omnipause neuron region. *J. Neurophysiol.* 98, 360-373.

862 Goffart, L., 2019. Kinematics and the neurophysiological study of visually-guided eye
863 movements. *Prog. Brain Res.* 249, 375–384.

864 Goffart, L., Burrelly, C., Quinton, J.-C., 2018. Neurophysiology of visually-guided eye
865 movements: critical review and alternative viewpoint. *J. Neurophysiol.* 120, 3234–3245.

866 Goffart, L., Burrelly, C., Quinet, J., 2017. Synchronizing the tracking eye movements with the
867 motion of a visual target: Basic neural processes. *Prog. Brain Res.* 236, 243–268.

868 Goffart, L., Chen, L.L., Sparks, D.L., 2003. Saccade dysmetria during functional perturbation of
869 the caudal fastigial nucleus in the monkey. *Ann. N.Y. Acad. Sci.* 1004, 220-228.

870 Goffart, L., Chen, L.L., Sparks, D.L., 2004. Deficits in saccades and fixation during muscimol
871 inactivation of the caudal fastigial nucleus in the rhesus monkey. *J. Neurophysiol.* 92, 3351-
872 3367.

873 Goffart, L., Cecala A.L., Gandhi, N.J., 2017. The superior colliculus and the steering of saccades
874 toward a moving visual target. *J. Neurophysiol.* 118, 2890-2901.

875 Goffart, L., Hafed Z.M., Krauzlis R.J., 2012. Visual fixation as equilibrium: evidence from
876 superior colliculus inactivation. *J. Neurosci.* 32, 10627–10636.

877 Goossens, H.H., Van Opstal, A.J., 2006. Dynamic ensemble coding of saccades in the monkey
878 superior colliculus. *J. Neurophysiol.* 95, 2326-2341.

879 Grandin A., Brandi A.M., Dubayle D., Graf W., Ugolini G., Hadjidimitrakis K., Moschovakis A.,
880 2002. Density gradients of trans-synaptically labeled collicular neurons after injections of
881 rabies virus in the lateral rectus muscle of the rhesus monkey. *J. Comp. Neurol.* 451, 346-361.

882 Guerrasio, L., Quinet, J., Büttner, U., Goffart, L., 2010. Fastigial oculomotor region and the
883 control of foveation during fixation. *J. Neurophysiol.* 103, 1988-2001.

884 Hafed Z.M., Goffart, L., Krauzlis R.J., 2009. A neural mechanism for microsaccade generation in
885 the primate superior colliculus. *Science* 323, 940-943.

886 Hafed Z.M., Goffart, L., Krauzlis R.J., 2008. Superior colliculus inactivation causes stable offsets
887 in eye position during tracking. *J. Neurosci.* 28, 8124-8137.

888 Hafed Z.M., Krauzlis R.J., 2012. Similarity of superior colliculus involvement in microsaccade
889 and saccade generation. *J. Neurophysiol.* 107, 1904–1916.

890 Helmchen, C., Rambold, H., Fuhry, L., Büttner, U., 1998. Deficits in vertical and torsional eye
891 movements after uni-and bilateral muscimol inactivation of the interstitial nucleus of Cajal of
892 the alert monkey. *Exp. Brain Res.* 119, 436-452.

893 Henn V., Lang W., Hepp K., Reisine H., 1984. Experimental gaze palsies in monkeys and their
894 relation to human pathology. *Brain* 107, 619–636.

895 Highstein, S.M., Holstein, G.R., 2006. The anatomy of the vestibular nuclei. *Prog. Brain Res.*
896 151, 157-203.

897 Hikosaka, O., Wurtz, R.H. (1985) Modification of saccadic eye movements by GABA-related
898 substances. I. Effect of muscimol and bicuculline in monkey superior colliculus. *J.*
899 *Neurophysiol.* 53, 266 –291.

900 Horn, A.K., Leigh, R.J., 2011. The anatomy and physiology of the ocular motor system.
901 *Handbook of clinical neurology* 102, 21-69.

902 Horn, A.K., Helmchen, C., Wahle, P., 2003. GABAergic neurons in the rostral mesencephalon
903 of the macaque monkey that control vertical eye movements. *Ann. N.Y. Acad. Sci.* 1004, 19-
904 28.

905 Hu X., Jiang H., Gu C., Li C., Sparks, D.L., 2007. Reliability of oculomotor command signals
906 carried by individual neurons. *Proc. Nat. Acad. Sci.* 104, 8137–8142.

907 Johansson, R., Holsanova, J., Holmqvist, K., 2013. Using eye movements and spoken discourse
908 as windows to inner space. In *The construal of spatial meaning: Windows into conceptual*
909 *space* (pp. 9-28). Oxford University Press.

910 Kaneko C.R., 1996. Effect of ibotenic acid lesions of the omnipause neurons on saccadic eye
911 movements in rhesus macaques. *J. Neurophysiol.* 75, 2229–2242.

- 912 Kaneko C.R., 1997. Eye movement deficits after ibotenic acid lesions of the nucleus prepositus
913 hypoglossi in monkeys. I. Saccades and fixation. *J. Neurophysiol.* 78, 1753-1768.
- 914 Kaneko, C.R., 1999. Eye movement deficits following ibotenic acid lesions of the nucleus
915 prepositus hypoglossi in monkeys II. Pursuit, vestibular, and optokinetic responses. *J.*
916 *Neurophysiol.* 81, 668-681.
- 917 Keller, E.L., 1974 Participation of medial pontine reticular formation in eye movement
918 generation in monkey. *J. Neurophysiol.* 37, 316–332.
- 919 Keller, E.L., Gandhi, N.J., Weir, P.T., 1996. Discharge of superior collicular neurons during
920 saccades made to moving targets. *J. Neurophysiol.* 76, 3573–3577.
- 921 Keller, E.L., Heinen, S.J., 1991. Generation of smooth-pursuit eye movements: neuronal
922 mechanisms and pathways. *Neurosc. Res.* 11, 79-107.
- 923 Keller, E.L., Khan, N.S., 1986. Smooth pursuit initiation in the presence of a textured
924 background in monkey. *Vision Res.* 26, 943-955.
- 925 Keller, E.L., McPeck, R.M., Salz, T., 2000 Evidence against direct connections to PPRF EBNs
926 from SC in the monkey. *J. Neurophysiol.* 84, 1303-1313.
- 927 Kojima, Y., Iwamoto, Y., Robinson, F.R., Noto, C.T., Yoshida, K., 2008. Premotor inhibitory
928 neurons carry signals related to saccade adaptation in the monkey. *J. Neurophysiol.* 99, 220–
929 230.
- 930 Krauzlis, R.J., 2004. Recasting the smooth pursuit eye movement system. *J. Neurophysiol.* 91,
931 591-603.
- 932 Krauzlis, R.J., Basso, M.A., Wurtz, R.H., 2000. Discharge properties of neurons in the rostral
933 superior colliculus of the monkey during smooth-pursuit eye movements. *J. Neurophysiol.* 84,
934 876–891.
- 935 Krauzlis, R.J., Goffart, L., Hafed, Z.M., 2017. Neuronal control of fixation and fixational eye
936 movements. *Philos. Trans. Roy. Soc. Lond. B Biol. Sci.* 372, 20160205.
- 937 Langer, T., Kaneko, C.R.S., Scudder, C.A., Fuchs, A.F., 1986. Afferents to the abducens nucleus
938 in the monkey and cat. *J. Comp. Neurol.* 245, 379–400.
- 939 Leigh, R.J., Zee, D.S., 2006. *The Neurology of Eye Movements.* Oxford, UK: Oxford University
940 Press.
- 941 Lennerstrand, G., 2007. Strabismus and eye muscle function. *Acta Ophthalm. Scand.* 85, 711-
942 723.
- 943 Lennerstrand, G., Tian, S., Zhao, T.X., 1993. Force development and velocity of human saccadic
944 eye movements. I: Abduction and adduction. *Clin. Vision Sci.* 8, 295–305.
- 945 Lestienne, F.G., Le Goff, B., Liverneaux, P.A., 1995. Head movement trajectory in three-
946 dimensional space during orienting behavior toward visual targets in rhesus monkeys. *Exp.*
947 *Brain Res.* 102, 393-406.

- 948 Lisberger, S.G., Morris, E.J., Tychsen, L., 1987. Visual motion processing and sensory-motor
949 integration for smooth pursuit eye movements. *Annu. Rev. Neurosci.* 10, 97-129, 1987.
- 950 Lisberger, S.G., Pavelko, T.A., Broussard, D.M., 1994. Responses during eye movements of
951 brain stem neurons that receive monosynaptic inhibition from the flocculus and ventral
952 paraflocculus in monkeys. *J. Neurophysiol.* 72, 909-927.
- 953 Luschei, E.S., Fuchs, A.F., 1972. Activity of brain stem neurons during eye movements of alert
954 monkeys. *J. Neurophysiol.* 35, 445–461.
- 955 May, J.G., Keller, E.L., Suzuki, D.A., 1988. Smooth-pursuit eye movement deficits with chemical
956 lesions in the dorsolateral pontine nucleus of the monkey. *J. Neurophysiol.* 59, 952-77.
- 957 May, P.J., Bohlen, M.O., Perkins, E., Wang, N., Warren, S., 2021. Superior colliculus projections
958 to target populations in the supraoculomotor area of the macaque monkey. *Vis. Neurosci.* 38.
- 959 May, P.J., Gamlin, P.D., Warren, S., 2022. A Novel Tectal/Pretectal Population of Premotor
960 Lens Accommodation Neurons. *Invest. Ophthalmol. Visual Sci.* 63, 35-35.
- 961 May, P.J., Hartwich-Young, R., Nelson, J., Sparks, D.L., Porter, J.D., 1990. Cerebellotectal
962 pathways in the macaque: implications for collicular generation of saccades. *Neuroscience* 36,
963 305-324.
- 964 May, P.J., Porter, J.D., Gamlin, P.D., 1992. Interconnections between the primate cerebellum
965 and midbrain near-response regions. *J. Comp. Neurol.* 315, 98–116.
- 966 May, P.J., Warren, S., Gamlin, P.D., Billig, I., 2018. An anatomic characterization of the
967 midbrain near response neurons in the macaque monkey. *Invest Ophthalmol Vis Sci* 59, 1486-
968 1502.
- 969 May, P.J., Billig, I., Gamlin, P.D., Quinet, J., 2019. Central mesencephalic reticular formation
970 control of the near response: lens accommodation circuits. *J. Neurophysiol.* 121, 1692-1703.
- 971 McCrea, R.A., Horn, A.K. Nucleus prepositus. *Prog. Brain Res.* 151, 205-230, 2006.
- 972 McCrea, R.A., Strassman, A., Highstein, S.M., 1987a. Anatomical and physiological
973 characteristics of vestibular neurons mediating the vertical vestibulo-ocular reflexes of the
974 squirrel monkey. *J. Comp. Neurol.* 264, 571–594.
- 975 McCrea, R.A., Strassman, A., May, E., Highstein, S.M., 1987b. Anatomical and physiological
976 characteristics of vestibular neurons mediating the horizontal vestibulo-ocular reflex of the
977 squirrel monkey. *J. Comp. Neurol.* 264, 547–570.
- 978 McDougal, D.H., Gamlin, P.D., 2015. Autonomic control of the eye. *Comprehens. Physiol.* 5,
979 439-473.
- 980 McFarland, J.L., Fuchs, A.F., 1992. Discharge patterns in nucleus prepositus hypoglossi and
981 adjacent medial vestibular nucleus during horizontal eye movement in behaving macaques. *J.*
982 *Neurophysiol.* 68, 319–332.
- 983 Mays, L.E., 1984. Neural control of vergence eye movements: convergence and divergence
984 neurons in midbrain. *J. Neurophysiol.* 51, 1091–1108.

- 985 Mays, L.E., Porter, J.D., Gamlin, P.D., Tello, C.A., 1986. Neural control of vergence eye
986 movements: neurons encoding vergence velocity. *J. Neurophysiol.* 56, 1007-1021.
- 987 Mays, L.E., Gamlin, P.D., 1995. A neural mechanism subserving saccade-vergence interactions.
988 In: *Eye movement research: Mechanisms, processes and applications*, edited by Findlay J,
989 Walker R, and Kentridge RW. Amsterdam, The Netherlands: Elsevier, p. 215–223.
- 990 Mays L.E., Gamlin, P.D., 1995. Neuronal circuitry controlling the near response. *Curr. Opin.*
991 *Neurobiol.* 5, 763-768.
- 992 Miller, J.M., Robins, D., 1992. Extraocular muscle forces in alert monkey. *Vision Res.* 32, 1099–
993 1113.
- 994 Miller, J.M., Bockisch, C.J., Pavlovski, D.S., 2002. Missing lateral rectus force and absence of
995 medial rectus co-contraction in ocular convergence. *J. Neurophysiol.* 87, 2421-2433.
- 996 Moschovakis, A.K., 1995. Are laws that govern behavior embedded in the structure of the
997 CNS? The case of Hering's law. *Vision Res.* 35, 3207-3216.
- 998 Moschovakis, A.K., 1997. The neural integrators of the mammalian saccadic system. *Front*
999 *Biosci* 2, d552-577.
- 1000 Moschovakis, A.K., Kitama, T., Dalezios, Y., Petit, J., Brandi, A.M., Grantyn, A.A., 1998. An
1001 anatomical substrate for the spatiotemporal transformation. *J. Neurosci.* 18, 10219-10229.
- 1002 Moschovakis, A.K., Scudder, C.A., Highstein, S.M. 1996. The microscopic anatomy and
1003 physiology of the mammalian saccadic system. *Prog. Neurobiol.* 50, 133–254.
- 1004 Moschovakis, A.K., Scudder, C.A., Highstein, S.M. 1991a. The structure of the primate
1005 oculomotor burst generator. I. Medium-lead burst neurons with upward on-directions. *J.*
1006 *Neurophysiol.* 65, 203-217.
- 1007 Moschovakis, A.K., Scudder, C.A., Highstein, S.M., Warren J.D.M., 1991b. The structure of the
1008 primate oculomotor burst generator. II. Medium-lead burst neurons with downward on-
1009 directions. *J. Neurophysiol.* 65, 218-229.
- 1010 Mustari, M.J., Fuchs, A.F., 1990. Discharge patterns of neurons in the pretectal nucleus of the
1011 optic tract (NOT) in the behaving macaque. *J. Neurophysiol.* 64, 77-90.
- 1012 Mustari, M.J., Fuchs, A.F., Wallman, J., 1988. Response properties of dorsolateral pontine units
1013 during smooth pursuit in the rhesus macaque. *J. Neurophysiol.* 60, 664-686.
- 1014 Mustari, M.J., Fuchs, A.F., Kaneko, C.R., Robinson, F.R., 1994. Anatomical connections of the
1015 primate pretectal nucleus of the optic tract. *J. Comp. Neurol.* 349, 111-128.
- 1016 Mustari, M.J., Ono, S., Das, V.E., 2009. Signal processing and distribution in cortical-brainstem
1017 pathways for smooth pursuit eye movements. *Ann. N. Y. Acad. Sci.* 1164, 147-154.
- 1018 Noda, H., Mikami, A., 1986. Discharges of neurons in the dorsal paraflocculus of monkeys
1019 during eye movements and visual stimulation. *J. Neurophysiol.* 56, 1129–1146.

- 1020 Noda, H., Suzuki, D.A., 1979. The role of the flocculus of the monkey in fixation and smooth
1021 pursuit eye movements. *J. Physiol.* 294, 335–348.
- 1022 Noda, H., Sugita, S., Ikeda, Y., 1990. Afferent and efferent connections of the oculomotor
1023 region of the fastigial nucleus in the macaque monkey. *J. Comp. Neurol.* 302, 330–348.
- 1024 Ohgaki, T., Markham, C.H., Schneider, J.S., Curthoys, I.S., 1989. Anatomical evidence of the
1025 projection of pontine omnipause neurons to midbrain regions controlling vertical eye
1026 movements. *J. Comp. Neurol.* 289, 610-625.
- 1027 Ono, S., Das, V.E., Mustari, M.J., 2003. Role of the dorsolateral pontine nucleus in short-term
1028 adaptation of the horizontal vestibuloocular reflex. *J. Neurophysiol.* 89, 2879-2885.
- 1029 Optican, L.M., 2009. Oculomotor system: models. In: *Encyclopedia of Neuroscience*, edited by
1030 Squire, L.R. Oxford, UK: Academic.
- 1031 Ostriker, G., Pellionisz, A., Llinás, R., 1985. Tensorial computer model of gaze--I. Oculomotor
1032 activity is expressed in non-orthogonal natural coordinates. *Neuroscience* 14, 483-500.
- 1033 Phillips, J.O., Ling, I., Siebold, C., Fuchs, A.F., 1996. Behavior of primate vestibulo-ocular reflex
1034 neurons and vestibular neurons during head-free gaze shifts. *Ann. N. Y. Acad. Sci.* 781, 276-
1035 291.
- 1036 Phillips, J.O., Ling, I., Fuchs, A.F., 1999. Action of the brain stem saccade generator during
1037 horizontal gaze shifts. I. Discharge patterns of omnidirectional pause neurons. *J. Neurophysiol.*
1038 81, 1284-1295.
- 1039 Pola, J., Wyatt, H.J., 1985. Active and passive smooth eye movements: effects of stimulus size
1040 and location. *Vision Res.* 25, 1063-1076.
- 1041 Porter, J. D., Guthrie, B. L., Sparks, D. L., 1983. Innervation of monkey extraocular muscles:
1042 localization of sensory and motor neurons by retrograde transport of horseradish peroxidase.
1043 *J. Comp. Neurol.* 218, 208-219.
- 1044 Quinet, J., Goffart, L., 2007. Head-unrestrained gaze shifts after muscimol injection in the
1045 caudal fastigial nucleus of the monkey. *J. Neurophysiol.* 98, 3269-3283.
- 1046 Quinet, J., Schultz, K, May, P.J., Gamlin, P.D., 2020. Neural control of rapid binocular eye
1047 movements: Saccade-vergence burst neurons. *Proc. Natl. Acad. Sci. USA* 117, 29123-29132.
- 1048 Raphan, T., Cohen, B., 1978. Brainstem mechanisms for rapid and slow eye movements. *Annu.*
1049 *Rev. Physiol.* 40, 527-552.
- 1050 Richmond, F.J., Singh, K., Corneil, B.D., 2001. Neck muscles in the rhesus monkey. I. Muscle
1051 morphometry and histochemistry. *J. Neurophysiol.* 86, 1717-1728.
- 1052 Roy, J.E., Cullen, K.E., 2002. Vestibuloocular reflex signal modulation during voluntary and
1053 passive head movements. *J. Neurophysiol.* 87, 2337-2357.
- 1054 Roy JE, Cullen, K.E., 2003. Brain stem pursuit pathways: dissociating visual, vestibular, and
1055 proprioceptive inputs during combined eye-head gaze tracking. *J. Neurophysiol.* 90, 271-290.

- 1056 Robinson, D.A., 1975. A quantitative analysis of extraocular muscle cooperation and squint.
1057 Invest. Ophthalmol. 14, 801–825.
- 1058 Robinson, D.A., 1970. Oculomotor unit behavior in the monkey. J. Neurophysiol. 33, 393–403,
1059 1970.
- 1060 Roy, J.E., Cullen, K.E., 2003. Brain stem pursuit pathways: dissociating visual, vestibular, and
1061 proprioceptive inputs during combined eye-head gaze tracking. J. Neurophysiol. 90, 271-290.
- 1062 Rucker, J.C., Lavin, P.J., 2021. Neuro-Ophthalmology: Ocular Motor System. Bradley's
1063 Neurology in Clinical Practice E-Book, 201.
- 1064 Schultheis, L.W., Robinson, D.A., 1981. Directional plasticity of the vestibulo-ocular reflex in
1065 the cat. Annals N. Y. Acad. Sci. 374, 504-512.
- 1066 Scudder, C.A., Fuchs, A.F., 1992. Physiological and behavioral identification of vestibular
1067 nucleus neurons mediating the horizontal vestibuloocular reflex in trained rhesus monkeys. J.
1068 Neurophysiol. 68, 244–264.
- 1069 Scudder, C.A., Fuchs, A.F., Langer, T.P., 1988. Characteristics and functional identification of
1070 saccadic inhibitory burst neurons in the alert monkey. J. Neurophysiol. 59, 1430–1454.
- 1071 Scudder, C.A., Kaneko, C.R.S., Fuchs, A.F., 2002. The brainstem burst generator for saccadic
1072 eye movements: a modern synthesis. Exp. Brain Res. 142, 439–462.
- 1073 Sharpe, J.A., 2008. Neurophysiology and neuroanatomy of smooth pursuit: lesion studies.
1074 Brain Cogn 68, 241–254.
- 1075 Sharpe, J., Wong, A.M., 2005. Anatomy and physiology of ocular motor systems. Walsh and
1076 Hoyt's Clinical Neuro-Ophthalmology 16, 809-885.
- 1077 Soetedjo, R., Kaneko, C.R., Fuchs, A.F., 2002. Evidence that the superior colliculus participates
1078 in the feedback control of saccadic eye movements. J. Neurophysiol. 87, 679–695.
- 1079 Soetedjo, R., Kojima, Y., Fuchs, A.F., 2019. How cerebellar motor learning keeps saccades
1080 accurate. J. Neurophysiol. 121, 2153-2162.
- 1081 Sparks, D.L., 2002. The brainstem control of saccadic eye movements. Nat. Rev. Neurosci. 3,
1082 952–964.
- 1083 Sparks, D.L., Barton, E.J., Gandhi, N.J., Nelson J., 2002. Studies of the role of the paramedian
1084 pontine reticular formation in the control of head restrained and head-unrestrained gaze
1085 shifts. Ann. N.Y. Acad Sci 956, 85–98.
- 1086 Sparks, D.L., Freedman, E.G., Chen, L.L., Gandhi, N.J., 2001. Cortical and subcortical
1087 contributions to coordinated eye and head movements. Vision Res. 41, 3295-3305.
- 1088 Sparks, D.L., Gandhi, N.J., 2003. Single cell signals: an oculomotor perspective. Prog. Brain Res.
1089 142, 35-53.
- 1090 Sparks, D.L., Holland, R., Guthrie, B.L., 1976. Size and distribution of movement fields in the
1091 monkey superior colliculus. Brain Res. 113, 21-34.

- 1092 Sparks, D.L., Mays L.E., 1990. Signal transformations required for the generation of saccadic
1093 eye movements. *Annu. Rev. Neurosci.* 13, 309–336.
- 1094 Sparks, D.L., Lee, C., & Rohrer, W. H., 1990. Population coding of the direction, amplitude, and
1095 velocity of saccadic eye movements by neurons in the superior colliculus. *Cold Spring Harb.*
1096 *Symp. Quant. Biol.* 55, 805-811.
- 1097 Strassman, A., Evinger C, McCrea, R.A., Baker RG, Highstein, S.M., 1987. Anatomy and
1098 physiology of intracellularly labelled omnipause neurons in the cat and squirrel monkey. *Exp.*
1099 *Brain Res.* 67, 436-440.
- 1100 Strassman, A., Highstein, S.M., McCrea, R.A., 1986a. Anatomy and physiology of saccadic burst
1101 neurons in the alert squirrel monkey. I. Excitatory burst neurons. *J. Comp. Neurol.* 249, 337–
1102 357.
- 1103 Strassman, A., Highstein, S.M., McCrea, R.A., 1986b. Anatomy and physiology of saccadic burst
1104 neurons in the alert squirrel monkey. II. Inhibitory burst neurons. *J. Comp. Neurol.* 249, 358–
1105 380.
- 1106 Straumann, D, Zee, D.S., Solomon, D., Lasker, A.G., Roberts, D.C., 1995. Transient torsion
1107 during and after saccades. *Vision Res.* 35, 3321-3334.
- 1108 Strohmaier, A.R., MacKay, K.J., Obersteiner, A., Reiss, K.M., 2020. Eye-tracking methodology
1109 in mathematics education research: A systematic literature review. *Educ. Stud. Math.* 104,
1110 147-200.
- 1111 Suzuki, D.A., May, J.G., Keller, E.L., Yee, R.D., 1990. Visual motion response properties of
1112 neurons in dorsolateral pontine nucleus of alert monkey. *J. Neurophysiol.* 63, 37-59.
- 1113 Suzuki, D.A., Yamada, T., Yee, R.D., 2003. Smooth-pursuit eye-movement-related neuronal
1114 activity in macaque nucleus reticularis tegmenti pontis. *J. Neurophysiol.* 89, 2146-2158.
- 1115 Sylvestre, P.A., Cullen, K.E., 1999. Quantitative analysis of abducens neuron discharge
1116 dynamics during saccadic and slow eye movements. *J. Neurophysiol.* 82, 2612–2632.
- 1117 Tanaka, M., 2005. Involvement of the central thalamus in the control of smooth pursuit eye
1118 movements. *J. Neurosci.* 25, 5866-5876.
- 1119 Takahashi, M., Shinoda, Y., 2018. Brain stem neural circuits of horizontal and vertical saccade
1120 systems and their frame of reference. *Neuroscience* 392, 281-328.
- 1121 Takahashi, M., Sugiuchi, Y., Na, J., Shinoda, Y., 2022. Brainstem circuits triggering saccades and
1122 fixation. *J. Neurosci.* 42, 789-803.
- 1123 Tweed, D., Glenn, B., Vilis, T., 1995. Eye-head coordination during large gaze shifts *J.*
1124 *Neurophysiol.* 73, 766-779.
- 1125 Ugolini, G., Klam, F., Doldan Dans, M., Dubayle, D., Brandi, A.M., Büttner-Ennever, J., Graf, W.,
1126 2006. Horizontal eye movement networks in primates as revealed by retrograde
1127 transneuronal transfer of rabies virus: differences in monosynaptic input to "slow" and "fast"
1128 abducens motoneurons. *J. Comp. Neurol.* 498, 762-785.

- 1129 Van Gisbergen, J.A.M., Robinson, D.A., Gielen, S. A., 1981. Quantitative analysis of generation
1130 of saccadic eye movements by burst neurons. *J. Neurophysiol.* 45, 417–442.
- 1131 Van Gisbergen, J.A., Van Opstal, A.J., 1989. The neurobiology of saccadic eye movements.
1132 *Models. Rev. Oculomot. Res.* 3, 69-101.
- 1133 Van Horn MR, Waitzman DM, Cullen, K.E., 2013. Vergence neurons identified in the rostral
1134 superior colliculus code smooth eye movements in 3D space. *J. Neurosci.* 33, 7274-7284.
- 1135 Wong, A. M., 2004. Listing's law: clinical significance and implications for neural control.
1136 *Survey Ophthalmol.* 49, 563-575.
- 1137 Yakushin, S.B., Gizzi, M., Reisine, H., Raphan, T., Büttner-Ennever, J., Cohen, B., 2000.
1138 Functions of the nucleus of the optic tract (NOT). II. Control of ocular pursuit. *Exp. Brain Res.*
1139 131, 433-447.
- 1140 Zhang H., Gamlin P.D., 1996. Single-unit activity within the posterior fastigial nucleus during
1141 vergence and accommodation in the alert primate. *Soc. Neurosci. Abstr.* 22: 2034.
- 1142 Zee, D.S., Yamazaki, A., Butler, P.H., Gücer, G., 1981. Effects of ablation of flocculus and
1143 paraflocculus of eye movements in primate. *J. Neurophysiol.* 46, 878–899.
- 1144

1145 **Acknowledgements**

1146

1147 The author is grateful to Drs Anja Horn, Richard J. Krauzlis, Paul J. May, Adonis Moschovakis,
1148 David L. Sparks and Catherine Vignal-Clermont for their corrections and suggestions to
1149 improve this article.

1150 This work was possible thanks to support from Centre National de la Recherche Scientifique
1151 and from the Fondation pour la Recherche Médicale.

1152

1153 Figure captions

1154

1155 **Figure 1:** Schematic representation of the extraocular muscles (A) and pulling directions
1156 engaged by the contraction of their fibers when the eyes are centered in the orbit (B). LR:
1157 lateral rectus, MR: medial rectus, SR: superior rectus, IR: inferior rectus, SO: superior oblique,
1158 IO: inferior oblique, tr: trochlea. The red arrows indicate the muscles whose contraction
1159 abducts the eye, the black arrows the muscles whose contraction adducts the eye. The
1160 turquoise arrows indicate the muscles whose contraction elevates the eye (supraduction), the
1161 yellow arrows those whose contraction infraducts the eye. The blue arrows indicate the
1162 muscles whose contraction causes an incyclotorsion (incycloduction) of the eye, the green
1163 arrows those whose contraction causes an excyclotorsion (excycloduction).

1164

1165 **Figure 2:** Neuronal network involved during horizontal saccades toward the left (A) and the
1166 right (B). Connecting lines ended by an arrow indicate excitatory connections; those ended by
1167 a circle indicate inhibitory synaptic connections. Blue color indicates the agonist neuronal
1168 elements, red color the antagonist ones. The thickness of connecting lines schematizes the
1169 strength with which the neurons fire. OPN: omnipause neurons, EBN: excitatory burst
1170 neurons, IBN: inhibitory burst neurons, MN: abducens motoneurons, AIN: abducens
1171 internuclear neurons. In the bottom, the thickness of arrows attached to the eyeballs
1172 schematizes the strength of muscle contraction. LR: lateral rectus, MR: medial rectus, SR:
1173 superior rectus, IR: inferior rectus, SO: superior oblique, IO: inferior oblique.

1174

1175 **Figure 3:** Neuronal network involved during vertical saccades toward the top (A) and the
1176 bottom (B). Connecting lines ended by an arrow indicate excitatory connections; those ended
1177 by a circle indicate inhibitory synaptic connections. Blue color indicates the agonist neuronal
1178 elements, red color the antagonist ones. The thickness of connecting lines schematizes the
1179 strength with which the neurons fire. uEBN: upward excitatory burst neurons, dEBN:
1180 downward excitatory burst neurons, uIBN: upward inhibitory burst neurons, MN:
1181 motoneurons. In the top, the thickness of arrows attached to the eyeballs schematizes the
1182 strength of muscle contraction. LR: lateral rectus, MR: medial rectus, SR: superior rectus, IR:
1183 inferior rectus, SO: superior oblique, IO: inferior oblique.

1184

1185 **Figure 4:** Neuronal network involved during combined horizontal eye and head movements
1186 toward the left. Connecting lines ended by an arrow indicate excitatory connections; those
1187 ended by a circle indicate inhibitory synaptic connections. Blue color indicates the agonist
1188 neuronal oculomotor elements, red color the antagonist ones. Green color indicates the
1189 cephalomotor elements. The thickness of connecting lines schematizes the strength with
1190 which the neurons fire. dSC: deep superior colliculus, RSN: reticulospinal neurons, EBN:
1191 excitatory burst neurons, IBN: inhibitory burst neurons, ABD: abducens motor and
1192 internuclear neurons, NMN: neck motoneurons, PVN: primary vestibular neurons, IVN:
1193 inhibitory vestibular neurons, EVN: excitatory vestibular neurons, VIII: eight cranial nerve.

1194

1195 **Figure 5:** Neuronal network involved in ending a leftward gaze shift while the head terminates
1196 its rotation (stabilized gaze direction). Connecting lines ended by an arrow indicate excitatory
1197 connections; those ended by a circle indicate inhibitory synaptic connections. Blue color
1198 indicates the agonist neuronal oculomotor elements, red color the antagonist ones. Green
1199 color indicates the neurons of the ascending tract of Deiters (ATDN). The thickness of
1200 connecting lines schematizes the strength with which the neurons fire. LR: lateral rectus, MR:
1201 medial rectus, VI, abducens nerve, MRMN: motor neurons innervating the medial rectus
1202 muscle, LRMN: motor neurons innervating the lateral rectus muscle, AIN: abducens
1203 internuclear neurons, NMN: neck motoneurons, PVN: primary vestibular neurons, IVN:
1204 inhibitory vestibular neurons, EVN: excitatory vestibular neurons, VIII: eight cranial nerve.

1205

1206 **Figure 6:** Neuronal network involved in ending a downward (A) and upward (B) gaze shift while
1207 the head terminates its movement (stabilized gaze direction). Connecting lines ended by an
1208 arrow indicate excitatory connections; those ended by a circle indicate inhibitory synaptic
1209 connections. Blue color indicates the agonist neuronal oculomotor elements, red color the
1210 antagonist ones. IOMN: motor neurons innervating the inferior oblique muscle, IRMN: motor
1211 neurons innervating the inferior rectus muscle, SRMN: motor neurons innervating the
1212 superior rectus muscle, SOMN: motor neurons innervating the superior oblique muscle, PVN:
1213 primary vestibular neurons, IVN: inhibitory vestibular neurons, EVN: excitatory vestibular
1214 neurons, VIII: eight cranial nerve.

1215

1216 **Figure 7:** Neuronal network involved in the near response. LR: lateral rectus, MR: medial
1217 rectus, SP: sphincter pupillae, DP: dilator pupillae, CM: ciliary muscle, CG: ciliary ganglion,
1218 MRMN: motor neurons innervating the medial rectus muscle, EWpg: Edinger-Westphal
1219 parasympathetic nucleus, cMRF: central mesencephalic reticular formation, SOA:
1220 supraoculomotor area, cFN: caudal fastigial nucleus, SC: superior colliculus.

1221

1222 **Figure 8:** Neuronal network involved in the generation of horizontal slow pursuit eye
1223 movements. LR: lateral rectus, MR: medial rectus, MN: motor neurons, AIN: abducens
1224 internuclear neurons, NHP/MV: complex nucleus prepositus hypoglossi-medial vestibular
1225 nucleus, ETN: excitatory tonic neurons, ITN: inhibitory tonic neurons, cFN: caudal fastigial
1226 nucleus, NOT: nucleus of the optic tract, rSC: rostral superior colliculus, rNRTP: rostral part of
1227 nucleus reticularis tegmenti pontis, DMPN: dorsomedian pontine nucleus, DLPN: dorsolateral
1228 pontine nucleus, VPL-Th: ventroposterolateral part of thalamus, FEF: frontal eye field, MST:
1229 medial superior temporal area of cerebral cortex.

Figure 1





Figure 2

A

LEFTWARD



B

RIGHTWARD




Figure 3




Figure 4




Figure 5





Figure 6

A



B




Figure 7




Figure 8

