



HAL
open science

Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques

Igor Lima Maldonado, Christophe Destrieux

► To cite this version:

Igor Lima Maldonado, Christophe Destrieux. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques. *LE STUDIUM Multidisciplinary Journal*, 2021, pp.64-88. <10.34846/le-studium.170.02.fr.01-2019>. <hal-03543714>

HAL Id: hal-03543714

<https://hal.science/hal-03543714v1>

Submitted on 9 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques

Igor Lima Maldonado^{1,5}, Christophe Destrieux^{1,3,6}.

¹UMR Inserm U1253, iBrain, Université de Tours, Tours, France

²Le Studium Loire Valley Institute for Advanced Studies, Orleans, France

³CHRU de Tours, Tours, France

⁴Departamento de Biomorfologia - Instituto de Ciências da Saúde, Universidade Federal da Bahia, Salvador, Brasil

⁵Programa de Pós-Graduação em Medicina e Saúde, Universidade Federal da Bahia, Salvador, Brasil

⁶Laboratory of Anatomy - Faculté de Médecine - 10 bd Tonnellé - 37032 Tours – France

REPORT INFO

Fellow: **Igor Lima Maldonado**

From: Brazil

Host laboratory in region Centre-Val de Loire: UMR Inserm U1253, iBrain, Université de Tours, Tours, France

Host scientist: **Christophe Destrieux**

Period of residence in region Centre-Val de Loire: January 2018 to December 2019.

Keywords: *limbic lobe, cingulum, cingulate gyrus, white matter, diffusion imaging, magnetic resonance, fiber dissection*

ABSTRACT

Although cortical elements of the limbic system are quite well defined, the underlying white matter pathways are not well detailed. This project aimed to better describe white matter tracts of the limbic system using both fiber dissection, and in and ex vivo Diffusion Weighted Imaging (DWI) techniques. In vivo data (from DWI and neuropsychological evaluations) were obtained from healthy subjects aged 82 and over previously enrolled in the FIBRATLAS project funded by the French Research Agency and run in the host laboratory. Correlations between neuropsychological and white matter characteristics on DWI were examined in this group to infer function of the limbic white matter tracts. Ex vivo data from the FIBRATLAS project database and from the Tours Body Donation Program were examined. The brains were extracted and used to study the anatomy of the limbic white matter tracts employing both DWI and fiber dissection according to a variant of the Klingler's method.

1- Introduction

In the nineteenth century, Paul Broca proposed the term “*great limbic lobe*” for a ring of convolutions around the corpus callosum in two seminal papers published in *Bulletins de la Société d'Anthropologie de Paris* and *Revue d'Anthropologie* (Broca, 1877, 1878). While the remaining cortex was regarded as “intellectual”, the limbic lobe was considered the “brutal” part of the brain. He thus introduced a dichotomy between the limbic lobe and surrounding cortex. Such a perspective, which relegated the limbic lobe to relatively primitive tasks, became very popular. That was the first attempt to group structures that are presently recognized as the **limbic system**. The identity of the limbic system is, however, a concept in evolution that continues to provoke

debate. There is, for instance, discussion surrounding which are the exact structures that should be considered as limbic or extra-limbic, given the richness of their connections (Olry & Nicolay, 1994).

The **cingulum** has regularly appeared as the main communication element of the limbic system. It was one of the central structures of the circuit described by Papez in 1937 as the “seat of dynamic vigilance by which environmental experiences are endowed with an emotional awareness” (Papez, 1937). We have however witnessed a very important development of related research over the last decades that has been at the origin of multiple updates and is pleading for a revolution in the way we think about the cingulum. This fasciculus crosses large distances within the frontal, parietal, occipital, and temporal lobes. Beyond

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

being an anatomical substrate of limbic connections, it is a major route for association fibers from cortical regions with distinct functional roles. This subject is of great importance for modeling of higher functions, with applications in cognitive neuroscience, neuropsychology, and a significant impact on how we interpret clinical manifestations of white matter diseases.

Studies on connectivity multiplied greatly in the 20th century. This was mainly related to methodological advances. In Switzerland, Joseph Klingler developed an anatomical method associating formaldehyde fixation with one or more freezing and defrosting cycles (Klingler, 1935; Klingler & Gloor, 1960; Ludwig & Klingler, 1956). He noticed that, curiously, freezing was able to change the texture of tissues and slightly separate structures. This phenomenon enabled the development of a method of **fiber dissection** and identification of white matter fasciculi that would be imperceptible in conventional preparations (Alarcon, et al., 2014; Arnts, Kleinnijenhuis, Kooloos, Schepens-Franke, & van Cappellen van Walsum, 2014; Bozkurt, et al., 2017; Costa, et al., 2018; Koutsarnakis, Liakos, Kalyvas, Sakas, & Stranjalis, 2015; I. Maldonado, E. Mandonnet, & H. Duffau, 2012; Maldonado, et al., 2013; I. L. Maldonado, E. Mandonnet, & H. Duffau, 2012; Mandonnet, et al., 2017; Martino, et al., 2013; Pascalau, Popa Stanila, Sfrangeu, & Szabo, 2018; Shah, Jhawar, & Goel, 2012; Zemmoura, Vons, Velut, & Destrieux, 2015). In animals, tract-tracing techniques developed intensively, and autoradiography became notable for its accuracy and high level of detail. Based on the axonal transport of injected radioactively labeled amino acids, this method required animal sacrifice and was impossible to apply to humans (Mufson & Pandya, 1984; Petrides & Pandya, 1984; Woolsey, 2016). In such a context of methodological challenge and absence of a research strategy that would be both effective *in vivo* and practicable in humans, significant effort came from magnetic resonance imaging (MRI) research. Magnetic resonance based **diffusion imaging** is the primary group of techniques used to infer tissue architecture from how it constrains diffusion of water molecules (Basser, Mattiello, & LeBihan, 1994a, 1994b; Conturo, et al., 1999; D. Le Bihan, et al., 1986). This is currently the most frequently used category of non-invasive methods to study the three-dimensional organization of the fasciculi that compose the cerebral white matter in both

clinical and research contexts (Basser, Pajevic, Pierpaoli, Duda, & Aldroubi, 2000; Catani & Thiebaut de Schotten, 2008; Denis Le Bihan, et al., 2001; Mukherjee, Berman, Chung, Hess, & Henry, 2008).

Examined together, data from the aforementioned techniques and an explosion in the number of publications on cerebral connectivity made opportune and necessary to rethink the main role of the cingulum in the human brain. In our works, we have been exposed to morphological and functional arguments that highlight a tangible shift in perspective. In this research project we studied the morphology of **white matter tracts** using fiber dissection, *in* and *ex vivo* Diffusion Weighted Imaging (DWI), with a special focus on the limbic system.

2- Experimental details

To carry out the project, the following steps were developed (results were presented in the same order):

- 1) A **literature review**, describing the state of the art and the up-to-date anatomy of limbic connectivity, especially that of the cingulum, the main fasciculus of the limbic system in human and non-human primates;
- 2) A **fiber dissection** study from the medial, ventral and lateral aspects of the human cerebral hemisphere, with special attention to the medial aspect and the limbic system. The dissected hemispheres were also subjected to MRI scanning;
- 3) A study in diffusion tensor imaging (DTI) through ***in vivo* DWI** of the connections ensured by this fasciculus, whether they are completely inside the limbic system or interconnecting extra-limbic areas using the cingulum as a route;
- 4) Compilation of the morphological results of the *in vivo* MRI with those of the **neuropsychological assessment**, checking for associations between white matter characteristics and the performance profiles;
- 5) ***Ex-vivo* DWI** scanning in high magnetic field (7T, 11.7T) of human cerebral hemispheres;
- 6) A theoretical and epistemological study of the role of the cingulum in humans for **synthetization** of the available data from resonance and fiber dissection studies, taking into account its limbic and extra-limbic connectivity, as recently evidenced.

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

2-1 Volunteers

In vivo and part of *ex vivo* data were collected in the frame of the FIBRATLAS project (<https://sites.google.com/site/fibratlas/>) funded by the French Research Agency and ran by the host laboratory. FIBRATLAS aims at producing a database containing *in vivo* (DWI MRI and neuropsychological testing) and *ex vivo* data (DWI and dissection) obtained for the same subjects involved in the body donation program of one of the following university body donation programs: Tours, Angers, Amiens, Brest, Lyon, Nancy. Part of this database will be available to the community for validation of tractography algorithms.

The volunteers, previously enrolled in the body donation program and aged 82 and over, were scanned *in vivo* for anatomical three-dimensional T1-weighted and DWI and underwent an extensive neuropsychological examination. The parameters of acquisition for *in vivo* MRI examination were as follows:

- B0 cartography - FOV 256x256; matrix 64x64; 4mm joined slices; TR 800ms; TE 4.92/7.38ms; NEX=1;
- 3D T1 anatomic images - matrix=256x256; voxel size=1x1x1mm ; TR 2300ms; TE 2.98ms; TI 900ms; NEX=1 ; angle 9°; phase direction A/P;
- DWI - FOV=256x256; voxel size : 1.7x1.7x1.7 mm]: 21 directions with b=300 mm/s², 25 directions with b=1200 mm/s², and 66 directions with b=1200 mm/s².

For DTI tractography, the tensor was computed in each voxel using fiber assignment by continuous tracking. Tracts were defined by regions of interest located according to published atlases. For image post-processing, the Trackvis (<http://www.trackvis.org/>) and Connectomist (CEA Neurospin, Saclay, France) packages were used.

2-2 Neuropsychological assessment

The level of education of each participant was systematically recorded and several psychometric fields were explored using standardized tests as follows.

- Manual dominance - Edinburgh test (Oldfield, 1971);
- Global evaluation – Mini Mental State Examination (Folstein et al., 1975);

- Language - DO80 (Deloche and Hannequin, 1997);
- Memory - RL/RI 16 test for verbal (Van der Linder et al, 2004) and DMS4 48 for visual memory (Barbeau et al, 2004);
- Visuo-spatial abilities: Rey's figure (copy) (Rey, 1959; Fastenau et al., 1999);
- Executive functions: verbal fluency (semantic and orthographic) (Godefroy and GREFEX, 2008), Stroop test (Grefex) (Stroop, 1935; Godefroy et al., 2008), Trail Making test (Reitan, 1955; Godefroy et al., 2008; Azouvi et al., 2017), Direct and indirect digit span (Wechsler, 2000);
- Attention and speed processing: Digit Symbol (Wechsler, 2011), TMTA (Godefroy and GREFEX 2008);
- Praxis: Mahieux-Laurent's battery (Mahieux-Laurent et al., 2009);
- Depressive symptoms: MADRS (Montgomery and Asberg, 1979).

For each of the studied tract, a mean value for each scalar parameter obtained from DWI acquisition was computed. The results of neuropsychological examinations were then examined together with those obtained mean values. Any potential association between tract morphology and neuropsychological performance was detailed and discussed.

2-3 Ex-vivo data

Ex vivo specimens were collected from two different sources: participants of the FIBRATLAS project, and extra volunteers of the Body Donation Program of the University of Tours School of Medicine. The brain was first extracted from the skull and fixed by immersion in a 5% formaldehyde solution in water for at least two to three months. Each body was treated in the first 36 hours after death. Both common carotid arteries were catheterized with the catheter directed upwards. One litter of a 10% formaldehyde buffered solution was injected in each carotid artery during 15 minutes thanks to a peristaltic pump. The day after, a circular craniotomy was performed; the encephalon was extracted after sectioning the falx, tentorium and medullary spinal junction, and suspended by the basilar artery in a 10 % buffered formalin solution for further fixation. Up to the end of the fellowship, a total of 12 hemispheres were scanned for post mortem DWI: 1 at

11.7T, 9 at 7T, and 2 at 3T with different protocols. The 11.7T scanner allowed high resolution imaging with voxels as small as 100µm, isotopically.

For fiber dissection, an adaption of the method described by Ludwig and Klingler was used. After freezing of the specimens for two weeks and defrosting, dissection of the white matter fibers was carried out using fine wooden spatulas. The grey substance, friable, was then gradually separated from the firmer white matter. Identifiable fibers revealed by ablation of the grey matter and neighboring structures were tracked as long as possible towards their cortical or subcortical origins and/or terminations. A surgical microscope (Leica OH4, Leica Microsystems, Wetzlar, Germany) was used as well as additional microsurgical instruments when the naked eye and the mono or binocular loupes were no longer sufficient.

2-4 Ethic issues

This work respected all Ethical Principles for Medical Research Involving Human Subjects stated in the Helsinki declaration. In addition, the authors warrant that was developed in respect to the current rules of good practices for research in all participating countries. It is in conformity with the Clinical Trial Regulation of the European Union number 536/2014 as well as the resolution number 466/2012 from the Brazilian Council of Health. The donors who gave their body to science signed a will by which they allow any research on their body after death. In addition, the FIBRATLAS project, from which part of the data was obtained, was approved by the ethical committee of Tours (Comité de protection des Personnes, 2015-R8) and by the ANSM (*Agence Nationale de Sûreté du Médicament et des Produits de Santé*, EudraCT/ID RCB: 2015-A00363-46). The confidentiality of the identity of all subjects that have donated their bodies to science were and will continue to be ensured in all stages, including in scientific communications and future publications.

3- Results and discussion

3-1 Results of the comparative connectivity review

For this part of the work, a total of **80 publications** was examined. In monkeys and chimpan-

zees, the structure of the white matter that composes the cingulum is complex and heterogeneous (Vogt *et al.* 1992). The bundle is organized as an associative complex, even though it has the appearance of a single compact fasciculus in gross anatomy. Charles Beever (1891) offered one of the first descriptions of the cingulum anatomy in the nineteenth century. He noted that in the Marmoset monkey, the bundle almost does not vary in diameter throughout its course, because of the continuous addition and exiting of fibers (Beever 1891; Jeremy D. Schmahmann 2006).

The **Rhesus monkey** has been the most frequent non-human primate used in investigations of cerebral connectivity historically. The cingulum subdivisions that have been proposed were mostly based on the results of radioisotope tracing and autoradiography, considered as a golden standard technique (Mufson and Pandya 1984; Jeremy D. Schmahmann 2006; Schmahmann *et al.* 2007). Such studies showed that fibers coming from the prefrontal cortex may travel large distances inside the cingulum, and some extend up to the retrosplenial cortex. Those from the parietal cortex could reach the parahippocampal gyrus of the temporal lobe through a different portion of the same bundle (Jeremy D. Schmahmann 2006). A population of small overlapping tracts responsible for regional connections, were also described with consideration of specific functional roles (Mogenson *et al.* 1980; Morgane *et al.* 2005; Catani *et al.* 2013). Consequently, the microstructural characteristics of the cingulum are not homogenous (Yeterian *et al.* 2011; Petrides *et al.* 2012) and its fibers may be grouped according to their anatomical terminations and functionalities. Two main types are considered: those that originate or terminate in the cingulate gyrus itself and those that travel inside the cingulum as a route for distant or adjacent cortical areas (Jeremy D. Schmahmann 2006).

Several authors agree that the cingulum is part of the route of several projection and association bidirectional pathways (Forkel *et al.*; Mufson and Pandya 1984; Petrides *et al.* 2012). Fibers from the anterior thalamic nuclei and the frontal lobe travel inside the fasciculus before reaching the retrosplenial cortex and to the cingulate gyrus itself (area 23) (Yeterian *et al.* 2012). Furthermore, those from the dorsolateral thalamic nuclei and the parietal cortex run in the cingulum before

reaching the presubiculum and the parahippocampal gyrus in the temporal lobe (Forkel *et al.*; Mufson and Pandya 1984; Jankowski *et al.* 2013; Heilbronner and Haber 2014). Other fibers connect post-Rolandic, parietal, and pre-occipital areas with the dorsal and ventrolateral prefrontal cortices (Aggleton *et al.* 2012; Petrides *et al.* 2012).

Association fibers spread to cortical areas at different points. While the rostral ones terminate in the prefrontal cortex, other groups leave the bundle in the frontal, parietal, and temporal lobes (Mufson and Pandya 1984). The longest fibers are afferent and travel along the periphery, and thalamic fibers adopt a more ventral position (Jeremy D. Schmahmann 2006).

Efferent pathways from the anterior cingulate gyrus, including monkey area 24, travel in the cingulum before reaching a large number of destinations in the same cerebral hemisphere: the orbitofrontal, the ventrolateral and dorsolateral prefrontal cortex, as well as the primary motor cortex, infero-rostral portions of the parietal lobe, the amygdala, the perirhinal cortex (Jeremy D. Schmahmann 2006), the extreme capsule, and insula (Petrides *et al.* 2012). Those from area 23 are dorsolateral and stream caudally to the parahippocampus and presubiculum (Mufson and Pandya 1984).

Most evidence from autoradiography suggests a general organization of the cingulum connectivity in two main components. The anterior (ventral) component connects most of the cingulate gyrus, orbitofrontal, septal, sub-callosal, and pericallosal regions (Mufson and Pandya 1984; Schmahmann *et al.* 2007). The posterior (dorsal) component connects the precuneus, presubiculum, parahippocampal gyrus, and hippocampus (Jeremy D. Schmahmann 2006; Aggleton *et al.* 2012).

Publications on findings of fiber dissection became more frequent after the seminal works of Ludwig and Klingler in 1956 (Ludwig and Klingler 1956) but received substantial impulse after the advent of methods for diffusion imaging and peri-operative subcortical stimulation in the last twenty years (Duffau 2016). Fiber dissection allows visualization of macroscopic organization of the fiber bundles and their three-dimensional disposition in the cerebral parenchyma (Fernández-Miranda *et al.* 2008; Insausti *et al.* 2017), goals that would be difficult to achieve using traditional anatomical preparations. Some variants

of the original technique were described, although the main methodological principle remains (i.e. the addition of freezing steps in the fixation protocol). The crystallization of the water and formalin increases volume and slightly separates groups of fibers, rendering them visible and dissectible (Zemmoura *et al.* 2015).

In fiber dissection studies, the cingulum appears as a compact and well-defined bundle travelling in the depth of the cingulate gyrus. After removal of the cingulate cortex, made friable by the preparation, a relatively firm layer of subcortical white matter is identified. This thin layer is roughly amorphous to the naked eye and accompanies the curvature of the medial aspect of the cingulate gyrus (Fernández-Miranda *et al.* 2008). Deeper to that, fibers are easily identifiable, grouped within a compact bundle, with a predominant antero-posterior direction. The lateral aspect of the white matter in the core of the cingulate gyrus is continuous with the medial portion of the centrum semiovale, where the compact subcortical layer is absent (Mollink *et al.* 2017).

The region of the centrum semiovale that is in direct anatomical relationship with the cingulum is rich in fibers from callosal and thalamic radiations (Fernández-Miranda *et al.* 2008). Inferiorly, the trunk of the cingulum is in contact with transversal fibers of the corpus callosum (Lawes *et al.* 2008; Maldonado *et al.* 2012). Lateral-most fibers are then in direct contact with the rostrum, genu, body, isthmus, and splenium without the interposition of cingulate cortex. The anterior extremity of the cingulum sharpens progressively as its fibers leave the main bundle to terminate in the subgenual gray matter and septal areas (Ribas 2010; Chaddad-neto *et al.* 2017).

In its trajectory around the corpus callosum, the anterior cingulum also has indirect anatomical relationships (through the cerebral cortex and the cingulate sulcus) with surrounding cortical areas, namely the superior frontal gyrus, the paracentral lobe, and the precuneus. There is evidence of extensions of the cingulate white matter with well-defined fibers penetrating the superior frontal gyrus (Li *et al.* 2013), the paracentral lobe, and precuneus (Cavanna and Trimble 2006; Wu *et al.* 2016).

Inferiorly and anteriorly to the splenium, cingulate fibers penetrate the parahippocampal gyrus. On their way, they maintain indirect relationships with the posterior column of the fornix (crus fornicis) and with the cistern spaces through the parahippocampal cortex (Concha *et al.* 2005).

The inferior extremity of the cingulum extends anteriorly in the depth of the parahippocampal gyrus and advances toward the hippocampal formation (Jones *et al.* 2013a).

The white matter of the parahippocampal gyrus and precuneus is a continuous structure. Inferiorly to the subparietal sulcus, the precuneus is connected with the cingulate and the parahippocampal gyrus (Ribas 2010). This is related to the presence of parahippocampal-precuneal direct association pathways, as a subcomponent of the cingulum. Similarly, irradiation of cingulate fibers to the upper frontal gyrus and the upper precuneus proposed the cingulum as a path for direct fronto-parietal associative connections (Burzynska *et al.* 2011; Maldonado *et al.* 2012; Ekman *et al.* 2016).

Diffusion imaging techniques have provided indirect information on the anatomy of fiber tracts in the last twenty years with increasing levels of confidence. The representation of the cingulum as a complex connective system, because of the assembling of smaller bundles, is a conception that was recently strengthened by imaging studies. Some authors have proposed systematization of its parts to characterize each of them (Catani and Thiebaut de Schotten 2008; Beckmann *et al.* 2009; Jones *et al.* 2013a; Heilbronner and Haber 2014).

In DTI, deterministic tractography of the cingulum may be obtained by placing regions of interest in specific anatomical sites (Jellison *et al.* 2004; Catani and Thiebaut de Schotten 2008; Jones *et al.* 2013b). In several studies, their locations were based on previous knowledge from autoradiography studies in monkeys. Using this technique, the general appearance of the human cingulum is similar to that observed in fiber dissection studies. The cingulum arches almost 180 degrees around the splenium and continues anteriorly. After traveling inside the cingulate gyrus, above the isthmus and trunk of the corpus callosum, it curves again around the genu. It terminates by spreading its fibers in the subcallosal gyrus and the paraolfactory area (Catani and Thiebaut de Schotten 2008; Sarkar *et al.* 2012). The human cingulum seems to contain **fibers with different lengths and with distinct connectivity profiles**, in the context of a complex population of pathways (Jones *et al.* 2013a). This was detailed through virtual dissection employing simultaneous regions of interest.

The middle portions of the cingulum contain long fibers that travel to the frontal lobe, terminating in the premotor or motor cortices (Gong *et al.* 2005; Xie *et al.* 2009; Bliss *et al.* 2016). The longest fibers seem to originate from the extremities of the bundle, as is the case of those arising from the subgenual cortex, amygdala, uncus, and parahippocampal gyrus (Thiebaut de Schotten *et al.* 2012). Additionally, some long fibers extend inferiorly, providing direct parieto-temporal connections (Jones *et al.* 2013a). Inferior portions of the cingulum may serve as a route for direct transcingular temporo-occipital connections.

Catani *et al.* proposed a systematization based on the anatomic division of the cingulum in two parts: the anterior-dorsal and the posterior-ventral (Catani and Thiebaut de Schotten 2008). The anterior-dorsal is composed of long fibers that follow the shape of the corpus callosum, go forward and around the genu, and continue in the subcallosal part towards the paraolfactory area. The posterior-ventral fibers travel along parahippocampal, retrosplenial and posterior precuneal areas (Catani *et al.* 2013).

Jones *et al.* (2013), using regions of interest based on the a priori anatomy of the Rhesus monkey, described three identifiable subdivisions with specific morphological and physiological features: the subgenual, the retrosplenial, and the parahippocampal (Jones *et al.* 2013a). Moreover, the authors described the parahippocampal subdivision in two different manners: restricted and unrestricted. In restricted reconstructions, the authors isolated transcingular temporo-parietal and temporo-occipital pathways. In unrestricted reconstructions, such fibers were depicted together with the “standard cingulum”. The subgenual subdivision is medial in relation to the others. The retrosplenial one is lateral to the subgenual in the portions in which they overlap. Finally, the parahippocampal subdivision contains the lateralmost fibers of the cingulum complex. Those subdivisions connect extensive areas around the corpus callosum. Few fibers run along the entire path. Indeed, further studies are required to detail the length, terminations, and functional roles of the smaller groups.

Using high-definition fiber tracking and a template from the Human Connectome Project, Wu *et al.* (2016) identified five potential subcomponents. The authors named them as CB I, II, III, IV, and V subcomponents (Wu *et al.* 2016). The CB-I subcomponent is anteromedial and extends from the orbitofrontal cortex to the precuneus,

running along most of the length of the corpus callosum, up to the splenium. The CB-II subcomponent travels from medial portions of the superior frontal gyrus to the retrosplenial, parahippocampal and perirhinal cortices. The CB-III subcomponent is the largest in terms of volume, originates in the medial aspect of the SFG, and ends in the precuneus, connecting many cortical areas. The CB-IV is the shortest, connecting the superior parietal lobe and precuneus with supplementary and premotor areas. It is located over the trunk of the corpus callosum and follows roughly the same trajectory of the dorsal component of the superior longitudinal fascicle (SFL-1). The author subdivided it into two parts: the paracingulate (paraCB) and supracingular (supraCB). Given these observations, the cingulum was suggested to comprise direct dorsal frontoparietal fibers, as it seems to be the case in fiber dissection. Lastly, the CB-V corresponds to the group of parahippocampal fibers that spread in the parietal and occipital lobes. The subdivision in five components also enabled some systematization of the three-dimensional disposition of different groups of fibers. While the CB-I subcomponent runs close to midline adjacent to the corpus callosum, CB-II, III, and IV occupy different portions of the cingulum above the corpus callosum, and most fibers of CB-IV have a predominant superior location in relation to the others. Finally, running exclusively in the temporal and occipital lobes, CB-V is the most inferior and laterally situated subcomponent (Wu *et al.* 2016).

3.2- Results from fiber dissection

3.2.1- Medial and ventral aspect

After midline section of each human cerebral specimen, the initial examination enabled identification the non-cortical structures. Among the main ones were the olfactory tract and bulb, internal carotid artery, optic nerve (II cranial nerve), optic chiasm, oculomotor nerve, midbrain (cerebral peduncle, tegmentum and tectum), and the splenium at the posterior-most part of the corpus callosum.

The next step was examination of the sulco-gyral architecture. The main sulci of the medial aspect were identified at this stage, namely the sulcus of the corpus callosum, the cingulate, subparietal, parieto-occipital and calcarine sulcus. In the frontobasal area, the main superficial anatomical landmarks visible were the olfactory sulcus

delimiting medially the gyrus rectus (or straight gyrus) and laterally the medial orbital gyrus. The orbital sulcus had the shape of a letter "H" delimiting four orbital gyri (anterior, posterior, lateral and medial). In the medial and inferior aspects of the temporal and occipital lobe, the hippocampus sulcus was identified as well as the collateral and occipitotemporal and calcarine sulcus. These landmarks were used references for the next steps of dissection, and for localization of the medial aspect cerebral gyri, the main being the superior frontal gyrus, paracentral lobule, cuneus, precuneus, lingula, cingulate, parahippocampal, fusiform, and inferior temporal gyrus.

On a second layer, the cortex rendered spongy by the preparation was removed from the sulci and short association fibers ("U" fibers) exposed. The fibers recover the depth of each sulcus and then reproduce the sulco-gyral anatomy studied in the previous step.

On a third layer we used delicate wooden spatulas and forceps to gently remove "U" fibers from the sulci. This made it possible to expose and follow the most superficial fasciculi such as the cingulum, the inferior longitudinal fasciculus, and the dorsal callosal radiations.

On a fourth layer, we used a scalpel blade to cut the ventricular septum and open the lateral ventricle. We were able to identify the ependyma which lines the ventricular cavity and observed the fornix which is located partly in the floor ventricular cavity adjacent to the choroid plexus. The dissection progressed into the walls of the lateral and third ventricles at the same time as it progressed inside the cingulum bundle. The procedure enabled exposition of the anterior columns of the fornix inside the hypothalamus, the mammillothalamic tract, anterior and dorsal thalamic radiations. Finally, the atrium and temporal horn of the lateral ventricle were widely opened for exposure of the tapetum and optic radiations around the temporal horn itself (temporal loop) and the sagittal stratum.

The optical radiations occupied the last layer of this dissection. At the end stages, structures of the optic pathways were visible: the optic nerve, optic chiasma, lateral geniculate body, pulvinar, optical radiations, and the dissected primary visual cortex.

3.2.3. Lateral aspect

This phase enabled further detailing of the anatomy of structures that were not completely exposed from the medial aspect, such as those in the walls of the lateral ventricle and the study of anatomical relationships. After study of surface landmarks such as done for the medial aspect, removal of the cerebral cortex and short "U" association fibers was performed, allowing dissection of deep portions of the cerebral white matter. This procedure exposed the long association fasciculi of the semioval center and the sagittal stratum around the lateral ventricle. Given the lack of detailed description of this region in the literature, special attention was given to its structures. Fibers of the inferior longitudinal fasciculus were observed in the inferior margin of the preparation, intermingled with other elements of the region. Although most of these fibers ran at the ventral aspect of the temporal and occipital lobe, some of them could indeed be dissected from the lateral aspect. At this stage, the temporal operculum was preserved not to eliminate the Middle Longitudinal Fasciculus (MdLF) located superficially. The MdLF penetrated the superior temporal gyrus and planum temporale as an organized fiber contingent progressively more compact from posterior to anterior. Posteriorly, albeit superficial, MdLF was clearly intermingled with the surrounding fibers.

Running at a deeper level than the MdLF, the Inferior Fronto Occipital Fasciculus (iFOF) reached the temporal stem and the insula just posterior and dorsal to the limen insulae. Consequently, the superior temporal gyrus including the planum temporale, and the middle temporal gyrus had to be removed prior to its dissection. The posterior segment of the iFOF was partially covered laterally by the dorsal part of the MdLF. As the dissection progressed from the posterior to the middle segment (trunk) of the iFOF, the latter ran deeper to the MdLF, reaching the antero-inferior portion of the external and extreme capsules. The ventral part of the external and extreme capsules also contained the UF, which was ventral to the anterior segment of the iFOF.

The next fiber contingent crossed by the dissection was the optic radiations that originated from the lateral geniculate body and ran in the lateral wall of the atrium and occipital horn prior to the lips of the calcarine fissure. As previously described by other authors, the optic radiations

can be subdivided in three segments from anterior to posterior. The first segment, or Meyer's loop, is the most anterior one and forms a posteriorly concave curve in a lateral oblique plane in close relationships with the anterior commissure and, more anteriorly, the UF. The second segment, or body of the optic radiations, runs along the lateral aspect of the atrium and occipital horn. Finally, the third segment joins the lips of the calcarine fissure. As a whole, the optic radiations – from the thalamus to the cortex – can be considered as specific posterior thalamic radiations and share the same general organization as other adjacent thalamic radiations to the occipital and parietal cortices.

Finally, the tapetum was located medial to the optic, containing fibers in a vertical organization. This layer, which is derived from the corpus callosum, covered the lateral aspect of the ventricular atrium and occipital horn. This was the deepest white matter layer of the lateral wall of the atrium, only limited from the cerebrospinal fluid by the ependyma.

As a summary, the lateral wall of the ventricular atrium comprised from medial to lateral: the ventricular ependyma, tapetum, optic radiations, iFOF, MdLF, posterior portion of the Superior Longitudinal Fasciculus / Arcuate Fasciculus complex, short "U" association fibers, and the cerebral cortex. Some of them also participated to the lateral wall of the occipital ventricular horn. The Inferior Longitudinal Fasciculus occupied only the ventro-lateral aspect of the dissection, its fibers contributing little to the ventral aspect of the superficial layer of the sagittal stratum. Most of these structures were partially superimposed on a sagittal plane. The MdLF, narrower than the iFOF, covered only its upper portion. The iFOF, in turn, partially covered the optic radiations, which overflows ventrally at the level of Meyer's loop. The optic radiations spilled over into the anterior-inferior portion of the iFOF as they are more ventral in location and extend around the temporal horn. As it progressed forward, the iFOF moved away from it to reach the temporal stem and the insula. Therefore, a space was created behind the directional change of the UF fibers, ahead of the Meyer's loop and deep to the iFOF. This space was occupied by fibers from the anterior commissure, opened in a fan of which the more caudal ones joined the SS. Although a large number of fibers were intermingled, this phenomenon varied according to the location considered. As a result, the layers can be separated at

certain sites, whereas in others, the delamination procedure causes the fibers to rupture. This was for instance the case: for the posterior segment of the iFOF, interspersed with the optic radiations as both approach the occipital cortex; for the anterior commissure intermingled with those of the Meyer's loop; and for the fibers of the anterior commissure intermingled with those of the posterior portion of the uncinate fasciculus.

3.3- Results from in vivo DWI

MRI examinations of 120 volunteers are comprised so far in the database of the FIBRATLAS project. Datasets of fifteen participants were assessed for study of limbic and extralimbic connections with special focus on the cingulum bundle. Six (42.8%) of them were male, and the mean age was 85 ± 3.5 years (mean \pm SD, range 82 - 93). Two male subjects (13.3%) reported more than twelve years of education, and three (21.4%) have completed primary school. All women and remaining four men have 12 years of education or less. Based on the ISCO-08 occupational map and the ISCED-97 classification of skill levels, a total of four (26.7%) participants had skill level I occupations (physical or manual activities), four (26.7%) level II (machinery or electronic equipment operation, driving, maintenance or manipulation of information), three (13.3%) level III (technical task with procedural knowledge in a specialized field), and four (26.7%) level IV (activities requiring complex problem-solving, creativity or decision-making based on extensive bodies of knowledge). In one case, no occupation was informed.

All thirty cingulum reconstructions were performed successfully. Direct frontoparietal transcingular tracks were observed in fifteen right and fourteen left cerebral hemispheres (not observed in the left hemisphere of one female subject). In these 29 tractograms, interconnections between the medial frontal cortex and the posterior precuneus were depicted. Interconnections involving frontal areas (the anterior middle frontal gyrus, the superior frontal gyrus, the precentral gyrus) and parietal areas (the postcentral gyrus and the precuneus, corresponding to the medial surface of the superior parietal lobe) were depicted. Pathways between the rostralateral and dorsal-lateral prefrontal cortex, Brodmann areas (BA) 10 and 8, 9, and the posterior medial parietal cortex (precuneus), BA 7, were observed in

twenty-four hemispheres; and between the primary motor cortex, BA 4, and the parietal medial areas in six right and seven left hemispheres. In the parietal lobe, most of interconnections (in 27 hemispheres) ended in the precuneus or the postcentral gyrus (in 13 hemispheres). Those between the anterior middle frontal gyrus (BA 10) and the precuneus were depicted in thirteen. Direct tracks between the posterior medial frontal cortex, supplementary motor area (BA 6), and the precuneus were observed in fourteen hemispheres. Inside the cingulum, tracks making direct connections between the superior frontal gyrus and precuneus assumed a more medial and inferior trajectory. In turn, fibers that interconnect the precuneus and supplementary motor area were more lateral in position.

The median fiber length and volume of the transcingular direct frontoparietal tracts were respectively 102.2 ± 30.8 mm (median \pm SD) and 2 ± 2.5 ml (median \pm SD) respectively for the right hemisphere, and 128.18 ± 19.4 (median \pm SD) and 5.6 ± 4.2 ml (median \pm SD) for the left hemisphere. The value of fractional anisotropy (FA) was 0.23 ± 0.049 (mean \pm SD, median 0.24) for the right, and 0.23 ± 0.031 (mean \pm SD, median of 0.24) for the left hemisphere. The number of tracks was 201.9 ± 223.22 (mean \pm SD, median of 81) in the right, and 424.9 ± 396.46 (mean \pm SD, median 369.50) in the left. The voxel count resulted in 3158.6 ± 2675.6 (mean \pm SD, median 2009.0) for the right and 4995.21 ± 3730.3 (mean \pm SD, median 3763.5) for the left hemisphere.

Sex was related to significant differences in the scalar measurements in our sample. Significantly higher FA was observed in women in the right hemispheres ($p=0.04$, $LI=-0.22$), and in males in the left hemisphere ($LI=0.17$) in conjunction with higher voxel count values ($p=0.01$). No other statistically significant difference of scalar measurements in the same sex was observed, but the volume laterality index indicated leftward lateralization ($LI = 0.70$) in men. Also, a total volume inferior or equal to the median (2 ml in the left hemisphere was more frequent among women ($p=0.03$)). Leftward lateralization was observed for FA in three men, and for fiber length in one man and four women. Right lateralization was observed in one female subject for FA and length. There was no association of volume or fiber length with profession or educational scores.

3.4- Results of neuropsychological assessment

Fourteen volunteers completed the neuropsychological assessment. The basic cognitive assessment using the Folstein's Mini-Mental State Examination showed a mean score of 26 (19 to 29). Three participants scored under 24.

Concerning the Rey Complex Figure Test, five subjects copied it using a type 1 strategy, four using a type 2, one using copy type 3 and five a type 4. Visuospatial abilities varied widely across participants. The mean immediate recall score was 11.1 ± 1.6 and the delayed recall score was 10.9 ± 1.3 (mean \pm SD), both significantly lower than the initial copy score (29.8 ± 0.4 , $p < 0.01$). No statistical difference was observed between immediate and delayed recall scores. Negative correlation was observed between age and copy score (mean correlation coefficient of $r = -0.559$, leading to an equivalent corrected alpha of 0.047). Positive correlation was observed between immediate recall score and delayed recall score ($r = 0.824$, $p=0.001$), copy time and immediate recall time (0.700 , $p=0.008$), copy time and delayed recall time ($r = 0.726$, $p=0.005$), and immediate recall and delayed recall time ($r = 0.600$, $p=0.03$).

For the RL/RI-16 free and cued recall task, Z-values and quantiles of the normalized results of the RL/RI-16 free and cued recall test were obtained. The total free recall and the sum of total recall scores were 22.07 ± 7.35 and 42.57 ± 5.27 (mean \pm SD, $p < 0.01$) respectively. Two participants presented with low Z scores, under the pathological threshold (quantile 5, $Z=-4.47$ in this sample).

In the Trail Making Test test the mean interval of time for completion of part A was 58.29 ± 16.6 (mean \pm SD) and the median 58 seconds. For part B this figure was 143.14 ± 41.9 seconds and the median 143.5 seconds. The difference of the time needed to complete parts A and B was $84.8 \pm 31.2s$ (mean \pm SD) and the median 94 seconds. Quantiles of the Z scores were obtained. A linear regression analysis did not show significant effect of age, sex and level of study on performance in parts A and B in our sample.

In the Stroop test, participants took longer to name the ink color word when the color name was incongruent with the ink color in all subjects. In such "Interference" situation, in which participants had to inhibit the written word in order to correctly name the color of the ink, the mean time was 167.83 ± 43.8 seconds. The mean difference

between the situation of interference and that of simple denomination was 80.87 seconds, with a median of 92 ± 47 seconds. The mean of total uncorrected errors was 2.08 ± 2.32 (mean \pm SD, range 0-8). In participants aged 85 years or older, the time necessary to complete the task in interference condition, mean time per word, and interference score were significantly higher.

Lastly, data from published studies were used to compare this sample to others of similar range of age (Supplementary Figure 1). For the Mini Mental State Examination Test, results were similar (Kalafat, 2003) or at the significance threshold (Kochhann et al. 2009, $p=0.05$). For the Rey Complex Figure Test (Fastenau et al. 1999; Foss et al. 2010), results were similar for both the copy and the delayed recall scores. Similarities were also observed in the Trail Making Test part A (Tombaugh et al. 2003, Amieva et al. 2009) and in the number of errors in the interference condition of the Stroop test (Bayard et al. 2011). Nevertheless, a significant difference was observed with Zimmermann et al. 2015 since this score was inferior in our sample.

3.5- Results of compilation of morphological and neuropsychological findings

For analysis, scalar data were distributed in two groups in function of the median scores. Associations between scalar values and neuropsychological performance were observed in women only. Women with a volume of transcingular frontoparietal tracks inferior to the median in the left hemisphere presented with a Mini Mental State score ≥ 25 ($p=0.02$). For the Rey figure, all women with median fiber length ≤ 101.93 mm in the right hemisphere were in the group above the mean copy normalized score ($p=0.02$). For the immediate recall, all women had scores inferior to the mean (11.11 ± 1.63 , mean \pm SD).

A Spearman correlation analysis of scalar variables was performed with the performance on Rey Complex Figure, RL/RI-16 Free and Cued Recall Task, Trail Making and the Stroop test. In the left hemisphere, FA ($r = -0.56$; $p=0.029$) and volume ($r = -0.53$; $p=0.028$), but not voxel count or fiber length, were negatively correlated with Stroop interference time. Voxel count was negatively correlated with the difference of time necessary to complete parts B and A of the Trail Making Test (B-A) ($r = -0.51$; $p=0.035$). In the right hemisphere, voxel count ($r = 0.58$; $p=0.014$)

and volume ($r = 0.51$; $p=0.031$) were correlated with memory reproduction, the immediate recall score of the Rey Complex Figure Test. Voxel count was also correlated with the delayed recall score ($r = 0.49$; $p=0.035$).

3.6- Results from ex vivo DWI

To date, a total of ten human brain specimens have been examined on MRI. Two were scanned at 3T (Martinos Center, Boston, USA). Other eight were scanned in high magnetic fields (CEA Neurospin, Saclay, France): seven at 7T, and one at 11.7T (with resolutions of 100, 150 and 200 μm).

These procedures resulted in a unique library of image datasets for future studies in detailed morphology of the human white matter. The datasets will be made available in the form of an open database on the FIBLATLAS project webpage (<https://fibratlas.univ-tours.fr/>), after an embargo period.

3.7- Synthesis of available data

In the present study, DTI tractography was used to detail the morphology of long association fibers interconnecting the frontal and parietal lobes through the dorsal cingulum. Exploration by two independent operators demonstrated transcingular tracks in 29 out of 30 cerebral hemispheres, which were anatomically detailed. These findings reinforce the idea of the cingulum as an important route of direct frontoparietal association pathways in parallel to those comprised in the Superior Longitudinal Fasciculus. A standardized neuropsychological assessment of the same volunteers suggested that faster planification and visuospatial processes were associated to the presence of longer tracks. To our best knowledge, this was the first study on diffusion imaging focusing on transcingular direct frontoparietal connections in the human brain. This is a subject of great importance in the field of modeling of mental functions. Beyond the participation of the cingulum in the limbic system, it seems to be an important element of highly functional dorsomedial circuits such as those related to motor activity initiation, executive functions, and the Default Mode Network.

Longer fibers were deeper (more lateral) inside the cingulum, an observation which is in line

with the findings in non-human primates (Mufson and Pandya 1984; Aggleton et al. 2012). The human cingulum is, however, larger in volume and looks richer in regional fibers. Connections to the Supplementary Motor Area (SMA) and precuneus were robust on left hemisphere. Hand preference and the importance of these connections to motor planning and visuospatial integration are possible explanations for this phenomenon. Asymmetries have been observed for other fascicles in humans. A classic example is the arcuate fasciculus (Parker et al. 2005; Rilling et al. 2008; Madhavan et al. 2014; Hecht et al. 2015; Keedwell et al. 2012; Fan et al. 2015) a phenomenon considered to be related to language dominance.

Our results investigated relevant relationships between scalar measurements of transcingular frontoparietal tracts and neuropsychometric parameters such as episodic memory (Rey complex figure delayed recall, RL/RI-16 free and cued recall), visuospatial processes and planification (Rey complex figure copy), cognitive inhibition (Stroop Test) and flexibility processes (Trail Making Test). Some structural particularities were identified in individuals with better results. Volunteers with better performance in immediate recall and delayed recall of the Rey Complex Figure Test presented with significantly larger volume and voxel count. A possible interpretation for this is the fact that the richness of interconnecting pathways tends to coexist with the increased activation of frontal and parietal areas that ensures better visuospatial abilities. In the Trail Making Test, individuals with less difference on time between parts A and B showed higher track voxel count in left transcingular pathways. In addition, more time needed to complete in Stroop test was related with smaller figures of fractional anisotropy and volume.

Cognitive inhibition (Stroop test) was associated with volume and FA on the left hemisphere, and memory reproduction (Rey complex figure test) was associated with volume and voxel count on the right. Lower microstructural organization of the cingulum was reported to be related to a decrease in performance in various cognitive domains and in the activation of the default mode network (Bathelt et al. 2019). In a study of 220 older adults without dementia, attention, and executive function performance correlated to the FA of the anterior cingulum. The same was observed in the posterior cingulum for memory, language, attention, executive and visuo spatial

function (Kantarci et al. 2011). The efficacy of cognitive control was also reported to be very sensitive to the cingulum microstructure (Metzler-Baddeley et al. 2012). Prior investigations that highlighted the link between a more developed functional connectivity and higher cognition have also pointed to the importance of frontoparietal connections in both adults (Charlton et al. 2010; Ritchie et al. 2015; Bettcher et al. 2016; Ekman et al. 2016) and children (Wendelken et al. 2016; Carter Wendelken, Emilio Ferrer, Simona Ghetti 2017). These findings are relevant to clinical neuroscience of both pathological and normal aging conditions. Indeed, executive functions are among the most commonly affected cognitive domain in older adults.

Anatomical and neuropsychological consequences of aging were certainly present in our series, such as the decline in white matter integrity and aging-functional modifications on episodic memory and executive functions. In a study with 94 older adults without dementia (90–103 years), loss of microstructural integrity within the cingulum was associated with chronological aging and global cognition (Bennett et al. 2017). These microstructural changes do not seem to occur homogeneously along the white matter tracks. An anterior-posterior gradient of FA decreases has been reported; the anterior portions of the dorsal cingulum being the most compromised portion of this bundle, particularly the left subgenual segment (Sibilia et al. 2017; Davis et al., 2009; Gunning et al. 2009). It is not clear whether white matter integrity losses are symmetrical or not. Comparisons with previously published results with samples with a similar age range showed similarity of most neuropsychological results.

Tract-tracing studies in non-human primates have demonstrated that the cingulum contains a very heterogeneous contingent of fibers, with at least three juxtaposed components with distinct cortical terminations (Mufson & Pandya, 1984). The first component emanates from in the anterior and lateral dorsal nucleus of the thalamus and occupies the ventral aspect of the bundle. Fibers from the anterior nuclei travel to the prefrontal, retrosplenial, and posterior cingulate cortices (monkey area 23), while those from the lateral dorsal nucleus terminate in the retrosplenial cortex, parahippocampal gyrus, and presubiculum. A second component originates from the cingulate cortex itself and has a dorsolateral position. It contains fibers from the anterior and ventral

cingulate cortex (area 24) that extend to the premotor cortex, area 23, and retrosplenial cortex, and fibers that leave area 23 to reach the prefrontal, parahippocampal gyrus, and presubiculum. Finally, a third component connects frontal and parietal associative areas. Fibers that originate in prefrontal areas travel to the retrosplenial cortex and those that originate with a posterior parietal topography extend to the prefrontal, parahippocampal, and presubicular cortices (Mufson & Pandya, 1984).

Inside the cingulate gyrus, fibers of different lengths make up the cingulum. The longest ones originate in the amygdala, uncus, and parahippocampal gyrus and travel to the subgenual area in the frontal lobe (Catani, et al., 2013; Husain & Schott, 2016; Nieuwenhuys, et al., 2008). On their way, they are joined by a large number of adjacent regional “U-shaped” association fibers interconnecting the temporal, occipital, parietal, and frontal lobes, and bringing them into relation with various parts of the cingulate gyrus (Bubb, et al., 2018; Catani & Thiebaut de Schotten, 2008).

A considerable amount of information on the functional aspects of the medial frontal cortex has been made available in the last years thanks to the multiplication in the number of studies in fMRI. In a recent term-based meta-analysis of 9721 publications, a division of the medial frontal cortex into three topographic zones was proposed (de la Vega, Chang, Banich, Wager, & Yarkoni, 2016). The activation peaks mostly reflected the functional anatomy described by previous authors. The anterior zone was significantly associated with episodic memory, social processes, decision making, reward and fear, the middle zone with aspects of cognitive control, decision making, fear and pain, and the posterior zone with motor aspects. In the middle zone itself, the anterior clusters were much more related to decision-making processes, those of the middle ventral cingulate cortex were related to affection (co-activating together with the amygdala and striatum), and the more dorsal ones were related to working memory (with greater co-activation of other regions associated to cognitive control). The posterior zone corresponded to regions previously known to be implicated on motor function in many studies, such as the SMA, the pre-SMA and the motor cingulate cortex. The SMA also showed association with pain processing, suggesting its involvement in the initiation of pain-induced movements, while the pre-SMA

was more related to cognitive control, co-activated with regions considered to be important for goal-driven cognitive processes (de la Vega, et al., 2016).

A recent discussion involved the significance of the cingulum for the connection of distant areas in the frontal and parietal lobes. In the Rhesus monkey, autoradiography studies in the 1980s indicated that dorsal frontoparietal interconnections were mainly ensured by the SLF-I component of the SLF/AF (Superior Longitudinal Fasciculus/Arcuate Fasciculus) complex, in the form of a long association bundle running parallel to the superior border of the cerebral hemisphere (Petrides & Pandya, 1984). Indeed, injected radioisotopes do migrate from the medial parietal cortex through the cingulum to reach distant regions of the frontal cortex (motor and premotor areas, areas 24 and 8) (Mufson & Pandya, 1984; Petrides & Pandya, 1984). In agreement with this, Schmahmann et al. (2007) depicted similar fibers extending to the medial (areas 8, 9, 14, 25, and 32) and dorsolateral (areas 8, 9, 11, and 46) aspects of the frontal and parietal (inferior parietal lobule and the retrosplenial cortex) lobes using diffusion spectrum imaging in non-human primates (Schmahmann, et al., 2007).

In humans, studies on this subject employing diffusion imaging or fiber dissection took place more than 20 years later than the initial descriptions with autoradiography (Jang & Hong, 2012; Kamali, Flanders, Brody, Hunter, & Hasan, 2014; Makris, et al., 2005; Yagmurlu, Middlebrooks, Tanriover, & Rhoton, 2016). Preferential anteroposterior diffusion was identified by means of diffusion imaging in a topography that was analogous to that of the monkey SLF-I (Makris, et al., 2005). However, fiber dissection studies did not isolate this bundle in a reproducible manner with less homogeneous results (Fernandez-Miranda, et al., 2008; I.L. Maldonado, E. Mandonnet, & H. Duffau, 2012; Pascalau, et al., 2018; Shah, et al., 2012; Wu, Sun, Wang, Wang, & Ou, 2016). A great number of regional connections were nonetheless described in that region and some controversy exists as they may be a confounding factor for diffusion imaging (I.L. Maldonado, et al., 2012). In this debate, the cingulum emerged as a potential additional source of long-distance pathways. Less affected by gyrification, it was estimated that this fasciculus may be the preferential route

of fibers directly connecting the dorsomedial cortex of the frontal and parietal lobes (I.L. Maldonado, et al., 2012; Wang, et al., 2016).

The presence of direct dorsal frontoparietal connections was suggested in the subgenual subdivision of the classification proposed by Jones et al. (2013) and in subcomponents III and IV in that proposed by Wu et al. (2016). Some authors have recently argued that fibers of the dorsal component of the superior longitudinal fasciculus (SLF-I) could actually be considered part of the cingulum (Wang, et al., 2016; Wu, et al., 2016). Once again, a debate arises - the very identity of the elements of the cerebral white matter is under discussion. Here, a discussion of nomenclature and the logic behind it is also pertinent, as detailed by Mandonnet et al. (2018). SLF-I was defined as the dorsal component of the SFL/AF complex, connecting areas of the cerebral cortex along the upper border of the cerebral hemisphere. The corona radiata physically separates this (laterally located) complex from the (more medial) cingulate fibers. This observation reinforces the importance of the detailed study of the arrangement of these association bundles in the coronal plane and the fact that cingulate fibers should not be labeled SLF-I (Mandonnet, Sarubbo, & Petit, 2018). Although they do indeed contribute to dorsal frontoparietal connectivity, they belong to a distinct anatomical structure.

4- Conclusions

In the study of limbic connectivity through fiber dissection, diffusion tensor and examination of the available literature data, the importance of the cingulum for long distance (even extra-limbic) interconnections stands out. Nevertheless, a higher proportion of locoregional fibers in the subgenual and parahippocampal segments are observed in humans than in non-human primates. This difference is probably enhanced by the significant evolutionary increase in the volume of the frontal and occipital lobes.

A high level of agreement between the results of diffusion imaging and fiber dissection in humans is noticed. The white matter of the dorsal cingulum contains fibers responsible for direct interconnections between highly specialized frontal and parietal areas of the medial aspect of the cerebral hemisphere such as the superior frontal gyrus, supplementary motor area, and precuneus. These routes connect regions that

have been acknowledged to be highly involved in the initiation of motor activity, executive functions and the default mode network. Therefore, the study of its functional aspects must go beyond its participation in the limbic system and in the cortical processing of emotions.

In the left hemisphere, fractional anisotropy was associated to cognitive inhibition, and voxel count was associated with flexibility processes. In the right hemisphere, voxel count and volume were associated with nonverbal memory reproduction. The findings suggest the participation of transcingular direct frontoparietal pathways on executive functions and visuospatial memory.

Our understanding of the structure and function of the white matter bundles has evolved considerably over the last two decades. In addition to its classical participation in the limbic system, the cingulum has been identified as the anatomical substrate of a connective system for several areas involved in higher functions. These recent observations have had a direct influence on how we have and will interpret disconnection syndromes. As the cingulum is a multilayered tract, an important avenue for future research is to distinguish the functional roles of the different layers. More than simply a fiber bundle, it should be now perceived as the primary interconnecting apparatus of the medial aspects of the cerebral lobes.

5- Perspectives of future collaborations with the host laboratory

The ex-fellow and the host researcher continue to work together on research projects, especially in the field of ex vivo imaging, fiber dissection, and cerebral vascularization. After his fellowship IL Maldonado was hired by the Tours University Hospital and remained working in the iBrain research unit, Inserm U1253. Fiber dissection and magnetic-resonance imaging continue to be among his most used methodological tools.

6- Scientific production in the framework of the fellowship

6.1- Published article

Maldonado IL, Parente de Matos V, Castro Cuesta TA, Herbet G, Destrieux C. The human cingulum: From the limbic tract to the connectionist paradigm. *Neuropsychologia*. 2020 Jul;144:107487.

6.2- Article in press

Maldonado IL, Destrieux C, Ribas ESC, Guimarães BSAB, Cruz PP, Duffau H. Composition and organization of the sagittal stratum in the human brain: a fiber dissection study. *Journal of Neurosurgery*. *In Press*.

6.3- Submitted articles

Cuesta TAC, Matos VP, Andersson F, Brachet M, Destrieux, Maldonado IL. Trans cingular dorsal frontoparietal connections: a study of diffusion imaging and neuropsychological testing in elderly. *Neuropsychology*. Under review.

Cuesta TAC, Matos VP, Destrieux C, Maldonado IL. Comparative connectivity in human and non-human primates. *Journal of Primatology*. Submitted.

6.4- Book chapters

Maldonado IL, Zemmoura I, Destrieux C. Middle longitudinal fasciculus in the human brain from fiber dissection. In: Turgut M, Yurttas C, Tubbs RS, editors. *Island of Reil (insula) in the human brain : anatomical, functional, clinical and surgical aspects*. Cham: Springer; 2018. p. 71–5.

Maldonado IL, Zemmoura I, Destrieux C. Gross anatomy of the Human Insula. In: Turgut M, Yurttas C, Tubbs RS, editors. *Island of Reil (insula) in the human brain : anatomical, functional, clinical and surgical aspects*. Cham: Springer; 2018. p. 15–22.

Destrieux C, Maldonado IL, Terrier L-M, Zemmoura I. Surgical anatomy of the insula. In: Turgut M, Yurttas C, Tubbs RS, editors. *Island of Reil (insula) in the human brain : anatomical, functional, clinical and surgical aspects*. Cham: Springer; 2018. p. 23–37.

6.5- Conference abstracts

Human Brain Mapping 2020

- HotRY, Popov A, Beaujoin J, Perez G, Poupon F, Maldonado IL, Mangin JF, Destrieux C and Poupon C. Ex vivo mapping of the cyto- and the myeloarchitecture of the human cerebral cortex using ultrahigh field MRI (7T and 11.7T).

- Smirnov M, Serres B, KERdilles G, Barantin L, Zhornyk V, Maldonado IL, Destrieux C. Towards an accurate identification of vascular territories in the human brain.

6-6- Lectures and workshops in international conferences

May 22-25, 2019. Surgery Follows Function. White Matter Dissection Course & Symposium. Graz, Austria.

- Cerebral Network System - Lateral Aspect
- Cerebral Network System - Medial Aspect
- White Matter Dissection Unit - Hands-On.

April 12-13, 2019. 6ème Journée de Neurochirurgie d'Oran, Oran, Algérie.

- *Anatomie intrinsèque de la substance blanche cérébrale*

January 30-31, 2019. World Federation of Neurosurgical Societies Anatomy Committee, Hands-On Cadaveric Course, Manipal, India.

- Intrinsic anatomy of the human brain
- Limbic lobe & related white matter tracts. Anatomy and surgery - 3D Lecture
- Medial Aspect white matter fiber tracts dissection

May 31- June 1, 2018 WMTours2018: Dissection and imaging of white matter tracts – Neurosurgical applications

Tours – France

- Autoradiography. Contributions to the understanding of white matter anatomy
- Step by step dissection of the arcuate fasciculus. Hands-On Workshop
- Step by step dissection of the ventral pathway. Hands-On Workshop.

September 21-23, 2018, The Cerebral White Matter – Functional Anatomy and Surgical Approaches, Cochabamba – Bolivia

- General organization of the hemispheric white matter.
- The cerebral white matter – Hands-On Cadaver Dissection.

6-7 Organization of a scientific meeting

5-7 December 2018. Frontiers in connectivity. Exploring and Dissecting the Cerebral White Matter

Lectures / workshops performed by the fellow:

- Tumor infiltration and white matter fibers
- General anatomy of the human cerebral white matter with fiber dissection
- Step-by-step dissection of the dorsal stream (Hands-on Workshop)
- Step-by-step dissection of the ventral stream (Hands-on Workshop)

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

- Step-by-step dissection of the optic radiations and tapetum (Hands-on Workshop)

- Morphology of the medial aspect of the cerebral hemisphere and limbic system

- Dissection of the medial aspect the cerebral hemisphere and limbic system (Hands-on Workshop)

6-8 Participation in educational website

Anthropotomia, e-manuel de dissection. Ed.: Destrieux C. Home page: <http://anthropotomia.univ-tours.fr/>; Central Nervous System page: <https://anthropotomia.univ-tours.fr/v2/en/central-nervous-system/>

6-9 Other publications during the period of residence

Lechanoine F, Janot K, Herbreteau D, Maldonado IL, Velut S. Surgical Thrombectomy Combined with Bilateral Decompressive Craniectomy in a Life-Threatening Case of Coma from Cerebral Venous Sinus Thrombosis: Case Report and Literature Review. *World Neurosurg.* 2018 Dec;120:485–9.

Dupuy-Bonafe I, Maldonado IL. “Please don’t move”: Cone-beam computed tomography and obstructive sleep apnea hypopnea syndrome. *Am J Orthod Dentofacial Orthop.* 2019 May;155(5):616–7.

Lechanoine F, Smirnov M, Armani GL, Carneiro P, Cottier P, Destrieux C, et al. Stereoscopic images from computed tomography angiograms. *World Neurosurg.* 2019 May 9;

Junior VA dos S, Santos M de S, Ribeiro NM da S, Maldonado IL. Combining proprioceptive neuromuscular facilitation and virtual reality for improving sensorimotor function in stroke survivors: a randomized clinical trial. *Journal of central nervous system disease.* 2019;11:1179573519863826.

Janot K, Charbonnier G, Boustia F, Lima Maldonado I, Bibi R, Pucheux J, et al. [Stroke prevention]. *Presse Med.* 2019 May 28;

Souza DCB, Santos MS, Ribeiro NMS, Maldonado IL. Inpatient trunk exercises after recent stroke: An update meta-analysis of randomized controlled trials. *NeuroRehabilitation.* 2019;Epub Ahead of Print.

Lessa KP, Vilasboas IGM, Aquino MA, Maldonado IL. Usage of a Translated Version of Terminologia Anatomica in Brazil: An Examination of the Neurosurgical Literature. *Brazilian Neurosurgery*. 2019; 38(2): 106-111.

7- Acknowledgements

We express our gratitude to the study volunteers and body donors, who make this research possible. We thank Gerald Deluermoz, Jean Paul da Silva, and Carine Rougeron from the Laboratory of Human Anatomy for their technical and administrative support.

This work was supported by the Le Studium, Loire Valley Institute for Advanced Studies, Orleans & Tours, France under Marie Sklodowska-Curie grand agreement no. 665790, European Commission.

We acknowledge the French Research Agency (Agency National de Recherche) for funding of the FIBRATLAS project.

We also thank the Planiol Foundation and the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil* (CAPES, Finance Code 001) for the financial support to our trainee Taryn Ariadna Castro-Cuesta.

8- References

Aggleton JP, Wright NF, Vann SD, Saunders RC (2012) Medial temporal lobe projections to the retrosplenial cortex of the macaque monkey. *Hippocampus* 22:1883–1900. doi: 10.1002/hipo.22024

Alarcon, C., de Notaris, M., Palma, K., Soria, G., Weiss, A., Kassam, A., & Prats-Galino, A. (2014). Anatomic study of the central core of the cerebrum correlating 7-T magnetic resonance imaging and fiber dissection with the aid of a neuronavigation system. *Neurosurgery*, 10 Suppl 2, 294-304; discussion 304.

Amieva H, Le Goff M, Stoykova R, et al (2009) Trail Making Test A et B (version sans correction des erreurs) : normes en population chez des sujets âgés, issues de l'étude des trois Cités. *Rev Neuropsychol* 1:210. doi: 10.3917/rne.013.0210

Amiez C, Sallet J, Hopkins WD, et al (2019) Sulcal organization in the medial frontal cortex provides insights into primate brain evolution. *Nat Commun* 10:. doi: 10.1038/s41467-019-11347-x

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

Amunts K, Zilles K (2015) Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron* 88:1086–1107. doi: 10.1016/j.neuron.2015.12.001

Arnts, H., Kleinnijenhuis, M., Kooloos, J. G., Schepens-Franke, A. N., & van Cappellen van Walsum, A. M. (2014). Combining fiber dissection, plastination, and tractography for neuroanatomical education: Revealing the cerebellar nuclei and their white matter connections. *Anat Sci Educ*, 7, 47-55.

Assaf Y, Johansen-Berg H, Thiebaut de Schotten M (2019) The role of diffusion MRI in neuroscience. *NMR Biomed* 32:e3762. doi: 10.1002/nbm.3762

Azouvi P. (2017). Neuropsychologie de l'attention : aspects théoriques, évaluation, rééducation.

Baker CM, Burks JD, Briggs RG, et al (2018) A Connectomic Atlas of the Human Cerebrum-Chapter 4: The Medial Frontal Lobe, Anterior Cingulate Gyrus, and Orbitofrontal Cortex. *Oper Neurosurg (Hagerstown, Md)* 15:S122–S174. doi: 10.1093/ons/opy257

Barbeau E, Tramon E, Joubert S, Mancini J, Ceccaldi M, Poncet M. (2004). Evaluation de la mémoire en reconnaissance visuelle : normalisation d'une nouvelle épreuve en choix forcé (DMS 48) et utilisé en neuropsychologie clinique. In: GREMEN VDLMa, editor(s). *L'évaluation des troubles de la mémoire*. Marseille: Solal; p. 85–101

Barona A, Reynolds CR, Chastain R (1984) A demographically based index of premorbid intelligence for the WAIS--R. *J Consult Clin Psychol* 52:885–887. doi: 10.1037/0022-006X.52.5.885

Basser, P. J., Mattiello, J., & LeBihan, D. (1994a). Estimation of the effective self-diffusion tensor from the NMR spin echo. *J Magn Reson B*, 103, 247-254.

Basser, P. J., Mattiello, J., & LeBihan, D. (1994b). MR diffusion tensor spectroscopy and imaging. *Biophys J*, 66, 259-267.

Basser, P. J., Pajevic, S., Pierpaoli, C., Duda, J., & Aldroubi, A. (2000). In vivo fiber tractography using DT-MRI data. *Magn Reson Med*, 44, 625-632.

Bathelt J, Johnson A, Zhang M, Astle DE (2019) The cingulum as a marker of individual

- differences in neurocognitive development. *Sci Rep* 9:1–16. doi: 10.1038/s41598-019-38894-z
- Baumstarck K, Boyer L, Boucekine M, et al (2013) Self-reported quality of life measure is reliable and valid in adult patients suffering from schizophrenia with executive impairment. *Schizophr Res* 147:58–67. doi: 10.1016/j.schres.2013.03.008
- Beckmann M, Johansen-Berg H, Rushworth MF (2009) Connectivity-based parcellation of human cingulate cortex and its relation to functional specialization. *J Neurosci* 29:1175–1190. doi: 10.1523/jneurosci.3328-08.2009
- Beevor CE (1891) On the Course of the Fibres of the Cingulum and the Posterior Parts of the Corpus Callosum and Fornix in the Marmoset Monkey. *Philos Trans R Soc London B* 182:135–199
- Bennett IJ, Greenia DE, Maillard P, et al (2017) Age-related white matter integrity differences in oldest-old without dementia. *Neurobiol Aging* 56:108–114. doi: 10.1016/j.neurobiolaging.2017.04.013
- Bettcher BM, Mungas D, Patel N, et al (2016) Neuroanatomical substrates of executive functions: Beyond prefrontal structures. *Neuropsychologia* 85:100–109. doi: 10.1016/j.neuropsychologia.2016.03.001
- Bliss TVP, Collingridge GL, Kaang B-K, Zhuo M (2016) Synaptic plasticity in the anterior cingulate cortex in acute and chronic pain. *Nat Rev Neurosci* 3–5. doi: 10.1038/nrn.2016.68
- Blyth R, Cook P, Alexander D (2003) Tractography with multiple fibre directions. *Methods* 10427:10427–10427
- Bonni S, Veniero D, Mastropasqua C, et al (2015) TMS evidence for a selective role of the precuneus in source memory retrieval. *Behav Brain Res* 282:70–75. doi: 10.1016/j.bbr.2014.12.032
- Bozkurt, B., Yagmurlu, K., Middlebrooks, E. H., Cayci, Z., Cevik, O. M., Karadag, A., Moen, S., Tanriover, N., & Grande, A. W. (2017). Fiber Connections of the Supplementary Motor Area Revisited: Methodology of Fiber Dissection, DTI, and Three Dimensional Documentation. *J Vis Exp*.
- Bozzali M (2002) White matter damage in Alzheimer's disease assessed in vivo using diffusion tensor magnetic resonance imaging. *J Neurol Neurosurg Psychiatry* 72:742–746. doi: 10.1136/jnnp.72.6.742
- Branco DM, Suarez RO, Whalen S, et al (2006) Functional MRI of memory in the hippocampus: Laterality indices may be more meaningful if calculated from whole voxel distributions. *Neuroimage* 32:592–602. doi: 10.1016/j.neuroimage.2006.04.201
- Broca, P. (1877). Sur la circonvolution limbique et la scissure limbique. *Bulletins de la Société d'anthropologie de Paris*, 12, 646-657.
- Broca, P. (1878). Anatomie comparée des circonvolutions cérébrales. Le grand lobe limbique et la scissure limbique dans la série des mammifères. *Revue d'Anthropologie*, 1, 385-498.
- Brown TL (1996) Attentional Selection and Word Processing in Stroop and Word Search Tasks: The Role of Selection for Action. *Am J Psychol* 109:265. doi: 10.2307/1423276
- Brown TL, Gore CL, Carr TH (2002) Visual attention and word recognition in Stroop color naming: Is word recognition “automatic?” *J Exp Psychol Gen* 131:220–240. doi: 10.1037/0096-3445.131.2.220
- Bruner E, Preuss TM, Chen X, Rilling JK (2017) Evidence for expansion of the precuneus in human evolution. *Brain Struct Funct* 222:1053–1060. doi: 10.1007/s00429-015-1172-y
- Bruner, E. (2018). Human Paleoneurology and the Evolution of the Parietal Cortex. *Brain, Behavior and Evolution*, 91(3), 136–147. doi:10.1159/000488889
- Bubb EJ, Metzler-Baddeley C, Aggleton JP (2018) The cingulum bundle: Anatomy, function, and dysfunction. *Neurosci Biobehav Rev*. doi: 10.1016/j.neubiorev.2018.05.008
- Budisavljevic S, Kawadler JM, Dell'Acqua F, et al (2015) Heritability of the limbic networks. *Soc Cogn Affect Neurosci* 11:746–57. doi: 10.1093/scan/nsv156
- Burzynska AZ, Nagel IE, Preuschhof C, et al (2011) Microstructure of frontoparietal connections predicts cortical responsivity and working memory performance. *Cereb Cortex* 21:2261–2271. doi: 10.1093/cercor/bhq293
- Carmichael ST, Price JL (1995) Limbic connections of the orbital and medial prefrontal cortex in macaque monkeys. *J Comp Neurol* 363:615–641. doi: 10.1002/cne.903630408

- Carnero-Pardo C (2014) Should the Mini-Mental State Examination be retired? *Neurol* (English Ed 29:473–481. doi: 10.1016/j.nrleng.2013.07.005
- Carter Wendelken, Emilio Ferrer, Simona Ghetti et al. (2017) Frontoparietal structural connectivity in childhood predicts development of functional connectivity and reasoning ability: A large-scale longitudinal investigation. *J Neurosci* 30:8549–8558. doi: 10.3389/fpsyg.2018.00265
- Castellanos FX, Margulies DS, Kelly C, et al (2008) Cingulate-Precuneus Interactions: A New Locus of Dysfunction in Adult Attention-Deficit/Hyperactivity Disorder. *Biol Psychiatry* 63:332–337. doi: 10.1016/j.biopsych.2007.06.025
- Castro-Costa É, Fuzikawa C, Uchoa E, et al (2008) Norms for the mini-mental state examination: adjustment of the cut-off point in population-based studies (evidences from the Bambuí health aging study). *Arq Neuropsiquiatr* 66:524–528. doi: 10.1590/S0004-282X2008000400016
- Catani M, Dell’Acqua F, Thiebaut de Schotten M (2013) A revised limbic system model for memory, emotion and behaviour. *Neurosci Biobehav Rev* 37:1724–1737. doi: 10.1016/j.neubiorev.2013.07.001
- Catani M, Mesulam M (2008) The arcuate fasciculus and the disconnection theme in language and aphasia: History and current state. *Cortex* 44:953–961. doi: 10.1016/j.cortex.2008.04.002
- Catani M, Thiebaut de Schotten M (2008) A diffusion tensor imaging tractography atlas for virtual in vivo dissections. *Cortex* 44:1105–1132. doi: 10.1016/j.cortex.2008.05.004
- Catani, M., Dell’Acqua, F., & Thiebaut de Schotten, M. (2013). A revised limbic system model for memory, emotion and behaviour. *Neuroscience & Biobehavioral Reviews*, 37, 1724–1737.
- Cavanna AE, Trimble MR (2006) The precuneus: A review of its functional anatomy and behavioural correlates. *Brain* 129:564–583. doi: 10.1093/brain/awl004
- Chaddad-neto F, Devanir M, Bozkurt B, et al (2017) Contralateral anterior interhemispheric-transcallosal-transrostral approach to the subcallosal region: a novel surgical technique. 1–7. doi: 10.3171/2017.4.JNS16951.
- Chai SC, Kung JC, Shyu BC (2010) Roles of the anterior cingulate cortex and medial thalamus in short-term and long-term aversive information processing. *Mol Pain* 6:1–10. doi: 10.1186/1744-8069-6-42
- Charlton RA, Barrick TR, Lawes INC, et al (2010) White matter pathways associated with working memory in normal aging. *Cortex* 46:474–489. doi: 10.1016/j.cortex.2009.07.005
- Concha L, Gross DW, Beaulieu C (2005) Diffusion tensor tractography of the limbic system. *AJNR Am J Neuroradiol* 26:2267–2274
- Conturo, T. E., Lori, N. F., Cull, T. S., Akbudak, E., Snyder, A. Z., Shimony, J. S., McKinstry, R. C., Burton, H., & Raichle, M. E. (1999). Tracking neuronal fiber pathways in the living human brain. *Proceedings of the National Academy of Sciences*, 96, 10422–10427.
- Cooper M, Thapar A, Jones DK (2015) ADHD severity is associated with white matter microstructure in the subgenual cingulum. *NeuroImage Clin* 7:653–660. doi: 10.1016/j.nicl.2015.02.012
- Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* 3:201–215. doi: 10.1038/nrn755
- Cornell DG, Roberts M, Oram G (1997) The Rey Osterrieth Complex Figure Test as a Neuropsychological Measure in Criminal Offenders. *Arch Clin Neuropsychol* 12:47–56
- Costa, M. D., Braga, V. L., Yagmurlu, K., Centeno, R. S., Cavalheiro, S., & Chaddad, F. (2018). A Technical Guide for Fiber Tract Dissection of the Internal Capsule. *Turk Neurosurg*.
- Davis SW, Dennis NA, Buchler NG, et al (2009) Assessing the effects of age on long white matter tracts using diffusion tensor tractography. *Neuroimage* 46:530–541. doi: 10.1016/j.neuroimage.2009.01.068
- De Benedictis A, Duffau H (2011) Brain hodotopy: From esoteric concept to practical surgical applications. *Neurosurgery* 68:1709–1723. doi: 10.1227/NEU.0b013e3182124690
- de la Vega, A., Chang, L. J., Banich, M. T., Wager, T. D., & Yarkoni, T. (2016). Large-Scale Meta-Analysis of Human Medial Frontal Cortex Reveals Tripartite Functional Organization. *J Neurosci*, 36, 6553–6562.

- Deloche G, Hannequin D. (1997). Test de dénomination orale d'images D0 80. Editions du Centre de Psychologie Appliquée.
- De Simoni S, Grover PJ, Jenkins PO, et al (2016) Disconnection between the default mode network and medial temporal lobes in post-traumatic amnesia. *Brain* 139:3137–3150. doi: 10.1093/brain/aww241
- Dion M, Potvin O, Belleville S, et al (2015) Normative data for the rappel libre/rappel indicé à 16 items (16-item free and cued recall) in the elderly quebec-french population. *Clin Neuropsychol* 28:1–19. doi: 10.1080/13854046.2014.915058
- Duffau H (2016) Stimulation Mapping of Myelinated Tracts in Awake Patients. *Brain Plast* 2:99–113. doi: 10.3233/BPL-160027
- Duley JF, Wilkins JW, Hamby SL, et al (1993) Explicit scoring criteria for the Rey-Osterrieth and Taylor complex figures. *Clin Neuropsychol* 7:29–38. doi: 10.1080/13854049308401885
- Dunlop K, Hanlon CA, Downar J (2017) Noninvasive brain stimulation treatments for addiction and major depression. *Ann N Y Acad Sci* 1394:31–54. doi: 10.1111/nyas.12985
- Ekman M, Fiebach CJ, Melzer C, et al (2016) Different Roles of Direct and Indirect Frontoparietal Pathways for Individual Working Memory Capacity. *J Neurosci* 36:2894–2903. doi: 10.1523/JNEUROSCI.1376-14.2016
- Etkin A, Egner T, Kalisch R (2011) Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends Cogn Sci* 15:85–93. doi: 10.1016/j.tics.2010.11.004
- Fan S, van den Heuvel OA, Cath DC, et al (2015) Mild White Matter Changes in Un-medicated Obsessive-Compulsive Disorder Patients and Their Unaffected Siblings. *Front Neurosci* 9:495. doi: 10.3389/fnins.2015.00495
- Fan S, van den Heuvel OA, Cath DC, et al (2015) Mild White Matter Changes in Un-medicated Obsessive-Compulsive Disorder Patients and Their Unaffected Siblings. *Front Neurosci* 9:495. doi: 10.3389/fnins.2015.00495
- Fastenau PS, Denburg NL, Hufford BJ (1999) Adult norms for the rey-osterrieth complex figure test and for supplemental recognition and matching trials from the extended complex figure test. *Clin Neuropsychol* 13:30–47. doi: 10.1076/clin.13.1.30.1976
- Fazekas F, Chawluk JB, Zimmerma A, June M (1987) MR Signal Abnormalities at 1.5 T in Alzheimer's Dementia and Normal Aging deficiency. *Am J Neuroradiol* 149:351–356
- Fernández-Miranda JC, Rhoton AL, Álvarez-Linera J, et al (2008) Three-dimensional microsurgical and tractographic anatomy of the white matter of the human brain. *Neurosurgery* 62:. doi: 10.1227/01.NEU.0000297076.98175.67
- Folstein MF, Folstein SE, McHugh PR. 'Minimal state'. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; 12: 189–98.
- Forkel SJ, Thiebaut de Schotten M, Kawadler JM, et al (2012) The anatomy of fronto-occipital connections from early blunt dissections to contemporary tractography. *Cortex* 56:73–84. doi: 10.1016/j.cortex.2012.09.005
- Foss MP, Bastos-formigheri DSM, Speciali JG (2010) Figuras complexas de Rey para idosos. *Aval Psicol*. doi: 10.5579/rnl.2016.0276
- Godefroy O, Jeannerod M, Allain P, Le Gall D (2008) Frontal lobe, executive functions and cognitive control. *Rev Neurol (Paris)* 164:. doi: 10.1016/S0035-3787(08)73302-2
- Gonçalves APB, Tarrasconi MA, Holz MR, et al (2019) Cognitive flexibility and inhibition in single-versus multiple-domain mild cognitive impairment: A comparative and discriminative analysis. *Psychol Neurosci* 12:209–223. doi: 10.1037/pne0000160
- Gong G, Jiang T, Zhu C, et al (2005) Asymmetry analysis of cingulum based on scale-invariant parameterization by diffusion tensor imaging. *Hum Brain Mapp* 24:92–98. doi: 10.1002/hbm.20072
- GREFEX (Azouvi P, Didic-Hamel Cooke M, Fluchaire I, Godefroy O, Hoclet E, Le Gall D, Marié RM, Meulemans T, Naegelé B, Peres B PB (2017) L'évaluation des fonctions executives en pratique clinique. *Rev Neuropsychol* 11:383–434
- Grèzes J, Valabrègue R, Gholipour B, Chevallier C (2014) A direct amygdala-motor pathway for emotional displays to influence action: A diffusion tensor imaging study. *Hum Brain Mapp* 35:5974–5983. doi: 10.1002/hbm.22598
- Gunning FM, Cheng J, Murphy CF, et al (2009) Anterior cingulate cortical volumes and treatment remission of geriatric depression. *Int J Geriatr Psychiatry* 24:829–836. doi: 10.1002/gps.2290

- Gunning-dixon FM, Brickman AM, Cheng JC, Alexopoulos GS (2009) Aging of cerebral white matter : a review of MRI findings. 109–117. doi: 10.1002/gps
- Haber SN (2016) Corticostriatal circuitry. *Neurosci 21st Century From Basic to Clin Second Ed* 1721–1741. doi: 10.1007/978-1-4939-3474-4_135
- Haggard P (2008) Human volition: Towards a neuroscience of will. *Nat Rev Neurosci* 9:934–946. doi: 10.1038/nrn2497
- Hamdan AC, Hamdan EMLR (2009) Effects of age and education level on the Trail Making Test in a healthy Brazilian sample. *Psychol Neurosci* 2:199–203. doi: 10.3922/j.psns.2009.2.012
- Hara T, Abo M, Kakita K, et al (2016) Does a combined intervention program of repetitive transcranial magnetic stimulation and intensive occupational therapy affect cognitive function in patients with post-stroke upper limb hemiparesis? *Neural Regen Res* 11:1932. doi: 10.4103/1673-5374.197134
- Hau, J., Aljawad, S., Baggett, N., Fishman, I., Carper, R. A., & Müller, R. (2019). The cingulum and cingulate U-fibers in children and adolescents with autism spectrum disorders. *Human Brain Mapping*. doi:10.1002/hbm.24586
- Hecht EE, Gutman DA, Bradley BA, et al (2015) Virtual dissection and comparative connectivity of the superior longitudinal fasciculus in chimpanzees and humans. *Neuroimage* 108:124–137. doi: 10.1016/j.neuroimage.2014.12.039
- Heilbronner SR, Haber SN (2014) Frontal Cortical and Subcortical Projections Provide a Basis for Segmenting the Cingulum Bundle: Implications for Neuroimaging and Psychiatric Disorders. *J Neurosci* 34:10041–10054. doi: 10.1523/JNEUROSCI.5459-13.2014
- Hilgetag C.C., Barbas H. (2006) Role of mechanical factors in the morphology of the primate cerebral cortex. *PLoS Comput. Biol.* 2:e22101371/journal.pcbi.0020022
- Hiroshima S, Anei R, Murakami N, Kamada K (2014) Functional localization of the supplementary motor area. *Neurol Med Chir (Tokyo)* 54:511–520. doi: 10.2176/nmc.oa2012-0321
- Hoffstaedter F, Grefkes C, Caspers S, et al (2014) The role of anterior midcingulate cortex in cognitive motor control: Evidence from functional connectivity analyses. *Hum Brain Mapp* 35:2741–2753. doi: 10.1002/hbm.22363
- Hoffstaedter F, Grefkes C, Zilles K, Eickhoff SB (2013) The “what” and “when” of self-initiated movements. *Cereb Cortex* 23:520–530. doi: 10.1093/cercor/bhr391
- Hoffstaedter F, Grefkes C, Zilles K, Eickhoff SB (2013) The “what” and “when” of self-initiated movements. *Cereb Cortex* 23:520–530. doi: 10.1093/cercor/bhr391
- Horn A, Ostwald D, Reiser M, Blankenburg F (2014) The structural-functional connectome and the default mode network of the human brain. *Neuroimage* 102:142–151. doi: 10.1016/j.neuroimage.2013.09.069
- Howells H, Thiebaut De Schotten M, Dell ’acqua F, et al (2018) Frontoparietal Tracts Linked to Lateralized Hand Preference and Manual Specialization. *Cereb Cortex* 1–13. doi: 10.1093/cercor/bhy040
- Husain, M., & Schott, J. M. (2016). *Oxford textbook of cognitive neurology and dementia*. Oxford, United Kingdom: Oxford University Press.
- Insausti R, Muñoz-López M, Insausti AM, Artacho-Pérula E (2017) The Human Periallocortex: Layer Pattern in Presubiculum, Parasubiculum and Entorhinal Cortex. A Review. *Front Neuroanat* 11:1–10. doi: 10.3389/fnana.2017.00084
- ISCO-08 (2008). INTERNATIONAL STANDARD CLASSIFICATION OF OCCUPATIONS
- Jang, S. H., & Hong, J. H. (2012). The anatomical characteristics of superior longitudinal fasciculus I in human brain: Diffusion tensor tractography study. *Neuroscience Letters*, 506, 146-148.
- Jankowski MM, Ronnqvist KC, Tsanov M, et al (2013) The anterior thalamus provides a subcortical circuit supporting memory and spatial navigation. *Front Syst Neurosci* 7:45. doi: 10.3389/fnsys.2013.00045
- Jbabdi S, Lehman JF, Haber SN, Behrens TE (2013) Human and Monkey Ventral Prefrontal Fibers Use the Same Organizational Principles to Reach Their Targets: Tracing versus Tractography. *J Neurosci* 33:3190–3201. doi: 10.1523/JNEUROSCI.2457-12.2013
- Jellison BJ, Field a. S, Medow J, et al (2004) Diffusion tensor imaging of cerebral white matter: a pictorial review of physics, fiber tract

- anatomy, and tumor imaging patterns. *Am J Neuroradiol* 25:356–69. doi: 10.1038/nrn2776
- Jeremy D. Schmahmann DNP (2006) Fiber pathways of the brain
- Jeurissen B, Descoteaux M, Mori S, Leemans A (2019) Diffusion MRI fiber tractography of the brain. *NMR Biomed* 32:e3785. doi: 10.1002/nbm.3785
- Jones DK (2008) Studying connections in the living human brain with diffusion MRI. *Cortex* 44:936–952. doi: 10.1016/j.cortex.2008.05.002
- Jones DK, Cercignani M (2010) Twenty-five pitfalls in the analysis of diffusion MRI data. *NMR Biomed* 23:803–820. doi: 10.1002/nbm.1543
- Jones DK, Cercignani M (2010) Twenty-five pitfalls in the analysis of diffusion MRI data. *NMR Biomed* 23:803–820. doi: 10.1002/nbm.1543
- Jones DK, Christiansen KF, Chapman RJ, Aggleton JP (2013) Distinct subdivisions of the cingulum bundle revealed by diffusion MRI fibre tracking: Implications for neuropsychological investigations. *Neuropsychologia* 51:67–78. doi: 10.1016/j.neuropsychologia.2012.11.018
- Jones DK, Christiansen KF, Chapman RJ, Aggleton JP (2013a) Distinct subdivisions of the cingulum bundle revealed by diffusion MRI fibre tracking: Implications for neuropsychological investigations. *Neuropsychologia* 51:67–78. doi: 10.1016/j.neuropsychologia.2012.11.018
- Jones DK, Knösche TR, Turner R (2013b) White matter integrity, fiber count, and other fallacies: The do's and don'ts of diffusion MRI. *Neuroimage* 73:239–254. doi: 10.1016/j.neuroimage.2012.06.081
- Kalafat M, Hugonot-Diener L, Poitrenaud J (2003) The Mini Mental State (MMS): French standardization and normative data [Standardisation et étalonnage français du “Mini Mental State” (MMS) version GRÉCO]
- Kamali A, Flanders AE, Brody J, et al (2014) Tracing superior longitudinal fasciculus connectivity in the human brain using high resolution diffusion tensor tractography. *Brain Struct Funct* 219:269–81. doi: 10.1007/s00429-012-0498-y
- Kamali, A., Flanders, A. E., Brody, J., Hunter, J. V., & Hasan, K. M. (2014). Tracing Superior Longitudinal Fasciculus Connectivity in the Human Brain using High Resolution Diffusion Tensor Tractography. *Brain Structure and Function*, 219, 1-21.
- Kantarci K, Senjem ML, Avula R, et al (2011) Diffusion tensor imaging and cognitive function in older adults with no dementia. *Neurology* 77:26–34. doi: 10.1212/WNL.0b013e31822313dc
- Keedwell PA, Chapman R, Christiansen K, et al (2012) Cingulum white matter in young women at risk of depression: The effect of family history and anhedonia. *Biol Psychiatry* 72:296–302. doi: 10.1016/j.biopsych.2012.01.022
- Keser Z, Hasan KM, Mwangi B, et al (2017a) Limbic Pathway Correlates of Cognitive Impairment in Multiple Sclerosis. *J Neuroimaging* 27:37–42. doi: 10.1111/jon.12381
- Keser Z, Hasan KM, Mwangi B, et al (2017b) Diffusion Tensor Imaging-Defined Sulcal Enlargement Is Related to Cognitive Impairment in Multiple Sclerosis. *J Neuroimaging* 27:312–317. doi: 10.1111/jon.12406
- Khalsa S, Mayhew SD, Chechlacz M, et al (2014) NeuroImage The structural and functional connectivity of the posterior cingulate cortex : Comparison between deterministic and probabilistic tractography for the investigation of structure – function relationships. *Neuroimage* 102:118–127. doi: 10.1016/j.neuroimage.2013.12.022
- Kim, S. J., Jeong, D.-U., Sim, M. E., Bae, S. C., Chung, A., Kim, M. J., ... Lyoo, I. K. (2006). Asymmetrically Altered Integrity of Cingulum Bundle in Posttraumatic Stress Disorder. *Neuropsychobiology*, 54(2), 120–125. doi:10.1159/000098262
- Klingler, J. (1935). Erleichterung der makroskopischen Praeparation des Gehirns durch den Gefrierprozess. *Schweizer Archiv für Neurologie und Psychiatrie*, 36, 247-256.
- Klingler, J., & Gloor, P. (1960). The connections of the amygdala and of the anterior temporal cortex in the human brain. *The Journal of comparative neurology*, 115, 333-369.
- Kochhann R, Cerveira MO, Godinho C, et al (2009) Evaluation of Mini-Mental State Examination scores according to different age and education strata, and sex, in a large Brazilian healthy sample. *Dement Neuropsychol* 3:88–93. doi: 10.1590/s1980-57642009dn30200004
- Koutsarnakis, C., Liakos, F., Kalyvas, A. V., Sakas, D. E., & Stranjalis, G. (2015). A

- Laboratory Manual for Stepwise Cerebral White Matter Fiber Dissection. *World Neurosurg*, 84, 483-493.
- Laver GD, Burke DM (1993) Why do semantic priming effects increase in old age? A meta-analysis. *Psychol Aging* 8:34–43. doi: 10.1037/0882-7974.8.1.34
- Lawes INC, Barrick TR, Murugam V, et al (2008) Atlas-based segmentation of white matter tracts of the human brain using diffusion tensor tractography and comparison with classical dissection. *Neuroimage* 39:62–79. doi: 10.1016/j.neuroimage.2007.06.041
- Lawton MP, Brody EM (1969) Assessment of Older People : Self-Maintaining and Instrumental Activities of Daily Living 1. *Gerontologist* 9:179–186
- Le Bihan, D., Breton, E., Lallemand, D., Grenier, P., Cabanis, E., & Laval-Jeantet, M. (1986). MR imaging of intravoxel incoherent motions: application to diffusion and perfusion in neurologic disorders. *Radiology*, 161, 401-417.
- Le Bihan, D., Mangin, J. F., Poupon, C., Clark, C. a., Pappata, S., Molko, N., & Chabriat, H. (2001). Diffusion tensor imaging: concepts and applications. *Journal of magnetic resonance imaging : JMIRI*, 13, 534-546.
- Lebel C, Beaulieu C (2011) Longitudinal Development of Human Brain Wiring Continues from Childhood into Adulthood. *J Neurosci* 31:10937–10947. doi: 10.1523/JNEUROSCI.5302-10.2011
- Leichnetz GR (2001) Connections of the Medial Posterior Parietal Cortex (Area 7m) in the Monkey. *Anat Rec* 236:215–236
- Lezak M (2012) *NEUROPSYCHOLOGICAL ASSESSMENT*. OXFORD
- Li W, Qin W, Liu H, et al (2013) Subregions of the human superior frontal gyrus and their connections. *Neuroimage* 78:46–58. doi: 10.1016/j.neuroimage.2013.04.011
- Liberati A, Altman DG, Tetzlaff J, et al (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 6:e1000100. doi: 10.1371/journal.pmed.1000100
- Lima CF, Krishnan S, Scott SK (2016) Roles of Supplementary Motor Areas in Auditory Processing and Auditory Imagery. *Trends Neurosci* 39:527–542. doi: 10.1016/j.tins.2016.06.003
- Locke S (1961) Limbic Nuclei of Thalamus and Connections of Limbic Cortex. *Arch Neurol* 4:355. doi: 10.1001/archneur.1961.00450100003002
- Ludwig, E., & Klingler, J. (1956). *Atlas Cerebri Humani*.
- Madhavan KM, McQueeney T, Howe SR, et al (2014) Superior longitudinal fasciculus and language functioning in healthy aging. *Brain Res* 1562:11–22. doi: 10.1016/j.brainres.2014.03.012
- Mahieux-Laurent F, Fabre C, Galbrun E, Dubrulle A, Moroni C. (2009). Validation d'une batterie brève d'évaluation des praxies gestuelles pour consultation Mémoire. Évaluation chez 419 témoins, 127 patients atteints de troubles cognitifs légers et 320 patients atteints d'une démence. *Rev. Neurol. (Paris)*; 165: 560-567.
- Maier S, Szalkowski A, Kamphausen S, et al (2012) Clarifying the Role of the Rostral dmPFC/dACC in Fear/Anxiety: Learning, Appraisal or Expression? *PLoS One* 7:. doi: 10.1371/journal.pone.0050120
- Maier-Hein KH, Neher PF, Houde J-C, et al (2017) The challenge of mapping the human connectome based on diffusion tractography. *Nat Commun* 8:1349. doi: 10.1038/s41467-017-01285-x
- Makris N, Kennedy DN, McInerney S, et al (2005) Segmentation of subcomponents within the superior longitudinal fascicle in humans: A quantitative, in vivo, DT-MRI study. *Cereb Cortex* 15:854–869. doi: 10.1093/cercor/bhh186
- Maldonado IL, Mandonnet E, Duffau H (2012) Dorsal Fronto-Parietal Connections of the Human Brain: A Fiber Dissection Study of Their Composition and Anatomical Relationships. *Anat Rec* 295:187–195. doi: 10.1002/ar.21533
- Maldonado IL, Parente de Matos V, Castro Cuesta TA, et al (2020) The human cingulum: From the limbic tract to the connectionist paradigm. *Neuropsychologia* 144:107487. doi: 10.1016/j.neuropsychologia.2020.107487
- Maldonado, I. L., de Champfleury, N. M., Velut, S., Destrieux, C., Zemmoura, I., & Duffau, H. (2013). Evidence of a middle longitudinal fasciculus in the human brain from fiber dissection. *J Anat*, 223, 38-45.
- Maldonado, I. L., Mandonnet, E., & Duffau, H. (2012). Dorsal fronto-parietal connections of the

- human brain: a fiber dissection study of their composition and anatomical relationships. *Anat Rec (Hoboken)*.
- Mandonnet E, Sarubbo S, Petit L, et al (2018) The Nomenclature of Human White Matter Association Pathways : Proposal for a Systematic Taxonomic Anatomical Classification. *Front Neuroanat* 12:. doi: 10.3389/fnana.2018.00094
- Mandonnet, E., Martino, J., Sarubbo, S., Corrivetti, F., Bouazza, S., Bresson, D., Duffau, H., & Froelich, S. (2017). Neuronavigated Fiber Dissection with Pial Preservation: Laboratory Model to Simulate Opercular Approaches to Insular Tumors. *World Neurosurg*, 98, 239-242.
- Mandonnet, E., Sarubbo, S., & Petit, L. (2018). The Nomenclature of Human White Matter Association Pathways: Proposal for a Systematic Taxonomic Anatomical Classification. *Front Neuroanat*, 12, 94.
- Mars RB, Jbabdi S, Reilly JXO, et al (2011) Diffusion-Weighted Imaging Tractography-Based Parcellation of the Human Parietal Cortex and Comparison with Human and Macaque Resting-State Functional Connectivity. *31:4087–4100*. doi: 10.1523/JNEUROSCI.5102-10.2011
- Mars RB, O’Muircheartaigh J, Folloni D, et al (2019) Concurrent analysis of white matter bundles and grey matter networks in the chimpanzee. *Brain Struct Funct* 224:1021–1033. doi: 10.1007/s00429-018-1817-8
- Mars RB, Sotiropoulos SN, Passingham RE, et al (2018) Whole brain comparative anatomy using connectivity blueprints. *Elife* 7:. doi: 10.7554/eLife.35237
- Marshal F. Folstein SEF and PRMch (1975) Mini-Mental State. A practical state method for grading the cognitive state of petients for the clinician. *J phychiat Res* 12:189–198
- Martino, J., da Silva-Freitas, R., Caballero, H., Marco de Lucas, E., Garcia-Porrero, J. A., & Vazquez-Barquero, A. (2013). Fiber dissection and diffusion tensor imaging tractography study of the temporoparietal fiber intersection area. *Neurosurgery*, 72, 87-97; discussion 97-88.
- Metzler-Baddeley C, Jones DK, Steventon J, et al (2012) Cingulum microstructure predicts cognitive control in older age and mild cognitive impairment. *J Neurosci* 32:17612–17619. doi: 10.1523/JNEUROSCI.3299-12.2012
- Metzler-Baddeley C, Jones DK, Steventon J, et al (2012)
- Migliaccio R, Bouhali F, Rastelli F, et al (2014) Damage to the medial motor system in stroke patients with motor neglect. *Front Hum Neurosci* 8:1–9. doi: 10.3389/fnhum.2014.00408
- Misdraji EL, Gass CS (2010) The Trail Making Test and its neurobehavioral components. *J Clin Exp Neuropsychol* 32:159–163. doi: 10.1080/13803390902881942
- Mogenson GJ, Jones DL, Yim CY (1980) From motivation to action: Functional interface between the limbic system and the motor system. *Prog Neurobiol* 14:69–97. doi: 10.1016/0301-0082(80)90018-0
- Mollink J, Kleinnijenhuis M, Cappellen van Walsum AM van, et al (2017) Evaluating fibre orientation dispersion in white matter: Comparison of diffusion MRI, histology and polarized light imaging. *Neuroimage* 157:561–574. doi: 10.1016/j.neuroimage.2017.06.001
- Morgane PJ, Galler JR, Mokler DJ (2005) A review of systems and networks of the limbic forebrain/limbic midbrain. *Prog Neurobiol* 75:143–160. doi: 10.1016/j.pneurobio.2005.01.001
- Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry* 1979; 134: 382-9.
- Mori S, Aggarwal M (2014) In vivo magnetic resonance imaging of the human limbic white matter. *Front Aging Neurosci* 6:1–6. doi: 10.3389/fnagi.2014.00321
- Mufson EJ, Pandya DN (1984) Some observations on the course and composition of the cingulum bundle in the rhesus monkey. *J Comp Neurol* 225:31–43. doi: 10.1002/cne.902250105
- Mufson EJ, Pandya DN (1984) Some observations on the course and composition of the cingulum bundle in the rhesus monkey. *J Comp Neurol* 225:31–43. doi: 10.1002/cne.902250105
- Mukherjee P, Berman JI, Chung SW, et al (2008) Diffusion tensor MR imaging and fiber tractography: Theoretic underpinnings. *Am J Neuroradiol* 29:632–641. doi: 10.3174/ajnr.A1051
- Nagae LM, Zarnow DM, Blaskey L, et al (2012) Elevated Mean Diffusivity in the Left Hemisphere Superior Longitudinal Fasciculus in Autism Spectrum Disorders Increases with More Profound Language Impairment. *Am J*

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

- Neuroradiol 33:1720–1725. doi: 10.3174/ajnr.A3037
- Nieuwenhuys, R., Voogd, J., & Huijzen, C. v. (2008). *The human central nervous system* (4th ed.). New York: Springer.
- Nilsson LG (2003) Memory function in normal aging. *Acta Neurol Scand Suppl* 107:7–13. doi: 10.1212/wnl.42.2.396
- O'Donnell, L. J., & Westin, C.-F. (2011). An Introduction to Diffusion Tensor Image Analysis. *Neurosurgery Clinics of North America*, 22(2), 185–196. doi:10.1016/j.nec.2010.12.004
- Oldfield RC (1971). The assessment the and analysis of handedness: edinburgh inventory. *Neuropsychologia*, 9:97–113
- Olry, R., & Nicolay, X. (1994). [From Paul Broca to the long-term potentiation: the difficulties in confirming a limbic identity]. *Hist Sci Med*, 28, 199-203.
- Opdebeeck C, Martyr A, Clare L (2015) Cognitive reserve and cognitive function in healthy older people: a meta-analysis. *Aging , Neuropsychol , Cogn* 37–41. doi: 10.1080/13825585.2015.1041450
- Palacios EM, Sala-Llonch R, Junque C, et al (2013) Resting-State Functional Magnetic Resonance Imaging Activity and Connectivity and Cognitive Outcome in Traumatic Brain Injury. *JAMA Neurol* 70:1–7. doi: 10.1001/jamaneurol.2013.38
- Pandya DN, Van Hoesen GW, Mesulam MM (1981) Efferent connections of the cingulate gyrus in the rhesus monkey. *Exp Brain Res* 42:319–330. doi: 10.1007/BF00237497
- Pandya MP and DN (2006) Efferent Association Pathways Originating in the Caudal Prefrontal Cortex in the Macaque Monkey. *J Comp Neurol* 498:227–251. doi: 10.1002/cne
- Papez, J. (1937). A proposed mechanism of emotion. *Archives of Neurology And Psychiatry*, 38, 725.
- Parker GJM, Luzzi S, Alexander DC, et al (2005) Lateralization of ventral and dorsal auditory-language pathways in the human brain. *Neuroimage* 24:656–666. doi: 10.1016/j.neuroimage.2004.08.047
- Pascalau, R., Popa Stanila, R., Sfrangeu, S., & Szabo, B. (2018). Anatomy of the Limbic White Matter Tracts as Revealed by Fiber Dissection and Tractography. *World Neurosurg*, 113, e672-e689.
- Pasquier F, Grymonprez L, Lebert F, Van der Linden M (2001) Memory Impairment Differs in Frontotemporal Dementia and Alzhemier's Disease. *Neurocase* 7:161–171. doi: 10.1093/neucas/7.2.161
- Petrides M, Pandya D (1984) Projections to the frontal lobe from the posterior-parietal region in the rhesus monkey. *J Comp Neurol* 228:105–116
- Petrides M, Tomaiuolo F, Yeterian EH, Pandya DN (2012) The prefrontal cortex: Comparative architectonic organization in the human and the macaque monkey brains. *Cortex* 48:46–57. doi: 10.1016/j.cortex.2011.07.002
- Petrides, M., & Pandya, D. N. (1984). Projections to the frontal lobe from the posterior-parietal region in the rhesus monkey. *The Journal of Comparative Neurology*, 228, 105-116.
- Procyk E, Wilson CRE, Stoll FM, et al (2016) Midcingulate Motor Map and Feedback Detection: Converging Data from Humans and Monkeys. *Cereb Cortex* 26:467–476. doi: 10.1093/cercor/bhu213
- Pruessner JC, Dedovic K, Khalili-Mahani N, et al (2008) Deactivation of the Limbic System During Acute Psychosocial Stress: Evidence from Positron Emission Tomography and Functional Magnetic Resonance Imaging Studies. *Biol Psychiatry* 63:234–240. doi: 10.1016/j.biopsych.2007.04.041
- Pruessner JC, Dedovic K, Khalili-Mahani N, et al (2008) Deactivation of the Limbic System During Acute Psychosocial Stress: Evidence from Positron Emission Tomography and Functional Magnetic Resonance Imaging Studies. *Biol Psychiatry* 63:234–240. doi: 10.1016/j.biopsych.2007.04.041
- Raichle ME, MacLeod AM, Snyder AZ, et al (2001) A default mode of brain function. *Proc Natl Acad Sci U S A* 98:676–82. doi: 10.1073/pnas.98.2.676
- Raj A, Hess C, Mukherjee P (2011) Spatial HARDI: Improved visualization of complex white matter architecture with Bayesian spatial regularization. *Neuroimage* 54:396–409. doi: 10.1016/j.neuroimage.2010.07.040
- Régis J., Mangin J.-F., Ochiai T., Frouin V., Rivière D., Cachia A., et al. . (2005). “Sulcal root” generic model: a hypothesis to overcome the variability of the human cortex folding patterns. *Neurol. Med. Chir. (Tokyo)* 45, 1–17.10.2176/nmc.45.1

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

- Reitan RM (1955) The relation of the Trail Making Test to organic brain damage. *J Consult Psychol* 19:393–394. doi: 10.1037/h0044509
- Rey A (1959) Manuel du test de copie d'une figure complexe de A. Rey. Paris Les Ed du Cent Psychol Appl
- Ribas GC (2010) The cerebral sulci and gyri. *Neurosurg Focus* 28:E2. doi: 10.3171/2009.11.FOCUS09245
- Rilling JK, Glasser MF, Preuss TM, et al (2008) The evolution of the arcuate fasciculus revealed with comparative DTI. *Nat Neurosci* 11:426–428. doi: 10.1038/nn2072
- Ritchie SJ, Bastin ME, Tucker-Drob EM, et al (2015) Coupled changes in brain white matter microstructure and fluid intelligence in later life. *J Neurosci* 35:8672–8682. doi: 10.1523/JNEUROSCI.0862-15.2015
- Rolls ET (2014) Limbic systems for emotion and for memory, but no single limbic system. *Cortex* 62:119–157. doi: 10.1016/j.cortex.2013.12.005
- Rolls ET (2015) Limbic systems for emotion and for memory, but no single limbic system. *Cortex* 62:119–157. doi: 10.1016/j.cortex.2013.12.005
- Roxo MR, Franceschini PR, Zubaran C, et al (2011) The limbic system conception and its historical evolution. *ScientificWorldJournal* 11:2428–41. doi: 10.1100/2011/157150
- Sarkar S, Craig MC, Catani M, et al (2012) Frontotemporal white-matter microstructural abnormalities in adolescents with conduct disorder: a diffusion tensor imaging study. *Psychol Med* 1–11. doi: 10.1017/S003329171200116X
- Scarpina F, Tagini S (2017) The stroop color and word test. *Front Psychol* 8:1–8. doi: 10.3389/fpsyg.2017.00557
- Schmahman JD, Pandya DN (2006) Fiber pathways of the brain
- Schmahmann JD, Pandya DN, Wang R, et al (2007) Association fibre pathways of the brain: Parallel observations from diffusion spectrum imaging and autoradiography. *Brain* 130:630–653. doi: 10.1093/brain/awl359
- Seghier ML (2008) Laterality index in functional MRI: methodological issues. *Magn Reson Imaging* 26:594–601. doi: 10.1016/j.mri.2007.10.010
- Shah A, Jhavar SS, Goel A (2012) Analysis of the anatomy of the Papez circuit and adjoining limbic system by fiber dissection techniques. *J Clin Neurosci* 19:289–298. doi: 10.1016/j.jocn.2011.04.039
- Shah A, Jhavar SS, Goel A (2012) Analysis of the anatomy of the Papez circuit and adjoining limbic system by fiber dissection techniques. *J Clin Neurosci* 19:289–298. doi: 10.1016/j.jocn.2011.04.039
- Shah, A., Jhavar, S. S., & Goel, A. (2012). Analysis of the anatomy of the Papez circuit and adjoining limbic system by fiber dissection techniques. *J Clin Neurosci*, 19, 289–298.
- Sibilia F, Kehoe EG, Farrell D, et al (2017) Aging-Related Microstructural Alterations Along the Length of the Cingulum Bundle. *Brain Connect* 7:366–372. doi: 10.1089/brain.2017.0493
- Skadorwa T, Kunicki J, Nauman P, Cizek B (2009) Image-guided dissection of human white matter tracts as a new method of modern neuroanatomical training. *Folia Morphol (Warsz)* 68:135–139
- Spreen, O., and Strauss E (1998) A compendium of neuropsychological tests: Administration, norms, and commentary, 2nd ed. New York: Oxford University Press.
- Spreng RN, Sepulcre J, Turner GR, et al (2013) Intrinsic Architecture Underlying the Relations among the Default, Dorsal Attention, and Frontoparietal Control Networks of the Human Brain. *J Cogn Neurosci* 25:74–86. doi: 10.1162/jocn_a_00281
- Stroop JR (1935) Studies of interference in serial verbal reactions. *J Exp Psychol* 18:643–662. doi: 10.1037/h0054651
- Strotzer M (2009) One century of brain mapping using Brodmann areas. *Clin Neuroradiol* 19:179–186. doi: 10.1007/s00062-009-9002-3
- Suarez RO, Whalen S, Nelson AP, et al (2009) Threshold-independent functional MRI determination of language dominance: A validation study against clinical gold standards. *Epilepsy Behav* 16:288–297. doi: 10.1016/j.yebeh.2009.07.034
- Tabassum Khan N (2017) Autoradiography: Detection and Analysis of Radioactive Entities. *J Biom Biostat* 08:1–2. doi: 10.4172/2155-6180.1000361
- Tamnes CK, Østby Y, Fjell AM, et al (2010) Brain Maturation in Adolescence and Young Adulthood: Regional Age-Related Changes in Cortical Thickness and White Matter Volume
- Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64–88
<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

- and Microstructure. *Cereb Cortex* 20:534–548. doi: 10.1093/cercor/bhp118
- Ten Donkelaar HJ, Kachlík D, Tubbs RS (2018) *An Illustrated Terminologia Neuroanatomica*. Springer International Publishing, Cham
- Thiebaut de Schotten M, Dell’Acqua F, Valabregue R, Catani M (2012) Monkey to human comparative anatomy of the frontal lobe association tracts. *Cortex* 48:82–96. doi: 10.1016/j.cortex.2011.10.001
- Thiebaut de Schotten M, Dell’Acqua F, Valabregue R, Catani M (2012) Monkey to human comparative anatomy of the frontal lobe association tracts. *Cortex* 48:82–96. doi: 10.1016/j.cortex.2011.10.001
- Thiebaut de Schotten M, ffytche DH, Bizzi A, et al (2011) Atlasing location, asymmetry and inter-subject variability of white matter tracts in the human brain with MR diffusion tractography. *Neuroimage* 54:49–59. doi: 10.1016/j.neuroimage.2010.07.055
- Thiebaut M, Schotten D, Dell F, et al (2012) Monkey to human comparative anatomy of the frontal lobe association tracts. *CORTEX* 48:82–96. doi: 10.1016/j.cortex.2011.10.001
- Tombaugh TN (2004) Trail Making Test A and B: Normative data stratified by age and education. *Arch Clin Neuropsychol* 19:203–214. doi: 10.1016/S0887-6177(03)00039-8
- Tournier J-D, Calamante F, Gadian DG, Connelly A (2004) Direct estimation of the fiber orientation density function from diffusion-weighted MRI data using spherical deconvolution. *Neuroimage* 23:1176–1185. doi: 10.1016/j.neuroimage.2004.07.037
- Tremblay M-P, St-Hilaire A, Hudon C, Macoir J (1999) Fiche descriptive de test : Epreuve de rappel libre/rappel indicé à 16items. *Réseau Québécois Rech sur le Vieillessement* 93:161–164. doi: 10.1016/S0035
- Tremblay MP, Potvin O, Callahan BL, et al (2015) Normative data for the rey-osterrieth and the taylor complex figure tests in Quebec-French people. *Arch Clin Neuropsychol* 30:78–87. doi: 10.1093/arclin/acu069
- Ugur Türe, M. Gazi Yasargil, Allan H. Friedman OA-M (2000) *Fiber Dissection Technique : Lateral Aspect of the Brain*. *Neurosurgery* 47:417–427
- van den Heuvel M, Mandl R, Luijckes J, Hulshoff Pol H (2008) Microstructural Organization of the Cingulum Tract and the Level of Default Mode Functional Connectivity. *J Neurosci* 28:10844–10851. doi: 10.1523/JNEUROSCI.2964-08.2008
- Van der Linden, M., Coyette, F., Poitrenaud, J., Kalafat, M., Calacis, F., Wyns C, Adam, S. et les membres du G (2004) (2004) L’épreuve de rappel libre/rappel indicé à 16 items (RL/RI 16). In: Van der Linden M, Adam S, Agniel A et les membres du GREMEM, editors. *L’évaluation des troubles de la mémoire*. *Neuropsychologie* 1–19
- Verhaeghen P, De Meersman L (1998) Aging and the Stroop effect: A meta-analysis. *Psychol Aging* 13:120–126. doi: 10.1037/0882-7974.13.1.120
- Vogt B (2009) in *Cingulate neurobiology and disease, Architecture, neurocytology, and comparative organization of monkey and human cingulate cortices*, ed Vogt B (Oxford UP, New York)
- Vogt BA, Finch DM, Olson CR (1992) Functional heterogeneity in cingulate cortex: The anterior executive and posterior evaluative regions. *Cereb Cortex* 2:435–443. doi: 10.1093/cercor/2.6.435-a
- Vogt BA, Finch DM, Olson CR (1992) Functional heterogeneity in cingulate cortex: The anterior executive and posterior evaluative regions. *Cereb Cortex* 2:435–443. doi: 10.1093/cercor/2.6.435-a
- Vogt BA, Vogt L, Farber NB, Bush G (2009) Architecture and Neurocytology of Monkey Cingulate Gyrus. *J Comp Neurol* 485:218–239. doi: 10.1002/cne.20512. Architecture
- Von Bonin and Percival Bailey (1947) *The neocortex of Macaca mulatta*. University of Illinois Press
- Vos SB, Aksoy M, Han Z, et al (2016) Trade-off between angular and spatial resolutions in in vivo fiber tractography. *Neuroimage* 129:117–132. doi: 10.1016/j.neuroimage.2016.01.011
- Vos SB, Viergever MA, Leemans A (2013) Multi-fiber tractography visualizations for diffusion MRI data. *PLoS One* 8:1–17. doi: 10.1371/journal.pone.0081453
- Wakana S, Jiang H, Nagae-Poetscher LM, et al (2004) Fiber tract-based atlas of human white matter anatomy. *Radiology* 230:77–87. doi: 10.1148/radiol.2301021640
- Wang Q, Li W, Liu XS, et al (2014) NIH Public Access. 27:380–392. doi: 10.1016/j.molcel.2007.05.041.A

Maldonado, I. L.; Destrieux, C. Characterization of limbic system connectivity through fiber dissection and diffusion imaging techniques, *LE STUDIUM Multidisciplinary Journal*, 2019, 3, 64-88

<https://doi.org/10.34846/le-studium.170.02.fr.01-2019>

- Wang Y, Hao L, Zhang Y, et al (2019) Entorhinal cortex volume, thickness, surface area and curvature trajectories over the adult lifespan. *Psychiatry Res Neuroimaging* 292:47–53. doi: 10.1016/j.psychresns.2019.09.002
- Wang, X., Pathak, S., Stefanescu, L., Yeh, F.-C., Li, S., & Fernandez-Miranda, J. C. (2016). Subcomponents and connectivity of the superior longitudinal fasciculus in the human brain. *Brain Structure and Function*, 221, 2075-2092.
- Wedeen VJ, Wang RP, Schmahmann JD, et al (2008) Diffusion spectrum magnetic resonance imaging (DSI) tractography of crossing fibers. *Neuroimage* 41:1267–1277. doi: 10.1016/j.neuroimage.2008.03.036
- Weschler D. (2000). Wechsler Adult Intelligence Scale (WAIS-III). Paris: Editions du Centre de Psychologie Appliquée.
- Wendelken C, Ferrer E, Whitaker KJ, Bunge SA (2016) Fronto-Parietal Network Reconfiguration Supports the Development of Reasoning Ability. *Cereb Cortex* 26:2178–2190. doi: 10.1093/cercor/bhv050
- Wirt RA, Hyman JM (2017) Integrating spatial working memory and remote memory: Interactions between the medial prefrontal cortex and hippocampus. *Brain Sci* 7:. doi: 10.3390/brainsci7040043
- Woolsey, T. A. (2016). Review on Cowan WM, Gottlieb DI, Hendrickson AE, Price JL, Woolsey TA. 1972. The autoradiographic demonstration of axonal connections in the central nervous system. *Brain Res* 37: 21–51. *Brain Research*, 1645, 25-27.
- Wu TC, Wilde EA, Bigler ED, et al (2010) Evaluating the relationship between memory functioning and cingulum bundles in acute mild traumatic brain injury using diffusion tensor imaging. *J Neurotrauma* 27:303–307. doi: 10.1089/neu.2009.1110
- Wu Y, Sun D, Wang Y, et al (2016) Segmentation of the Cingulum Bundle in the Human Brain: A New Perspective Based on DSI Tractography and Fiber Dissection Study. *Front Neuroanat* 10:84. doi: 10.3389/fnana.2016.00084
- Wu, Y., Sun, D., Wang, Y., Wang, Y., & Ou, S. (2016). Segmentation of the Cingulum Bundle in the Human Brain: A New Perspective Based on DSI Tractography and Fiber Dissection Study. *Frontiers in neuroanatomy*, 10, 84.
- Xie Y, Huo F, Tang J (2009) Cerebral cortex modulation of pain. *Acta Pharmacol Sin* 30:31–41. doi: 10.1038/aps.2008.14
- Yagmurlu, K., Middlebrooks, E. H., Tanriover, N., & Rhoton, A. L. (2016). Fiber tracts of the dorsal language stream in the human brain. *Journal of Neurosurgery*, 124, 1396-1405.
- Yeterian EH, Pandya DN, Tomaiuolo F, Petrides M (2012) The cortical connectivity of the prefrontal cortex in the monkey brain. *Cortex* 48:68–81. doi: 10.1016/j.cortex.2011.03.004
- Zemmoura I, Blanchard E, Raynal P, et al (2015) How Klingler’s dissection permits exploration of brain structural connectivity? An electron microscopy study of human white matter. *Brain Struct Funct* 221:2477–2486. doi: 10.1007/s00429-015-1050-7
- Zemmoura, I., Vons, J., Velut, S., & Destrieux, C. (2015). From Vesalius to tractography. *Journal of neurosurgical sciences*, 59, 309-325.
- Zhang S, Li CR (2012) NeuroImage Functional connectivity mapping of the human precuneus by resting state fMRI. *Neuroimage* 59:3548–3562. doi: 10.1016/j.neuroimage.2011.11.023
- Zhang Y, Schuff N, Jahng G-H, et al (2007) Diffusion tensor imaging of cingulum fibers in mild cognitive impairment and Alzheimer disease. *Neurology* 68:13–19. doi: 10.1212/01.wnl.0000250326.77323.01
- Zimmermann N, de Cardoso CO, Trentini CM, et al (2015) Normas Brasileiras Preliminares E A Investigação Dos Efeitos De Idade E Escolaridade No Desempenho Dos Testes Wisconsin De Classificação De Cartas Modificado, Stroop De Cores E Palavras E Dígitos Em Adultos. *Dement e Neuropsychol* 9:120–127. doi: 10.1590/1980-57642015DN92000006