



**HAL**  
open science

**Low-valent molybdenum carbonyl complexes as an entry to octahedral MoI<sub>3</sub>L<sub>3</sub> complexes. Synthesis and x-ray molecular structure of triiodotris(tetrahydrofuran)molybdenum**

F. Albert Cotton, Rinaldo Poli

► **To cite this version:**

F. Albert Cotton, Rinaldo Poli. Low-valent molybdenum carbonyl complexes as an entry to octahedral MoI<sub>3</sub>L<sub>3</sub> complexes. Synthesis and x-ray molecular structure of triiodotris(tetrahydrofuran)molybdenum. *Inorganic Chemistry*, 1987, 26 (10), pp.1514-1518. 10.1021/ic00257a011 . hal-03545065

**HAL Id: hal-03545065**

**<https://hal.science/hal-03545065>**

Submitted on 27 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Low-Valent Molybdenum Carbonyl Complexes as an Entry to Octahedral $\text{MoI}_3\text{L}_3$ Complexes. Synthesis and X-ray Molecular Structure of $\text{MoI}_3(\text{THF})_3$

F. Albert Cotton\* and Rinaldo Poli

Received November 13, 1986

$\text{MoI}_3(\text{THF})_3$  has been synthesized by iodination at room temperature of either  $\text{Mo}(\text{CO})_3(\eta^6\text{-C}_6\text{H}_5\text{CH}_3)$  or  $\text{Mo}_2\text{I}_4(\text{CO})_8$  in tetrahydrofuran (THF) as solvent. Crystals are orthorhombic, space group *Pbcn*, with  $a = 9.252$  (3) Å,  $b = 14.541$  (5) Å,  $c = 14.273$  (3) Å,  $V = 1920$  (2) Å<sup>3</sup>,  $Z = 4$ ,  $d_{\text{calcd}} = 2.397$  g cm<sup>-3</sup>, and  $R = 0.0501$  for 926 data having  $F_o^2 > 3\sigma(F_o^2)$ . The molecule is octahedral, in the meridional configuration. Substitution reactions have been carried out with  $\text{PR}_3$  ( $R = \text{Me, Et}$ ) to produce  $\text{MoI}_3(\text{PR}_3)_3$ . The direct interaction between  $\text{MoX}_3$  ( $X = \text{Cl, I}$ ) and  $\text{PR}_3$  ( $R = \text{Me, Et}$ ) in toluene has been studied. Reduction competes with coordinative addition to produce mixtures of  $\text{MoX}_3(\text{PR}_3)_3$  and  $\text{Mo}_2\text{X}_4(\text{PR}_3)_3$  for  $X = \text{Cl, I}$  and  $R = \text{Me}$  and mixtures of  $[\text{PHR}_3][\text{MoI}_4(\text{PR}_3)_2]$  and  $\text{Mo}_2\text{X}_4(\text{PR}_3)_4$  for  $X = \text{I}$  and  $R = \text{Et}$ . A possible mechanism for these reactions is discussed.

### Introduction

In relation to our recent work on the synthesis of  $\text{Mo}_2\text{I}_4\text{L}_4$  and  $\text{Mo}_2\text{I}_4(\text{LL})_2$  ( $L = \text{phosphine}$ ;  $\text{LL} = \text{diphosphine}$ ) complexes by decarbonylation of molybdenum(II) carbonyl derivatives,<sup>1-4</sup> where molybdenum(III) iodide complexes were also sometimes obtained as a result of a competitive disproportionation reaction, we became interested in finding alternative and better routes to derivatives of molybdenum(III) iodide. This interest was reinforced upon realizing that such derivatives were not known, although the chemistry of the corresponding chlorides and bromides had been developed. The  $[\text{MoX}_6]^{3-}$  and  $[\text{Mo}_2\text{X}_6]^{3-}$  ( $X = \text{Cl, Br}$ )<sup>5,6</sup> anions are well-known.  $\text{MoX}_3\text{L}_3$  ( $X = \text{Cl, Br}$ ) compounds with  $L = \text{pyridine}$  were prepared for the first time<sup>7</sup> by high-temperature reaction of  $\text{MoX}_3$  and the ligand and later<sup>8,9</sup> from  $(\text{NH}_4)_2\text{MoX}_5 \cdot \text{H}_2\text{O}$ . Compounds with other hard-donor-based ligands were obtained<sup>10</sup> by disproportionation reactions from  $\text{Mo}_2\text{X}_4(\text{CO})_8$ ; the unavailability of a good preparative method for

$\text{Mo}_2\text{I}_4(\text{CO})_8$ <sup>11a</sup> may have prevented such chemistry from being performed on the iodide system.<sup>11b</sup> San Filippo and Schaefer King were able to prepare the  $\text{MoX}_3(\text{py})_3$  ( $X = \text{Cl, Br}$ ;  $\text{py} = \text{pyridine}$ ) compounds by interaction of the  $\text{Mo}_2\text{X}_4(\text{py})_4$  dimers of  $\text{Mo}(\text{II})$  with pyridine at high temperatures.<sup>12</sup>

The cleanest route to  $\text{MoX}_3\text{L}_3$  complexes, which also has the advantage of being applicable to the preparation of derivatives containing soft-donor-based ligands,<sup>13</sup> seems to be the substitution reaction on preformed  $\text{MoX}_3(\text{THF})_3$  ( $\text{THF} = \text{tetrahydrofuran}$ ). Such a method has been applied so far only to the chloride system, the corresponding bromide and iodide being unknown.  $\text{MoCl}_3(\text{THF})_3$  has been conveniently obtained by reduction of  $\text{MoCl}_4(\text{THF})_2$  with zinc<sup>14</sup> or tin<sup>15</sup> or more recently via  $\text{MoCl}_3(\text{MeCN})_3$ , which in turn had been obtained from  $\text{MoCl}_4(\text{MeCN})_2$ .<sup>16</sup> The

- (1) Cotton, F. A.; Poli, R. *J. Am. Chem. Soc.* **1986**, *108*, 5628.
- (2) Cotton, F. A.; Dunbar, K. R.; Poli, R. *Inorg. Chem.* **1986**, *25*, 3700.
- (3) Cotton, F. A.; Poli, R. *Inorg. Chem.* **1986**, *25*, 3703.
- (4) Cotton, F. A.; Poli, R. *Inorg. Chem.* **1986**, *25*, 3624.
- (5) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 4th ed.; Wiley: New York, 1980, p 864.
- (6) Cotton, F. A.; Ucko, D. A. *Inorg. Chim. Acta* **1972**, *6*, 161 and references therein.
- (7) Rosenheim, A.; Abel, G.; Lewy, R. *Z. Anorg. Allg. Chem.* **1931**, *197*, 189.
- (8) Brencic, J. V. *Z. Anorg. Allg. Chem.* **1974**, *403*, 218.
- (9) Brencic, J. V.; Leban, I. *Z. Anorg. Allg. Chem.* **1978**, *445*, 251.
- (10) Westland, A. D.; Muriithi, N. *Inorg. Chem.* **1972**, *11*, 2971.

- (11) (a) Colton, R.; Rix, C. *J. Aust. J. Chem.* **1969**, *22*, 305. (b) A good-yield synthesis of  $\text{Mo}_2\text{I}_4(\text{CO})_8$  is now available,<sup>11c,d</sup> and the disproportionation reaction with pyridine to afford  $\text{MoI}_3(\text{py})_3$  has been reported.<sup>4</sup> (c) Poli, R. "Tesi di perfezionamento", Scuola Normale Superiore: Pisa, Italy, 1985. (d) Calderazzo, F.; Poli, R.; Zanazzi, P. F., manuscript in preparation.
- (12) San Filippo, J., Jr.; Schaefer King, M. A. *Inorg. Chem.* **1976**, *15*, 1228.
- (13) (a) Atwood, J. L.; Hunter, W. E.; Carmona-Guzman, E.; Wilkinson, G. *J. Chem. Soc., Dalton Trans.* **1980**, 467. (b) Carmona, E.; Marin, J. M.; Poveda, M. L.; Atwood, J. L.; Rogers, R. D. *J. Am. Chem. Soc.* **1983**, *105*, 3014.
- (14) (a) Anker, M. W.; Chatt, J.; Leigh, G. J.; Wedd, A. G. *J. Chem. Soc., Dalton Trans.* **1975**, 2639. (b) Dilworth, J. R.; Richards, R. L. *Inorg. Synth.* **1980**, *20*, 121.
- (15) Dilworth, J. R.; Zubieta, J. A. *J. Chem. Soc., Dalton Trans.* **1983**, 397.
- (16) Roh, S.-Y.; Bruno, J. W. *Inorg. Chem.* **1986**, *25*, 3105.

unavailability of molybdenum(IV) bromide and iodide starting materials may have prevented the development of such a chemistry for the heavier halides. We therefore tried to find alternative strategies for the preparation of MoI<sub>3</sub>(THF)<sub>3</sub>. The direct interaction between MoI<sub>3</sub><sup>17</sup> and THF is not a good one: no reaction occurs at room temperature, while at the reflux temperature I-(CH<sub>2</sub>)<sub>4</sub>-I and a highly insoluble material, presumably a molybdenum oxide or oxo-iodide, are obtained.<sup>18</sup> This is presumably due to the inertness of the MoX<sub>3</sub> material. MoCl<sub>3</sub> has two polymorphs, both of which contain extended networks of chloride ions with molybdenum ions embedded in adjacent octahedral cavities with strong metal-metal interactions.<sup>19</sup> On the other hand, at higher temperatures the oxophilicity of Mo(III) becomes the dominant feature and extraction of oxygen from the solvent occurs.

Since reduction (because of unavailability of starting materials) and direct interaction of MoI<sub>3</sub> and THF (because of inertness) are not feasible routes, we decided to try the oxidation of lower-valent molybdenum derivatives.

We wish to report here the successful synthesis of MoI<sub>3</sub>(THF)<sub>3</sub> by iodination of two different complexes, i.e. Mo(CO)<sub>3</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>) and Mo<sub>2</sub>I<sub>4</sub>(CO)<sub>8</sub>, and its conversion to MoI<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub> (R = Me, Et). The direct interaction between MoX<sub>3</sub> (X = Cl, I) and PR<sub>3</sub> (R = Me, Et) will also be discussed.

### Experimental Section

All operations were performed under an atmosphere of prepurified argon by using standard Schlenk tube techniques. Solvents were carefully dried and distilled under dinitrogen prior to use. IR spectra were recorded on a Perkin-Elmer 783 spectrophotometer. Elemental analyses were by Galbraith Laboratories Inc., Knoxville, TN. MoCl<sub>3</sub>,<sup>20</sup> MoI<sub>3</sub>,<sup>17</sup> Mo(CO)<sub>3</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>)<sup>21</sup> and Mo<sub>2</sub>I<sub>4</sub>(CO)<sub>8</sub><sup>11c</sup> were prepared as described elsewhere. PMe<sub>3</sub> (Strem Chemicals) and PEt<sub>3</sub> (Aldrich) were used as received.

**Synthesis of MoI<sub>3</sub>(THF)<sub>3</sub>. A. From Mo(CO)<sub>3</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>) and I<sub>2</sub> in THF.** Mo(CO)<sub>3</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>) (3.18 g, 11.7 mmol) was dissolved in THF (50 mL) and stirred for a few minutes. The IR bands at 1917 s and ca. 1775 vs (br) cm<sup>-1</sup> of the resulting solution indicated that conversion to Mo(CO)<sub>3</sub>(THF)<sub>3</sub> had occurred. This solution was then slowly added to a well-stirred solution of I<sub>2</sub> (4.46 g, 17.6 mmol) in THF (50 mL). At the end of the addition the clear dark red solution showed IR bands at 2066 s, 2006 vs, ca. 1970 w, and 1932 s cm<sup>-1</sup>. Stirring was continued at room temperature for 48 h, during which time the pale brown microcrystalline product precipitated out of the solution. The material was recovered by filtration, washed with THF, and dried in vacuo; yield 5.18 g (64%). This material had spectroscopic properties (IR in Nujol mull) identical with those of the analytically pure material, and it was used directly for the subsequent substitution reactions (vide infra).

**B. From Mo<sub>2</sub>I<sub>4</sub>(CO)<sub>8</sub> and I<sub>2</sub> in THF.** Mo<sub>2</sub>I<sub>4</sub>(CO)<sub>8</sub> (0.67 g, 0.72 mmol) was dissolved in 20 mL of THF. Gas evolution occurred, and the complete conversion<sup>11c</sup> to MoI<sub>2</sub>(CO)<sub>3</sub>(THF)<sub>2</sub> was confirmed by IR (bands at 2023 s, ca. 1945 sh, and 1929 vs cm<sup>-1</sup>). I<sub>2</sub> (0.19 g, 0.76 mmol) was added and the resulting mixture stirred at room temperature for about 20 h. IR monitoring showed the intensity of the bands of MoI<sub>2</sub>(CO)<sub>3</sub>(THF)<sub>2</sub> to decrease and new bands at 2066, 2006, and 1932 cm<sup>-1</sup> first to emerge and then decrease again while the pale brown product precipitated out of the solution. The solid was collected and characterized as described above in procedure A; yield 0.56 g (56%).

Analytically pure material in a crystalline form that was also suitable for the X-ray diffractometric analysis was obtained by simply setting aside without stirring a solution of MoI<sub>2</sub>(CO)<sub>3</sub>(THF)<sub>2</sub> and I<sub>2</sub> in equimolar amounts prepared as described above. Anal. Calcd for C<sub>12</sub>H<sub>24</sub>I<sub>3</sub>MoO<sub>3</sub>: C, 20.8; H, 3.5. Found: C, 21.1; H, 3.8. IR (Nujol mull, cm<sup>-1</sup>): 1340 m, 1295 w, 1245 w, 1180 w, 1045 m, 1005 s, 955 w, 925 sh, 915 m, 865 m, 840 vs, 685 w, 590 w.

**Reaction between MoI<sub>3</sub>(THF)<sub>3</sub> and PR<sub>3</sub> (R = Me, Et). Synthesis of MoI<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub>.** A. R = Me. MoI<sub>3</sub>(THF)<sub>3</sub> (0.49 g, 0.71 mmol) was suspended in THF (10 mL) and treated with PMe<sub>3</sub> (0.25 mL, 2.5 mmol). The resulting mixture was stirred at room temperature for ca. 2 h, during

which time all the solid dissolved to produce a clean bright red solution. The solvent was removed under reduced pressure and the residue crystallized by dissolution in toluene (2 mL) and reprecipitation by addition of 10 mL of *n*-hexane. After it was cooled to ca. -20 °C overnight, the brick red product was filtered off, washed with *n*-hexane, and dried in vacuo; yield 0.26 g (52%). Anal. Calcd for C<sub>9</sub>H<sub>27</sub>I<sub>3</sub>MoP<sub>3</sub>: C, 15.3; H, 3.9. Found: C, 15.7; H, 3.9. IR (Nujol mull, cm<sup>-1</sup>): 1420 m, 1300 m, 1280 m, 1120 m (br), 950 s, 850 w, 730 m, 670 w.

B. R = Et. By a procedure identical with that described above for R = Me, from 0.80 g of MoI<sub>3</sub>(THF)<sub>3</sub> and 0.6 mL of PEt<sub>3</sub> in 15 mL of THF, 0.70 g of the compound MoI<sub>3</sub>(PEt<sub>3</sub>)<sub>3</sub> was obtained (yield 73%). Anal. Calcd for C<sub>18</sub>H<sub>45</sub>I<sub>3</sub>MoP<sub>3</sub>: C, 26.0; H, 5.5. Found: C, 25.6; H, 5.3. IR (Nujol mull, cm<sup>-1</sup>): 1415 m, 1250 w, 1110 w (br), 1035 s, 995 w, 975 w, 760 s, 745 s, 730 m-s, 710 m, 685 w-m, 675 w.

**Reaction between MoX<sub>3</sub> and PR<sub>3</sub> in Toluene. A. MoI<sub>3</sub> and PMe<sub>3</sub>.** MoI<sub>3</sub> (0.52 g, 1.09 mmol) was suspended in 15 mL of toluene and treated with 0.45 mL of PMe<sub>3</sub> (4.5 mmol). Stirring at room temperature slowly developed a pale green color in the solution. The mixture was warmed to about 80 °C for 48 h to produce a deep emerald green solution and a pale solid. The latter was filtered off, washed with toluene, and dried in vacuo. It exhibited an IR band (Nujol mull) at 2405 cm<sup>-1</sup>. Anal. Calcd for C<sub>7</sub>H<sub>10</sub>I<sub>3</sub>P: C, 17.7; H, 5.0. Found: C, 17.7; H, 4.9. The volume of the filtered solution was reduced to ca. 5 mL, and 20 mL of *n*-hexane was added to cause the precipitation of brick red microcrystalline material. After it was cooled to ca. 5 °C overnight, the solid was filtered off, washed with *n*-hexane, and dried in vacuo. It was shown to be MoI<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub> by comparison of its spectroscopic properties with those of the material obtained from MoI<sub>3</sub>(THF)<sub>3</sub> as described above; yield 0.31 g (41%). From the mother solution, now having a blue color, 0.17 g of blue crystals were obtained after evaporation to dryness and recrystallization of the residue from 15 mL of hot *n*-hexane. The spectroscopic properties of this sample (IR and UV/vis) are in accord with its formulation as the known<sup>1</sup> quadruply bonded Mo(II) dimer Mo<sub>2</sub>I<sub>4</sub>(PMe<sub>3</sub>)<sub>4</sub>.

B. MoCl<sub>3</sub> and PMe<sub>3</sub>. MoCl<sub>3</sub> (0.50 g, 2.47 mmol) was treated in toluene (20 mL) with PMe<sub>3</sub> (0.80 mL, 8.0 mmol). The mixture was briefly (ca. 15 min) refluxed and then filtered. The red solution was concentrated to about 5 mL under reduced pressure and then treated with *n*-hexane (10 mL). The resulting yellow solid was filtered off, washed with *n*-hexane, and dried in vacuo. Analytical and spectroscopic properties indicated this compound to be the known<sup>16</sup> MoCl<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub>, yield 0.52 g (48%). The mother solution of this solid was dichroic red/blue; its UV/visible properties confirmed the presence of Mo<sub>2</sub>Cl<sub>4</sub>(PMe<sub>3</sub>)<sub>4</sub>.<sup>22</sup>

C. MoI<sub>3</sub> and PEt<sub>3</sub>. MoI<sub>3</sub> (0.47 g, 0.98 mmol) was suspended in 20 mL of toluene and treated with 0.45 mL of PMe<sub>3</sub> (3.0 mmol). A pale blue-green color slowly developed upon stirring at room temperature. The mixture was refluxed for about 5 h, after which time it consisted of a blue-green solution and some yellow solid. Following the filtration of the hot mixture, a yellow solid crystallized upon cooling to room temperature; yield 0.13 g. Anal. Calcd for C<sub>18</sub>H<sub>46</sub>I<sub>4</sub>MoP<sub>3</sub>: C, 22.5; H, 4.8. Found: C, 22.9; H, 5.2. IR (Nujol mull, cm<sup>-1</sup>): 2395 w, 1415 m, 1260 m, 1245 w, 1050 m, 1035 s, 995 w, 970 w, 885 w, 870 m, 755 s, 740 w, 715 s, 700 w, 660 w, 620 w. The mother solution was evaporated to dryness and the blue residue recrystallized from 5 mL of hot *n*-hexane. A 0.08-g amount of blue crystals was recovered from the solution after cooling to -20 °C. Anal. Calcd for C<sub>24</sub>H<sub>50</sub>I<sub>4</sub>Mo<sub>2</sub>P<sub>4</sub>: C, 24.6; H, 5.2. Found: C, 24.8; H, 5.1. IR (Nujol mull, cm<sup>-1</sup>): 1410 m, ca. 1260 m (br), ca. 1100 m (br, with shoulder at ca. 1140), 1040 s, 985 w, 780 sh, 765 s, 750 s, 730 sh, 685 w, 670 w, 610 w, 490 w, 450 w. UV/vis (CH<sub>2</sub>Cl<sub>2</sub>; λ<sub>max</sub>, nm (ε, mol L<sup>-1</sup> cm<sup>-1</sup>): 640 (1.2 × 10<sup>3</sup>), 383 (2.6 × 10<sup>3</sup>).

**Thermal Treatment of MoI<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub> (R = Me, Et) in Toluene. A. R = Me.** A toluene solution of MoI<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub> (ca. 1.4 × 10<sup>-2</sup> M) was refluxed overnight. No significant change was observed during the treatment.

B. R = Et. MoI<sub>3</sub>(PEt<sub>3</sub>)<sub>3</sub> (0.24 g, 0.29 mmol) was refluxed in toluene (10 mL) for ca. 5 h. The resulting mixture contained some yellow solid in a green solution. From the filtered solution yellow crystals deposited upon cooling to room temperature. They were decanted, washed with *n*-hexane, and dried in vacuo. They were shown to be of [PHEt<sub>3</sub>]-[MoI<sub>4</sub>(PEt<sub>3</sub>)<sub>2</sub>] by comparison of their spectroscopic properties with those of the material obtained from MoI<sub>3</sub> and PEt<sub>3</sub> (vide supra). UV/vis spectroscopy on the mother solution confirmed the presence of Mo<sub>2</sub>I<sub>4</sub>(PEt<sub>3</sub>)<sub>4</sub>.

**X-ray Crystallography for MoI<sub>3</sub>(THF)<sub>3</sub>.** A parallelepiped-shaped single crystal was glued to the inside of a thin-walled glass capillary, which was then sealed under nitrogen and mounted on a Nicolet PI diffractometer with its longest dimension nearly parallel to the φ axis. The cell dimensions were determined by a least-squares calculation on

(17) Westland, A. D.; Muriithi, N. *Inorg. Chem.* **1973**, *12*, 2356.

(18) Poli, R., unpublished observations.

(19) Schäfer, H.; von Schnering, H.-G.; Tillack, J.; Kuhn, F.; Wöhrle, H.; Baumann, H. Z. *Anorg. Allg. Chem.* **1967**, *353*, 281.

(20) Mallock, A. *Inorg. Nucl. Chem. Lett.* **1967**, *3*, 441.

(21) Strohmeier, W. *Chem. Ber.* **1961**, *94*, 3337.

(22) Cotton, F. A.; Extine, M. W.; Felthouse, T. R.; Kolthammer, B. W. S.; Lay, D. G. *J. Am. Chem. Soc.* **1981**, *103*, 4040.

**Table I.** Crystal Data for MoI<sub>3</sub>(THF)<sub>3</sub>

formula	C <sub>12</sub> H <sub>24</sub> I <sub>3</sub> MoO <sub>3</sub>
fw	692.88
space group	<i>Pbcn</i>
systematic absences	0 <i>kl</i> , <i>k</i> ≠ 2 <i>n</i> ; <i>h</i> 0 <i>l</i> , <i>l</i> ≠ 2 <i>n</i> ; <i>hk</i> 0, <i>h</i> + <i>k</i> ≠ 2 <i>n</i>
<i>a</i> , Å	9.252 (3)
<i>b</i> , Å	14.541 (5)
<i>c</i> , Å	14.273 (3)
α, deg	90
β, deg	90
γ, deg	90
<i>V</i> , Å <sup>3</sup>	1920 (2)
<i>Z</i>	4
<i>d</i> <sub>calcd</sub> , g cm <sup>-3</sup>	2.397
cryst size, mm	0.25 × 0.40 × 0.50
μ(Mo Kα), cm <sup>-1</sup>	54.44
data collec instrument	Nicolet Pī
radiation (monochromated in incident beam)	Mo Kα (λ <sub>K</sub> = 0.71073 Å)
orientation reflcns:	15; 20–30
no.; range (2θ), deg	
temp, °C	20
scan method	ω
data collec range, 2θ, deg	3–50.7
no. of unique data, total with <i>F</i> <sub>o</sub> <sup>2</sup> > 3σ( <i>F</i> <sub>o</sub> <sup>2</sup> )	1429, 926
no. of parameters refined	88
transmission factors: max, min	1.000, 0.769
<i>R</i> <sup>a</sup>	0.0501
<i>R</i> <sub>w</sub> <sup>b</sup>	0.0642
quality-of-fit indicator <sup>c</sup>	1.336
largest shift/esd, final cycle	0.24
largest peak, e Å <sup>-3</sup>	1.346

<sup>a</sup>*R* = Σ||*F*<sub>o</sub>|| - ||*F*<sub>c</sub>|| / Σ||*F*<sub>o</sub>||. <sup>b</sup>*R*<sub>w</sub> = [Σ*w*(||*F*<sub>o</sub>|| - ||*F*<sub>c</sub>||)<sup>2</sup> / Σ*w*||*F*<sub>o</sub>||<sup>2</sup>]<sup>1/2</sup>; *w* = 1/σ<sup>2</sup>(||*F*<sub>o</sub>||). <sup>c</sup>Quality of fit = [Σ*w*(||*F*<sub>o</sub>|| - ||*F*<sub>c</sub>||)<sup>2</sup> / (N<sub>observ</sub> - N<sub>params</sub>)]<sup>1/2</sup>.

**Table II.** Atomic Positional Parameters and Equivalent Isotropic Displacement Parameters (Å<sup>2</sup>) and Their Estimated Standard Deviations for MoI<sub>3</sub>(THF)<sub>3</sub><sup>a</sup>

atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i>
Mo	0.500	0.37167 (7)	0.250	2.17 (3)
I1	0.500	0.18281 (8)	0.250	4.06 (4)
I2	0.3162 (1)	0.37131 (7)	0.09507 (8)	3.76 (2)
O1	0.500	0.5238 (8)	0.250	3.1 (2)
O2	0.3103 (8)	0.3830 (9)	0.3414 (6)	3.3 (2)
C1	0.491 (2)	0.5820 (8)	0.334 (1)	8.7 (9)
C2	0.478 (2)	0.671 (1)	0.303 (2)	7.2 (6)
C3	0.303 (2)	0.338 (2)	0.430 (1)	6.1 (4)
C4	0.141 (2)	0.371 (1)	0.469 (1)	5.7 (5)
C5	0.109 (2)	0.454 (1)	0.408 (1)	5.6 (5)
C6	0.198 (2)	0.441 (1)	0.319 (2)	5.7 (3)

<sup>a</sup>Anisotropically refined atoms are given in the form of the equivalent isotropic displacement parameter defined as  $\frac{1}{3}[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$ .


the setting angles of 15 reflections with 20° < 2θ < 30° and were subsequently confirmed by axial photographs. Data collection was performed with a scan rate of 3.0° min<sup>-1</sup> for reflections having 3.0° < 2θ < 39.7° and 2.0° min<sup>-1</sup> for those having 39.7° < 2θ < 50.7°. The periodic monitoring of six standard reflections showed no significant variation of intensity. Data were corrected for Lorentz and polarization effects, and an empirical absorption correction<sup>23</sup> based on azimuthal (ψ) scans of five reflections having an Eulerian angle χ near 90° was also applied.

The structure was solved by SHELX-76 direct methods and then refined by subsequent least-squares full-matrix cycles and difference Fourier maps with the Enraf-Nonius SDP software. Anisotropic refinement converged at *R* = 0.0501 (*R*<sub>w</sub> = 0.0642). The highest peak in the last difference Fourier map (1.336 e Å<sup>-3</sup>) was located at 0.88 Å from I(2), and it was considered to be the result of series termination errors. Pertinent crystal data are reported in Table I. Fractional atomic coordinates

**Table III.** Selected Bond Distances (Å) and Angles (deg) and Their Estimated Standard Deviations

Mo-I1	2.746 (2)	Mo-O1	2.212 (12)
Mo-I2	2.790 (1)	Mo-O2	2.193 (8)
I1-Mo-I2	89.89 (3)	I2-Mo-O2	89.1 (2)
I1-Mo-O1	180.00 (0)	I2-Mo-O2'	90.9 (2)
I1-Mo-O2	94.3 (3)	O1-Mo-O2	85.7 (3)
I2-Mo-I2	180.00 (0)	O2-Mo-O2'	171.4 (5)
I2-Mo-O1	90.11 (3)		





<sup>a</sup>Numbers in parentheses are estimated standard deviations in the least significant digits.

**Figure 1.** ORTEP view of MoI<sub>3</sub>(THF)<sub>3</sub> with the atomic numbering scheme employed.

and equivalent isotropic displacement parameters are listed in Table II, while Table III contains selected bond distances and angles. Full tables of bond distances and angles, as well as anisotropic displacement parameters and a listing of observed and calculated structure factors, have been deposited as supplementary material.

## Results and Discussion

Interaction of THF adducts of low-valent molybdenum carbonyl derivatives proves to be a convenient method for the synthesis of MoI<sub>3</sub>(THF)<sub>3</sub>. Mo(CO)<sub>3</sub>(THF)<sub>3</sub>, obtained in situ from Mo(CO)<sub>3</sub>(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>CH<sub>3</sub>) (eq 1), and MoI<sub>2</sub>(CO)<sub>3</sub>(THF)<sub>2</sub>, obtained in situ from Mo<sub>2</sub>I<sub>4</sub>(CO)<sub>8</sub> (eq 3), were employed. The stoichiometries of the iodination reactions are outlined in eq 2 and 4.



It is important to perform reaction 2 by adding the molybdenum complex to the iodine solution so that an excess of iodine will always be present. This will minimize the side reaction of carbonylation of the starting material according to eq 5.




IR monitoring of reactions 2 and 4 shows the intermediacy of a carbonyl complex whose IR spectrum exhibits CO stretching vibrations at 2066, 2006, and 1932 cm<sup>-1</sup>. Such a spectrum is strikingly similar to that of the [MoI<sub>3</sub>(CO)<sub>4</sub>]<sup>-</sup> ion (bands at 2072 s, 2010 s, and 1945 m cm<sup>-1</sup> in CH<sub>2</sub>Cl<sub>2</sub> solution for the *n*-Bu<sub>4</sub>N<sup>+</sup>

salt)<sup>24</sup> and would therefore suggest such an intermediate to contain a MoI<sub>3</sub>(CO)<sub>4</sub> moiety of a molybdenum(II) center, possibly bridged by one or more iodide ligands to one or more molybdenum(III) centers. This is, however, pure speculation based only on this IR evidence. We have been so far unsuccessful in isolating this intermediate.

We attribute the success of these syntheses of MoI<sub>3</sub>(THF)<sub>3</sub> to the mild conditions employed. When we performed the iodination directly on Mo(CO)<sub>6</sub>, which required higher temperatures for the labilization of the CO ligands, only oxygenated molybdenum products and I-(CH<sub>2</sub>)<sub>4</sub>-I were obtained.<sup>18</sup>


The structure of the MoI<sub>3</sub>(THF)<sub>3</sub> molecule is shown in Figure 1. The molecule resides on a crystallographic twofold axis, this passing through the Mo, I(1), and O(1) atoms. The coordination geometry is octahedral, the ligands having the meridional arrangement. The same geometry has been observed in other MCl<sub>3</sub>(THF)<sub>3</sub> complexes of different metals (M = Sc,<sup>25</sup> Ti,<sup>26</sup> V,<sup>27</sup> Cr<sup>27</sup>) and in other Mo(III) complexes, such as MoCl<sub>3</sub>(py)<sub>3</sub>,<sup>8</sup> MoBr<sub>3</sub>(4-pic)<sub>3</sub>,<sup>9</sup> (4-pic = 4-picoline), and MoI<sub>3</sub>(PMe<sub>2</sub>Ph)<sub>2</sub>(POMe<sub>2</sub>Ph).<sup>4</sup> The Mo-I distance compares well with those found<sup>4</sup> in MoI<sub>3</sub>(PMe<sub>2</sub>Ph)<sub>2</sub>(POMe<sub>2</sub>Ph) (ax, 2.746 (2) vs. 2.756 (2) Å; eq, 2.790 (1) vs. 2.762 (2) and 2.768 (2) Å; ax refers to the iodine atoms trans to the THF ligand, whereas eq refers to the iodine atoms trans to each other) and with those in the *trans*-(MoI<sub>4</sub>(PET<sub>3</sub>Ph)<sub>2</sub>)<sup>-</sup> anion (2.784 (1), 2.789 (1), 2.754 (1), and 2.797 (1) Å).<sup>4</sup> In particular, the Mo-I<sub>ax</sub> bond length is shorter than the Mo-I<sub>eq</sub> one, a feature that is observed also in all the other MX<sub>3</sub>L<sub>3</sub> compounds so far structurally characterized and that is in agreement with the accepted order of trans influence (I<sup>-</sup> > Cl<sup>-</sup> > OR<sub>2</sub>).<sup>28</sup> The observed trend in the Mo-O distances is also expected on trans-influence grounds: the Mo-O<sub>ax</sub> (trans to iodide) distance is longer (2.21 (1) Å) than the Mo-O<sub>eq</sub> one (2.193 (8) Å). Again, this is also observed in the MCl<sub>3</sub>(THF)<sub>3</sub> complexes (M = Sc, Ti, V, Cr).<sup>27</sup> The thermal ellipsoids of the THF carbon atoms, particularly those attached to the oxygen atoms, are strongly elongated perpendicularly to the ideal THF plane. This indicates a high thermal motion. The bond distances and angles related to the THF molecules are therefore a time-averaged result over all the possible conformations assumed in the crystal and are not particularly meaningful from the chemical point of view.

Interaction of MoI<sub>3</sub>(THF)<sub>3</sub> with 3 equiv of PR<sub>3</sub> (R = Me, Et) cleanly and rapidly affords solutions of MoI<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub> (eq 6).




This represents the first clean synthetic route to octahedral MoI<sub>3</sub> derivatives.

The direct interaction between MoI<sub>3</sub> and PR<sub>3</sub> has been carried out in toluene as solvent. In the case of the trimethylphosphine reaction, MoI<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub> has been obtained, but a considerable amount of reduction to Mo<sub>2</sub>I<sub>4</sub>(PMe<sub>3</sub>)<sub>4</sub> simultaneously takes place. PHMe<sub>3</sub><sup>+</sup>I<sup>-</sup> is also produced under those conditions (see eq 7).




The reduction process to the quadruply bonded Mo(II) dimer, with concomitant protonation of PMe<sub>3</sub>, presumably involves oxidative C-C coupling of solvent molecules. Since toluene has unusually reactive (benzylic) C-H bonds, the reaction might proceed differently (or not at all) in other solvents. The corresponding reaction starting from the molybdenum(III) chloride

Scheme I




gives similar results. The only noticeable difference was the reaction time necessary to consume the starting material completely (eq 8).



Another point concerning both these reactions in toluene is that dibenzyl might well be a byproduct. We have not attempted to detect it, however, so this aspect of the overall process remains uncertain. It is not impossible, however, that adventitious H<sub>2</sub>O is the source of protons, with the oxygen atom being captured by PMe<sub>3</sub>. Again, we do not have any evidence to rule this in, or out.

It is worth mentioning that the reaction between MoCl<sub>3</sub> and phosphine had been already reported by Edwards, Walton, and co-workers.<sup>29</sup> They performed the reaction in ethanol and were able to obtain only dinuclear derivatives of molybdenum(II). While we still obtain some amounts of Mo(II) materials, our main products are the complexes of Mo(III), indicating that the reducing power of the solvent plays an important role in this reaction.

Interaction of MoI<sub>3</sub> with PET<sub>3</sub> again gave rise to some reduction process, affording Mo<sub>2</sub>I<sub>4</sub>(PET<sub>3</sub>)<sub>4</sub>, but the Mo(III) product this time was the anionic [MoI<sub>4</sub>(PET<sub>3</sub>)<sub>2</sub>]<sup>-</sup> complex, isolated as its PHEt<sub>3</sub><sup>+</sup> salt (eq 9). This exhibits the characteristic P-H stretching



vibration at 2395 cm<sup>-1</sup> in the IR spectrum. We suggest for the anion a *trans* configuration, since this was found for the very similar [MoI<sub>4</sub>(PET<sub>2</sub>Ph)<sub>2</sub>]<sup>-</sup> anion.<sup>4</sup>

We propose Scheme I to rationalize these results. The interaction between MoX<sub>3</sub> and PR<sub>3</sub> would presumably proceed stepwise through the intermediates MoX<sub>3</sub>(PR<sub>3</sub>) and MoX<sub>3</sub>(PR<sub>3</sub>)<sub>2</sub>. The latter can then take up the last PR<sub>3</sub> molecule to produce MoX<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub>. If, however, the intermediate MoX<sub>3</sub>(PR<sub>3</sub>)<sub>2</sub> undergoes a one-electron reduction at the expense of one electron of the solvent, we can then generate the molybdenum(II) product, along with some HX that would then be trapped by the phosphine to form PHR<sub>3</sub><sup>+</sup>X<sup>-</sup>. We have then rationalized the formation of all the products observed in the PMe<sub>3</sub> reaction. The formation of the Mo(III) compound observed in the PET<sub>3</sub> reaction, i.e. [PHEt<sub>3</sub>][MoI<sub>4</sub>(PET<sub>3</sub>)<sub>2</sub>], can also be rationalized if we suppose the reaction between intermediate MoX<sub>3</sub>(PR<sub>3</sub>)<sub>2</sub> and PR<sub>3</sub> to be reversible and to be more so for PET<sub>3</sub> than for PMe<sub>3</sub>. The intermediate MoX<sub>3</sub>(PR<sub>3</sub>)<sub>2</sub> would therefore be present in solution in substantial amounts to react with the PHR<sub>3</sub><sup>+</sup>X<sup>-</sup> species generated in the reductive pathway to eventually produce the observed Mo(III) anion. Independent observations that are in agreement with this view are the following: (i) MoI<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub> is stable in refluxing toluene for at least several hours, giving no indication of reduction to Mo<sub>2</sub>I<sub>4</sub>(PMe<sub>3</sub>)<sub>4</sub>; (ii) MoCl<sub>3</sub>(PMe<sub>3</sub>)<sub>3</sub> is also rather

(24) Barbati, A.; Calderazzo, F.; Poli, R.; Zanazzi, P. F. *J. Chem. Soc., Dalton Trans.* **1986**, 2569.

(25) Atwood, J. L.; Smith, K. D. *J. Chem. Soc., Dalton Trans.* **1974**, 921.

(26) Handlowic, M.; Miklos, D.; Zikmund, M. *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.* **1981**, *37*, 811.

(27) Cotton, F. A.; Duraj, S. A.; Powell, G. L.; Roth, W. J. *Inorg. Chim. Acta* **1986**, *113*, 81.

(28) Pidcock, A.; Richards, R. E.; Venanzi, L. M. *J. Chem. Soc. A* **1966**, 1707.

(29) Armstrong, J. E.; Edwards, D. A.; Maguire, J. J.; Walton, R. A. *Inorg. Chem.* **1979**, *18*, 1172.

stable in refluxing toluene, but to a lesser extent than its iodide analogue, since slow appearance of the typical  $\delta \rightarrow \delta^*$  electronic transition of  $\text{Mo}_2\text{Cl}_4(\text{PMe}_3)_4$  can be detected in the visible spectrum; (iii) thermal treatment of  $\text{MoI}_3(\text{PEt}_3)_3$  as obtained from  $\text{MoI}_3(\text{THF})_3$  by the procedure described in eq 6 gives rise to a mixture of  $\text{Mo}_2\text{I}_4(\text{PEt}_3)_4$  and  $[\text{PHEt}_3][\text{MoI}_4(\text{PEt}_3)_2]$ .

It would therefore seem that dissociation of a phosphine molecule from  $\text{MoI}_3(\text{PR}_3)_3$  is a much more facile process for  $\text{PEt}_3$  than for  $\text{PMe}_3$ . This is in agreement with the relative basicity of the two ligands and also with their steric requirements. A similar trend was found for the Mo(II) quadruply bonded dimers  $\text{Mo}_2\text{Cl}_4(\text{PR}_3)_3$  in their reaction with CO: kinetic studies indicated that the dissociation of a phosphine is the necessary primary step, and the dimer with  $\text{PMe}_3$  was found to be much less reactive than those containing phosphines with longer alkyl substituents.<sup>30</sup>

Finally, we would like to remark that the chemistry depicted in Scheme I may be related to other phosphine-molybdenum carbonyl reactions that were found to lead to salts of phosphonium cations.<sup>4,11a,31</sup>

## Conclusions

A convenient route to  $\text{MoI}_3(\text{THF})_3$  has been developed by diiodine oxidation of molybdenum(0) and molybdenum(II) carbonyl starting materials. This route represents a new strategy for the preparation of THF adducts of transition-metal halides, the commonly used ones being direct interaction of the metal(III) halide with THF or reduction of higher valent starting materials. We anticipate that other unknown  $\text{MX}_3(\text{THF})_3$  complexes, particularly  $\text{MoBr}_3(\text{THF})_3$  and  $\text{WX}_3(\text{THF})_3$  ( $\text{X} = \text{Br}, \text{I}$ ), could possibly be obtained in the same way.

**Acknowledgment.** We thank the National Science Foundation for support of this work, Dr. Cynthia Day of Crystallitics, Lincoln, NB, for collection of intensity data on  $\text{MoI}_3(\text{THF})_3$ , and Professor H. D. Kaesz for helpful discussion.

**Registry No.**  $\text{MoI}_3(\text{THF})_3$ , 107680-52-4;  $\text{Mo}(\text{CO})_3(\eta^6\text{-C}_6\text{H}_5\text{CH}_3)$ , 12083-34-0;  $\text{Mo}_2\text{I}_4(\text{CO})_8$ , 22547-54-2;  $\text{MoI}_2(\text{CO})_3(\text{THF})_2$ , 103904-36-5;  $\text{MoI}_3(\text{PMe}_3)_3$ , 107680-53-5;  $\text{MoI}_3(\text{PEt}_3)_3$ , 107680-54-6;  $\text{Mo}_2\text{Cl}_4(\text{PMe}_3)_4$ , 67619-17-4;  $\text{MoCl}_3(\text{PMe}_3)_3$ , 73534-30-2;  $\text{Mo}_2\text{I}_4(\text{PEt}_3)_4$ , 107680-55-7;  $[\text{MoI}_4(\text{PEt}_3)_2][\text{PHEt}_3]$ , 107680-57-9;  $\text{Mo}_2\text{I}_4(\text{PMe}_3)_4$ , 89637-15-0;  $\text{I}_2$ , 7553-56-2.

**Supplementary Material Available:** Full tables of bond distances and angles and a listing of anisotropic displacement parameters for  $\text{MoI}_3(\text{THF})_3$  (3 pages); a listing of observed and calculated structure factors for  $\text{MoI}_3(\text{THF})_3$  (5 pages). Ordering information is given on any current masthead page.

- (30) (a) Cotton, F. A.; Darensbourg, D. J.; Kolthammer, B. W. S. *J. Organomet. Chem.* **1981**, *217*, C14. (b) Daniels, L. M. Ph.D. Dissertation, Texas A&M University, 1984.
- (31) (a) Lewis, J.; Whyman, R. *J. Chem. Soc. A* **1967**, 77. (b) Tsang, W.-S.; Meek, D. W.; Wojcicki, A. *Inorg. Chem.* **1968**, *7*, 1263. (c) Baker, P. K.; Fraser, S. G. *Inorg. Chim. Acta* **1986**, *116*, L1.