



The tick tock of odontogenesis

Li Zheng, Lauren Ehardt, Blake Mcalpin, Imad About, Doohak Kim, Silvana Papagerakis, Petros Papagerakis

► To cite this version:

Li Zheng, Lauren Ehardt, Blake Mcalpin, Imad About, Doohak Kim, et al.. The tick tock of odontogenesis. Experimental Cell Research, 2014, 325 (2), pp.83-89. <10.1016/j.yexcr.2014.02.007>. <hal-03547658>

HAL Id: hal-03547658

<https://hal.science/hal-03547658v1>

Submitted on 5 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

The tick tock of odontogenesis

Li Zheng^{a,b}, Lauren Ehardt^a, Blake McAlpin^a, Imad About^c, Doohak Kim^a,
Silvana Papagerakis^{b,d}, Petros Papagerakis^{a,e,f,*}

^aDepartment of Pediatric Dentistry, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

^bDepartment of Otolaryngology, Medical School, University of Michigan, Ann Arbor, MI, USA

^cAix-Marseille Université, CNRS, ISM UMR 7287, 13288, Marseille cedex 09, France

^dDepartment of Periodontics and Oral Medicine, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

^eCenter for Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

^fCenter for Organogenesis, University of Michigan, Ann Arbor, MI, USA

ABSTRACT

Although a big deal of dental research is being focused to the understanding of early stages of tooth development, a huge gap exist on our knowledge on how the dental hard tissues are formed and how this process is controlled daily in order to produce very complex and diverse tooth shapes adapted for specific functions. Emerging evidence suggests that clock genes, a family of genes that controls circadian functions within our bodies, regulate also dental mineralized tissues formation. Enamel formation, for example, is subjected to rhythmical molecular signals that occur on short (24 h) periods and control the secretion and maturation of the enamel matrix. Accordingly, gene expression and ameloblast functions are also tightly modulated in regular daily intervals. This review summarizes the current knowledge on the circadian controls of dental mineralized tissues development with a special emphasis on amelogenesis.

Contents

Introduction to the circadian clock	2
Expression and potential roles of the circadian clock in mineralized tissues	2
Bone	2
Enamel	3
Dentin	5
Cementum, and periodontal ligament	5
Clock genes and stem cells	5
Conclusions	5

Conflict of interest	6
Acknowledgments	6
References	6

Introduction to the circadian clock

Circadian rhythms are generated within an organism by endogenous biological clocks driven by cyclic events [1]. The key requirement for the description of a rhythm as *circadian* is to show that the rhythm persists under constant conditions (i.e. the rhythm must continue within an approximate 24 h period when all external time cues are removed from the environment), thus demonstrating the endogenous existence of a time-keeping mechanism. Circadian rhythms are controlled by the body's "central" clock located in the brain at the suprachiasmatic nucleus (SCN). Body functions are also regulated directly by "peripheral" clocks located in several tissues. The central clock is light responsive and can be entrained by light/dark cycles. The peripheral clocks can be entrained by the central clock or independently by other physiological stimuli such as feeding [2].

At the molecular level, the circadian clock is regulated by differential expression of ~20 transcription factors called clock genes. The central clock located in the brain is composed of about 20,000 neurons, all of which express clock genes that oscillate in synchrony [3,4]. Clock genes are defined by a set of criteria that include rhythm in activity or amount as well as molecular evidence of a feedback mechanism [5,6]. The key mammalian clock genes are named Aryl hydrocarbon receptor nuclear translocator-like (*Arntl* or *Bmal1*), circadian locomotor output cycles kaput (*Clock*), Period1 (*Per1*), Period2 (*Per2*), Period 3 (*Per3*), and Cryptochrome 1 and 2 (*Cry1*, *Cry2*). Clock genes functions are regulated by feedback mechanisms consisting of an auto-regulatory ability in which specific clock proteins negatively regulate their own synthesis. Clock genes transmit output signals that drive rhythms of gene expression in central and peripheral tissues.

Evidence for a role of clock genes in circadian physiology and clock controlled gene expression comes from the studies of clock gene knockout mice. Wild-type mice will increase their locomotor activity at night as measured by wheel-running activity. This characteristic is used to assess circadian phenotypes of knockout mice. For example, mice deficient in *Per1* lose wheel-running rhythmicity when placed in constant darkness [7]. This observation is much more pronounced in *Per1/Per2* double knockout mice, which display an immediate loss of wheel-running rhythmicity when placed in constant darkness. Moreover, *Per1/Per2* double knockout mice have no circadian rhythms in central and peripheral (i.e. hepatic) clocks, or clock-controlled gene expression [8,9]. On the other hand, deletion of *Clock* and *Bmal1* genes results not only in circadian disturbances, but also in metabolic abnormalities of lipid and glucose homeostasis [10–12]. Thus, clock gene knockout mice have profound changes in circadian rhythmicity and offer a unique experimental genetic model to analyze the link between circadian gene networks and organ physiology.

The most direct mechanism by which clock genes drive circadian gene expression is through regulation of promoter activity of clock-controlled genes (CCG) [13]. In a given tissue, approximately 10–20% of the organ specific genes are under circadian control [14]. Recent

studies in humans confirm and further specify the role of clock genes in human diseases [1]. Exciting links between peripheral organs (such as the gut) and the brain are being discovered [2]. Regulation of stem cells behavior is also being linked to clock genes opening a whole new area of fascinating research [15]. Most importantly, understanding how the circadian systems work may facilitate innovative treatment options for patients with untreatable diseases such as autoimmune diseases and cancer as well as for psychiatric conditions [16].

In oral health, the possibilities of clock genes involvement into patho-physiology of oral and craniofacial tissues remain largely unexplored [17]. This review summarizes the potential roles of clock genes in dental tissues formation with a special focus on enamel development. We also present hypotheses regarding the potential connections between dysregulated clock gene expression and mineralized tissues formation in general. The recently hypothesized role of clock genes in regulating stem cells properties is also briefly mentioned.

Expression and potential roles of the circadian clock in mineralized tissues

Bone

The diurnal variation in the synthesis of type I collagen and osteocalcin are well known [18,19] supporting the hypothesis that synthesis and secretion of the matrix proteins are under circadian control. More recent studies have confirmed that osteoblast contain a peripheral clock mechanism that regulates bone volume [20,21]. *Per1*, *Per2*, *Cry1*, *Bmal1*, and *Clock* all showed robust, rhythmic expression over a 24 h period in osteoblasts, and their expression was decreased and became arrhythmic in *Per1^{-/-}* and *Per2m/m* bones, indicating that bone has a peripheral clock. It has also been documented that the molecular clock, specifically the *Per* genes, inhibits bone formation by preventing osteoblast proliferation [21]. It has been also found that *Per2* mutant mice and mice lacking *Cry2*($-/-$) displayed significantly increased bone volume at 12 weeks of age, when bone turnover is high [22]. Of interest, *Cry2* influences mostly the osteoclastic cellular component of bone while *Per2* acts on osteoblast parameters [22]. Furthermore, it is recently shown that mineralization in developing bone tissue is regulated by a local circadian oscillator mechanism [23]. Thus, we can postulate that bone deposition and mineralization are under direct circadian controls. It remains to clarify how the potential interplay between the "central" and "peripheral" clocks regulates bone homeostasis.

The clinical relevance of these cyclic events controlled by clock genes is currently being studied for several bone related diseases and this new knowledge may influence novel treatment options [24]. For example, the circadian clock plays a key role in obesity and diabetes which may be controlled by osteoblasts released hormones [25]. In dentistry, studies elucidating the role of clock genes in alveolar bone homeostasis warrant novel exciting

discoveries in the field of orthodontics tooth movement and periodontal tissue regeneration.

Enamel

Dental enamel is formed mainly during two distinct developmental stages: the secretory stage (when the full thickness of enamel is completed) and the maturation stage (when the enamel organic material is replaced by hydroxyapatite crystals) [26–28]. Ameloblasts, the cells responsible for making enamel, are specialized epithelial cells with distinct morphological characteristics that change at each developmental stage. Secretory and maturation ameloblasts are characterized by expression of enamel stage-specific genes that perform stage-specific functions. Besides the stage-dependent developmental control of enamel formation several observations of the enamel matrix suggest that enamel is formed incrementally. In human dental enamel for instance, there are two regularly occurring incremental markers: daily

cross-striations and long-period striae of Retzius (SR). These lines correspond to what was, at precise points in time during the secretory stage of amelogenesis, the enamel surface. Between the SR and running parallel with them are distinct growth lines known as cross-striations (Fig. 1). Cross-striations delineate the amount of enamel deposited in a single day [29,30]. Enamel cross-striations occur because of a circadian rhythm of ameloblast activity [27,31]. Similar growth lines have been showed recently in mouse enamel as well [32]. The distance between daily incremental lines averages 12 μm in rodents [27].

At the cellular level, we have shown that clock proteins are strongly expressed in pre-ameloblasts and differentiated ameloblasts throughout amelogenesis both in human embryonic teeth (Fig. 1) and mouse embryonic and post-natal teeth [8] (Fig. 1). Clock genes products are detected specifically in ameloblasts and their expression levels and localization alternates following circadian rhythms. In addition, melatonin receptors, key regulators of the circadian functions, are also differentially expressed in ameloblasts [33].




Fig. 1 – PER2 expression in mouse molar at PN21 stage (A). Dental pulp is devoid of staining in both incisors and molars (A). Odontoblasts in first molars are strongly stained for PER2 ((A) black arrowheads). PER2 shows relatively weak expression in periodontal dental ligament (PDL) cells when compared with odontoblast expression ((A) and (C)). Epithelial rests of Malassez (ERM) show strong expression of PER2 within the PDL space ((C) black arrows). PER2 proteins are also detected in the nucleus of osteoblasts and osteoclasts in the alveolar bone ((C) white arrow). On P4, both ameloblast and odontoblast show strong expression of PER2 in the nucleus (B). In human embryonic teeth, PER2 is detected in inner enamel epithelium (IEE) and outer enamel epithelium (OEE) but not in stellate reticulum (SR) and dental pulp ((D) and (E)). Black lines in figure F indicate the enamel cross striations in human tooth enamel. Ten cross striations are visualized between two successive Striae of Retzius (perpendicular lines). Am, ameloblast; DP, dental pulp; ERM, epithelial rests of Malassez; od, odontoblasts; Scale bars=200 μm in (A), 100 μm in (B)–(E), 10 μm in (F).

We have also recently published that there are daily variations in the rate of production and secretion of enamel proteins between early morning and late afternoon suggesting that enamel protein secretion is under circadian control [27] (Fig. 2). Enamel proteins secretion peaks at around 8 pm and follows a circadian pattern. We have also evaluated RNA oscillations of clock genes and amelogenin and enamelin in teeth of mice kept under normal

light and dark (LD) cycles at two time points: 10 am and 10 pm (Fig. 2). This preliminary data shows that *Per2* and amelogenin expression peaks at 10 pm with minimum expression at 10 am. In contrast, we have not detected any significant difference for enamelin expression. It is possible that enamelin is not under circadian control. Of significance, amelogenin expression continues to oscillate under constant dark (DD) conditions, however




Fig. 2 – Different amounts of proteins secreted in rat enamel are measured at different time points (A). Amelogenin RNA expression levels in developing mouse molars are higher at the night time compared to the day time. In contrast, enamelin RNA levels do not show significant differences between night and day time (B). Immunohistochemistry results show that BMAL1 (C) and PER2 (D) are differentially localized in ameloblasts at a given time point (black arrows). On serial sections, BMAL1 is mostly expressed in the cytoplasm (E) while PER2 is localized mainly in the nucleus (F). Most of BMAL1 proteins are localized in the cytoplasm at 8 am and then in the nucleus of ameloblasts at 8 pm (G). In contrast, PER2 proteins are found in the nucleus at 8 am and in the cytoplasm of ameloblasts at 8 pm (H). Relative mRNA levels are evaluated by qRT-PCR between wild type and knock-out mice kidneys ((I)–(J)) and in transfected HAT-7 cells (K). *Calbindin-D28k* (CABD 28K) RNA expression is altered in *Per2* knockout mice (I). Similar trend is be found in *Car2* RNA expression level (J). Over-expression of *Cry1* results in the down-regulation of *Amelx* and *Klk4* expression levels in HAT-7 cells (K). Am, ameloblast; od, odontoblast; nu, nucleus; cp, cytoplasm. Scale bars=20 μm in (A), 200 μm in (C) and (D), 40 μm in (E) and (F), 10 μm in (G) and (H).

in less extend and with a slight shift in peak times [31]. In vitro studies have also confirmed that amelogenin expression is modulated by clock genes [31,34,35] (Fig. 2). Consistent with the above findings, our preliminary analysis of *Per2* knock-out teeth shows abnormal enamel matrix formation (unpublished). Our data also suggest that Kallikrein 4 (*Klk4*), a key gene that regulates enamel maturation, might be under circadian control [31]. Taken together the above-discussed data, we postulate that enamel matrix production and maturation are closely controlled by selectively regulating key enamel matrix proteins encoding genes.

Clock genes expression is also synchronized in two other epithelial tissues with high similarities with ameloblasts, i.e. duodenum, and kidney. We found that genes involved in the calcium and phosphate metabolism such as calbindin-D28K (a calcium binding protein), carbonic anhydrase 2 (*Car2*), and alkaline phosphatase oscillate in regular daily intervals in kidney cells suggesting that these genes might be functional targets of clock genes. Furthermore, expression of calbindin-D28K and alkaline phosphatase is altered in *Per2* knock-out mice kidney (Fig. 2). It is then logical to hypothesize that these genes may also be targets of the circadian clock (CCG) in ameloblasts. Consistently with our hypothesis clock genes knockout mice (*Per2* KO) show also defects of enamel mineralization (unpublished data). However, the precise mechanisms of calcium regulation in ameloblasts remain to be elucidated.

CCG control of enamel related genes is suggested to be direct or indirect through the circadian regulation of key transcription factors as we showed for *Runx2* [34]. Recently, another group showed that *Runx2* is under circadian control in the suprachiasmatic nucleus and functions in the control of rhythmic behavior [36]. More studies are needed to fully comprehend the complex circadian controls that regulate enamel formation. Nevertheless, we can postulate that the ameloblast circadian clock plays significant roles in enamel secretion, maturation and mineralization.

Dentin

Similar to enamel, dentin is formed incrementally suggesting the involvement of a circadian clock mechanism during dentinogenesis. Daily incremental lines in dentin are termed von Ebner's lines. von Ebner's lines delineate the amount of mineral deposited in a single day [37]. The numbers of short-period incremental lines observed in enamel or dentin between sequentially administered dyes correspond to the number of days between successive administrations [38]. Circadian rhythms have also been demonstrated using ³H-proline tracers that label collagen during dentin formation [39]. Twice as much collagen is secreted during the daylight 12 h as during the night time 12 h.

At the cellular level, we recently showed evidence that odontoblasts express clock proteins [8]. Consistently, Ohtsuka-Isoya et al. [40] presented convincing evidence that SCN is related to the production of daily lines in dentin, as rats in which the SCN is fully or partially lesioned show respective permanent or temporary loss of circadian dentin increments. These studies suggest that dentin, similar to bone and enamel, is controlled by a circadian clock mechanism. However, very little is known on the molecular circadian mechanisms that regulate dentin formation. It will be interesting to elucidate the differential controls of dentin

formation by the master clock in the brain and/or a peripheral clock located in odontoblasts.

Cementum, and periodontal ligament

It has been described that human cementum is formed incrementally [41] and we found that periodontal ligament (PDL) cells differentially express clock proteins [8] (Fig. 1). Mouse cementoblasts lack of staining (Fig. 1) but we do not know if this is a generalized characteristic of cementoblasts. It is possible, that once cementum formation is completed clock expression is turned off in cementoblasts. Alternatively, species or teeth differences may exist. Of interest, Epithelial Rest of Mallasez (ERMs) strongly express clock proteins (Fig. 1). ERM are hypothesized to play a role in periodontal ligament regeneration. Further studies are needed to elucidate the potential links between clock genes and periodontal maintenance and turn over.

Clock genes and stem cells

A surprisingly finding of our analysis of clock gene expression in tooth is a strong expression of clock genes in areas rich in stem cells. For example, ERM cells, which are considered as dormant progenitor cells [42], show the strongest staining among all other cells in PDL (Fig. 1). Clock proteins are also strongly expressed by dental epithelium stem cells at the incisor cervical root area [8]. Of interest, clock genes mutants show variations in incisor size, being always smaller in *Per2* knockout mice (unpublished).

Similar to dental epithelia, hair cells express clock genes [43]. Indeed, clock genes modulate the human hair cycle clock independently of the brain suggesting that a peripheral hair clock also exists [44,45]. Clock genes and their products are also co-expressed with stem cells markers in hair and perturbations of circadian clocks result in changes in hair follicle renewal [46]. Of great interest, this study also reveals that disruption of clock equilibrium, through deletion of *Bmal1* or *Per1/2*, resulted in a progressive accumulation or depletion of dormant stem cells, respectively [46]. Hair and dental epithelium share common pathways [47]. We postulate that at least part of the developmental alterations observed in the enamel and hair of clock genes mutant mice are due to changes of stem cells behavior in these tissues. However, the potential role of the circadian clock in regulating stem cells behavior in tooth and hair remains elusive.

Conclusions

Both, the secretory and maturation stages of enamel formation are characterized by cyclic events that control cell morphology, gene expression and protein secretion and removal. In order to understand how enamel development is overall controlled, studying the circadian control of ameloblast differentiation is necessary. The concrete benefits of this research will provide the basis for a better understanding of enamel abnormalities and differences among the population. Once such knowledge becomes available, there is a promise that diseases associated with enamel formation can be better understood, diagnosed and treated. The time spent each day by ameloblasts to perform unique functions as well as the time spent into the secretory versus the maturation

stages are critical for the final product, the enamel. Enamel may vary considerably between individuals, due to differences in circadian profiles. Therefore, alteration of the circadian network would result in significant differences of enamel morphology, thickness and mineral content, predisposing teeth to disease or protecting them against it.

Equally important emerging evidence confirms that, bone, dentin and cementum are also under circadian controls. In the *Per2* ko mice both bone and teeth are affected. These data suggest that a synchronized formation of teeth and bone may be controlled by the central and/or peripheral circadian body clocks. This perspective opens new avenues of research that would study the potential communications between bone and teeth and/or their coordinated regulation by the circadian central and peripheral clocks. It is anticipated that circadian clock research will contribute to broader our understanding of how mineralized tissues are formed. A system's biology approach having as "skeleton" the circadian system may help us understand how mineralized tissues are developed in synchrony [17]. New inputs such as mastication, nutrition or bone load could be established in the circadian system equation increasing our understanding of this complex communication between the brain and the peripheral organs.

Although recent studies have shown that the behavior of some types of stem cells is regulated by the circadian clock, we are still far from understanding how circadian rhythmicity contributes to their function. Indeed, additional emerging evidence [15] and our own unpublished data suggests an important role of the circadian clock in regulating aging, cancer, and potentially tissue repair. Future studies on the relationship between clock genes and stem cells are expected to elucidate the contribution of the timing as a "fourth dimension" on the study of stem cells [48] in the years to come.

- [1] J. Richards, M.L. Gumz, Mechanism of the circadian clock in physiology, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 304 (2013) R1053–1064.
- [2] P.C. Konturek, T. Brzozowski, S.J. Konturek, Gut clock: implication of circadian rhythms in the gastrointestinal tract, *J. Physiol. Pharmacol.* 62 (2011) 139–150.
- [3] M. Ikeda, Calcium dynamics and circadian rhythms in supra-chiasmatic nucleus neurons, *Neuroscientist* 10 (2004) 315–324.
- [4] D.K. Welsh, D.E. Logothetis, M. Meister, S.M. Reppert, Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms, *Neuron* 14 (1995) 697–706.
- [5] E.M. Myers, J. Yu, A. Sehgal, Circadian control of eclosion: interaction between a central and peripheral clock in *Drosophila melanogaster*, *Curr. Biol.* 13 (2003) 526–533.
- [6] J.A. Williams, A. Sehgal, Molecular components of the circadian system in *Drosophila*, *Annu. Rev. Physiol.* 63 (2001) 729–755.
- [7] S. Masubuchi, N. Kataoka, P. Sassone-Corsi, H. Okamura, Mouse Period1 (mPER1) acts as a circadian adaptor to entrain the oscillator to environmental light/dark cycles by regulating mPER2 protein, *J. Neurosci.* 25 (2005) 4719–4724.
- [8] B. Zheng, U. Albrecht, K. Kaasik, M. Sage, W. Lu, S. Vaishnav, Q. Li, Z.S. Sun, G. Eichele, A. Bradley, C.C. Lee, Nonredundant roles of the mPer1 and mPer2 genes in the mammalian circadian clock, *Cell* 105 (2001) 683–694.
- [9] Y.S. Jang, Y.J. Kang, T.J. Kim, K. Bae, Temporal expression profiles of ceramide and ceramide-related genes in wild-type and mPer1/mPer2 double knockout mice, *Mol. Biol. Rep.* 39 (2012) 4215–4221.
- [10] J. Husse, S.C. Hintze, G. Eichele, H. Lehnert, H. Oster, Circadian clock genes Per1 and Per2 regulate the response of metabolism-associated transcripts to sleep disruption, *PLoS One* 7 (2012) e52983.
- [11] Y.J. Lee, D.H. Han, Y.K. Pak, S.H. Cho, Circadian regulation of low density lipoprotein receptor promoter activity by CLOCK/BMAL1, Hes1 and Hes6, *Exp. Mol. Med.* 44 (2012) 642–652.
- [12] G.K. Paschos, S. Ibrahim, W.L. Song, T. Kunieda, G. Grant, T.M. Reyes, C.A. Bradfield, C.H. Vaughan, M. Eiden, M. Masoodi, J.L. Griffin, F. Wang, J.A. Lawson, G.A. Fitzgerald, Obesity in mice with adipocyte-specific deletion of clock component Arntl, *Nat. Med.* 18 (2012) 1768–1777.
- [13] R.V. Kondratov, R.K. Shamanna, A.A. Kondratova, V.Y. Gorbacheva, M.P. Antoch, Dual role of the CLOCK/BMAL1 circadian complex in transcriptional regulation, *FASEB J.* 20 (2006) 530–532.
- [14] Y. Kumaki, M. Ukai-Tadenuma, K.D. Uno, J. Nishio, K.H. Masumoto, M. Nagano, T. Komori, Y. Shigeyoshi, J.B. Hogenesch, H.R. Ueda, Analysis and synthesis of high-amplitude cis-elements in the mammalian circadian clock, *Proc. Nat. Acad. Sci. U.S.A.* 105 (2008) 14946–14951.
- [15] P. Janich, K. Toufighi, G. Solanas, N.M. Luis, S. Minkwitz, L. Serrano, B. Lehner, S.A. Benitah, Human epidermal stem cell function is regulated by circadian oscillations, *Cell Stem Cell* (2013).
- [16] A.M. Schroeder, C.S. Colwell, How to fix a broken clock, *Trends Pharmacol. Sci.* (2013).
- [17] S. Papagerakis, L. Zheng, S. Schnell, M.A. Sartor, E. Somers, W. Marder, B. McAlpin, D. Kim, J. McHugh, P. Papagerakis, The circadian clock in oral health and diseases, *J. Dent. Res.* (2013).
- [18] C.M. Gundberg, M.E. Markowitz, M. Mizruchi, J.F. Rosen, Osteocalcin in human serum: a circadian rhythm, *J. Clin. Endocrinol. Metab.* 60 (1985) 736–739.
- [19] D.J. Simmons, G. Nichols Jr., Diurnal periodicity in the metabolic activity of bone tissue, *Am. J. Physiol.* 210 (1966) 411–418.
- [20] L. Fu, M.S. Patel, A. Bradley, E.F. Wagner, G. Karsenty, The molecular clock mediates leptin-regulated bone formation, *Cell* 122 (2005) 803–815.
- [21] L. Fu, M.S. Patel, G. Karsenty, The circadian modulation of leptin-controlled bone formation, *Prog. Brain Res.* 153 (2006) 177–188.
- [22] E. Maronde, A.F. Schilling, S. Seitz, T. Schinke, I. Schmutz, G. van der Horst, M. Amling, U. Albrecht, The clock genes Period 2 and Cryptochrome 2 differentially balance bone formation, *PLoS One* 5 (2010) e11527.
- [23] J.D. McElderry, G. Zhao, A. Khmaladze, C.G. Wilson, R.T. Franceschi, M.D. Morris, Tracking circadian rhythms of bone mineral deposition in murine calvarial organ cultures, *J. Bone Miner. Res.* 28 (2013) 1846–1854.
- [24] T. Iimura, A. Nakane, M. Sugiyama, H. Sato, Y. Makino, T. Watanabe, Y. Takagi, R. Numano, A. Yamaguchi, A fluorescence spotlight on the clockwork development and metabolism of bone, *J. Bone Miner. Metab.* 30 (2012) 254–269.

- [25] D.J. DiGirolamo, T.L. Clemens, S. Kousteni, The skeleton as an endocrine organ, *Nat. Rev. Rheumatol.* 8 (2012) 674–683.
- [26] J.C. Hu, Y.H. Chun, T. Al Hazzazi, J.P. Simmer, Enamel formation and amelogenesis imperfecta, *Cells Tissues Organs* 186 (2007) 78–85.
- [27] J.P. Simmer, P. Papagerakis, C.E. Smith, D.C. Fisher, A.N. Rountrey, L. Zheng, J.C. Hu, Regulation of dental enamel shape and hardness, *J. Dent. Res.* 89 (2010) 1024–1038.
- [28] C.E. Smith, Cellular and chemical events during enamel maturation, *Crit. Rev. Oral Biol. Med.* 9 (1998) 128–161.
- [29] J.A. Dean, D.R. Avery, M.L. Swartz, Attachment of anterior tooth fragments, *Pediatr. Dent.* 8 (1986) 139–143.
- [30] S. Risnes, C.B. Moinichen, D. Septier, M. Goldberg, Effects of accelerated eruption on the enamel of the rat lower incisor, *Adv. Dent. Res.* 10 (1996) 261–269.
- [31] L. Zheng, Y.J. Seon, M.A. Mourao, S. Schnell, D. Kim, H. Harada, S. Papagerakis, P. Papagerakis, Circadian rhythms regulate amelogenesis, *Bone* 55 (2013) 158–165.
- [32] A. Sehic, M. Nirvani, S. Risnes, Incremental lines in mouse molar enamel, *Arch. Oral Biol.* 58 (2013) 1443–1449.
- [33] S. Kumasaka, M. Shimosuma, T. Kawamoto, K. Mishima, R. Tokuyama, Y. Kamiya, P. Davaadorj, I. Saito, K. Satomura, Possible involvement of melatonin in tooth development: expression of melatonin 1a receptor in human and mouse tooth germs, *Histochem. Cell Biol.* 133 (2010) 577–584.
- [34] M. Athanassiou-Papaefthymiou, D. Kim, L. Harbron, S. Papagerakis, S. Schnell, H. Harada, P. Papagerakis, Molecular and circadian controls of ameloblasts, *Eur. J. Oral Sci.* 119 (Suppl. 1) (2011) 35–40.
- [35] R.S. Lacruz, J.G. Hacia, T.G. Bromage, A. Boyde, Y. Lei, Y. Xu, J.D. Miller, M.L. Paine, M.L. Snead, The circadian clock modulates enamel development, *J. Biol. Rhythms* 27 (2012) 237–245.
- [36] M.E. Reale, I.C. Webb, X. Wang, R.M. Baltazar, L.M. Coolen, M.N. Lehman, The transcription factor Runx2 is under circadian control in the suprachiasmatic nucleus and functions in the control of rhythmic behavior, *PLoS One* 8 (2013) e54317.
- [37] M.C. Dean, Comparative observations on the spacing of short-period (von Ebner's) lines in dentine, *Arch. Oral Biol.* 43 (1998) 1009–1021.
- [38] M.C. Dean, A.E. Scandrett, The relation between long-period incremental markings in dentine and daily cross-striations in enamel in human teeth, *Arch. Oral Biol.* 41 (1996) 233–241.
- [39] M. Ohtsuka, S. Saeki, K. Igarashi, H. Shinoda, Circadian rhythms in the incorporation and secretion of ^3H -proline by odontoblasts in relation to incremental lines in rat dentin, *J. Dent. Res.* 77 (1998) 1889–1895.
- [40] M. Ohtsuka-Isoya, H. Hayashi, H. Shinoda, Effect of suprachiasmatic nucleus lesion on circadian dentin increment in rats, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 280 (2001) R1364–1370.
- [41] S.I. Kvaal, T. Solheim, D. Bjerketvedt, Evaluation of preparation, staining and microscopic techniques for counting incremental lines in cementum of human teeth, *Biotech. Histochem.* 71 (1996) 165–172.
- [42] K. Oka, M. Morokuma, K. Imanaka-Yoshida, Y. Sawa, K. Isokawa, M.J. Honda, Cellular turnover in epithelial rests of Malassez in the periodontal ligament of the mouse molar, *Eur. J. Oral Sci.* 120 (2012) 484–494.
- [43] M. Tanioka, H. Yamada, M. Doi, H. Bando, Y. Yamaguchi, C. Nishigori, H. Okamura, Molecular clocks in mouse skin, *J. Invest. Dermatol.* 129 (2009) 1225–1231.
- [44] Y. Al-Nuaimi, J.A. Hardman, T. Biro, I.S. Haslam, M.P. Philpott, B.I. Toth, N. Farjo, B. Farjo, G. Baier, R.E. Watson, B. Grimaldi, J.E. Kloepper, R. Paus, A meeting of two chronobiological systems: circadian proteins period1 and BMAL1 modulate the human hair cycle clock (2013) *J. Invest. Dermatol.* (2013).
- [45] Y. Watabe, M. Tomioka, A. Watabe, M. Aihara, S. Shimba, H. Inoue, The clock gene brain and muscle Arnt-like protein-1 (BMAL1) is involved in hair growth, *Arch. Dermatol. Res.* 305 (2013) 755–761.
- [46] P. Janich, G. Pascual, A. Merlos-Suarez, E. Batlle, J. Ripperger, U. Albrecht, H.Y. Cheng, K. Obrietan, L. Di Croce, S.A. Benitah, The circadian molecular clock creates epidermal stem cell heterogeneity, *Nature* 480 (2011) 209–214.
- [47] J. Pispas, I. Thesleff, Mechanisms of ectodermal organogenesis, *Dev. Biol.* 262 (2003) 195–205.
- [48] J.M. Gimble, Z.E. Floyd, B.A. Bunnell, The 4th dimension and adult stem cells: can timing be everything?, *J. Cell Biochem.* 107 (2009) 569–578.