



HAL
open science

The magnetopause position in the case of an expanding ionosphere

Jean-Mathias Griessmeier, E. Johansson, H. Lammer, U. Motschmann

► To cite this version:

Jean-Mathias Griessmeier, E. Johansson, H. Lammer, U. Motschmann. The magnetopause position in the case of an expanding ionosphere. "European Planetary Science Congress 2010, 2010, à renseigner, Unknown Region. pp.236. <hal-03548990>

HAL Id: hal-03548990

<https://hal.science/hal-03548990v1>

Submitted on 31 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



The magnetopause position in the case of an expanding ionosphere

J.-M. Grießmeier (1), E. Johansson (2), **H. Lammer** (3) and U. Motschmann (2)

(1) Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E) & Observatoire des Sciences de l'Univers en région Centre (OSUC), Orleans, France (griessmeier@astron.nl), (2) Technical University of Braunschweig, Germany, (3) Space Research Institute, Graz, Austria

Abstract

The magnetosphere of a planet provides protection against the planetary environment. Its size and location determine how well the planet is shielded against the stellar wind, stellar Coronal Mass Ejections, or stellar and galactic cosmic rays. The question of magnetospheric shielding is particularly relevant for close-in extrasolar planets, where the magnetosphere can be strongly compressed by the enhanced stellar wind. The size of the magnetosphere and the position of the magnetopause are determined by the pressure equilibrium between the pressure exerted by the stellar wind and the pressure contributions from within the magnetosphere (see e.g. [1]). The standoff distance R_S is defined as the planetocentric distance of the magnetopause along the line connecting the planet and the star where the pressure equilibrium is satisfied. In the case of a magnetized planet, the magnetospheric pressure is usually dominated by the contribution caused by the planetary magnetic field. Here, we discuss the case of a non-magnetized or weakly magnetized planet, where the magnetospheric pressure is provided by a strongly expanding planetary ionosphere. Such ionospheres are expected for close-in exoplanets where strong stellar heating leads to intense photoionization, but may also be of relevance for the early terrestrial evolution during the Hadean epoch. Both analytical and numerical results from a hybrid simulation [2] are shown and compared.

References

- [1] Grießmeier, J.-M., Stadelmann, A., Grenfell, J. L., Lammer, H. and Motschmann, U.: On the protection of extrasolar Earth-like planets around K/M stars against galactic cosmic rays, *Icarus*, 199, 526–535, 2009.
- [2] Johansson, E. P. G., Bagdonat, T., and Motschmann, U.: Consequences of expanding exoplanetary atmospheres for magnetospheres, *Astron. Astrophys.*, 496, pp. 869–877, 2009.