



HAL
open science

Inverse identification of bi-material soft tissue parameters

Aflah Elouneg, Danas Sutula, Arnaud Lejeune, Jérôme Chambert, Franz Chouly, Stéphane Bordas, Emmanuelle Jacquet

► **To cite this version:**

Aflah Elouneg, Danas Sutula, Arnaud Lejeune, Jérôme Chambert, Franz Chouly, et al.. Inverse identification of bi-material soft tissue parameters. 14th World Congress on Computational Mechanics (WCCM) ECCOMAS 2020, Jan 2021, Paris, France. hal-03549361

HAL Id: hal-03549361

<https://hal.science/hal-03549361>

Submitted on 31 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INVERSE IDENTIFICATION OF BI-MATERIAL SOFT TISSUE PARAMETERS

A. Eloune^{1,*}, D. Sutula¹, A. Lejeune¹, J. Chambert¹, F. Chouly²,
S.P.A. Bordas³, E. Jacquet¹

¹ Univ. Bourgogne Franche-Comté, FEMTO-ST institute, UFC/CNRS/ENSMM/UTBM, Department of Applied Mechanics, Besançon, France, aflah.eloune@femto-st.fr

² Univ. Bourgogne Franche Comté, Institut de Mathématiques de Bourgogne, Dijon, France.

³ University of Luxembourg, Institute of Computational Engineering, Luxembourg.

Key Words: Inverse Identification, Hyperelasticity, Digital Image Correlation, Skin Tumors.

Keloid scars are considered to be benign tumors of the skin that can grow beyond the initial limits of a wound. The preferential appearance of keloids in certain anatomical sites and their pattern of growth attests to the influence of the mechanical stress as one of the driving factors [1]. Thus, a mechanical characterization of a keloid scar is to be done in order to better understand the keloid and, ultimately, to prevent its growth. A uniaxial tensile experiment was performed *in vivo* on a young caucasian female with a keloid scar located on the upper part of her left arm. The reaction force measurements were acquired by a force sensor in the loading cell and the surface displacement field was captured using the Digital Image Correlation technique [2]. The scarred skin was modelled as a bi-material 2D-structure. The non-linear forward solver used artificially generated measurements with an additive white Gaussian noise [3]. By using the simulated data, the material parameters of the both the keloid and the healthy skin media were identified through an FEMU-based open-source framework [4, 5]. The results show that it is possible to identify accurately at least 4 material parameters from a uniaxial test. Different uncertainty analyses including noise levels and the number of snapshots have been performed to explore the computational limits of model parameter identifiability. Our next steps consist of using the real experimental data to identify the material parameters.

REFERENCES

- [1] R. Ogawa, “Keloid and hypertrophic scarring may result from a mechanoreceptor or mechanosensitive nociceptor disorder,” *Med. Hypotheses*, vol. 71, pp. 493–500, 2008.
- [2] J. Chambert, T. Lihoreau, S. Joly, B. Chatelain, P. Sandoz, P. Humbert, E. Jacquet, and G. Rolin, “Multimodal investigation of a keloid scar by combining mechanical tests *in vivo* with diverse imaging techniques,” *J. Mech. Behav. Biomed. Mater.*, vol. 99, pp. 206–215, 2019.
- [3] H. Rappel, L. A. A. Beex, and S. P. A. Bordas, “Bayesian inference to identify parameters in viscoelasticity,” *Mech. Time-Depend. Mater.*, vol. 22, pp. 221–258, May 2018.
- [4] H.-P. Chen and Y. Ni, “Finite element model updating,” in *Structural Health Monitoring of Large Civil Engineering Structures*, ch. 6, pp. 123–154, John Wiley & Sons, Ltd, 2018.
- [5] P. Hauseux, J. S. Hale, S. Cotin, and S. P. Bordas, “Quantifying the uncertainty in a hyperelastic soft tissue model with stochastic parameters,” *Appl. Math. Model.*, vol. 62, pp. 86–102, 2018.