

The experimental emergence of convention in a non-human primate

Anthony Formaux, Dany Paleressompoulle, Joël Fagot, Nicolas Claidière

► To cite this version:

Anthony Formaux, Dany Paleressompoulle, Joël Fagot, Nicolas Claidière. The experimental emergence of convention in a non-human primate. Philosophical Transactions of the Royal Society B: Biological Sciences, 2021, 377, 10.1098/rstb.2020.0310. hal-03549748

HAL Id: hal-03549748 https://hal.science/hal-03549748

Submitted on 31 Jan2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Formaux, A., Paleressompoulle, D., Fagot, J., & Claidière, N. (2022). The experimental emergence of convention in a non-human primate. *Philosophical Transactions of the Royal Society B: Biological Sciences, 377*(1843), 20200310. doi:10.1098/rstb.2020.0310

The experimental emergence of convention in a non-

human primate

Anthony Formaux^{1,2}, Dany Paleressompoulle¹, Joël Fagot^{1,2,3}, Nicolas

Claidière^{1,2,3*}

1 – Aix Marseille Univ, CNRS, LPC, Marseille, France

2 – Station de Primatologie-Celphedia, CNRS UAR846, Rousset, France

3 - Institute for Language, Communication and the Brain, Université Aix-Marseille, CNRS, Aix-en-

Provence, France.

* - Correspondance to : Nicolas Claidière, Laboratoire de Psychologie Cognitive, UMR7290, Université Aix-Marseille/CNRS, 13331 Marseille, France. Email: <u>nicolas.claidiere@normalesup.org</u>

OrcidID : Anthony Formaux (0000-0001-7857-2349), Joël Fagot (0000-0002-9824-9685), Nicolas Claidière (0000-0002-4472-6597)

1

2 **1. Summary**

Conventions form an essential part of human social and cultural behaviour and may also be important to other animal societies. Yet, despite the wealth of evidence that has accumulated for culture in non-human animals, we know surprisingly little about non-human conventions beyond a few rare examples. We follow the literature in behavioural ecology and evolution and define conventions as systematic behaviours that solve a coordination problem in which two or more individuals need to display complementary behaviour to obtain a mutually beneficial outcome. We

9 start by discussing the literature on conventions in non-human primates from this perspective and 10 conclude that all the ingredients for conventions to emerge are present and therefore that they ought to be more frequently observed. We then probe emergence of conventions by using a unique 11 12 novel experimental system in which pairs of Guinea baboons (Papio papio) can voluntarily 13 participate together in touch-screen based cognitive testing and we show that conventions readily 14 emerge in our experimental setup and that they share three fundamental properties of human 15 conventions (arbitrariness, stability and efficiency). These results question the idea that observational learning, and imitation in particular, is necessary to establish conventions, they 16 17 suggest that positive reinforcement is enough.

Keywords: primate social cognition, social learning, animal culture, language evolution,
 imitation

20 **2. Introduction**

21 Conventions form an essential part of human social and cultural behaviour and may also be 22 important to other animal societies. Yet only a few rare examples of conventions have been 23 documented (see Stephens & Heinen, 2018 for a discussion of non-human conventions), despite the 24 existence of evolutionary precursors (Kappeler, Fichtel, & van Schaik, 2019). The first part of this 25 article discusses current evidence of conventions in non-human primates and concludes that 26 conventions ought to be more frequently observed. The scarcity of convention has been explained 27 by the limited social learning capacities of non-human primates, especially regarding imitation. 28 However, in the second part of the article we present a set of experiments showing the emergence 29 of convention in a non-human primate using a unique novel experimental system. These 30 experiments show that Guinea baboons (Papio papio) can readily develop conventions that share at 31 least three properties of human conventions (arbitrariness, stability and efficiency) through simple 32 reinforcement learning. We start by clarifying the meaning and properties of convention.

Intuitively conventions are rule-like behaviour such as shaking hands as a greeting. Beyond this intuitive understanding however, conventions have proven difficult to define and have been much debated. Here, we will follow a 'Lewisian' approach (Lewis, 1969) and define conventions as systematic behaviours that solve a coordination problem in which two or more individuals need to display complementary behaviour to obtain a mutually beneficial outcome.

38 A classic example of convention, used by Hume (1888) and taken up by Lewis, is that of two 39 rowers on a boat. If both rows are synchronized, the boat will move forward smoothly, whereas if they row out of sync, it will behave erratically. In Hume's (then Lewis's) use, this example illustrates 40 41 the main properties of convention: 1) rowers do not keep promises to act in concert, 2) they may 42 express their shared interest explicitly, but not necessarily, 3) to row in rhythm, there must be a 43 gradual installation of synchronous behaviour, 4) they may know that what is being set up between 44 them is a convention, but not know how to describe it verbally ("We start at the 90° angle and give 45 a stroke every 2 seconds") and 5) they have adopted a rhythm that they keep because it works, but there are alternatives. We discuss some of these characteristics in more detail below. 46

47 Individuals can find *ad hoc* solutions to coordination problems when they arise, as when 48 finding seating arrangements around the table for instance. However, when coordination problems 49 happen regularly and/or impose a high cost for a failure to coordinate, individuals may benefit from 50 establishing a convention (Hawkins & Goldstone, 2016). Notice that compared to other types of 51 positive interactions such as helping (one individual benefits at the expense of another) or 52 cooperation (individuals can benefit more by being selfish), individuals mutually benefit from 53 coordination: it is in their own best interest and in the interest of the partner(s) to coordinate. 54 Therefore, there is no conflict of interest between individuals when solving a coordination problem 55 or when establishing a convention. Rather, conventions are important because they help stabilize 56 interactions by creating mutual expectations between individuals. Someone picking up his/her car 57 in France expects other individuals to drive on the right side of the road and so adjusts his/her 58 behaviour accordingly. These expectations need not be explicit nor mutually acknowledged for a

59 convention to exist (Moore, 2013). In fact, sometimes we are barely aware that a convention exists: 60 you might lunch at a certain time simply because you expect other people to lunch around that time. According to Hawkins' theoretical model (2019), three processes allow the emergence and 61 62 maintenance of a convention: the population level (social network management and cultural 63 transmission), the dyad level (coordination and social interaction) and the individual level (executive functions and expectations). A convention is not necessarily created as a result of a problem 64 65 encountered by the whole population at one point. If enough dyads in the population repeatedly 66 encounter a coordination problem and each individual belongs to several dyads, a population-wide 67 convention can emerge.

68 Of course, a paradigmatic example of convention in humans is language because to 69 communicate individuals need to coordinate on word-meaning relationships. Studies have shown 70 how humans can develop new conventions to communicate during the natural emergence of new 71 languages such as pidgin (which occurs when individuals with no common language need to 72 communicate, Bickerton, 1981) or sign language (for instance the Nicaraguan sign language for 73 example, Kegl, 1994) and these studies have been complemented by experiments and models 74 showing how conventions develop during the emergence of artificial communication systems (Fay, 75 Garrod, Roberts, & Swoboda, 2010; Galantucci, 2005). Conventions are traditionally distinguished 76 from habits and norms. Habits are patterns of individual behaviour that do not (strongly) depend on 77 other individuals, but they can be shared and cultural: taking a shower vs. taking a bath for instance. 78 In contrast, norms may be seen as solving coordination problems when the interest of individuals are not aligned. They are associated to a sense of "oughtness" and punishment that conventions 79 80 and habits lack, e.g. "crossing at a red light" (e.g. Bicchieri, 2005). Clearly, given the diversity of 81 human social behaviour, there are no strict boundaries between habits, conventions, and norms. 82 What was once a habit, "spitting in the street", can become a norm "do not spit" (Nichols, 2002). Conventions, e.g. "crossing at pedestrian crossings" can be felt to some like norms that ought to be 83 84 followed at all times and by others as mere conventions. There are also habits, "eating cheese

before dessert", that, although not a coordination problem *per se*, are easier to deal with when
everyone follows the same rules (e.g. to have a meal together). Therefore, conventions are not an
"all or nothing" kind of thing but a "more or less" one: behaviours are more or less conventional
depending on how closely they match the prototypical example of convention.

89 In the framework of game theory (Deming, Neumann, & Morgenstern, 1944), a coordination 90 problem is a problem with multiple Nash equilibriums (a strategy is a Nash equilibrium when no 91 player can do better by changing his strategy alone; (Nash, 1951)) and a convention is a Nash 92 equilibrium to such problem (see Lewis (1969) for the original discussion and Stephens and Heinen 93 (2018) for a more detailed discussion in the context of animal behaviour). The Nash equilibrium, i.e. 94 the convention, is governed by initial conditions and dynamics (influenced by the way individuals 95 choose to act and the environment). Note that conventions require multiple equilibriums because if 96 there are no alternative options, there would be no coordination problem to solve in the first place 97 and two individuals would necessarily converge on the unique solution. However, different 98 equilibriums need not be equally rewarding, one may lead to higher rewards for instance (called 99 payoff dominant strategies) and some equilibriums may be more rewarding than others when 100 individuals deviate from the equilibrium (risk dominant equilibrium). Note that in the framework of 101 game theory a convention is 'arbitrary' because there are multiple Nash equilibriums to a 102 coordination problem, so a population could have ended up using another convention. It does not 103 mean that individuals are arbitrarily choosing one behaviour over another, they may strategically 104 choose a payoff dominant strategy for instance. Importantly, compared to their one-shot 105 counterpart, repeated economic games can help create conventions because with repetition 106 multiple equilibriums often appear (Fudenberg & Maskin, 2008).

107 To summarise, when individuals regularly face coordination problems, they may develop 108 conventions as solutions to these problems. Conventions are systematic behaviours that are 109 mutually beneficial and provide stable expectations about other individuals. Thus conventions, like 110 norms, help individuals stabilize their social environment and navigate their social world. It is important to note that conventions do not require individuals to do the same action (one could pull

112 while the other pushes), they can concern as few as two individuals, and they do not necessarily

113 lead to the optimal solution (other more profitable equilibria may exist). Conventions are usually

associated with three important properties (Hawkins & Goldstone, 2016):

- Arbitrariness: at least one alternative exists and would be equally acceptable if everyone
 coordinated on it.
- 117 2. Efficiency: individuals benefit from establishing a convention.
- 118 3. **Stability:** over multiple games behaviour converges towards an attractor.

119 Conventions are the natural outcome of individuals repeatedly facing coordination problems

120 together and are therefore expected in group-living animals and especially in tightly bonded non-

121 human primates. Recently, Stephens et al. (2018) reviewed evidence of conventions in non-human

animals using a game-theoretic perspective but did not discuss primates, and Kappeler et al. (2019)

123 provided an important analysis of the evolutionary behavioural, emotional and cognitive precursors

124 of norms and conventions in non-human primates. Here, we complement this work by discussing

125 specific cases of non-human primate behaviour in the context just discussed that is graded and

126 grounded in a game-theoretic perspective.

3. Conventions in non-human primates?

128 Susan Perry and colleagues' description of social conventions in white-faced capuchin 129 monkeys (Cebus capucinus) is one of the best description of conventions we have in non-human 130 primates (Perry et al., 2003). Using 13 years of data collection in their field sites in Costa Rica, Perry 131 et al. describe the fashion-like emergence of new behaviour. During hand-sniffing for instance (first 132 reported by Fedigan, 1993) two individuals are resting together and one individual holds the hand 133 or foot of another over its own face. These behaviours are similar in many respects to human 134 conventions such as shaking hands: they require coordination between pairs of individuals because 135 individuals need to be together and to perform complementary behaviour, emerge in some groups

136 and not others and can spread to a large proportion of individuals within the group. They can also 137 stay stable for a long time, in one group hand-sniffing was observed during seven years before 138 disappearing. Finally, these behaviours could have a function close to our handshakes because they 139 may strengthen the bonds between individuals. These behaviours therefore appear to have all the 140 properties of conventions. Similarly, the hand-clasp grooming (McGrew, Marchant, Scott, & Tutin, 2001; van Leeuwen, 2021; van Leeuwen, Cronin, Haun, Mundry, & Bodamer, 2012) and social-141 142 scratch grooming of chimpanzees (Nakamura, McGrew, Marchant, & Nishida, 2000) also appear to 143 meet the requirements of conventions: they are stable behaviour that solve coordination problems 144 for which alternative behaviour exist. In addition, the social structure may vary between groups of 145 the same species and some behaviours may be considered conventional under the definition 146 proposed here. For example, van de Waal (2018) showed that neighbouring groups of vervet 147 monkeys differ in the structure of their social networks as well as how they handle conflict. When 148 conflicts arise within a group, alternative behaviours can be adopted to bring back stability, some 149 groups adopt affiliative behaviours (conciliation or reconciliation) while others adopt agonistic 150 behaviours (coalition or redirection). Another similar example is the emergence and transmission of 151 a "pacific culture" within a troop of wild olive baboons (Papio anubis) (Sapolsky, 2006; Sapolsky & 152 Share, 2004). Due to unfortunate circumstances, a large proportion of the most aggressive males of 153 the troop died of tuberculosis, the few remaining males were untypically affiliative and non-154 aggressive, creating a troop with high rates of grooming and relaxed dominance hierarchy. The low 155 level of aggression, compared to a neighbouring group residing in the same reserve, was stable and 156 maintained in the group for over 20 years despite the migration of new individuals. These are good 157 examples of convention, because different behaviours can be stable solutions to coordination 158 problems (how to respect the dominance hierarchy for instance) and individuals benefit from a 159 stable social organisation (social stability has important fitness benefits (Gilby et al., 2013; Schülke, 160 Bhagavatula, Vigilant, & Ostner, 2010; Silk, 2007; Silk et al., 2009).

161 Notably, most social learning experiments (see Heyes, 1994 for a definition of social 162 learning) do not result in the establishment of conventions, mostly because they do not involve 163 coordination problems but individual solving abilities. For instance, in the classical study of Whiten, 164 Horner, and de Waal (2005), chimpanzees can solve a puzzle box using two different techniques and 165 they learn to do so by watching other individuals. Since only one individual solves the problem at a 166 time, there is no coordination between them regarding the technique they use. It also emphasizes the difficulty of defining precisely what a convention is, and the fine line between social learning 167 168 and convention studies.

169 Some social learning experiments however show suggestive evidence of conventionality. In 170 the studies of van de Waal, Borgeaud, and Whiten (2013) for instance, groups of vervet monkeys 171 (Chlorocebus aethiops) learned to eat food of one colour (e.g. pink) and avoid food from another 172 colour (e.g. blue). Once this preference had been learned, results showed that monkeys continued to prefer to eat food of only one colour even when they were exposed to edible versions of both 173 174 colours. Even more surprisingly, monkeys that migrated from one group (e.g. pink) to another (e.g. 175 blue) sometimes adopted the behaviour of members of the group against their own previous 176 preference (e.g. they switch from pink to blue). One possible explanation for the vervet monkeys' behaviour is that they needed to compete for food to establish their dominance within the group¹. 177 178 Indeed, dominance hierarchies are examples of conventions, they are stable solutions to 179 coordination problems (to access food and/or reproduction) and provide stable expectations

¹ We thank Kate Arnold (University of St Andrews) for pointing this out to us.

regarding others' behaviour (see Bergman, Beehner, Cheney, & Seyfarth, 2003). In a particular group there certainly are several stable solutions to organise the dominance between individuals, that organisation is therefore conventional to some extent. However, the dominance hierarchy is also typical of a species and constrained by genetic dispositions and although it could be tempting to extend the concept of conventions to behaviour that are largely genetically determined

185 (Stephens & Heinen, 2018), here we will limit ourselves to learned conventions.

186 This consideration is also important to determine if non-human primate communication is 187 conventional. Non-human primates' vocalisations and gestures come from a largely fixed repertoire 188 and are not conventional (e.g. Arnold & Zuberbühler, 2008; Kemp et al., 2017; Ouattara, Lemasson, 189 & Zuberbühler, 2009; Tomasello & Zuberbühler, 2002). Although, note that conventions, in the form 190 of geographical dialects, have been observed in other vocal-learning non-human species (e.g. Slater, 191 1986; Weilgart & Whitehead, 1997) and that there is some limited evidence of vocal flexibility in 192 non-human primates (Snowdon & Elowson, 1999; Watson et al., 2015). Watson et al. (2015) in 193 particular observed the convergence of food calls during the integration of two chimpanzee groups 194 over a three-year period.

195 In non-human primates, some flexible gestural communications can emerge through 196 "ontogenetic ritualisation" (Tomasello & Call, 1997), a process through which an action at the 197 beginning of an action sequence, such as raising an arm before play-hitting another individual, 198 progressively becomes a signal to initiate the action sequence, raising an arm to initiate rough-and-199 tumble play (Tomasello, 2008). However, ontogenetic ritualisation does not systematically give rise 200 to conventions because there is a direct link between the sequence of actions to perform and the 201 signal that is used to initiate it, there is therefore no alternative signal through which this could be 202 initiated (Moore, 2013). Chimpanzee and bonobo gestures overlap extensively in form and meaning 203 for instance (Graham, Hobaiter, Ounsley, Furuichi, & Byrne, 2018). However, in one group of 204 mandrills (Mandrillus sphinx) Laidre (2011) described the emergence of a novel eye-covering 205 gesture probably used to signal that the individual was resting. This gesture was unique to this

group and had been observed for nearly a decade in the group and never seen elsewhere where
other behaviours are used to signal 'resting' (such as lying down or turning one's back). Therefore,
this signal may be seen as conventional to some extent, even if it is iconic. Interestingly, during the
emergence of sign language when signals are created, they are initially iconic, motivated by features
of the situated interaction. Later, iconic signals become arbitrary symbols as they are stylised and
ritualised. This happens to the point that new learners use them without awareness of the initial
motivation, and, in many cases, the initial iconic motivation is undetectable (Garrod et al. 2007).

213 Of course, the absence of conventions in non-human primates' communication contrasts 214 sharply with human language that is often considered to be a paradigmatic example of convention 215 since the word-meaning association is largely arbitrary (Lewis, 1969). Noteworthily, Tomasello 216 (2008) has proposed that linguistic and other human conventions are learned by imitation and that 217 the lack of a capacity for imitation in non-human primates could explain both their apparent lack of 218 conventions and their lack of language (see also Moore, 2013). Tamariz (2019) goes further and 219 argues that it is not imitation (copying goals and behaviour) but copying without paying attention to 220 goals, that makes language uniquely human (Tamariz, 2019). For example, children first learn to 221 reproduce the sound of a word before they learn the conventional way to use them.

222 The experiments we present below come to question the role of imitation in the 223 establishment of conventions since we observe the emergence of convention in a non-human 224 primate without imitation or other forms of observational learning. In our experiments two 225 baboons are seated next to each other facing computer touchscreens and they can see each other's 226 screen. When the experiment starts one baboon is randomly selected and must choose between 227 two stimuli, then the next individual is presented with the same two stimuli and must choose one 228 too. When the same two stimuli are selected, the two baboons are rewarded. Using this 229 coordination task, we wanted to ask the following three main questions:

- Will the baboons solve this coordination problem by developing a convention?
 Compared to choosing the stimuli independently or to responding by watching what the
 other individual selected.
- 233 2. If the baboons develop a convention, is observational learning necessary or important?
- 3. If the baboons develop a convention, to what extent are they arbitrary and depend onthe group social structure?
- 236

4. Method

238 Section 4.1 Participants

Nineteen Guinea baboons (*Papio papio*, 13 females) from the CNRS primatology center in Rousset-sur-Arc were tested in this study (mean age: 12.2 years [min=3.25; max=25.3]). They belonged to a social group of 19 individuals living in a 25x30m outdoor enclosure connected to a 6x4m indoor enclosure and two 8x4m trailers (Fig.1a). The monkeys had *ad libitum* access to five S-ALDMs (Social - Automated Computer Learning Devices for Monkeys; see below) four in the first trailer and six in the second.

245 Social Automated Computer Learning Devices for Monkeys (S-ALDMs)

246 The ALDMs (Fagot & Bonté, 2010; Fagot & Paleressompoulle, 2009) are operant conditioning test 247 systems that can be used for testing non-human primates in social setting. ALDMs use an automatic radio frequency identification device implanted in each forearm of the monkeys to detect the 248 249 location and identity of specific individuals. This device makes it possible to test the animals without 250 having to capture and isolate them (additional information of the setup can be found in Fagot, 251 Gullstrand, Kemp, Defilles, & Mekaouche, 2014; Fagot, Marzouki, Huguet, Gullstrand, & Claidière, 252 2015). The ALDMs presented in Fagot et al. (2015) were modified in 2018 and connected in pairs to allow an individual in one ALDM to both see its partner and its touchscreen of the neighboring 253 254 ALDM (see Fig.1b). We dubbed this new version of ALDMs the Social-ALDMs or S-ALDMs for short.

Two individuals could therefore see each other and their responses on the screen when a transparent partition between the S-ALDM was used or they had no visual access when an opaque partition was used. All the monkeys had previously participated in studies using this S-ALDM testing system but had never performed the present experiment.



259

Figure 1: (a) 19 baboons living in an enclosure have *ad libitum* access to two trailers, each containing several S-ALDMs. The diagram shows the two bungalows and the five units. (b) Detail of the organization of a workstation, where two monkeys can work side by side, while seeing each other.

264 **Computerised task**

We used two different tasks, one when two individuals are present, the "dual task", and one when an individual is alone, the "filler task". These two tasks are presented below. When a monkey entered an S-ALDM, a blue screen was displayed in its monitor with a 4-second delay. If another individual was detected in the neighboring S-ALDM during this delay, a blue screen was also displayed on the adjoining monitor, announcing the synchronization of the two machines and the start of the dual task (Fig.2 and video in ESM). Otherwise, if the neighboring S-ALDM stayed empty

- for 4 secs, the filler task started. The function of the filler task was mainly to keep the baboons
- 272 occupied within the S-ALDM while waiting for another individual to arrive.
- 273 Dual task



274

Figure 2: Schematic representation of the dual task on touch screen. A to E: succession of events.

277 During the dual task (Fig. 2), which is the focus of this paper, the test program identified the two monkeys when they entered an S-ALDM, synchronized the two computers, and displayed a blue 278 279 fixation cross at the bottom center of both screens (Fig.2B). The test started once both monkeys 280 had pressed the fixation cross and within a delay of 4-secs (otherwise the trial was aborted and re-281 presented). One monkey of the pair was then randomly selected. This first monkey, that we are 282 going to call "proposer", had to choose between two randomly placed visual stimuli on the screen 283 (Fig.2C). These stimuli were located on the side of the screen so that the second monkey, the 284 "responder", had the best possible view. Once the stimulus was selected by the proposer, it would 285 flash twice (Fig.2D), then both stimuli would disappear and appear on the responder's screen. The 286 responder then had to choose one of the two stimuli according to what the neighbor had chosen 287 (Fig.2E). If both proposer and responder selected the same target the trial was considered a success,

otherwise, a miss. The success of a test was coded as a binary variable: correct response = 1 and incorrect response = 0. A success triggered the delivery of a reward to the two individuals of the pair, a miss resulted in a 3 secs timeout for both. We presented experiments with seven or five different stimuli, therefore 21 or 10 possible different combinations. To balance all possible combinations the trials were organized in randomly ordered block of 42 (for seven stimuli) or 40 (for five) trials.

294 Filler task

If a monkey presented itself at an S-ALDM and no other monkey participated in the task in the neighboring station (at the time of synchronization of the blue screens, see Fig. 2A), a version of the Wisconsin Card Sorting Task (WCST, Berg, 1948) was presented. The WSCT is commonly used in executive control studies to assess cognitive flexibility and has been adapted to the ALDM system by Bonté, Flemming, and Fagot (2011). In the present study, the WCST was used as a filler task therefore these data will not be analyzed in the context of this paper.

301 Statistical analysis

302 We used two measures to understand the emergence of conventions in our experiments. 303 The first variable of interest is the score of individuals (i.e. whether they succeeded at the task and 304 were rewarded) because it indicates if the monkeys are solving the coordination task. To analyse 305 the score, we consider independently the score when an individual is in the role of the proposer, 306 from the score when in the role of the responder. This is because the roles are very different, the 307 proposer cannot respond based on visual information, the receiver can. Note that individuals can in 308 principle have very different scores as proposer and responder, if one individual uses visual 309 information and not another for instance. To understand the evolution of score over time we 310 analyse the average score on a block of trials (40 or 42 trials, see above) performed by an individual. 311 To understand how the conventions emerge, we also used the Elo-score rating technique 312 (Elo, 1978). The Elo-score is used to rank chess players and to analyse dominance hierarchies in

animals for instance (Strauss & Holekamp, 2019). The advantage is that it is a dynamic index, when

a contest takes place, the Elo-score of the winner increases, while that of the looser decreases. In our case, we used the Elo-score package (Sánchez-Tójar, Schroeder, & Farine, 2018; see ESM and the analysis code for details) to represent the dominance between stimuli: every time a monkey chooses one stimulus over another, that stimulus 'wins', the other loses (the k parameter was set to 1 and the initial elo-score was set at 0). The Elo-score technique therefore allows us to see if there is a hierarchy appearing during the experiment and to track changes in the hierarchy if they appear, which is important to assess stability.

321 We performed several experiments in which we manipulated the partition between 322 individuals (either transparent or opaque) and the stimuli (all stimuli are freely available with the 323 data and code to analyse the results). Since these experiments are based on the voluntary partition 324 of monkeys, we could not control the number of trials done per individual or by pairs of individuals, 325 so the number of blocks performed by individuals vary and we analyse only complete blocks. The 326 evolution of Score with the number of blocks was evaluated using a generalized mixed linear model 327 (GLMM, Baayen, Davidson, & Bates, 2008) with binomial error distribution and a logit link function. 328 Models were analysed using the stan glmer function of R's rstanarm package (Goodrich, Gabry, Ali, 329 & Brilleman, 2020) with default weakly informative priors as recommended. Convergence was 330 checked using the shinystan package (Muth, Oravecz, & Gabry, 2018) and we found no convergence 331 issues. To consider inter-individual variability and repeated measurement, we used mixed models 332 including a random intercept and slope (represented by the number of blocks of 42 trials 333 performed) for each individual. Thus, we limit our analysis to complete blocks of trials (all the model 334 details are presented in the ESM). Additional analyses use nonparametric tests due to the non-335 independent nature of the data.

336

All analyses were done with the statistical software R (R Core Team, 2021).

5. Study 1: stability of convention

338 In this first study, the baboons had to choose between pairs of coloured stimuli from seven 339 possible colours (see video 1 in ESM). During the first period of the experiment, monkeys could see 340 each other's responses through the transparent partition between them. We start by focusing on 341 the proposer whose role is to choose which of the two colours is to be selected. We found that the 342 proposer's score was initially high (mean score for the first block = 0.75, SEM = 0.02, min = 0.55, 343 max = 0.95) and steadily increased at the beginning of the experiment with the number of blocks (β 344 = 0.03, 95%CI = [0.02,0.04]; Fig.3A). This increase in score was linked to the establishment of a 345 hierarchy between colours (Fig.3B), some colours (such as yellow and orange) became progressively 346 more chosen by the proposer than others (such as light and dark blue). We found that at the end of 347 the experiment (we used the last two blocks of trials for each proposer to get a reliable measure of 348 score for every pair of colours) the difference in dominance rank between colours was highly 349 correlated to the average score for that pair (Kendall, N = 21, τ = 0.71, z = 4.53, p < 0.001), for 350 instance the probability of success when orange and dark blue were presented together was on 351 average 99% (a difference in Elo-score of 316) while it was 60% for light vs. dark blue (an Elo-score 352 difference of 38).

353





355 Figure 3: Establishment of a colour convention. A: evolution of proposers' score with the 356 number of blocks per individual (grey) and on average (in black, error bars represent standard error 357 of the mean). B: Evolution of Elo-score between colours based on proposers' behaviour (the seven 358 colours presented are an approximate match to the colours seen by the baboons). C: change in 359 score for the responders with the change in partition, from transparent to opaque. D: Average 360 responder score for every combination of colour pair depending on the colour pair difference in Elo-361 score for three performance groups (Low, Medium and High performance). The fitted curve and 362 standard errors are based on a quasibinomial fit.

363

The maintenance of the colour hierarchy can be achieved through two non-exclusive processes, the responder could watch the choice of the proposer and respond accordingly, or the responder could have learned the hierarchy and respond without watching. To tease apart these two explanations, we introduced an opaque partition between the two participants. We found that the responder's score decreased for 13/15 individuals (Binomial test, p = 0.004), but that the

369 average score remained high (Fig. 4C). Most of the decisions taken by the responders are therefore 370 based on the knowledge of the hierarchy and do not require visual access. However, we noticed 371 that all the monkeys with very high score with the transparent partition (above 85%) suffered from 372 the introduction of the opaque partition, suggesting these individuals watched the proposer. To test 373 this hypothesis, we divided the group of 19 individuals into three equally sized groups with Low, 374 Medium and High performing individuals during the transparent condition. For each group we 375 pooled the data and calculated the average score for every pair of colours. Since these data are 376 non-independent (the same colour appears in multiple pairs), we used a non-parametric correlation 377 test to evaluate the relationship between the average score of every pair of colours with the 378 difference in Elo-score depending on the transparent and opaque conditions (Fig.4D). We found 379 that high performing monkeys used their knowledge of the hierarchy of colours when the Elo-score 380 difference between the stimuli was high and resorted to visual checking of their social partner when 381 the difference was small (Kendall, N = 5, τ = -0.42, z = -2.66, p = 0.008). In contrast, medium or low 382 performing individuals seem to have only learned the dominance hierarchy of colours, and 383 therefore have a lower score when the rank difference is small (Medium: Kendall, N = 5, τ = -0.13, z 384 = -0.85, p = 0.40; Low: Kendall, N = 5, τ = -0.06, z = -0.36, p = 0.72).

385 The results of study 1 therefore show that when baboons had to coordinate by choosing the 386 same colour among a pair, they quickly created a hierarchy between colours that improved their 387 performance (the score increased). All the monkeys that participated regularly learned at least part 388 of the hierarchy. In addition, some monkeys also watched their partner to respond appropriately 389 when the pairs were difficult to discriminate. The hierarchical organisation of colours, although 390 stable, does not seem to be arbitrary because the score at the beginning of the experiment was 391 already initially high, suggesting that baboons did not choose colours arbitrarily but based on pre-392 existing preferences. To test the generality of our results, the arbitrariness and stability of 393 conventions we conducted two additional sets of experiments.

394

6. Study 2: generalisation of the results

396 We replicated the results of study 1 with a new set of five black and white images to test the 397 generality of our conclusions (Fig.4, experiment 2.1; see video 2 in ESM). We found equivalent 398 results to our first study, the score of the proposers increased in a similar fashion (β = 0.02, 95%CI = 399 [0.01,0.04]) and was linked to the dominance hierarchy that emerged during the experiment 400 (Kendall, N = 10, τ = 0.91, z = 3.67, p < 0.001). Although the deleterious effect of the opaque 401 partition was less widespread than previously (the responder's score decreased for 11/16 402 individuals, Binomial test, p =0.11), we found a similar relationship between the monkeys' 403 performance in the transparent condition and the effect of the opaque partition. The performance 404 of high performing monkeys decreased more with the introduction of the opaque partition when 405 the pairs of stimuli were close in Elo-scores than when they were far apart (High performance: 406 Kendall, N = 5, τ = -0.56, z = -2.24, p = 0.025; Medium: Kendall, N = 5, τ = -0.28, z = -1.16, p = 0.24; 407 Low: Kendall, N = 6, τ = 0.33, z = 1.34, p = 0.18).

408







416

Monkeys could therefore establish a convention with different types of stimuli such as coloured squares, or black and white images. Next, in experiment 2.2, we questioned the importance of the transparent condition to establish the convention. Originally, we thought the monkeys would use the opportunity to see the action of their neighbour to choose the correct response, but the results showed that for a majority the introduction of the opaque partition had limited effect or no effect at all. We decided to question the necessity of having visual access to establish a convention by introducing five new black and white stimuli and performing the same experiment but with an opaque partition from the start. Visual inspection of figure 4 did not reveal qualitative differences
between the experiment with a transparent or an opaque partition (Fig.4 and ESM). We noticed
that the average score was lower and the evolution of the dominance less noisy in the opaque
condition, this is consistent with the fact that monkeys sometime rely on visual access, but it could
also be due to differences in the stimuli we presented since we used different stimuli for the two
experiments. Nonetheless, it appeared that individuals could rapidly establish a convention with or
without visual access to their partner's responses.

431

432 **7. Study 3: Arbitrariness of convention**

433 A distinctive feature of conventions is their arbitrariness. In experiment 3.1, we tested the 434 possibility of establishing an arbitrary convention by individually training the same monkeys as in 435 experiment one and two to choose the five black and white stimuli of experiment 2.1 in the 436 opposite order of dominance (we used an opaque partition so that monkeys had no visual access to 437 their neighbours' behaviour; see video 3 and additional details can be found in the ESM). For 438 instance, in experiment 2.1 we found that the "dog paw" was more dominant than the "leaf", we 439 therefore trained the monkeys on the opposite choice by rewarding the "leaf" and not the "paw". 440 This is a strong test of arbitrariness because presumably monkeys ordered the pairs according to 441 their shared preference, therefore, if an opposite convention is established and remain stable, this 442 shows that opposite conventions can be maintained (a less stringent test would simply try to 443 inverse two stimuli for instance). 444 All individuals were trained in the same reversed order. During training, we found, as expected, that the score was initially very low (for the first two blocks of trial, mean = 0.30, SE = 445 446 0.02, min = 0.19, max = 0.43). However, the performance rapidly improved as the monkeys learned

to choose the correct stimuli for each pair (for the last two blocks of each individual, mean score =

448 0.72, SE = 0.05, min = 0.28, max = 0.99; see also Fig.S19). After the training phase we performed the

same coordination experiment with the transparent partition as previously and found that the score remained high and showed no sign of decreasing ($\beta = 0.03$, 95%CI = [-0.03,0.08]). The hierarchy was also stable (Fig.S20) and in the opposite order of the one previously established (Table 1). This shows that the established conventions are to some extent arbitrary.

453

	First exposition	After reversed training
¥	140	-258
	63	-93
n far	8	13
	-77	112
Ø	-133	225

454

Table 1: Inversion of hierarchy between stimuli after training. First exposition refers to the spontaneous Elo-scores obtained during experiment 2.1 and "after reversed training" to the Eloscores obtained after each individual had been trained on the reversed order.

458 Finally, in experiment 3.2, we wanted to test the impact that each baboon could have on the 459 establishment of convention based on their social relationships with others. We reasoned that well-460 connected individuals could either be influential and change others or, on the contrary, be more 461 flexible and adapt to others. We selected 11 individuals that had taken part in previous experiments 462 and used data from experiment 2.1 (generalisation of the results, transparent condition, N= 38409) 463 to calculate the number of times two individuals performed trials together. Using the modularity 464 classes algorithm available in the software Gephi, we identified three clusters (Fig. 5). From each 465 cluster, we selected the individual with the highest degree (Mako, Violette, Ewine) and individually

- 466 trained them on five new stimuli organised in an arbitrary hierarchy. The remaining eight individuals
- 467 were trained using the same procedure but in the reverse order (Fig.6 and ESM for details on
- 468 training). Once trained, we tested the establishment of a new convention with a transparent
- 469 partition as before.



470

Figure 5: Social network of the 11 individuals tested in experiment 3.2. Individuals with highest degree in their cluster (Mako, Violette, Ewine) were trained on the opposite convention compared to other individuals. The social network was obtained using the Force Atlas algorithm of Gephi (Bastian, Heymann, & Jacomy, 2009). Colours indicate different clusters, determined using Modularity classes. The thickness of the links represents the weight of the links between two individuals (equivalent to their number of trials performed together). The size of the nodes represents the weighted degree of the node (i.e. the sum of the link weights).

478

Individuals rapidly learned to choose the correct stimuli during training (for the last two
blocks of each individual, mean score = 0.86, SE = 0.03, min = 0.66, max = 0.96; see also Fig.S24 for
the evolution through time). During the coordination experiment, we found that the group
converged on the convention of the low-degree individuals (Fig.6A) because high-degree individuals
changed their behaviour more than did low-degree individuals. For instance, as can be seen on

- 484 Fig.6B, the most dominant and least dominant stimuli were reversed for Violette (High degree) and
- did not change for Fana (Low degree; see Fig.S27 for the plots of each individual).



486

487 Figure 6: Evolution of Elo-scores trajectories for the entire group (A) and for the first 1000
488 trials of two individuals (B), Violette (high degree group) and Fana (low degree group).

489

490 **8. Discussion and conclusions**

In this series of experiments, we showed that when monkeys were faced with a coordination problem, they could spontaneously develop efficient arbitrary stable conventions. Conventions were efficient because the score improved with the emergence of conventions and because we found a direct relationship between the score and the hierarchy among the stimuli (Study 1 and 2.1). We showed that conventions were arbitrary and stable because after training to choose stimuli according to a new hierarchy, monkeys could establish and maintain a new convention based on this new hierarchy (Study 3.1). These three properties are defining features of human conventions and our results therefore complement a small number of field studies (discussed previously)
showing that non-human primates and maybe other animals can establish conventions akin to
humans.

501 Our results also speak more generally to our understanding of conventions and how they are 502 established. Moore (2013) for instance raises two important questions in the context of linguistic 503 conventions: (1) which manner of learning would allow the acquisition of convention? And what 504 must we know in order to participate in convention? Further specifying that:

505"Where a coordinating tool acquires its functional properties through convention, those who506learn it must be particularly attentive to the nature of the action that others perform—that507is, to the means R that an observed agent employs in pursuit of her goal E. That's because in508conventional coordination, unlike in the case of nut-cracking described at the outset, one509couldn't (unless one was very lucky) figure out the appropriate means to a goal

510 independently of copying the actions performed by others." (Moore, 2013; R stands for

511 regularity in behaviour and E for end goal)

522

2004).

512 In our study, monkeys established a new convention without any visual access to each other's

513 behaviour (experiment 2.2), therefore showing that imitation, or more generally observational 514 learning, is not necessary to establish conventions: in the opaque condition baboons were able to 515 establish a convention through positive reinforcement and knew nothing about the behaviour of 516 their conspecifics. However, we also found that the most successful monkeys were able to copy the 517 proposer when the task became difficult due to a small Elo-score difference between pairs of stimuli 518 (Fig.3D: when the difference in elo-score is small the effect of the opaque partition on the score of 519 high performing monkeys is stronger). This is in agreement with the growing literature on social 520 learning showing that individuals can learn from the observation of conspecifics (Whiten, 2021) and 521 with another study showing this effect with touch-screen (Subiaul, Cantlon, Holloway, & Terrace,

523 According to Lewis (1969), there are three main sources explaining the emergence of 524 conventions: agreement, saliency and precedence. Agreement occurs when individuals can 525 communicate and form mutual expectations (i.e. agree on a course of action), for instance when 526 making an appointment. Conventions can also emerge when a solution is so salient that most 527 individuals tend towards that solution (i.e a preference system). Finally, conventions can emerge 528 through precedence: if a solution becomes more salient and remarkable because it was previously 529 successfully chosen (i.e. a regularity of behaviour). In the context of our study, agreement is unlikely to explain the emergence of conventions because monkeys (1) cannot explicitly agree on which 530 531 colour to choose and moreover (2) continued to follow convention when the opaque partition 532 prevented observation of their partner. On the other hand, monkeys can create conventions 533 through a combination of saliency and precedence. Monkeys may all perceive certain stimuli as 534 more salient than others (saliency) and the result of each trial also influences the stimuli future 535 saliency (precedence).

536 In humans, studies have described tipping points and the role of a determined minority in 537 changing an established convention (Andreoni, Nikiforakis, & Siegenthaler, 2021; Centola, Becker, 538 Brackbill, & Baronchelli, 2018). We attempted to study the importance of well-connected 539 individuals (with a high degree within their cluster), reasoning that if well-connected individuals 540 were to change their behaviour this could change the whole group. Contrary to our expectations we 541 found that well-connected individuals were more likely to adapt to others rather than influence 542 them. Given that high-degree individuals were also a minority (three vs. eight), several factors could 543 explain this result. Individuals may have been conformist and followed the majority for instance 544 (e.g. Haun, Rekers, & Tomasello, 2012), or they may be well-connected because they are flexibly 545 adapting to others. Additional experiments will be able to provide a more detailed understanding of 546 the influence of network parameters and other biases by manipulating the proportion and identity 547 of individuals that are trained on alternative conventions.

548	Conventions are an important aspect of human culture because like norms they stabilise
549	cultural diversity. Surprisingly, although the evolutionary precursors of conventions and norms exist
550	in non-human primates (Kappeler et al., 2019), conventions do not seem to be widespread and their
551	contribution to non-human primate culture seems almost anecdotal. One might therefore be
552	tempted to think that conventions require elaborate social learning mechanisms. Using a freely
553	accessible touch-screen system in which pairs of individuals can perform experiments together, we
554	have shown, through five experiments, that non-human primates can develop conventions using
555	positive reinforcement learning and that although copying can be used by some individuals, it is not
556	necessary.
557	Acknowledgments
558	We thank Julie Gullstrand and Sebastien Barniaud as well as the staff at the Rousset-sur-Arc
559	Primate Center (CNRS-UPS846, France) for technical supports.
560	Ethics
561	This research was carried out in accordance with European Union and French ethical
562	standards and received approval from the French Ministère de l'Education Nationale et de la
563	Recherche (approval no. APAFIS-2717-2015111708173794-V3).
564	Data Accessibility
565	The data and analysis code used in this article are available here:
566	https://osf.io/b68x2/
567	Authors' Contributions
568	A.F., N.C., J.F. planned the experiment, D.P., N.C., J.F. planned and constructed the S-ALDMs,
569	A.F., N.C. performed the experiment, A.F., N.C. analysed the results, A.F., N.C., J.F. wrote the
570	manuscript.
571	Competing Interests
572	We have no competing interests.
573	Funding

- 574 This project received funding from the Institute for Language Communication and the Brain
- 575 (ILCB) through the Agence Nationale de la Recherche grant ANR-16-CONV-0002

576 **9. References**

- 577 Andreoni, J., Nikiforakis, N., & Siegenthaler, S. (2021). Predicting social tipping and norm change in
- 578 controlled experiments. *Proceedings of the National Academy of Sciences, 118*(16),
- 579 e2014893118. doi:10.1073/pnas.2014893118
- Arnold, K., & Zuberbühler, K. (2008). Meaningful call combinations in a non-human primate. *Current Biology, 18*(5), R202-R203. doi:<u>https://doi.org/10.1016/j.cub.2008.01.040</u>
- 582 Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random
- 583 effects for subjects and items. *Journal of Memory and Language, 59*(4), 390-412.
- 584 doi:<u>http://dx.doi.org/10.1016/j.jml.2007.12.005</u>
- 585 Bastian, M., Heymann, S., & Jacomy, M. (2009). Gephi: an open source software for exploring and
- 586 *manipulating networks*. Paper presented at the International AAAI Conference on Weblogs
- 587 and Social Media, San Jose, California, USA.
- 588 Berg, E. A. (1948). A Simple Objective Technique for Measuring Flexibility in Thinking. *The Journal of* 589 *General Psychology*, *39*(1), 15-22. doi:10.1080/00221309.1948.9918159
- Sos Ceneral Psychology, 55(1), 15 22: 40.110.1000/00221505.15 10.5510155
- 590 Bergman, T. J., Beehner, J. C., Cheney, D. L., & Seyfarth, R. M. (2003). Hierarchical Classification by
- 591Rank and Kinship in Baboons. Science, 302(5648), 1234-1236. doi:10.1126/science.1087513
- 592 Bicchieri, C. (2005). *The grammar of society: The nature and dynamics of social norms*: Cambridge
- 593 University Press.
- 594 Bickerton, D. (1981). *Roots of language*. Ann Arbor: Karoma Publishers.
- 595 Bonté, E., Flemming, T., & Fagot, J. (2011). Executive control of perceptual features and abstract
- relations by baboons (Papio papio). *Behavioural Brain Research, 222*(1), 176-182.
- 597 doi:<u>https://doi.org/10.1016/j.bbr.2011.03.034</u>

- 598 Centola, D., Becker, J., Brackbill, D., & Baronchelli, A. (2018). Experimental evidence for tipping
- points in social convention. *Science*, *360*(6393), 1116. doi:10.1126/science.aas8827
- Deming, W., Neumann, J., & Morgenstern, O. (1944). Theory of Games and Economic Behavior.
- 601 Journal of the American Statistical Association, 40, 263.
- 602 Elo, A. E. (1978). *The rating of chessplayers, past and present*: Arco Pub.
- 603 Fagot, J., Gullstrand, J., Kemp, C., Defilles, C., & Mekaouche, M. (2014). Effects of freely accessible
- 604 computerized test systems on the spontaneous behaviors and stress level of Guinea
- 605 baboons (*Papio papio*). *American Journal of Primatology, 76*(1), 56-64.
- 606 doi:10.1002/ajp.22193
- 607 Fagot, J., Marzouki, Y., Huguet, P., Gullstrand, J., & Claidière, N. (2015). Assessment of Social
- Cognition in Non-human Primates Using a Network of Computerized Automated Learning
 Device (ALDM) Test Systems. *JoVE*(99), e52798. doi:doi:10.3791/52798
- 610 Fay, N., Garrod, S., Roberts, L., & Swoboda, N. (2010). The Interactive Evolution of Human
- 611 Communication Systems. *Cognitive Science*, *34*(3), 351-386.
- 612 doi:<u>https://doi.org/10.1111/j.1551-6709.2009.01090.x</u>
- 613 Fedigan, L. (1993). Sex differences and intersexual relations in adult white-faced capuchins (Cebus
- 614 capucinus). International Journal of Primatology, 14(6), 853-877. doi:10.1007/BF02220256
- 615 Fudenberg, D., & Maskin, E. (2008). THE FOLK THEOREM IN REPEATED GAMES WITH DISCOUNTING
- 616 OR WITH INCOMPLETE INFORMATION. In *A Long-Run Collaboration on Long-Run Games* (pp.
 617 209-230): WORLD SCIENTIFIC.
- 618 Galantucci, B. (2005). An Experimental Study of the Emergence of Human Communication Systems.
- 619 *Cognitive Science, 29*(5), 737-767. doi:<u>https://doi.org/10.1207/s15516709cog0000_34</u>
- 620 Gilby, I. C., Brent, L. J. N., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey, A. E.
- 621 (2013). Fitness benefits of coalitionary aggression in male chimpanzees. *Behavioral Ecology*
- 622 and Sociobiology, 67(3), 373-381. doi:10.1007/s00265-012-1457-6

- 623 Goodrich, B., Gabry, J., Ali, I., & Brilleman, S. (2020). rstanarm: Bayesian applied regression
- 624 modeling via Stan (Version 2.21.1). Retrieved from <u>https://mc-stan.org/rstanarm</u>
- 625 Graham, K. E., Hobaiter, C., Ounsley, J., Furuichi, T., & Byrne, R. W. (2018). Bonobo and chimpanzee
- 626 gestures overlap extensively in meaning. *PLOS Biology*, *16*(2), e2004825.
- 627 doi:10.1371/journal.pbio.2004825
- Haun, D. B. M., Rekers, Y., & Tomasello, M. (2012). Majority-Biased Transmission in Chimpanzees
- and Human Children, but Not Orangutans. *Current biology*, 22(8), 727-731. Retrieved from
 http://linkinghub.elsevier.com/retrieve/pii/S0960982212002631
- 631 Hawkins, R. X. D., & Goldstone, R. L. (2016). The Formation of Social Conventions in Real-Time
- 632 Environments. *PLoS ONE, 11*(3), e0151670. doi:10.1371/journal.pone.0151670
- Hawkins, R. X. D., Goodman, N. D., & Goldstone, R. L. (2019). The Emergence of Social Norms and
 Conventions. *Trends in Cognitive Sciences*, 23(2), 158-169.
- 635 doi:<u>https://doi.org/10.1016/j.tics.2018.11.003</u>
- 636 Heyes, C. M. (1994). Social learning in animals: categories and mechanisms. Biological Reviews,
- 637 69(2), 207-231. doi:10.1111/j.1469-185X.1994.tb01506.x
- Hume, D. (1888). A treatise of human nature: Oxford Univ Press.
- 639 Kappeler, P. M., Fichtel, C., & van Schaik, C. P. (2019). There Ought to Be Roots: Evolutionary
- 640 Precursors of Social Norms and Conventions in Non-Human Primates. In *The Normative*
- 641 *Animal?* New York: Oxford University Press.
- 642 Kegl, J. (1994). The Nicaraguan sign language project: An overview. *Signpost, 7,* 24-31.
- 643 Kemp, C., Rey, A., Legou, T., Boë, L.-J., Berthommier, F., Becker, Y., & Fagot, J. (2017). Vocal
- 644 Repertoire of Captive Guinea Baboons (Papio papio). *Origins of Human Language:*
- 645 *Continuities and Discontinuities with Nonhuman Primates*, 15.
- Laidre, M. E. (2011). Meaningful Gesture in Monkeys? Investigating whether Mandrills Create Social
- 647 Culture. *PLoS ONE, 6*(2), e14610. Retrieved from
- 648 http://dx.doi.org/10.1371%2Fjournal.pone.0014610

- 649 Lewis, D. (1969). *Convention: A philosophical study*: Harvard Univ Pr.
- 650 McGrew, W. C., Marchant, L. F., Scott, S. E., & Tutin, C. E. G. (2001). Intergroup Differences in a
- 651 Social Custom of Wild Chimpanzees: The Grooming Hand Clasp of the Mahale Mountains.
- 652 *Current Anthropology, 42*(1), 148-153. Retrieved from
- 653 http://www.journals.uchicago.edu/doi/abs/10.1086/318441
- Moore, R. (2013). Imitation and conventional communication. *Biology & Philosophy, 28*(3), 481-500.
- 655 doi:10.1007/s10539-012-9349-8
- 656 Muth, C., Oravecz, Z., & Gabry, J. (2018). User-friendly Bayesian regression modeling: A tutorial with 657 rstanarm and shinystan. *Quantitative Methods for Psychology*, *14*(2), 99-119.
- 658 Nakamura, M., McGrew, W. C., Marchant, L. F., & Nishida, T. (2000). Social scratch: Another custom
- 659 in wild chimpanzees? *Primates, 41*(3), 237-248. doi:10.1007/BF02557594
- 660 Nash, J. (1951). Non-cooperative games. *Annals of mathematics*, 286-295.
- Nichols, S. (2002). On The Genealogy Of Norms: A Case For The Role Of Emotion In Cultural

662 Evolution. *Philosophy of Science, 69*(2), 234-255. doi:doi:10.1086/341051

663 Ouattara, K., Lemasson, A., & Zuberbühler, K. (2009). Campbell's Monkeys Use Affixation to Alter

664 Call Meaning. *PLoS ONE, 4*(11), e7808. doi:10.1371/journal.pone.0007808

- 665 Perry, S., Baker, M., Fedigan, L., Grosa Louis, J., Jack, K., MacKinnon, Katherine C., . . . Rose, L.
- 666 (2003). Social conventions in wild white faced Capuchin monkeys: Evidence for traditions in
- a neotropical primate. *Current Anthropology*, 44(2), 241-268. doi:10.1086/345825
- 668 R Core Team. (2021). R: A language and environment for statistical computing. Vienna, Austria: R
- 669 Foundation for Statistical Computing. Retrieved from <u>http://www.R-project.org/</u>.
- 670 Sánchez-Tójar, A., Schroeder, J., & Farine, D. R. (2018). A practical guide for inferring reliable
- 671 dominance hierarchies and estimating their uncertainty. *Journal of Animal Ecology*, 87(3),

672 594-608. doi:<u>https://doi.org/10.1111/1365-2656.12776</u>

- 673 Sapolsky, Robert M. (2006). Social Cultures among Nonhuman Primates. Current Anthropology,
- 674 47(4), 641-656. doi:10.1086/504162

- 675 Sapolsky, R. M., & Share, L. J. (2004). A pacific culture among wild baboons: Its emergence and
- 676 transmission. *PLoS Biology*, 2(4), 534-541. Retrieved from <Go to ISI>://000221194700020
- 677 Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social Bonds Enhance Reproductive
- 578 Success in Male Macaques. *Current Biology, 20*(24), 2207-2210.
- 679 doi:<u>https://doi.org/10.1016/j.cub.2010.10.058</u>
- 680 Silk, J. B. (2007). Social Components of Fitness in Primate Groups. *Science*, *317*(5843), 1347-1351.
- 681 doi:10.1126/science.1140734
- 682 Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., . . . Cheney, D. L.
- 683 (2009). The benefits of social capital: close social bonds among female baboons enhance
- 684 offspring survival. Proceedings of the Royal Society B: Biological Sciences, 276(1670), 3099-
- 685 3104. doi:10.1098/rspb.2009.0681
- 686 Slater, P. J. B. (1986). The cultural transmission of bird song. *Trends in Ecology & Evolution, 1*(4), 94-
- 687 97. doi:<u>https://doi.org/10.1016/0169-5347(86)90032-7</u>
- 688 Snowdon, C. T., & Elowson, A. M. (1999). Pygmy Marmosets Modify Call Structure When Paired.
- 689 *Ethology*, *105*(10), 893-908. doi:<u>https://doi.org/10.1046/j.1439-0310.1999.00483.x</u>
- 690 Stephens, D. W., & Heinen, V. K. (2018). Modeling nonhuman conventions: the behavioral ecology
- 691 of arbitrary action. *Behavioral Ecology, 29*(3), 598-608. doi:10.1093/beheco/ary011
- 692 Strauss, E. D., & Holekamp, K. E. (2019). Inferring longitudinal hierarchies: Framework and methods
- for studying the dynamics of dominance. *Journal of Animal Ecology*, 88(4), 521-536.
- 694 doi:<u>https://doi.org/10.1111/1365-2656.12951</u>
- 695 Subiaul, F., Cantlon, J. F., Holloway, R. L., & Terrace, H. S. (2004). Cognitive Imitation in Rhesus
- 696 Macaques. *Science*, *305*(5682), 407-410. doi:10.1126/science.1099136
- 697 Tomasello, M. (2008). Origins of human communication: MIT press.
- 698 Tomasello, M., & Call, J. (1997). *Primate cognition*: Oxford University Press.

- 699 Tomasello, M., & Zuberbühler, K. (2002). Primate vocal and gestural communication. In M. Bekoff,
- C. Allen, & G. Burghardt (Eds.), *The cognitive animal: empirical and theoretical perspectives on animal cognition*. Cambridge: MIT Press.
- van de Waal, E. (2018). On the neglected behavioural variation among neighbouring primate
- 703 groups. *Ethology*, *124*(12), 845-854. doi:<u>https://doi.org/10.1111/eth.12815</u>
- van de Waal, E., Borgeaud, C., & Whiten, A. (2013). Potent social learning and conformity shape a
 wild primate's foraging decisions. *Science*, *340*(6131), 483-485.
- van Leeuwen, E. J. C. (2021). Temporal stability of chimpanzee social culture. *Biology Letters, 17*(5),
 20210031. doi:10.1098/rsbl.2021.0031
- van Leeuwen, E. J. C., Cronin, K. A., Haun, D. B. M., Mundry, R., & Bodamer, M. D. (2012).
- 709 Neighbouring chimpanzee communities show different preferences in social grooming
- behaviour. *Proceedings of the Royal Society B: Biological Sciences, 279*(1746), 4362-4367.
- 711 doi:doi:10.1098/rspb.2012.1543
- 712 Watson, S. K., Townsend, S. W., Schel, A. M., Wilke, C., Wallace, E. K., Cheng, L., . . . Slocombe, K. E.
- 713 (2015). Vocal Learning in the Functionally Referential Food Grunts of Chimpanzees. *Current*714 *Biology*.
- 715 Weilgart, L., & Whitehead, H. (1997). Group-specific dialects and geographical variation in coda
- 716 repertoire in South Pacific sperm whales. Behavioral Ecology and Sociobiology, 40(5), 277-
- 717 285. doi:10.1007/s002650050343
- 718 Whiten, A. (2021). The burgeoning reach of animal culture. *Science*, *372*(6537), eabe6514.
- 719 doi:10.1126/science.abe6514
- 720 Whiten, A., Horner, V., & de Waal, F. B. (2005). Conformity to cultural norms of tool use in
- 721 chimpanzees. Nature, 437(7059), 737-740. Retrieved from
- 722 <u>http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation</u>
- 723 <u>&list_uids=16113685</u>

724