



HAL
open science

Nanoscale 3D distribution of low melt and fluid fractions in mantle rocks

Emmanuel Gardes, Luiz Morales, Wilhelm Heinrich, David Sifre, Leila Hashim, Fabrice Gaillard, Marquardt Katharina

► **To cite this version:**

Emmanuel Gardes, Luiz Morales, Wilhelm Heinrich, David Sifre, Leila Hashim, et al.. Nanoscale 3D distribution of low melt and fluid fractions in mantle rocks. EGU General Assembly 2016, 2016, Vienne, Austria. hal-03551122

HAL Id: hal-03551122

<https://hal.science/hal-03551122v1>

Submitted on 1 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Nanoscale 3D distribution of low melt and fluid fractions in mantle rocks

Emmanuel Gardes (1), Luiz Morales (2), Wilhelm Heinrich (2), David Sifre (3), Leila Hashim (3), Fabrice Gaillard (3), and Marquardt Katharina (4)

(1) CIMAP, Caen, France (gardes@ganil.fr), (2) GFZ, Potsdam, Germany, (3) ISTO, Orléans, France, (4) BGI, Bayreuth, Germany

The presence of melts or fluids in the intergranular medium of rocks strongly influences their bulk physico-chemical properties (e.g. mass transport and chemical reactivity, electrical conductivity, seismic wave velocity, etc). Actually, the effects can be so large that only small melt or fluid fractions must sometimes be involved for explaining mantle geophysical discontinuities and anomalies. The investigation of the distribution of such small fractions in the intergranular medium of mantle rocks is therefore crucial for relating them to bulk and large scale properties. However, it involves submicrometric structures which are hardly characterizable using conventional techniques. Here we present how the FIB-SEM-STEM microscope can be used to produce 3D imaging at unequalled resolution. We show that low melt and fluid fractions can form films as thin as 20 nm at olivine grain boundaries, and that they can modify the physico-chemical properties of mantle rocks by orders of magnitude. The fine relationships between films at grain boundaries, tubules at triple junctions and pockets at grain corners can be explored, and appear to be complex and to differ from usual visions.