



HAL
open science

**Establishing the geometry and nature of sediments
trapped in either natural or artificial dam lakes in
contrasted drainage basins from Western Europe
(French Massif Central and Pyrenees)**

Emmanuel Chapron, Léo Chassiot, Claude Zouzou, Anaëlle Simonneau,
Didier Galop, Christian Di Giovanni

► **To cite this version:**

Emmanuel Chapron, Léo Chassiot, Claude Zouzou, Anaëlle Simonneau, Didier Galop, et al.. Establishing the geometry and nature of sediments trapped in either natural or artificial dam lakes in contrasted drainage basins from Western Europe (French Massif Central and Pyrenees). EGU General Assembly 2016, 2016, Vienne, Austria. hal-03551938

HAL Id: hal-03551938

<https://hal.science/hal-03551938>

Submitted on 2 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Establishing the geometry and nature of sediments trapped in either natural or artificial dam lakes in contrasted drainage basins from Western Europe (French Massif Central and Pyrenees)

Emmanuel Chapron (1), Léo Chassiot (2), Claude Zouzou (1), Anaëlle Simonneau (2), Didier Galop (1), and Christian Di Giovanni (2)

(1) Université Fédérale de Toulouse, GEODE UMR 5602, Toulouse, France (emmanuel.chapron@univ-tlse2.fr; hermann.zouzou@aiesec.net; didier.galop@univ-tlse2.fr), (2) Université d'Orléans, ISTO UMR 7325, Orleans, France (leo.chassiot@cirs-orleans.fr; anaëlle.simonneau@univ-orleans.fr; christian.digiovanni@univ-orleans.fr)

Lacustrine sedimentary archives from artificial dam lakes are poorly documented both in terms of basin fill geometries and dominating sedimentary processes. In order to better understand their sensitivities to regional environmental changes, we performed a similar multidisciplinary study of French natural and artificial dam lakes in contrasted drainage basins from the volcanic Massif Central (lakes Aydat and Crégut) and two granitic sectors of the northern Pyrenees (lakes Fourcat and Orédon). Our approach combined high-resolution sub bottom profiling (14 kHz and 4 kHz chirp) and a detailed study of sediment cores based on qualitative and quantitative analysis (radiographies, sediment physical and chemical properties) together with radionuclide and radiocarbon dates. In all cases either changes in land uses within the drainage basin or the flooding of natural lakes by dams and the production of hydroelectricity induced changes in sedimentation rates and modes. Human activities affecting either the catchment or the lake itself favored enhanced clastic sediment supply in the lake basins and/or higher and fluctuating lake levels. Subaquatic slopes failures are also identified in Lake Aydat formed by a lava flow 8.5 kYrs ago and in glacial lakes Crégut (Massif Central) and Orédon (Pyrenees) now used to produce hydroelectricity and suggest that lake level changes and ground accelerations during earthquakes can remobilize distinct sectors of the basin fills and not only deltaic environments.