



**HAL**  
open science

## Role of litter decomposition sensitivity to water content in non-additive litter mixture effect: theoretical demonstration and validation with a peatland litter experiment

Sébastien Gogo, Fabien Leroy, Renata Zoccatelli, Léonard Bernard-Jannin,  
Fatima Laggoun-Défarge

### ► To cite this version:

Sébastien Gogo, Fabien Leroy, Renata Zoccatelli, Léonard Bernard-Jannin, Fatima Laggoun-Défarge. Role of litter decomposition sensitivity to water content in non-additive litter mixture effect: theoretical demonstration and validation with a peatland litter experiment. 19th EGU General Assembly, 2017, Vienne, Austria. pp.12297. hal-03552755

**HAL Id: hal-03552755**

**<https://hal.science/hal-03552755v1>**

Submitted on 3 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## **Role of litter decomposition sensitivity to water content in non-additive litter mixture effect: theoretical demonstration and validation with a peatland litter experiment**

Sébastien Gogo (1,2,3), Fabien Leroy (1,2,3), Renata Zoccatelli (1,4), Léonard Bernard-Jannin (1,2,3), Fatima Laggoun-Défarge (1,2,3)

(1) Université d'Orléans, Institut des Sciences de la Terre d'Orléans, France (sebastien.gogo@univ-orleans.fr), (2) CNRS, ISTO, UMR 7327, 45071 Orléans, France, (3) BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France, (4) CETRAHE, Cellule d'Expertise et de transfert en TRAçages appliqués à l'Hydrogéologie et à l'Environnement

In this work, we showed theoretically that differences in litter water content, evaporation rate and reaction rate sensitivity to water content can give account of non-additive litter mixture effect. More specifically two litters with the same dependence to litter water content and contrasted water content, and 2 litters with contrasted decomposition sensitivity to litter water content can exert synergistic mixture effect on decomposition when the 2 litters interact. In these situations, water can flow from the wettest to the driest litter, changing the whole reaction rate without changing the whole litter water content. The reaction rate increase of the litter receiving the water was relatively more important than the reaction rate decrease of the litter supplying the water. These theoretical considerations were validated with experimental data. *Sphagnum rubellum* and *Molinia caerulea* decompose faster in measured mixture than expected from the rates obtained in monoculture incubation. *Sphagnum rubellum* litter can contain more water, which evaporates at a slower rate than *Molinia caerulea*. It is thus proposed that water flowed from *Sphagnum rubellum* litter to the *Molinia caerulea* litter, with a substantial increase of the decomposition of the latter. The physical and biochemical litter characteristics towards water explains a fraction of the synergistic effect of mixing the 2 litters, which suggests that other factors intervene in this effect, such as the carbon substrate.