



# Evidence of phase nucleation during olivine diffusion creep: a new perspective for mantle strain localisation

Jacques Précigout, Holger Stünitz

## ► To cite this version:

Jacques Précigout, Holger Stünitz. Evidence of phase nucleation during olivine diffusion creep: a new perspective for mantle strain localisation. 19th EGU General Assembly, 2017, Vienne, Austria. pp.4520. hal-03552760

**HAL Id: hal-03552760**

**<https://hal.science/hal-03552760>**

Submitted on 3 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## **Evidence of phase nucleation during olivine diffusion creep: a new perspective for mantle strain localisation**

Jacques Précigout (1) and Holger Stünitz (1,2)

(1) Institut des Sciences de la Terre d'Orléans (ISTO), Université d'Orléans UMR/CNRS 7327, Orléans, France (jacques.precigout@univ-orleans.fr), (2) Department of Geology, University of Tromsø, Tromsø, Norway

Mantle strain localisation is of great importance for lithosphere dynamics, but the cause for this phenomenon remains very elusive, particularly in conditions of the strong and ductile uppermost mantle. In these latter, grain size reduction leading to diffusion creep in olivine is believed to be one of the best candidates to account for strain localisation. However, the mechanisms of grain size reduction in this regime are still poorly understood. Here we show the results of Griggs-type experiments that document grain size reduction and material weakening during wet olivine diffusion creep at 900 °C and 1.2 GPa. While occurring for both, mono-phase and two-phase aggregates, grain size reduction is coeval with strain localisation and local phase mixing in olivine–pyroxene aggregates. Based on evidence of fluid inclusions and cracks filled with a fine-grained phase mixture, we conclude that grain size reduces as a result of fluid-assisted nucleation. Cavitation induced by grain boundary sliding (creep cavitation) can be inferred, and may play a critical role for olivine grain size reduction. Amongst their implications for rock rheology in general, our findings highlight a key process for strain localisation in the ductile uppermost mantle.

This study has been published under the reference: “Précigout, J., and Stünitz, H. (2016) Evidence of phase nucleation during olivine diffusion creep: a new perspective for mantle strain localisation. *Earth and Planetary Science Letters* 455: 94–105, doi:10.1016/j.epsl.2016.09.029”.