



HAL
open science

Pegmatite emplacement: insights from two-phase flow numerical modelling.

Alexis Plunder, Laetitia Le Pourihet, Ludovic Räss, Charles Gumiaux

► **To cite this version:**

Alexis Plunder, Laetitia Le Pourihet, Ludovic Räss, Charles Gumiaux. Pegmatite emplacement: insights from two-phase flow numerical modelling.. 20th EGU General Assembly, 2018, Vienne, Austria. pp.13245. hal-03552835

HAL Id: hal-03552835

<https://hal.science/hal-03552835>

Submitted on 2 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Pegmatite emplacement: insights from two-phase flow numerical modelling.

Alexis Plunder (1), Laetitia Le Pourihet (1), Ludovic Räss (2), and Charles Gumiaux (3)

(1) Sorbonne Université, Institut des Sciences de la Terre de Paris, Paris, France (alex.plunder@gmail.com), (2) Faculté des géosciences et de l'environnement, Institut des Sciences de la Terre, University of Lausanne, Lausanne, Switzerland, (3) Institut des Sciences de la Terre d'Orléans, Université d'Orléans, Orléans, France

Amongst the fluids present in lithospheric processes, silicate rich melts with their range of composition are one of the prime sources of ore on Earth. Understanding how such fluids propagate and interact with the porosity of the lithosphere is therefore of prime interest both for the long-term lithospheric deformation in relation with the weakening properties of fluids or to predict ore concentration linked or not to the chemical evolution of the melts.

Pegmatites are igneous rocks, generally with a granitic composition and characterised by a crystal growth dominated texture. They are often enriched in rare elements such as Lithium, Tantalum or Beryllium (amongst others) offering valuable ore deposit. In nature they are observed as pegmatite fields showing an internal organisation, with respect to the host rock (either magmatic or metamorphic) and the tectonic structures. Here, we focus on the geometry of pegmatite fields using a numerical modelling approach. The numerical model allows us to resolve the formation and propagation of pipe structures in a viscous porous media also called porosity waves. We use the finite difference approach to solve the two-phase flow problem that couples a Stokes solver to predict the deformation of the porous matrix to a non-linear Darcy flow representing the pore fluid displacement (Räss et al., in prep).

We present here the first results of our study and discuss the effects heterogeneities in the host porous media and compare the statistical distribution of the modelled pipes with natural examples for the Montagne Noire and the Massif Central.

Räss, L., Duretz, T., and Y. Podladchikov. Hydro-mechanical coupled flow in deforming porous media.