



**HAL**  
open science

## Dissolution of a single mineral grain: comparison of microfluidic experiments with pore-scale simulations

Piotr Szymczak, Filip Dutka, Vitaliy Starchenko, Florian Osselin, Silvana Magni, Anthony J. C. Ladd

► **To cite this version:**

Piotr Szymczak, Filip Dutka, Vitaliy Starchenko, Florian Osselin, Silvana Magni, et al.. Dissolution of a single mineral grain: comparison of microfluidic experiments with pore-scale simulations. EGU General Assembly 2020, May 2020, Online, France. 10.5194/egusphere-egu2020-12253 . hal-03554428

**HAL Id: hal-03554428**

**<https://hal.science/hal-03554428v1>**

Submitted on 4 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU2020-12253

<https://doi.org/10.5194/egusphere-egu2020-12253>

EGU General Assembly 2020

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



## Dissolution of a single mineral grain: comparison of microfluidic experiments with pore-scale simulations

**Piotr Szymczak**<sup>1</sup>, Filip Dutka<sup>1</sup>, Vitaliy Starchenko<sup>2</sup>, Florian Osselin<sup>3</sup>, Silvana Magni<sup>1,4</sup>, and Anthony J.C. Ladd<sup>5</sup>

<sup>1</sup>Faculty of Physics, University of Warsaw, Poland (piotr.szymczak@fuw.edu.pl)

<sup>2</sup>Chemical Sciences Division, Oak Ridge National Laboratory, USA

<sup>3</sup>Institut des Sciences de la Terre d'Orléans, France

<sup>4</sup>Institute of Geophysics, Polish Academy of Sciences, Warsaw, Poland

<sup>5</sup>Department of Chemical Engineering, University of Florida, Gainesville, USA

We investigate the dissolution of a single grain of soluble mineral by microfluidic experiments and numerical simulations. The experiments use gypsum cylinders (10 mm radius, 0.5 mm thick) cast from rehydrated CaSO<sub>4</sub> hemihydrate. The numerical simulations used a finite-volume discretization of the reactive-transport equations with a mesh that conforms to the evolving shape of the mineral. Using the coefficients for dilute aqueous ions, we overpredict the dissolution rate by about 25%. However, including the Debye-Huckel correction for the ion activity gives a substantial reduction in diffusion across the boundary layer at the dissolving solid surface and brings the simulation time scale into quantitative agreement with experiment.

The asymmetry introduced by the flow causes the initially cylindrical sample to take on a shape resembling one half of a figure eight, with the tip pointing in the downstream direction. The simulations give a near perfect match to the experimental size and shape. We quantify the evolution of the volume of the grain and its surface area, as well as its overall shape as the function of the Peclet number. Next we discuss the differences between the geometric surface area and the reactive surface area of a dissolving grain and explore a potential use of these results to upscale the reactive transport problem and obtain the effective reaction rates in a multi-grain system.