

Sb, As and W contents of magmas: insights from geochemical databases

Valentin Mollé, Giada Iacono-Marziano, Eric Gloaguen, Johann Tuduri, Anthony Pochon, Héctor Campos

► To cite this version:

Valentin Mollé, Giada Iacono-Marziano, Eric Gloaguen, Johann Tuduri, Anthony Pochon, et al.. Sb, As and W contents of magmas: insights from geochemical databases. vEGU21: Gather Online, Apr 2021, Online, France. 10.5194/egusphere-egu21-12365. hal-03554519

HAL Id: hal-03554519 https://hal.science/hal-03554519v1

Submitted on 3 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



EGU21-12365, updated on 03 Feb 2022 https://doi.org/10.5194/egusphere-egu21-12365 EGU General Assembly 2021 © Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Sb, As and W contents of magmas: insights from geochemical databases

Valentin Mollé¹, Giada Iacono-Marziano¹, Eric Gloaguen^{1,2}, Johann Tuduri^{1,2}, Anthony Pochon¹, and Héctor Campos¹ ¹ISTO, UMR7327, Université d'Orléans, CNRS, BRGM, F-45071 Orléans, France ²BRGM, F-45060 Orléans, France

Magmatic heat sources allow hydrothermal fluids to transport and deposit various types of metals and metalloids. For instance, antimony (Sb) is frequently spatially associated with mafic intrusions, and may be associated with various amounts of other elements, such as Hg, As, W, Au and Ag. However, source-sink relationships in those settings remain poorly constrained. Whether mafic magmas contribute fluids, metals and metalloids to hydrothermal systems at the origin of Sb mineralisation remains uncertain. Spatial and chronological correlations between Sb ore deposits and mafic magmatism have been acknowledged in Variscan settings (i.e. the Armorican Massif and the Central Iberian Zone), but no causal processes have been established yet (Pochon *et al.*, 2019).

We herein investigated the Sb, As and W contents of magmatic rocks available in the literature, to assess mafic magmas as a potential source for metals and metalloids. The GEOROC database covers most of the rock types occurring in a TAS diagram (n = 7215, whole rock), whereas the Jenner and O'Neill (2012) database focuses on MORBs (n = 601, glass).

Sb, As and W are highly covariant, suggesting a common behaviour during magmatic processes. Sb, As and W concentrations in oceanic magmatic rocks increase with increasing K₂O content. This increase is up to two orders of magnitude in mafic compositions, and one order of magnitude through intermediate and differentiated compositions. Differentiated alkaline magmas therefore generally yield higher Sb, As and W concentrations. Variations in Sb, As and W with major and trace elements composition suggest a major role of mantle source processes, and a minor contribution from fractional crystallisation. In particular, Sb, As and W contents in mafic compositions show a good correlation with the La/Sm ratio, and an absence of correlation with the 87 Sr/ 86 Sr and the 143 Nd/ 144 Nd isotopic ratios, suggesting a crucial control of partial melting processes.

Continental magmatic rocks show a strong Sb, As and W variability, with values up to 3 orders of magnitude higher than oceanic rocks, suggesting the occurrence of crustal contamination. Post-magmatic alteration does not seem to have any effect on Sb, As and W concentrations.

We finally investigate mafic rocks spatially and temporally associated with Sb-Hg \pm As-W-Au-Ag ore deposits, and discuss the possible processes at the origin of their enrichment.

GEOROC, 2020. Geochemical rock database., accessed: 02/10/2020.

Jenner F. E., O'Neill H. St. C., 2012. *Analysis of 60 elements in 616 ocean floor basaltic glasses: TECHNICAL BRIEF.* Geochemistry, Geophysics, Geosystems 13 (2), 11 p.

Pochon A., Branquet Y., Gloaguen E., Ruffet G., Poujol M., Boulvais P., Gumiaux C., Cagnard F., Baele J.-M., Kéré I., Gapais D., 2019. *A Sb* ± *Au mineralizing peak at 360 Ma in the Variscan belt.* BSGF – Earth Sciences Bulletin 190 (4), 12 p.