



HAL
open science

Mechanoregulation of PDZ Proteins, An Emerging Function

Elsa Bazellères, André Le Bivic

► **To cite this version:**

Elsa Bazellères, André Le Bivic. Mechanoregulation of PDZ Proteins, An Emerging Function. Mechanoregulation of PDZ Proteins, An Emerging Function, pp.257 - 275, 2021, 10.1007/978-1-0716-1166-1_15 . hal-03557734

HAL Id: hal-03557734

<https://hal.science/hal-03557734>

Submitted on 4 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mechanoregulation of PDZ Proteins, An Emerging Function

Elsa Bazellières and André Le Bivic

Abstract

Mechanical forces have emerged as essential regulators of cell organization, proliferation, migration, and polarity to regulate cellular and tissue homeostasis. Changes in forces or loss of the cellular response to them can result in abnormal embryonic development and diseases. Over the past two decades, many efforts have been put in deciphering the molecular mechanisms that convert forces into biochemical signals, allowing for the identification of many mechanotransducer proteins. Here we discuss how PDZ proteins are emerging as new mechanotransducer proteins by altering their conformations or localizations upon force loads, leading to the formation of macromolecular modules tethering the cell membrane to the actin cytoskeleton.

Key words PDZ proteins, Forces, Tight junctions, Adherens junctions, Actomyosin

1 A Brief Introduction on Mechanotransduction/Mechanoregulation in Biology

All the cells and tissues of the body are subject to external and internal forces. These forces can affect the shape and intracellular organization of cells, their proliferation, their migration, and their intercellular interactions. Forces influence the development of embryos [1] as well as cell functions and homeostasis in the adult [2]. Moreover, many disease states are characterized by changes in these forces and/or a loss of the normal cellular response to them [3]. Over the last decade, mechanotransduction has emerged as a key process in development and diseases. Mechanotransduction can be defined as a cellular event that converts a mechanical input such as fluid shear stress (blood vessels), stretch (lung, intestine), osmotic forces (urinary tract), mechanical load (bone, muscle) [4, 5] as well as the impact of the stiffness of the extracellular matrix (ECM) that surrounds most cells leading to a biochemical response [6]. In well-studied examples of mechanotransduction, proteins can undergo force-induced changes into conformations that lead to modification in affinity for binding partners or catalytic activity. The mechanical load triggers biochemical changes that can

propagate via the activation of specific signaling pathways [7]. In a simple context, the mechanical load can accelerate the association or dissociation of protein–ligand bonds [8]. In the case of adhesion proteins, that exhibit catch–bond behavior, mechanical loads can produce changes in the conformation of the proteins that lead to high affinity for a binding partner [9–11].

Deciphering how fundamental physical principles are controlled by protein interactions is central to understand how forces are converted into biochemical signals. Elucidation of the specificity, selectivity, and regulatory mechanisms involved in protein–protein interactions can therefore provide important insights into many biological processes such as cell proliferation, cell migration, and cell polarity. Structural studies have revealed that mechanosensitive proteins with multiple domains and flexible interdomain interfaces can pass through multiple conformations [12, 13]. For example, a bent conformation can open up to a straighter arrangement of domains along the direction of the applied force. These changes in conformation result in modification of the affinity for the binding partners, exposure or hiding of different catalytic domains that will trigger differential intracellular signaling responses or even redistribution of the protein within the cell [14–16]. Among the proteins that mediate protein–protein interactions, there is the large PDZ (postsynaptic density 95/Disc large/Zonula occludens) family. Most of the PDZ proteins are multimodular scaffold proteins and often contain multiple PDZ domains which can interact with various binding partners and thereby assemble supramolecular signaling complexes [17].

As PDZ domains interact with motifs present in many proteins, understanding the regulatory mechanism of PDZ mediated interactions is important to gain insight into biological processes. So far, posttranslational modifications, autoinhibition, and allosteric interactions have been proposed to regulate PDZ-mediated interactions and thus intracellular signaling [18], but little is known about the impact of mechanical inputs on PDZ proteins. In this review, we will focus on the current knowledge on the mechanoregulation of PDZ proteins, focusing on few recent findings.

2 PDZ Proteins as Mechanotransducers

Biological mechanotransducers can be defined as a single protein or a protein complex that produce or enable a chemical signal in response to mechanical stimuli. These mechanotransducers can participate in mechanoreception and mechanotransmission as direct mechanosensitive structure, which respond by altering their conformation upon force loads, or as second line mechanotransducers. Among the mechanotransducer proteins studied so far, the PDZ protein family is emerging as a new pool of protein sensitive to

forces, but only few examples are known so far. In epithelial cells, the adhesion machineries at the cell–cell or cell–substrate interfaces are known to play an important role in mechanotransduction [10, 19]. These machineries are composed of several proteins, and many of them are part of the PDZ family. Being integrated in a complex modular array tethering the cell membrane to the cell cytoskeleton, the adhesion scaffolding PDZ proteins might constitute an important link to transduce the mechanical signals into an appropriate cell response to maintain cellular homeostasis. We will thus focus on PDZ proteins from the adhesion machinery that have been recently shown to be regulated by forces in a direct or indirect manner (Fig. 1).

3 Force Regulated PDZ Proteins

3.1 ZO1

ZO1 is a tight junction scaffolding protein that belongs to the MAGUK (membrane associated guanylated kinase homolog) family. It is characterized by the occurrence of three PDZ, one SH3 and a GUK (guanylate kinase homologous domain) domains [21]. ZO1 is a cytoplasmic protein, that anchors actin filaments through its actin binding domain at its C terminal region [22, 23]. At the N-terminus, the PDZ1 and PDZ3 domains bind to the tight junction membrane associated proteins, Claudins and JAM-A, and to TAZ, a member of Hippo pathway, while the PDZ2 domain promotes heterodimerization between ZO1 and either ZO2 or ZO3 [22, 24–26]. A larger region encompassing the PDZ3, SH3, U5, and GUK domains (ZPSG-1) interacts with and recruits to junctions the transmembrane TJ protein occludin, but also DbpA/ZONAB (DNA binding protein A/ZO1 associated Nucleic Acid-binding protein) [27, 28]. The sequestration of DbpA by ZO1 and ZO2 at the junctions of confluent monolayers inhibits its nuclear activity that regulates gene expression and cell proliferation [27–30].

Recently, Spadaro et al. demonstrated that ZO1 stretches upon mechanical forces [31]. In 2011, Then et al. have already proposed an impact of membrane tension on the localization of ZO1 [32]. Under hyperosmotic stress that generates an increase in membrane tension, the actin cytoskeleton is reorganized with the appearance of a dense F-actin cortical ring. In this condition, ZO1 expression is increased, and it colocalizes with the newly formed actin ring [32]. As ZO1 is anchored to the actin filaments through its C-terminal region, this change in actin organization exerts direct pulling forces on the ZO1 protein. When present as a heterodimer with ZO2 within cells, ZO1 is in a stretched configuration that allows binding to DpbA. However, the depletion of ZO2 together with an inhibition of myosin contractility promotes a folding of the N-terminal and C-terminal end of ZO-1, and releases the




Fig. 1 Hypothetical role of the PDZ protein in the strengthening and stabilization of intercellular adhesions. (a) At the level of the tight junctions (TJs), under low tension the mechanotransducer ZO1 is in an inactive

interaction with DpbA and Occludin. By applying linearly increasing forces to purified full-length single ZO1 molecules with magnetic tweezers, Spadaro et al. could demonstrate that both the C-terminal region and the ZPSG-1 module of ZO1 that comprises the PDZ3, SH3, U5, and GUK domains unfold at forces ranging from 5 to 20 pN, releasing the autoinhibition interaction between ZPSG and C terminal domain of ZO1 [31]. ZPSG-1 domain is not only important for the junctional localization of ZO1 but also for its interaction with Occludin, DbpA, and thus for barrier formation and epithelial polarization [33–38]. Forces act as an allosteric effector by stretching ZO1 protein to promote the interaction of ZPSG-1 with its ligands, occludin and DbpA. While the stretched conformation of ZO1 is the active conformation, the folded conformation is the inactive form in which the ZPSG domain is autoinhibited. In its inactive form, junctional ZO1 remains anchored to the membrane through binding of its N-terminal domain with interactors such as TAZ or Claudins, while the C-terminal half is intramolecularly autoinhibited. The release of ZPSG-1 domains may also regulate the interaction of other proteins such as α -Catenin, Afadin, JAM-A, Vinculin and Shroom2 [39].

3.2 MUPP1

MUPP1 belongs to the family of multi-PDZ proteins and contains a L27 domain at its N-terminal region followed by 13 PDZ domains [40]. MUPP1 is a structural paralog of PATJ, and both share several binding partners such as PALS1, PAR-6, AMOT, Jeap, ZO3, Claudins, or Nectins [41]. Lanaspa et al. by the use of osmolarity changes demonstrated that both acute and chronic hyperosmolarity in inner medullary collecting duct 3 cells induce an increase in the expression of MUPP1, ZO1, and Afadin [42]. As MUPP1 expression increases, it localizes to the apical side of the membrane at the level of the Tight Junctions (TJs). To survive hyperosmotic stresses and to maintain the integrity of the cell sheet with efficient barrier functions, cells have to adapt through

Fig. 1 (continued) conformation, and the second line mechanotransducers Afadin and Mupp1 are in the cytoplasm. The actin filaments are exerting low mechanical loads. Under high tension, generated by actomyosin bundles, ZO1 unfolds and un.masks several domains that leads to the binding with Occludin and JAM together with the recruitment of DbpA and Afadin. Afadin is then able to recruit Rap2c generating a positive feedback loop on RHO allowing for actin contractions and thus force generation. This increase in forces results in the recruitment of Mupp1 at the TJs, that will bind to ZO1, ZO2, and JAM but also potentially to the CRB3A/Pals1 polarity complex. The formation of this macromolecular complex that tethers transmembrane proteins (Claudins, Occludins, JAM) to the actin cytoskeleton is key for the strengthening and the stabilization of the TJs. **(b)** In the lateral domain of the cells, at the level of the lateral Adherens Junctions (LAJs), the actin filaments that binds to the E-cadherin–catenin complex, exert low tension and Scribble is in the cytoplasm. Above the lateral junctions, the E-cadherin–catenin complex is link to vinculin and binds to actomyosin bundles that are under high mechanical loads. SCRIBBLE is also enriched at the AJ level, and plays a key role in the stabilization of the AJs

reorganization of the actin cytoskeleton as mentioned previously but also through differential expression of proteins, that move to the junctions to counterbalance the changes in membrane tension. Depletion of MUPP1, results in a disruption of the TJ barrier, with a drop of transepithelial resistance of about 25% [42]. Molecularly, depletion of MUPP1, in hyperosmotic conditions, triggers a decreased expression and loss of membrane localization of Claudin4, a TJ protein [42]. These data clearly show that MUPP1 is important in the maintenance of the epithelial homeostasis and confirm previous study that pointed out a predominant role of MUPP1 in the disruption of both TJs barrier and apicobasal polarity [43–45]. A more recent study shows that MUPP1 stability and degradation in endothelial cells depends on the regulation of PDZRN3, an E3 ubiquitin ligase PDZ domain containing ring finger 3 protein [46]. The interactions between PDZRN3, MUPP1, PAR3, and aPKC regulate the stabilization of TJs in endothelial cells. Perturbation of PDZRN3 expression induces degradation of MUPP1 that correlates with destabilization of the actin cytoskeleton and disruption of the TJs. All these studies point toward a role of MUPP1 in the stabilization and maintenance of actin cytoskeleton and TJs in cells.

20 years ago, it was suggested that MUPP1 could change its conformation through the interaction of its PDZ10 domain with the 5-hydroxytryptamine type 2C receptors [47]. This interaction induces a conformational change in MUPP1 and triggers a clustering of the 5-hydroxytryptamine type 2C receptors at the cell membrane, triggering downstream signal transduction pathways. It is of interest that different PDZ domains of MUPP1 can bind to CADM1, a transmembrane cell adhesion protein, with different affinities. The PDZ2 of MUPP1, for example, presents the highest affinity for CADM1, but when this PDZ is absent, CADM1 can still interact with other PDZ domains of MUPP1. These data could point to a change in conformation of MUPP1 that would unmask the PDZ2 domain leading to a strong binding to CADM1 [48, 49]. In the case of CADM1 and maybe of other interactors of MUPP1, the interactions with multiple domains can be an alternative when some domains are either occupied by other ligands or hidden by an autoinhibitory conformation of MUPP1. These multi-low-affinity interactions might then serve as transient interaction contacts before MUPP1 reaches its stretched configuration. However, the unfolding/folding regulation of MUPP1 is still under debate as only indirect evidence has been obtained to date.

3.3 PAR3

PAR3 is part of the Partitioning defective proteins and presents a CRI oligomerization domain, 3 PDZ domains and aPKC binding domain. A recent study pointed out that cortical forces are responsible for the clustering of PAR3 on the cell cortex. In *C. elegans*, inhibition of actin contraction with blebbistatin or actin filament

polymerization causes a reduction of the PAR3 clusters on the cortex, demonstrating that higher cortical tension can drive PAR3 cluster formation at the cortex [50]. The actomyosin membrane associated network generates flows [51], which promote symmetry breaking along the anterior posterior axis through the advection of polarity components [52–54]. The polarized actomyosin contractions pull actin networks along the membrane toward the anterior region while triggering local disassembly and turnover via increased local tension, resulting in flow of material [55]. Based on the advective flow model proposed by Goerhing et al. [53] that predicts that diffusivity and turnover of cortical polarity proteins should be slow and stable enough to be passively transport by the advective cortical flow, Wang et al. [50] postulate a “clustering and stabilization” hypothesis for PAR3. By using fluorescence recovery after photobleaching, they demonstrate that the half-life of PAR3 and PKC are shorter before symmetry breaking. During the early to middle establishment phase, when the clusters are formed, the half-life increases. Molecularly, during this phase the activity of Cdc42 is reduced allowing PKC3 and PAR6 to associate with PAR3 clusters at the cortex and thus facilitates effective transport by advective cortical flows. During the maintenance phase, the half-lives of PAR3 and PKC are shorter, when the clusters disassemble, concomitant with an increased activity of Cdc42 that prevents PKC and PAR6 association with PAR3. As cortical tensions trigger a dynamic equilibrium between an unclustered and clustered form of PAR3 correlating with its association with PAR6 and PKC, it is tempting to speculate that PAR3 may undergo structural changes by mechanical stretching, which could relieve the amino-terminal CRI domain from intramolecular inhibitions, thus promoting PAR3 oligomerization [56]. These mechanically induced conformational changes may directly activate the ability of the CRI domain to facilitate its oligomerization, and proper localization in cells [56]. Furthermore, similar cortical flows are observed during cell division or cell migration, and then could, if proven, also explain the interaction between the different members of the PAR complex.

204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240

4 PDZ Proteins “Second Line” Mechanotransducers

241

In this part, we will describe how some PDZ proteins interact with proteins that are activated upon forces. These interactions may trigger a change in the conformation or in the localization of the “second line” mechanotransducers, thereby regulating different signal pathways.

242
243
244
245
246

4.1 PAR6

PAR6 is part of the Partitioning defective proteins in *C. elegans*, and encodes a protein with PKC binding domain (PB1), a semi-Cdc42/Rac1 interactive binding (CRIB) and a PDZ domain [57]. Recently a PDZ binding motif (PBM) was identified at its C-terminal region [58]. PAR6 in many species is involved in cell division, cell polarization, and cell migration, processes that are characterized by changes in cellular forces [59–61].

PAR6 function is able to segregate to the apical or leading edge of epithelial cells upon activation by Cdc42-GTP [57, 62]. Accumulating evidence indicates that Cdc42 responds to and is activated upon different mechanical loads such as hyperosmolarity, shear stress, and intercellular increased tension [63]. Upon exposure to mechanical loads, Cdc42 translocates from cytosol to the membrane and is concomitantly activated [64]. Activation of Cdc42 by exchange of GDP for GTP triggers actin polymerization and generation of tension [65, 66]. Binding of activated Cdc42GTP to the CRIB domain of PAR6 favors the interaction with CRUMBS over the interaction with PALS1 through a change in conformation of the CRIB-PDZ domains of PAR6 [67]. Upon Cdc42-GTP binding, a portion of the flexible CRIB motif folds into a stable conformation with the PDZ domain [68], triggering an increase affinity for the interaction with CRUMBS, and releasing the interaction with PALS1. The modular nature of PAR6 may allow the mutually exclusive interaction of CRUMBS and PALS1 with its PDZ domains, regulating the assembly and localization of different polarity complexes depending on the cellular context.

The localization of PAR6 to the plasma membrane depends on both the PDZ and the PBM domains, as both deletions caused a strong mislocalization of PAR6 to the cytosol. The PBM domain binds with different affinities to the PDZ1 and PDZ3 of PAR3, whose clustering is mediated by an increase in cortical tension as previously mentioned [50]. The weak but multivalent interaction of one PAR3 molecule with two PAR6 molecules might allow the assembly a large cluster of PAR complexes at the cell membrane. The formation of the large-scale cluster of PAR complexes can facilitate the actomyosin dependent advective transport of several PAR6 proteins during the establishment of polarity but also the formation of large clusters made with several proteins such as aPKC, PALS1, or CRUMBS that can serve different functions.

4.2 DLG

DLG is a MAGUK protein that presents an L27 domain, 3 PDZ domains, an SH3 domain, a HOOK domain and a Guanylate like domain. DLG is a member of the basolateral polarity complex SCRIBBLE, and is involved in processes such as cell division, cell migration and cell polarity [69]. The N-terminal L27 domain interacts with the two L27 domains of calcium/calmodulin-dependent serine protein kinase (CASK) [70]. CASK is a membrane-associated guanylate kinase and a scaffolding protein, that has been

demonstrated to recruit or organize other proteins at the plasma membrane to coordinate signal transduction pathways within the cytoplasm and nucleus [71]. A study done in 2016, using a cell mechanical stretch device shows that CASK expression and localization to the basal membrane is needed for the inhibition of proliferation of cells under cyclic stretch [72]. The authors show that CASK interacts with $\beta 1$ -integrin, however, the tension-driven interaction between these two proteins still has to be formally proven. Depletion of CASK results in aberrant proliferation of cells under mechanical stress demonstrating that CASK localization under tension is important for the mechanoregulation of the cell division [72]. In 2019, Porter et al. demonstrated that the direct interaction of CASK with DLG is required for normal cortical recruitment of NUMA, a key component of spindle orientation machinery [73]. Disruption of this interaction affects the integrity of epithelial architecture and results in misoriented cell division that give rise to multilumen cyst in 3D. Since the formation of 3D cysts generates an increase in tension at the cell–cell interface [74], it is tempting to speculate that CASK is recruited at the cell–cell interface in a force-dependent manner and then will recruit DLG allowing the formation of well polarized 3D cysts.

DLG interacts with the C terminal PDZ binding domain of CD97 and together these proteins are part of the adherens junctional signaling complex composed of E-cadherin and catenins [75]. This macromolecular complex is linked to the F actin cortex and is thus submitted to cellular forces. The localization of CD97 at the plasma membrane is actin dependent as blocking actin polymerization and elongation prevents its membrane localization. When present at the cell membrane CD97 strengthens the adherens junction (AJs) [76] while deletion of its PDZ binding domain results in the loss of cell–cell contacts [77] upon mechanical shear stress. Taken together these data show a role of CD97 in the mechanoregulation of cell–cell contacts. In a recent study, mechanical stimulation of epithelial cells applied by using shear stress or wound assay results in a rapid phosphorylation of Ser740 of CD97. This phosphorylation disrupts the binding of DLG1 to the PDZ binding domain of CD97, and correlates with a disorganization of the actin cytoskeleton. PKC contributes to the mechanical force induced cellular responses [78] and is a potential candidate to phosphorylate CD97 at S740. PKC α interacts via its PBM with PDZ3 domain of DLG1 [79]. Upon mechanical stress, PKC α might be recruited to the cell membrane via DLG1 and will then trigger phosphorylation of CD97 causing F-actin depolymerization, loss of cell–cell contact, and DLG1 detachment. This signaling pathway when activated induces depolymerization of actin and loss of cell–cell contacts, that will result in a relaxation of the tension at this particular cellular junction avoiding tissue breaking.

4.3 Afadin

Afadin is a filamentous actin binding protein with two Ras domains, a forkhead association domain, a dilute domain, a PDZ domain, and three proline-rich domains. Afadin is implicated in many cellular processes from cell survival, cell proliferation to cell migration and formation of the apical junctions in epithelial cells [80–83]. Afadin can interact both directly or indirectly with the actin cytoskeleton through several partners such as JAM-A, ZO1/ZO2, vinculin, and α -actinin [84–88]. Afadin is thus a strong candidate to be involved in mechanoregulated processes as many of its partners have been already linked to mechanotransduction. Interestingly, depletion of Afadin, JAM-A or double depletion of ZO1/ZO2 results in similar phenotypes with increased contractility triggered by activation of RhoA and phosphorylation of myosin-light chain [89–91]. In response to ZO1/ZO2 double depletion, Afadin is recruited to the cellular cell contacts, and inhibiting contractility perturbs the homogeneous localization of Afadin. Thus, myosin contractility is essential for maintaining Afadin uniform distribution along the zonula adherens (ZA). Removing Afadin in ZO1/ZO2 depleted cells specifically altered actomyosin architecture at the ZA of tricellular junctions where actin cables are anchored. This perturbation is accompanied by discontinuities in the E-cadherin, claudin, and occludin stainings [92]. Taken together, this data shows a synergy between ZO and Afadin depletions in disrupting tissue integrity under tension. Afadin may therefore act as a robust protein scaffold that maintains ZA architecture at tricellular junctions.

JAM-A, a component of the TJs, via its PDZ binding motif, can associate with signaling molecules such as scaffold PDZ proteins, ZO1/2, and Afadin, as well as with the guanine exchange factor PDZ-GEF2 [83, 93]. Monteiro et al. demonstrate that JAM-A interacts with Afadin, ZO2, and PDZ-GEF1 to activate the small GTPase Rap2c [89]. The activation of Rap2c controls the contraction of the apical cytoskeleton regulating the epithelial permeability to prevent cell injury. In this study, the authors also mention that Afadin is able to immunoprecipitate a doublet of JAM-A that might represent phosphorylated forms of JAM-A, although they did not investigate this phosphorylation but another team did. Scott et al show that mechanical stimulation can trigger phosphorylation of JAM-A [94]. Forces were applied by using fluid shear stress or by using magnetic tweezer. After applying forces, several biochemical analyses were performed that demonstrate that tension induces rapid phosphorylation of JAM-A S284. This phosphorylation of JAM-A induces activation of RhoA by PKC ζ triggering cell stiffening by modifying actin cytoskeleton. These results clearly demonstrate that JAM-A is a direct transducer of mechanical forces. Interestingly, when JAM-A is localized at the TJs which are under high levels of tension [95] it is phosphorylated [96]. It is thus tempting to speculate that mechanical tension, above a certain

threshold might induce the phosphorylation of JAM-A. This phosphorylation will control the affinity for Afadin thereby controlling the contraction of apical cytoskeleton that allows for the maintenance of ZA and TJs.

4.4 SCRIBBLE

SCRIBBLE is a classical multimodular scaffold protein that contains 16 leucine-rich repeats (LRR) and four PDZ domains. N-terminal LRR domain can associate with LGL and is required for the association with the lateral cortex and the establishment of apicobasal polarity [97, 98]. The C-terminal PDZ domains of SCRIBBLE can interact with a diverse range of proteins such as β -Pix [99], Vangl2 [100], and LGN [101], allowing for cell polarization, cell migration [102], establishment of planar cell polarity, and cell division, respectively. In epithelial cells, SCRIBBLE is required for normal intercellular adhesion by stabilizing E-cadherin-p120 catenin at the plasma membrane [103]. 10 years ago, E-cadherin complex was tagged as a mechanosensor complex. Since then many studies using different approaches such as FRET sensor, magnetic tweezer, and cell stretching have confirmed that E-cadherin can sense changes in mechanical forces within an epithelial monolayer [104–106]. The role of SCRIBBLE in mechanotransduction has, however, not been investigated so far. Some hints based on the function of SCRIBBLE in the stabilization of E-cadherin point to a role in the mechanoregulation of AJs. E-cadherin receptors form adhesive clusters that are coupled to the contractile actomyosin cortex [107, 108]. At the cellular level, E-cadherin adhesion not only binds cells together but also mechanically couples the contractile modules of neighboring cells together to generate junctional tension [109]. This junctional tension has been mainly treated as homogeneous along the cell–cell interface. However, E-cadherin forms clusters of different sizes along the junctions between cells [108–110]. Interestingly, despite overall basolateral localization, SCRIBBLE is enriched at the ZAs, where it stabilizes E-cadherin-p120 catenin adhesions [111, 112]. At the level of the ZAs, clusters of stabilized E-cadherin are linked to large actomyosin bundles [113–115]. Below the AJs, at the lateral adherens junctions (LAJs), E-cadherin is coupled to less aligned actomyosin network. Contractile tension is thereby greater at the ZA than at the LAJ. SCRIBBLE could be enriched at the ZA in a tension-dependent manner, and thereby by stabilizing E-cadherin could be implicated in a feedback positive loop in the generation of forces at the ZAs. This tension driven localization of SCRIBBLE to E-cadherin can also take place during the orientation of the mitotic spindle to align epithelial cell division. In 2017, Hart et al., by applying uniaxial stretch on epithelial cells, showed that cell division aligned with the stretch axis [116]. The orientation of this cell division axis requires trans-engagement of E-cadherin adhesions and involves tension-dependent recruitment of LGN to

E-cadherin. LGN can directly interact with the cytosolic tail of E-cadherin that localizes LGN at the cell–cell contacts [117]. Increase in tension does not trigger a polarized distribution of E-cadherin, while LGN and myosin IIA are polarized suggesting that an additional intermediate, LGN interacting protein might be involved in its polarized localization. This protein could be SCRIBBLE as other ones (DLG, Afadin, ERM proteins) were ruled out in this study. SCRIBBLE is indeed able to interact with LGN, and the interaction between LGN and E-cadherin is reduced in cells depleted for SCRIBBLE. The formation of this ternary complex is also important for proper cell division [101]. Taken together, all these data point to SCRIBBLE as a likely “second line” mechanotransducer to trigger signaling pathways downstream of the mechanosensitive E-cadherin units.

5 Concluding Remarks

In the last two decades, mechanical stimuli have emerged as essential regulators of several biological processes. Mechanotransduction has revealed a new layer of control of the interaction between proteins, and will potentially lead to global guiding principles for the organization of complex living systems. The convergent interests from physicists and biologists to understand the complexity of integrated systems have led to the development of new biophysical tools. These new technological advances such as stretching devices, optical/magnetic tweezers, or atomic force microscopy have enabled to measure and apply controlled forces on cells. Applying different forces on cells trigger several cellular responses and activate different signaling pathways that have led to the identification of an emerging mechanotransducing function for PDZ proteins. These mechanotransducers can participate in mechanoreception and mechanotransmission as direct mechanosensitive components or as second line mechanotransducers.

Pulling forces generated by the actomyosin network can stretch molecules exposing their cryptic binding sites or cryptic phosphorylation sites, triggering specific signaling pathways [5, 118]. In the case of ZO1, physical forces are responsible for the stretching of the molecule that is needed for its correct localization while allowing for the interaction with its different partners. This activated unfolded ZO1 protein is important for the maintenance of the TJs. The actomyosin network can thus directly by its contractions pull on proteins but also on the actin cytoskeleton, resulting in flow of material that leads to the formation of polarized clusters of proteins at the cell cortex that allow for adaptation of cell cytoskeleton [51, 119]. Under higher cortical tension PAR3 clusters and MUPP1 is recruited to the cell cortex. The tension-dependent clustering of PAR3 could promote its oligomerization at the cell

cortex where it could recruit PAR6 and aPKC, leading to the establishment of anteroposterior or apicobasal polarity. The recruitment of the PDZ protein MUPP1 to the TJs is key to maintaining the integrity of the TJ barrier during hyperosmotic stress. Finally, loading forces can regulate cell adhesion machineries [105, 106, 120]. The adhesive contacts can adapt to external forces, by modifying their binding to specific proteins, their structural conformations, their stability but also their links to the actin cytoskeleton. Some PDZ proteins such as SCRIBBLE, Afadin, or ZO1 can, upon loading forces, strengthen the cell adhesion machinery at the ZA and TJs localization respectively to preserve epithelial homeostasis by allowing for the proper polarization of the tissue and the tethering of the adhesions to the actin cytoskeleton. In these cases, the connections between adhesions are strengthened by assembly of new components, however in other cases the connections can be severed [121]. DLG is located at cell–cell contacts and upon forces generation will recruit aPKC at the ZA. The recruitment of aPKC by DLG at the ZA can then regulate the load of tension at the cell–cell contact by controlling the depolymerization of actin and disruption of cell–cell contact leading to a relaxation of the tissue when the forces are too elevated.

The active nature of cell adhesion machineries implies that the time from the mechanosensing to the mechanoreponse is slow and occurs within seconds to minutes allowing the occurrence of many different interactions between proteins [122–124]. In the case of PDZ scaffolds proteins, this could trigger specific interactions allowing for the cells to have a multimodal adaptive “repertoire” to properly adapt their responses to the change in forces.

Acknowledgments

Elsa Bazellières and André Le Bivic are supported by CNRS. This project was developed in the context of the LabEx INFORM (ANR-11-LABX-0054) and of the A*MIDEX project (ANR-11-IDEX-0001-02) funded by the “Investissements d’Avenir” French Government program.

References

- Petridou NI, Heisenberg C (2019) Tissue rheology in embryonic organization. *EMBO J* 38:e102497
- Salvi AM, DeMali KA (2018) Mechanisms linking mechanotransduction and cell metabolism. *Curr Opin Cell Biol* 54:114–120
- Jaalouk DE, Lammerding J (2009) Mechanotransduction gone awry. *Nat Rev Mol Cell Biol* 10:63–73
- Sidhwani P, Yelon D (2019) Fluid forces shape the embryonic heart: insights from zebrafish. *Curr Top Dev Biol* 132:395–416
- Pruitt BL, Dunn AR, Weis WI, Nelson WJ (2014) Mechano-transduction: from molecules to tissues. *PLoS Biol* 12:e1001996
- Dzamba BJ, DeSimone DW (2018) Extracellular matrix (ECM) and the sculpting of

539 embryonic tissues. *Curr Top Dev Biol* 130:245–274

540

541 7. Hoffman BD, Grashoff C, Schwartz MA (2011) Dynamic molecular processes mediate cellular mechanotransduction. *Nature* 475:316–323

542

543

544

545 8. Depoil D, Dustin ML (2014) Force and affinity in ligand discrimination by the TCR. *Trends Immunol* 35:597–603

546

547

548 9. Chen W, Lou J, Evans EA, Zhu C (2012) Observing force-regulated conformational changes and ligand dissociation from a single integrin on cells. *J Cell Biol* 199:497–512

549

550

551

552 10. Hu X, Margadant FM, Yao M, Sheetz MP (2017) Molecular stretching modulates mechanosensing pathways: molecular stretching modulates mechanosensing pathways. *Protein Sci* 26:1337–1351

553

554

555

556

557 11. Pannekoek W-J, de Rooij J, Gloerich M (2019) Force transduction by cadherin adhesions in morphogenesis. *F1000Res* 8:1044

558

559

560 12. Cailliez F, Lavery R (2005) Cadherin mechanics and complexation: the importance of calcium binding. *Biophys J* 89:3895–3903

561

562

563 13. Springer TA, Dustin ML (2012) Integrin inside-out signaling and the immunological synapse. *Curr Opin Cell Biol* 24:107–115

564

565

566 14. Rutherford SL, Zuker CS (1994) Protein folding and the regulation of signaling pathways. *Cell* 79:1129–1132

567

568

569 15. Papagrigoiriou E, Gingras AR, Barsukov IL, Bate N, Fillingham IJ, Patel B, Frank R, Ziegler WH, Roberts GC, Critchley DR, Emsley J (2004) Activation of a vinculin-binding site in the talin rod involves rearrangement of a five-helix bundle. *EMBO J* 23:2942–2951

570

571

572

573

574

575 16. Smith ML, Gourdon D, Little WC, Kubow KE, Eguiluz RA, Luna-Morris S, Vogel V (2007) Force-induced unfolding of fibronectin in the extracellular matrix of living cells. *PLoS Biol* 5:e268

576

577

578

579

580 17. Manjunath GP, Ramanujam PL, Galande S (2018) Structure function relations in PDZ-domain-containing proteins: implications for protein networks in cellular signalling. *J Biosci* 43:155–171

581

582

583

584

585 18. Lee H-J, Zheng JJ (2010) PDZ domains and their binding partners: structure, specificity, and modification. *Cell Commun Signal* 8:8

586

587

588 19. Charras G, Yap AS (2018) Tensile forces and mechanotransduction at cell–cell junctions. *Curr Biol* 28:R445–R457

589

590

591 20. Heinemann U, Schuetz A (2019) Structural features of tight-junction proteins. *Int J Mol Sci* 20:6020

592

593

594 21. Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA (1986) Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biol* 103:755–766

595

596

597

598

599

600 22. Fanning AS, Jameson BJ, Jesaitis LA, Anderson JM (1998) The tight junction protein ZO-1 establishes a link between the transmembrane protein occludin and the actin cytoskeleton. *J Biol Chem* 273:29745–29753

601

602

603

604

605 23. Itoh M, Nagafuchi A, Moroi S, Tsukita S (1997) Involvement of ZO-1 in cadherin-based cell adhesion through its direct binding to α catenin and actin filaments. *J Cell Biol* 138:181–192

606

607

608

609

610 24. Itoh M, Furuse M, Morita K, Kubota K, Saitou M, Tsukita S (1999) Direct binding of three tight junction-associated Maguks, Zo-1, Zo-2, and Zo-3, with the CooH termini of claudins. *J Cell Biol* 147:1351–1363

611

612

613

614

615 25. Wittchen ES, Haskins J, Stevenson BR (1999) Protein interactions at the tight junction. Actin has multiple binding partners, and ZO-1 forms independent complexes with ZO-2 and ZO-3. *J Biol Chem* 274:35179–35185

616

617

618

619

620

621 26. Remue E, Meerschaert K, Oka T, Boucherie C, Vandekerckhove J, Sudol M, Gettemans J (2010) TAZ interacts with zonula occludens-1 and -2 proteins in a PDZ-1 dependent manner. *FEBS Lett* 584:4175–4180

622

623

624

625

626

627 27. Balda MS, Matter K (2000) The tight junction protein ZO-1 and an interacting transcription factor regulate ErbB-2 expression. *EMBO J* 19:2024–2033

628

629

630

631 28. Spadaro D, Tapia R, Jond L, Sudol M, Fanning AS, Citi S (2014) ZO proteins redundantly regulate the transcription factor DbpA/ZONAB. *J Biol Chem* 289:22500–22511

632

633

634

635

636 29. Balda MS, Garrett MD, Matter K (2003) The ZO-1-associated Y-box factor ZONAB regulates epithelial cell proliferation and cell density. *J Cell Biol* 160:423–432

637

638

639

640 30. Sourisseau T, Georgiadis A, Tsapara A, Ali RR, Pestell R, Matter K, Balda MS (2006) Regulation of PCNA and cyclin D1 expression and epithelial morphogenesis by the ZO-1-regulated transcription factor ZONAB/DbpA. *Mol Cell Biol* 26:2387–2398

641

642

643

644

645

646

647 31. Spadaro D, Le S, Laroche T, Mean I, Jond L, Yan J, Citi S (2017) Tension-dependent stretching activates ZO-1 to control the

648

649

- 650 junctional localization of its interactors. *Curr*
651 *Biol* 27:3783–3795.e8
- 652 32. Then C, Bergler T, Jeblick R, Jung B,
653 Banas B, Krämer BK (2011) Hypertonic stress
654 promotes the upregulation and phosphoryla-
655 tion of zonula occludens 1. *Nephron Physiol*
656 119:p11–p21
- 657 33. Fanning AS, Ma TY, Anderson JM (2002)
658 Isolation and functional characterization of
659 the actin binding region in the tight junction
660 protein ZO-1. *FASEB J* 16:1835–1837
- 661 34. Fanning AS, Little BP, Rahner C,
662 Utepergenov D, Walther Z, Anderson JM
663 (2007) The unique-5 and -6 motifs of ZO-1
664 regulate tight junction strand localization and
665 scaffolding properties. *Mol Biol Cell*
666 18:721–731
- 667 35. Umeda K, Ikenouchi J, Katahira-Tayama S,
668 Furuse K, Sasaki H, Nakayama M, Matsui T,
669 Tsukita S, Furuse M, Tsukita S (2006) ZO-1
670 and ZO-2 independently determine where
671 claudins are polymerized in tight-junction
672 strand formation. *Cell* 126:741–754
- 673 36. Ikenouchi J, Umeda K, Tsukita S, Furuse M,
674 Tsukita S (2007) Requirement of ZO-1 for
675 the formation of belt-like adherens junctions
676 during epithelial cell polarization. *J Cell Biol*
677 176:779–786
- 678 37. Rodgers LS, Beam MT, Anderson JM, Fan-
679 ning AS (2013) Epithelial barrier assembly
680 requires coordinated activity of multiple
681 domains of the tight junction protein ZO-1.
682 *J Cell Sci* 126:1565–1575
- 683 38. Odenwald MA, Choi W, Buckley A,
684 Shashikanth N, Joseph NE, Wang Y, Warren
685 MH, Buschmann MM, Pavlyuk R,
686 Hildebrand J, Margolis B, Fanning AS,
687 Turner JR (2017) ZO-1 interactions with
688 F-actin and occludin direct epithelial polariza-
689 tion and single lumen specification in 3D cul-
690 ture. *J Cell Sci* 130:243–259
- 691 39. Fanning AS, Anderson JM (2009) Zonula
692 occludens-1 and -2 are cytosolic scaffolds
693 that regulate the assembly of cellular junc-
694 tions. *Ann N Y Acad Sci* 1165:113–120
- 695 40. Ullmer C, Schmuck K, Figge A, Lübbert H
696 (1998) Cloning and characterization of
697 MUPP1, a novel PDZ domain protein.
698 *FEBS Lett* 424:63–68
- 699 41. Assémat E, Bazellières E, Pallesi-Pocachard E,
700 Le Bivic A, Massey-Harroche D (2008) Polar-
701 ity complex proteins. *Biochim Biophys Acta*
702 1778:614–630
- 703 42. Lanaspa M, Andres-Hernando A, Rivard C,
704 Dai Y, Berl T (2008) Hypertonic stress
705 increases claudin-4 expression and tight junc-
706 tion integrity in association with MUPP1 in
IMCD3 cells. *Proc Natl Acad Sci U S A*
105:15797–15802
- 707 43. Latorre IJ, Roh MH, Frese KK, Weiss RS,
708 Margolis B, Javier RT (2005) Viral
709 oncoprotein-induced mislocalization of select
710 PDZ proteins disrupts tight junctions and
711 causes polarity defects in epithelial cells. *J*
712 *Cell Sci* 118:4283–4293
- 713 44. Shin K, Straight S, Margolis B (2005) PATJ
714 regulates tight junction formation and polar-
715 ity in mammalian epithelial cells. *J Cell Biol*
716 168:705–711
- 717 45. Straight SW, Shin K, Fogg VC, Fan S, Liu C-J,
718 Roh M, Margolis B (2004) Loss of PALS1
719 expression leads to tight junction and polarity
720 defects. *Mol Biol Cell* 15:1981–1990
- 721 46. Sewduth RN, Kovacic H, Jaspard-Vinassa B,
722 Jecko V, Wavasseur T, Fritsch N, Pernot M,
723 Jeaningros S, Roux E, Dufourcq P,
724 Couffinhal T, Duplâa C (2017) PDZRN3
725 destabilizes endothelial cell-cell junctions
726 through a PKC ζ -containing polarity complex
727 to increase vascular permeability. *Sci Signal*
728 10:464
- 729 47. Bécamel C, Figge A, Poliak S, Dumuis A,
730 Peles E, Bockaert J, Lübbert H, Ullmer C
731 (2001) Interaction of serotonin
732 5-hydroxytryptamine type 2C receptors with
733 PDZ10 of the multi-PDZ domain protein
734 MUPP1. *J Biol Chem* 276:12974–12982
- 735 48. Fujita E, Tanabe Y, Imhof BA, Momoi MY,
736 Momoi T (2012) A complex of synaptic adhe-
737 sion molecule CADM1, a molecule related to
738 autism spectrum disorder, with MUPP1 in
739 the cerebellum. *J Neurochem* 123:886–894
- 740 49. Baliova M, Juhasova A, Jursky F (2014) Using
741 a collection of MUPP1 domains to investigate
742 the similarities of neurotransmitter transpor-
743 ters C-terminal PDZ motifs. *Biochem Bio-*
744 *phys Res Commun* 454:25–29
- 745 50. Wang S-C, Low TYF, Nishimura Y, Gole L,
746 Yu W, Motegi F (2017) Cortical forces and
747 CDC-42 control clustering of PAR proteins
748 for *Caenorhabditis elegans* embryonic polar-
749 ization. *Nat Cell Biol* 19:988–995
- 750 51. Munro E, Nance J, Priess JR (2004) Cortical
751 flows powered by asymmetrical contraction
752 transport PAR proteins to establish and main-
753 tain anterior-posterior polarity in the early
754 *C. elegans* embryo. *Dev Cell* 7:413–424
- 755 52. Goehring NW, Hoegge C, Grill SW, Hyman
756 AA (2011) PAR proteins diffuse freely across
757 the anterior-posterior boundary in polarized
758 *C. elegans* embryos. *J Cell Biol* 193:583–594
- 759 53. Goehring NW, Trong PK, Bois JS,
760 Chowdhury D, Nicola EM, Hyman AA,
761 Grill SW (2011) Polarization of PAR proteins
762 763

- 764 by advective triggering of a pattern-forming
765 system. *Science* 334:1137–1141
- 766 54. Mayer M, Depken M, Bois JS, Jülicher F, Grill
767 SW (2010) Anisotropies in cortical tension
768 reveal the physical basis of polarizing cortical
769 flows. *Nature* 467:617–621
- 770 55. McFadden WM, McCall PM, Gardel ML,
771 Munro EM (2017) Filament turnover tunes
772 both force generation and dissipation to control
773 long-range flows in a model actomyosin
774 cortex. *PLoS Comput Biol* 13:e1005811
- 775 56. Benton R, Johnston DS (2003) *Drosophila*
776 PAR-1 and 14-3-3 inhibit Bazooka/PAR-3
777 to establish complementary cortical domains
778 in polarized cells. *Cell* 115:691–704
- 779 57. Joberty G, Petersen C, Gao L, Macara IG
780 (2000) The cell-polarity protein Par6 links
781 Par3 and atypical protein kinase C to Cdc42.
782 *Nat Cell Biol* 2:531–539
- 783 58. Renschler FA, Bruekner SR, Salomon PL,
784 Mukherjee A, Kullmann L, Schütz-Stoffregen
785 MC, Henzler C, Pawson T, Krahn MP, Wies-
786 ner S (2018) Structural basis for the interac-
787 tion between the cell polarity proteins Par3
788 and Par6. *Sci Signal* 11:517
- 789 59. Uroz M, Wistorf S, Serra-Picamal X, Conte V,
790 Sales-Pardo M, Roca-Cusachs P, Guimerà R,
791 Trepas X (2018) Regulation of cell cycle pro-
792 gression by cell-cell and cell-matrix forces.
793 *Nat Cell Biol* 20:646–654
- 794 60. Ladoux B, Mège R-M, Trepas X (2016)
795 Front-rear polarization by mechanical cues:
796 from single cells to tissues. *Trends Cell Biol*
797 26:420–433
- 798 61. Trepas X, Wasserman MR, Angelini TE,
799 Millet E, Weitz DA, Butler JP, Fredberg JJ
800 (2009) Physical forces during collective cell
801 migration. *Nat Phys* 5:426–430
- 802 62. Etienne-Manneville S, Hall A (2003) Cell
803 polarity: Par6, aPKC and cytoskeletal cross-
804 talk. *Curr Opin Cell Biol* 15:67–72
- 805 63. Lewis A, Di Ciano C, Rotstein OD, Kapus A
806 (2002) Osmotic stress activates Rac and
807 Cdc42 in neutrophils: role in hypertonicity-
808 induced actin polymerization. *Am J Phys Cell*
809 *Phys* 282:C271–C279
- 810 64. Li S, Chen BPC, Azuma N, Hu Y-L, Wu SZ,
811 Sumpio BE, Shyy JY-J, Chien S (1999) Dis-
812 tinct roles for the small GTPases Cdc42 and
813 Rho in endothelial responses to shear stress. *J*
814 *Clin Invest* 103:1141–1150
- 815 65. Tang DD, Gunst SJ (2004) The small GTPase
816 Cdc42 regulates actin polymerization and
817 tension development during contractile stim-
818 ulation of smooth muscle. *J Biol Chem*
819 279:51722–51728
- 820 66. Plutoni C, Bazellieres E, Le Borgne-Rochet-
821 M, Comunale F, Brugues A, Séveno M,
822 Planchon D, Thuault S, Morin N, Bodin S,
823 Trepas X, Gauthier-Rouvière C (2016)
824 P-cadherin promotes collective cell migration
825 via a Cdc42-mediated increase in mechanical
826 forces. *J Cell Biol* 212:199–217
- 827 67. Whitney DS, Peterson FC, Kittell AW, Egner
828 JM, Prehoda KE, Volkman BF (2016) Bind-
829 ing of crumbs to the Par-6 CRIB-PDZ mod-
830 ule is regulated by Cdc42. *Biochemistry*
831 55:1455–1461
- 832 68. Garrard SM, Capaldo CT, Gao L, Rosen MK,
833 Macara IG, Tomchick DR (2003) Structure
834 of Cdc42 in a complex with the GTPase-
835 binding domain of the cell polarity protein,
836 Par6. *EMBO J* 22:1125–1133
- 837 69. Su W-H, Mruk DD, Wong EWP, Lui W-Y,
838 Cheng CY (2012) Polarity protein complex
839 Scribble/Lgl/Dlg and epithelial cell barriers.
840 *Adv Exp Med Biol* 763:149–170
- 841 70. Lee S, Fan S, Makarova O, Straight S, Margolis
842 B (2002) A novel and conserved protein-
843 protein interaction domain of mammalian
844 Lin-2/CASK binds and recruits SAP97 to
845 the lateral surface of epithelia. *Mol Cell Biol*
846 22:1778–1791
- 847 71. Zhu Z-Q, Wang D, Xiang D, Yuan Y-X, Wang
848 Y (2014) Calcium/calmodulin-dependent
849 serine protein kinase is involved in exendin-
850 4-induced insulin secretion in INS-1 cells.
851 *Metabolism* 63:120–126
- 852 72. Jiang M, Qiu J, Zhang L, Lü D, Long M,
853 Chen L, Luo X (2016) Changes in tension
854 regulates proliferation and migration of fibro-
855 blasts by remodeling expression of ECM pro-
856 teins. *Exp Ther Med* 12:1542–1550
- 857 73. Porter AP, White GRM, Mack NA, Malliri A
858 (2019) The interaction between CASK and
859 the tumour suppressor Dlg1 regulates mitotic
860 spindle orientation in mammalian epithelia. *J*
861 *Cell Sci* 132:jcs230086
- 862 74. Hannezo E, Prost J, Joanny J-F (2014) The-
863 ory of epithelial sheet morphology in three
864 dimensions. *Proc Natl Acad Sci U S A*
865 111:27–32
- 866 75. Takeichi M (2014) Dynamic contacts: rear-
867 ranging adherens junctions to drive epithelial
868 remodelling. *Nat Rev Mol Cell Biol*
869 15:397–410
- 870 76. Becker S, Wandel E, Wobus M, Schneider R,
871 Amasheh S, Sittig D, Kerner C, Naumann R,
872 Hamann J, Aust G (2010) Overexpression of
873 CD97 in intestinal epithelial cells of trans-
874 genic mice attenuates colitis by strengthening
875 adherens junctions. *PLoS One* 5:e8507

- 876 77. Hilbig D, Sittig D, Hoffmann F, 933
877 Rothmund S, Warnt E, Quaas M, 934
878 Stürmer J, Seiler L, Liebscher I, Hoang NA, 935
879 Käs JA, Banks L, Aust G (2018) Mechano- 936
880 dependent phosphorylation of the 937
881 PDZ-binding Motif of CD97/ADGRE5 938
882 modulates cellular detachment. *Cell Rep* 939
883 24:1986–1995
- 884 78. Sivaramakrishnan S, Schneider JL, Sitikov A, 940
885 Goldman RD, Ridge KM (2009) Shear stress 941
886 induced reorganization of the keratin inter- 942
887 mediate filament network requires phosphor- 943
888 ylation by protein kinase C ζ . *Mol Biol Cell* 944
889 20:2755–2765
- 890 79. O'Neill AK, Gallegos LL, Justilien V, Garcia 945
891 EL, Leitges M, Fields A, Hall RA, Newton AC 946
892 (2011) PKC α interacts with its novel substrate 947
893 discs large homolog (DLG) 1 to promote 948
894 cellular migration. *J Biol Chem jbc.* 949
895 M111.294603
- 896 80. Kanzaki N, Ogita H, Komura H, Ozaki M, 950
897 Sakamoto Y, Majima T, Ijuin T, Takenawa T, 951
898 Takai Y (2008) Involvement of the nectin- 952
899 afadin complex in PDGF-induced cell sur- 953
900 vival. *J Cell Sci* 121:2008–2017
- 901 81. Sato T, Fujita N, Yamada A, Ooshio T, 954
902 Okamoto R, Irie K, Takai Y (2006) Regula- 955
903 tion of the assembly and adhesion activity of 956
904 E-cadherin by nectin and afadin for the for- 957
905 mation of adherens junctions in Madin-Darby 958
906 canine kidney cells. *J Biol Chem* 959
907 281:5288–5299
- 908 82. Ooshio T, Kobayashi R, Ikeda W, Miyata M, 960
909 Fukumoto Y, Matsuzawa N, Ogita H, Takai Y 961
910 (2010) Involvement of the interaction of afa- 962
911 din with ZO-1 in the formation of tight junc- 963
912 tions in Madin-Darby canine kidney cells. *J 964
913 Biol Chem* 285:5003–5012
- 914 83. Severson EA, Lee WY, Capaldo CT, Nusrat A, 965
915 Parkos CA (2009) Junctional adhesion mole- 966
916 cule A interacts with afadin and PDZ-GEF2 967
917 to activate Rap1A, regulate β 1 integrin levels, 968
918 and enhance cell migration. *Mol Biol Cell* 969
919 20:1916–1925
- 920 84. Asada M, Irie K, Morimoto K, Yamada A, 970
921 Ikeda W, Takeuchi M, Takai Y (2003) ADIP, 971
922 a novel afadin- and α -actinin-binding protein 972
923 localized at cell-cell adherens junctions. *J Biol 973
924 Chem* 278:4103–4111
- 925 85. Mandai K, Nakanishi H, Satoh A, Obaishi H, 974
926 Wada M, Nishioka H, Itoh M, Mizoguchi A, 975
927 Aoki T, Fujimoto T, Matsuda Y, Tsukita S, 976
928 Takai Y (1997) Afadin: a novel actin 977
929 filament-binding protein with one PDZ 978
930 domain localized at cadherin-based cell-to- 979
931 cell adherens junction. *J Cell Biol* 980
932 139:517–528
86. Mandai K, Nakanishi H, Satoh A, 933
Takahashi K, Satoh K, Nishioka H, 934
Mizoguchi A, Takai Y (1999) Ponsin/ 935
SH3P12: an I-afadin- and vinculin-binding 936
protein localized at cell-cell and cell-matrix 937
adherens junctions. *J Cell Biol* 938
144:1001–1018 939
87. Ooshio T, Irie K, Morimoto K, Fukuhara A, 940
Imai T, Takai Y (2004) Involvement of 941
LMO7 in the association of two cell-cell adhe- 942
sion molecules, nectin and E-cadherin, 943
through afadin and alpha-actinin in epithelial 944
cells. *J Biol Chem* 279:31365–31373 945
88. Takai Y, Nakanishi H (2003) Nectin and afa- 946
din: novel organizers of intercellular junc- 947
tions. *J Cell Sci* 116:17–27 948
89. Monteiro AC, Sumagin R, Rankin CR, 949
Leoni G, Mina MJ, Reiter DM, Stehle T, 950
Dermody TS, Schaefer SA, Hall RA, 951
Nusrat A, Parkos CA (2013) JAM-A associ- 952
ates with ZO-2, afadin, and PDZ-GEF1 to 953
activate Rap2c and regulate epithelial barrier 954
function. *Mol Biol Cell* 24:2849–2860 955
90. Fanning AS, Van Itallie CM, Anderson JM 956
(2012) Zonula occludens-1 and -2 regulate 957
apical cell structure and the zonula adherens 958
cytoskeleton in polarized epithelia. *Mol Biol 959
Cell* 23:577–590 960
91. Miyata M, Rikitake Y, Takahashi M, 961
Nagamatsu Y, Yamauchi Y, Ogita H, 962
Hirata K, Takai Y (2009) Regulation by afadin 963
of cyclical activation and inactivation of Rap1, 964
Rac1, and RhoA small G proteins at leading 965
edges of moving NIH3T3 cells. *J Biol Chem* 966
284:24595–24609 967
92. Choi W, Acharya BR, Peyret G, Fardin M-A, 968
Mège R-M, Ladoux B, Yap AS, Fanning AS, 969
Peifer M (2016) Remodeling the zonula 970
adherens in response to tension and the role 971
of afadin in this response. *J Cell Biol* 972
213:243–260 973
93. Ebnet K, Schulz CU, Zu Brickwedde M-KM, 974
Pendl GG, Vestweber D (2000) Junctional 975
adhesion molecule interacts with the PDZ 976
domain-containing proteins AF-6 and ZO-1. 977
J Biol Chem 275:27979–27988 978
94. Scott DW, Tolbert CE, Burrige K (2016) 979
Tension on JAM-A activates RhoA via 980
GEF-H1 and p115 RhoGEF. *Mol Biol Cell* 981
27:1420–1430 982
95. Priya R, Yap AS (2015) Active tension: the 983
role of cadherin adhesion and signaling in 984
generating junctional contractility. *Curr Top 985
Dev Biol* 112:65–102 986
96. Iden S, Misselwitz S, Peddibhotla SSD, 987
Tuncay H, Rehder D, Gerke V, Robenek H, 988
Suzuki A, Ebnet K (2012) aPKC 989

990 phosphorylates JAM-A at Ser285 to promote
 991 cell contact maturation and tight junction for-
 992 mation. *J Cell Biol* 196:623–639

993 97. Albertson R (2004) Scribble protein domain
 994 mapping reveals a multistep localization
 995 mechanism and domains necessary for estab-
 996 lishing cortical polarity. *J Cell Sci*
 997 117:6061–6070

998 98. Zeitler J, Hsu CP, Dionne H, Bilder D (2004)
 999 Domains controlling cell polarity and prolifer-
 1000 eration in the *Drosophila* tumor suppressor
 1001 Scribble. *J Cell Biol* 167:1137–1146

1002 99. Audebert S, Navarro C, Nourry C, Chasserot-
 1003 Golaz S, Lécine P, Bellaïche Y, Dupont J-L,
 1004 Premont RT, Sempéré C, Strub J-M, Van
 1005 Dorsaelaer A, Vitale N, Borg J-P (2004)
 1006 Mammalian Scribble forms a tight complex
 1007 with the betaPIX exchange factor. *Curr Biol*
 1008 14:987–995

1009 100. Kallay LM, McNickle A, Brennwald PJ, Hub-
 1010 bard AL, Braiterman LT (2006) Scribble
 1011 associates with two polarity proteins, *lgl2*
 1012 and *vangl2*, via distinct molecular domains. *J*
 1013 *Cell Biochem* 99:647–664

1014 101. Wang X, Dong B, Zhang K, Ji Z, Cheng C,
 1015 Zhao H, Sheng Y, Li X, Fan L, Xue W, Gao
 1016 W-Q, Zhu HH (2018) E-cadherin bridges
 1017 cell polarity and spindle orientation to ensure
 1018 prostate epithelial integrity and prevent carci-
 1019 nogenesis in vivo. *PLoS Genet* 14:e1007609

1020 102. Osmani N, Vitale N, Borg J-P, Etienne-Man-
 1021 neville S (2006) Scrib controls Cdc42 locali-
 1022 zation and activity to promote cell
 1023 polarization during astrocyte migration.
 1024 *Curr Biol* 16:2395–2405

1025 103. Lohia M, Qin Y, Macara IG (2012) The
 1026 Scribble polarity protein stabilizes E-cad-
 1027 herin/p120-catenin binding and blocks
 1028 retrieval of E-cadherin to the golgi. *PLoS*
 1029 *One* 7:e51130

1030 104. le Duc Q, Shi Q, Blonk I, Sonnenberg A,
 1031 Wang N, Leckband D, de Rooij J (2010)
 1032 Vinculin potentiates E-cadherin mechanosen-
 1033 soring and is recruited to actin-anchored sites
 1034 within adherens junctions in a myosin
 1035 II-dependent manner. *J Cell Biol*
 1036 189:1107–1115

1037 105. Borghi N, Sorokina M, Shcherbakova OG,
 1038 Weis WI, Pruitt BL, Nelson WJ, Dunn AR
 1039 (2012) E-cadherin is under constitutive
 1040 actomyosin-generated tension that is
 1041 increased at cell–cell contacts upon externally
 1042 applied stretch. *Proc Natl Acad Sci U S A*
 1043 109:12568–12573

1044 106. Bazellières E, Conte V, Elosegui-Artola A,
 1045 Serra-Picamal X, Bintanel-Morcillo M, Roca-
 1046 Cusachs P, Muñoz JJ, Sales-Pardo M,
 Guimerà R, Trepast X (2015) Control of cell-
 cell forces and collective cell dynamics by the
 intercellular adhesion. *Nat Cell Biol*
 17:409–420

107. Wu SK, Lagendijk AK, Hogan BM, Gomez
 GA, Yap AS (2015) Active contractility at
 E-cadherin junctions and its implications for
 cell extrusion in cancer. *Cell Cycle*
 14:315–322

108. Engl W, Arasi B, Yap LL, Thiery JP, Viasnoff
 V (2014) Actin dynamics modulate mechan-
 osensitive immobilization of E-cadherin at
 adherens junctions. *Nat Cell Biol*
 16:584–591

109. Wu SK, Gomez GA, Michael M, Verma S,
 Cox HL, Lefevre JG, Parton RG, Hamilton
 NA, Neufeld Z, Yap AS (2014) Cortical
 F-actin stabilization generates apical-lateral
 patterns of junctional contractility that inte-
 grate cells into epithelia. *Nat Cell Biol*
 16:167–178

110. Lugo-Martínez V-H, Petit CS, Fouquet S, Le
 Beyec J, Chambaz J, Pinçon-Raymond M,
 Cardot P, Thenet S (2009) Epidermal growth
 factor receptor is involved in enterocyte anoikis
 through the dismantling of E-cadherin-
 mediated junctions. *Am J Physiol Gastrointest*
Liver Physiol 296:G235–G244

111. Choi J, Troyanovsky RB, Indra I, Mitchell BJ,
 Troyanovsky SM (2019) Scribble, Erbin, and
 Lano redundantly regulate epithelial polarity
 and apical adhesion complex. *J Cell Biol*
 218:2277–2293

112. Yoshihara K, Ikenouchi J, Izumi Y, Akashi M,
 Tsukita S, Furuse M (2011) Phosphorylation
 state regulates the localization of Scribble at
 adherens junctions and its association with E-
 cadherin-catenin complexes. *Exp Cell Res*
 317:413–422

113. Meng W, Mushika Y, Ichii T, Takeichi M
 (2008) Anchorage of microtubule minus
 ends to adherens junctions regulates epithelial
 cell-cell contacts. *Cell* 135:948–959

114. Priya R, Yap AS, Gomez GA (2013)
 E-cadherin supports steady-state Rho signal-
 ing at the epithelial zonula adherens. *Differ-*
entiation 86:133–140

115. Truong Quang B-A, Mani M, Markova O,
 Lecuit T, Lenne P-F (2013) Principles of
 E-cadherin supramolecular organization
 in vivo. *Curr Biol* 23:2197–2207

116. Hart KC, Tan J, Siemers KA, Sim JY, Pruitt
 BL, Nelson WJ, Gloerich M (2017)
 E-cadherin and LGN align epithelial cell divi-
 sions with tissue tension independently of cell
 shape. *Proc Natl Acad Sci U S A* 114(29):

- 1103 E5845–E5853. [https://doi.org/10.1073/](https://doi.org/10.1073/pnas.1701703114)
1104 [pnas.1701703114](https://doi.org/10.1073/pnas.1701703114)
- 1105 117. Gloerich M, Bianchini JM, Siemers KA,
1106 Cohen DJ, Nelson WJ (2017) Cell division
1107 orientation is coupled to cell–cell adhesion by
1108 the E-cadherin/LGN complex. *Nat Commun* 8:1–11
1109
- 1110 118. del Rio A, Perez-Jimenez R, Liu R, Roca-
1111 Cusachs P, Fernandez JM, Sheetz MP
1112 (2009) Stretching single talin rod molecules
1113 activates vinculin binding. *Science*
1114 323:638–641
- 1115 119. Mitrossilis D, Fouchard J, Pereira D, Postic F,
1116 Richert A, Saint-Jean M, Asnacios A (2010)
1117 Real-time single-cell response to stiffness.
1118 *PNAS* 107:16518–16523
- 1119 120. Choquet D, Felsenfeld DP, Sheetz MP
1120 (1997) Extracellular matrix rigidity causes
1121 strengthening of integrin–cytoskeleton lin-
1122 kages. *Cell* 88:39–48
121. Stehbens SJ, Paszek M, Pemble H, 1123
Ettinger A, Gierke S, Wittmann T (2014) 1124
CLASPs link focal-adhesion-associated 1125
microtubule capture to localized exocytosis 1126
and adhesion site turnover. *Nat Cell Biol* 1127
16:561–573 1128
122. Sackmann E, Smith A-S (2014) Physics of cell 1129
adhesion: some lessons from cell-mimetic sys- 1130
tems. *Soft Matter* 10:1644–1659 1131
123. Riveline D, Zamir E, Balaban NQ, Schwarz 1132
US, Ishizaki T, Narumiya S, Kam Z, Geiger B, 1133
Bershadsky AD (2001) Focal contacts as 1134
mechanosensors externally applied local 1135
mechanical force induces growth of focal con- 1136
tacts by an mdial-dependent and rock- 1137
independent mechanism. *J Cell Biol* 1138
153:1175–1186 1139
124. Kaverina I, Krylyshkina O, Beningo K, 1140
Anderson K, Wang Y-L, Small JV (2002) Ten- 1141
sile stress stimulates microtubule outgrowth 1142
in living cells. *J Cell Sci* 115:2283–2291 1143