



HAL
open science

A Short Note On Some Novel Applications of Semi Module Homomorphisms

Mohammad Abobala, Mikail Bal, Malath Aswad

► **To cite this version:**

Mohammad Abobala, Mikail Bal, Malath Aswad. A Short Note On Some Novel Applications of Semi Module Homomorphisms. International Journal of Neutrosophic Science, 2022, pp.08 - 13. 10.54216/ijns.180101 . hal-03559359

HAL Id: hal-03559359

<https://hal.science/hal-03559359v1>

Submitted on 18 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A Short Note On Some Novel Applications of Semi Module Homomorphisms

Mohammad Abobala, Tishreen University, Department of Mathematics, Syria

Mikail Bal, Gaziantep University, Department of Mathematics, Turkey

Malath Aswad, Albaath University, Department of Mathematics, Syria

Co-: mohammadabobala777@gmail.com

Abstract

In this paper, we study the algebraic relationships between n -refined neutrosophic modules by using semi-module homomorphisms.

On the other hand, this work shows the relationship between neutrosophic geometrical AH-isometry and semi-module isomorphisms.

Keywords: AH-isometry, Semi homomorphism, semi isomorphism, n -refined neutrosophic module

1.Introduction

Neutrosophy is a new kind of philosophy founded by Smarandache [1] to study the uncertainty and the lack of information in many areas of science and life.

The indeterminacy element I and its refinements, were useful in algebra where we find some generalizations of classical algebraic structures such as neutrosophic modules, n -refined neutrosophic matrices [2-5].

The concept of semi-modules homomorphism /isomorphism was presented in [6] as a tool to study the relations between modules which are defined over different rings.

In this work, we show a new application of semi homomorphisms in the study of n -refined neutrosophic modules. Also, we present a novel application in neutrosophic Euclidean geometry [9] based on semi-isomorphisms, where we show that algebraic isometries used in the study of neutrosophic Euclidean geometrical shapes can be considered as semi-module isomorphisms.

Main discussion

First of all, we show that the concept of semi-homomorphism (semi-isomorphism) is essential in the study of neutrosophic Euclidean geometry.

Neutrosophic Euclidean geometry was built over the idea of AH-isometry, where the AH-isometry is a function that preserves distances between neutrosophic points in $R(I) \times R(I)$.

Definition 1.

(a). Let $R(I) \times R(I)$ be the neutrosophic plane with two N-dimensions, then.

$$g: R(I) \times R(I) \rightarrow R^2 \times R^2$$

$$g(a + bI, c + dI) = ((a, a + b), (c, c + d))$$

is called the two dimensional AH-isometry.

(b). the function $T: R(I) \rightarrow R \times R; T(a + bI) = (a, a + b)$ is called the one dimensional AH-isometry.

Theorem 2

g, T preserve operations and distances.

In the following, we prove that the two dimensional AH-isometry is a semi-module isomorphism.

Definition 3:

Let $(M, +, \cdot)$ be a module over the ring R , then $(M(I), +, \cdot)$ is called a weak neutrosophic module over the ring R , and it is called a strong neutrosophic module if it is a module over the neutrosophic ring $R(I)$.

Elements of $M(I)$ have the form $x + yI; x, y \in M$, i.e $M(I)$ can be written as $M(I) = M + MI$.

Definition 4:

- (a) Let $(M, +, \cdot)$ be a module over the ring R , we say that $M_n(I) = M + MI_1 + \dots + MI_n = \{x_0 + x_1I_1 + \dots + x_nI_n; x_i \in M\}$ is a weak n-refined neutrosophic module over the ring R . Elements of $M_n(I)$ are called n-refined neutrosophic vectors, elements of R are called scalars.
- (b) If we take scalars from the n-refined neutrosophic ring $R_n(I)$, we say that $M_n(I)$ is a strong n-refined neutrosophic module over the n-refined neutrosophic ring $M_n(I)$. Elements of $M_n(I)$ are called n-refined neutrosophic scalars.

Definition 5:

Let M be a module over a ring R , N be a module over a ring T , $\varphi: M \rightarrow N$ be a well defined map, we say that φ is an f -semi module homomorphism if and only if the following conditions are true:

- (a) $\varphi(x + y) = \varphi(x) + \varphi(y)$ for all $x, y \in M$.
- (b) There is a ring homomorphism $f: R \rightarrow T$ such $\varphi(r \cdot x) = f(r) \cdot \varphi(x)$ for all $r \in R, x \in M$.

Example 6:

Let M be a module over a ring R , $M(I)$ be the corresponding strong neutrosophic module over $R(I)$, $M(I_1, I_2)$ be the corresponding strong refined neutrosophic module over $R(I_1, I_2)$. Then

$\varphi: M(I_1, I_2) \rightarrow M(I)$; $\varphi(a, bI_1, cI_2) = a + (b + c)I$ is a semi homomorphism.

Theorem 7.

Let $g: R(I) \times R(I) \rightarrow R^2 \times R^2$ be the two dimensional AH-isometry, and $T: R(I) \rightarrow R \times R$ be the one dimensional AH-isometry, then, g is a semi-module isomorphism.

Proof.

we have $R(I) \times R(I)$ is a module over the ring $R(I)$, and $R^2 \times R^2$ is a module over the ring $R \times R$.

According to [14], T is a ring isomorphism.

On the other hand, g preserves addition between $R(I) \times R(I)$ and $R^2 \times R^2$, and g is a bijection.

Now, we prove that g has the following property.

$$g[(m + nI). (a + bI, c + dI)] = T(m + nI)g(a + bI, c + dI)$$

$$l_1 = [(m + nI). (a + bI, c + dI)] = g[(m + nI)(a + bI), (m + nI)(c + dI)]$$

$$= g[(ma + I(ab + na + nb)), (mc + I(amd + nc + nd))]$$

$$= ((m, m + n). (a, a + b), (m, m + n). (c, c + d)) = (m, m + n)((a, a + b), (c, c + d))$$

$$= T(m + nI)g(a + bI, c + dI).$$

Thus g is a semi isomorphism.

Result 8.

According to the previous theorem, the neutrosophic Euclidean geometry can be understood as a semi isomorphic image of a neutrosophic module.

Now, we study semi homomorphisms between n -refined neutrosophic modules.

Theorem 9

Let R be a ring, $R_n(I)$ be the corresponding n -refined neutrosophic ring.

Let M be a module over R , $M_n(I)$ be the corresponding n -refined neutrosophic module over $R_n(I)$, then.

(a). $M_{n-1}(I)$ is a semi homomorphic image of $M_n(I)$.

(b). M is a semi homomorphic image of $M_n(I)$.

Proof.

(a). $g: R_n(I) \rightarrow R_{n-1}(I)$; $g(a_0 + a_1I_1 + \dots + a_nI_n) = a_0 + a_1I_1 + \dots + (a_{n-1} + a_n)I_{n-1}$

is a ring homomorphism.

We define $f: M_n(I) \rightarrow M_{n-1}(I); f(x_0 + x_1I_1 + \dots + x_nI_n) = x_0 + x_1I_1 + \dots + (x_{n-1} + x_n)I_{n-1}$.

It is clear that f is well defined and preserves addition.

Now, Let's compute:

$$f[(a_0 + a_1I_1 + \dots + a_nI_n). (x_0 + x_1I_1 + \dots + x_nI_n)] = f \left[\sum_{i,j=0}^n a_i x_j I_i I_j \right]$$

The coefficient of I_n is $a_0x_n + a_nx_0 + a_nx_n$.

The coefficient of I_{n-1} is $a_0x_{n-1} + a_{n-1}x_0 + a_nx_{n-1} + a_{n-1}x_n + a_{n-1}x_{n-1}$.

This implies that

$$f \left[\sum_{i,j=0}^n a_i x_j I_i I_j \right] = f[A + I_{n-1}(a_0x_{n-1} + a_{n-1}x_0 + a_nx_{n-1} + a_{n-1}x_n + a_{n-1}x_{n-1}) + I_n(a_0x_n + a_nx_0 + a_nx_n)]$$

$$= A + I_{n-1}(a_0x_{n-1} + a_{n-1}x_0 + a_nx_{n-1} + a_{n-1}x_n + a_{n-1}x_{n-1}) + I_n(a_0x_n + a_nx_0 + a_nx_n)$$

Now, we compute.

$$g(a_0 + a_1I_1 + \dots + a_nI_n). f(x_0 + x_1I_1 + \dots + x_nI_n) = (a_0 + \dots + a_{n-2}I_{n-2} + I_{n-1}(a_n + a_{n-1})). (x_0 + \dots + x_{n-2}I_{n-2} + I_{n-1}(x_n + x_{n-1}))$$

Where $B = a_0x_{n-1} + a_0x_n + a_nx_0 + a_{n-1}x_0 + a_nx_{n-1} + a_nx_n + a_{n-1}x_{n-1} + a_{n-1}x_n$

Which is exactly equal to the coefficient I_{n-1} in $f[(\sum_{i=0}^n a_i I_i. \sum_{i=0}^n x_i I_i)]$.

This means that $f[(\sum_{i=0}^n a_i I_i. \sum_{i=0}^n x_i I_i)] = g(\sum_{i=0}^n a_i I_i). f(\sum_{i=0}^n x_i I_i)$, hence f is a semi module homomorphism.

(b). since $M_{n-i}(I)$ is a semi homomorphic image of $M_{n-i+1}(I)$, we get the following sequence:

$$M_n(I) \rightarrow M_{n-1}(I) \rightarrow \dots \rightarrow M_2(I) \rightarrow M_1(I) \rightarrow M(I)$$

Thus M is a semi homomorphic image of $M_n(I)$.

Example 10

Let $R = Z, M = Z_3$. M is a module over R .

Let $M_3(I) = \{a_0 + a_1I_1 + a_2I_2 + a_3I_3; a_i \in Z_3\}$ be the corresponding 3-refined neutrosophic module over $R_3(I)$

$Z_3(I) = \{b_0 + b_1I_1 + b_2I_2 + b_3I_3; b_i \in Z\}$.

$$(a). g: Z_3(I) \rightarrow Z_2(I); g(a_0 + a_1I_1 + a_2I_2 + a_3I_3) = a_0 + a_1I_1 + (a_2 + a_3)I_2$$

Is a ring homomorphism.

$$(b). f: M_3(I) \rightarrow M_2(I); f(x_0 + x_1I_1 + x_2I_2 + x_3I_3) = x_0 + x_1I_1 + (x_2 + x_3)I_2$$

is a semi homomorphism, that is because:

f preserves addition clearly.

$$\begin{aligned} & f[(a_0 + a_1I_1 + a_2I_2 + a_3I_3) \cdot (x_0 + x_1I_1 + x_2I_2 + x_3I_3)] \\ &= f[a_0x_0 + I_1(a_0x_1 + a_1x_0 + a_1x_1 + a_1x_2 + a_1x_3 + a_2x_1 + a_3x_1) + I_2(a_0x_2 + a_2x_2 + a_2x_3 + a_3x_2) \\ &\quad + I_3(a_0x_3 + a_3x_0 + a_3x_3)] + a_2x_0 \\ &= a_0x_0 + I_1(a_0x_1 + a_1x_0 + a_1x_1 + a_1x_2 + a_1x_3 + a_2x_1 + a_3x_1) \\ &\quad + I_2(a_0x_2 + a_2x_2 + a_2x_3 + a_3x_2 + a_2x_0 + a_0x_2 + a_3x_0 + a_3x_3) \end{aligned}$$

On the other hand.

$$\begin{aligned} & g(a_0 + a_1I_1 + a_2I_2 + a_3I_3) \cdot f(x_0 + x_1I_1 + x_2I_2 + x_3I_3) = (a_0 + a_1I_1 + I_2(a_3 + a_2))(x_0 + x_1I_1 + I_2(x_3 + x_2)) \\ &= a_0x_0 + I_1(a_0x_1 + a_1x_0 + a_1x_1 + a_1x_2 + a_1x_3 + a_2x_1 + a_3x_1) \\ &\quad + I_2(a_2x_0 + a_3x_0 + a_2x_2 + a_2x_3 + a_3x_2 + a_0x_2 + a_3x_3 + a_0x_3) \\ &= f[(a_0 + a_1I_1 + a_2I_2 + a_3I_3) \cdot (x_0 + x_1I_1 + x_2I_2 + x_3I_3)] \end{aligned}$$

$$(c). \ker f = \{a_2I_2 - a_2I_3 \in M_3(I)\} = \{a_2(I_2 - I_3); a_2 \in M\}$$

$$M_3(I)/\ker f \cong_s f(M_3(I)) = M_2(I).$$

The symbol \cong_s means semi isomorphic property.

Conclusion

In this paper, we have presented some novel applications of semi module homomorphisms/isomorphisms, where we proved that every module is a semi homomorphic image of it corresponding n-refined neutrosophic module.

Also, we have shown that the AH-isometry used in the theory of neutrosophic Euclidean geometry is a semi-module isomorphism.

References

- [1] Smarandache, F., " A Unifying Field in Logics: Neutrosophic Logic, Neutrosophy, Neutrosophic Set, Neutrosophic Probability", American Research Press. Rehoboth, 2003.
- [2] Abobala, M., Hatip, A., and Bal, M., " A Review On Recent Advantages In Algebraic Theory Of Neutrosophic Matrices", IJNS, Vol.17, 2021.
- [3] Agboola, A.A.A., "On Refined Neutrosophic Algebraic Structures", NSS, Vol.10, pp. 99-101. 2015.
- [4] Abobala, M., " A Study Of Nil Ideals and Kothe's Conjecture In Neutrosophic Rings", International Journal of Mathematics and Mathematical Sciences", Hindawi, 2021

- [5] Olgun, N., Hatip, A., Bal. M., and Abobala, M., "A novel approach to necessary and sufficient conditions for the diagonalization of refined neutrosophic matrices", IJNS, 2021.
- [6] Abobala, M., " Semi Homomorphisms and Algebraic Relations Between Strong Refined Neutrosophic Modules and Strong Neutrosophic Modules", Neutrosophic Sets and Systems, Vol. 39, 2021.
- [7] Abobala, M., "On Some Algebraic Properties of n-Refined Neutrosophic Elements and n-Refined Neutrosophic Linear Equations", Mathematical Problems in Engineering, Hindawi, 2021
- [8] Abobala, M., "Neutrosophic Real Inner Product Spaces", NSS, Vol. 43, 2021.
- [9] Abobala, M., and Hatip, A., "An Algebraic Approach To Neutrosophic Euclidean Geometry", NSS, Vol. 43, 2021.
- [10] Abobala, M., "On The Characterization of Maximal and Minimal Ideals In Several Neutrosophic Rings", NSS , Vol. 45, 2021.