



HAL
open science

G protein-coupled receptor-effector macromolecular membrane assemblies (GEMMAs)

Sergi Ferré, Francisco Ciruela, Carmen W Dessauer, Javier González-Maeso, Terence E Hébert, Ralf Jockers, Diomedes E Logothetis, Leonardo Pardo

► To cite this version:

Sergi Ferré, Francisco Ciruela, Carmen W Dessauer, Javier González-Maeso, Terence E Hébert, et al.. G protein-coupled receptor-effector macromolecular membrane assemblies (GEMMAs). *Pharmacology & Therapeutics*, 2021, pp.107977. <10.1016/j.pharmthera.2021.107977>. <hal-03560153>

HAL Id: hal-03560153

<https://hal.science/hal-03560153v1>

Submitted on 22 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License



Contents lists available at ScienceDirect

Pharmacology & Therapeutics

journal homepage: www.elsevier.com/locate/pharmthera

G protein-coupled receptor-effector macromolecular membrane assemblies (GEMMAs)

Sergi Ferré ^{a,*}, Francisco Ciruela ^b, Carmen W. Dessauer ^c, Javier González-Maeso ^d, Terence E. Hébert ^e, Ralf Jockers ^f, Diomedes E. Logothetis ^g, Leonardo Pardo ^h

^a Integrative Neurobiology Section, National Institute on Drug Addiction, Intramural Research Program, NIH, DHHS, Baltimore, MD, USA

^b Department of Pathology and Experimental Therapeutics, School of Medicine and Health Sciences, Institute of Neurosciences, IDIBELL, University of Barcelona, L'Hospitalet de Llobregat, Spain

^c Department of Integrative Biology and Pharmacology, McGovern Medical School, University of Texas Health Science Center at Houston, Houston, TX, USA

^d Department of Physiology and Biophysics, School of Medicine, Virginia Commonwealth University, Richmond, VA, USA

^e Department of Pharmacology and Therapeutics, McGill University, Montréal, Québec, Canada

^f Université de Paris, Institut Cochin, INSERM, CNRS, Paris, France

^g Laboratory of Electrophysiology, Departments of Pharmaceutical Sciences, Chemistry and Chemical Biology and Center for Drug Discovery, School of Pharmacy at the Bouvé College of Health Sciences and College of Science, Northeastern University, Boston, MA, USA

^h Laboratory of Computational Medicine, Biostatistics Unit, Faculty of Medicine, Autonomous University of Barcelona, Bellaterra, Spain

ARTICLE INFO

Available online xxxx

Editor: P. Holzer

Keywords:

G protein-coupled receptors
G protein subunits
Plasma membrane effector
GPCR oligomerization
GPCR allosterism

ABSTRACT

G protein-coupled receptors (GPCRs) are the largest group of receptors involved in cellular signaling across the plasma membrane and a major class of drug targets. The canonical model for GPCR signaling involves three components – the GPCR, a heterotrimeric G protein and a proximal plasma membrane effector – that have been generally thought to be freely mobile molecules able to interact by ‘collision coupling’. Here, we synthesize evidence that supports the existence of GPCR–effector macromolecular membrane assemblies (GEMMAs) comprised of specific GPCRs, G proteins, plasma membrane effector molecules and other associated transmembrane proteins that are pre-assembled prior to receptor activation by agonists, which then leads to subsequent rearrangement of the GEMMA components. The GEMMA concept offers an alternative and complementary model to the canonical collision-coupling model, allowing more efficient interactions between specific signaling components, as well as the integration of the concept of GPCR oligomerization as well as GPCR interactions with orphan receptors, truncated GPCRs and other membrane-localized GPCR-associated proteins. Collision-coupling and pre-assembled mechanisms are not exclusive and likely both operate in the cell, providing a spectrum of signaling modalities which explains the differential properties of a multitude of GPCRs in their different cellular environments. Here, we explore the unique pharmacological characteristics of individual GEMMAs, which could provide new opportunities to therapeutically modulate GPCR signaling.

Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations¹

¹GPCR nomenclature used in the current review follows the guidelines of the International Union of Basic and Clinical Pharmacology Committee on Receptor Nomenclature and Drug Classification (<https://www.guidetopharmacology.org/nomenclature.jsp>) and adds an ‘R’ at the end as an abbreviation of ‘receptor’: AC, adenylate cyclase; AKAP, A-kinase anchoring protein; AMPAR, ionotropic glutamate AMPA receptors; BiBC or BiFC, bimolecular bioluminescence or fluorescence complementation; BRET or FRET, bioluminescence or fluorescence resonance energy transfer; CT or NT, carboxy terminus or amino terminus; ER, endoplasmic reticulum; FRAP, fluorescence recovery after photobleaching; GEMMA, GPCR-effector macromolecular membrane assembly; GPCR, G protein-coupled receptor; Kir, inwardly rectifying potassium channel; PDZ, PSD-95/Discs large/Zona occludens-1; PKA, protein kinase A; PM-effector, plasma membrane-effector; PIP₂, phosphatidylinositol 4,5-bisphosphate; PLC, phospholipase C; RAMP, Receptor activity-modifying protein; Rluc, *Renilla* luciferase; TM, transmembrane; YFP, yellow fluorescence protein.

* Corresponding author at: Integrative Neurobiology Section, National Institute on Drug Abuse, IRP, NIH, DHHS, Triad Technology Building, 333 Cassell Drive, Baltimore, MD 21224, USA. E-mail address: sferre@intra.nida.nih.gov (S. Ferré).

¹ GPCR nomenclature used in the current review follows the guidelines of the International Union of Basic and Clinical Pharmacology Committee on Receptor Nomenclature and Drug Classification (<https://www.guidetopharmacology.org/nomenclature.jsp>) and adds an ‘R’ at the end as an abbreviation of ‘receptor’.

<https://doi.org/10.1016/j.pharmthera.2021.107977>

0163-7258/Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: S. Ferré, F. Ciruela, C.W. Dessauer, et al., G protein-coupled receptor-effector macromolecular membrane assemblies (GEMMAs), *Pharmacology & Therapeutics*, <https://doi.org/10.1016/j.pharmthera.2021.107977>

Contents

1. Introduction: the GEMMA concept	0
2. Evidence for pre-assembly and ligand-induced rearrangement of specific GPCR, G proteins and effector complexes	0
3. Compartmentalization of signaling molecules at the plasma membrane by membrane nanodomains and scaffold proteins	0
4. GPCR oligomerization	0
5. Targeting specific GEMMAs	0
6. Concluding remarks and future directions	0
Declaration of Competing Interest	0
Acknowledgements	0
References.	0

1. Introduction: the GEMMA concept

G protein-coupled receptors (GPCRs) form the largest group of receptors involved in cellular signaling across the plasma membrane. Heterotrimeric ($G\alpha\beta\gamma$) guanine nucleotide-binding proteins (G proteins) are the initial transducers that convey information from agonist-occupied GPCRs to a variety of effector proteins, many of them localized to the plasma membrane (PM-effectors). The canonical model thus involves three components, the GPCR, a heterotrimeric G protein and a proximal PM-effector, thought initially to be freely mobile molecules able to interact by 'collision coupling'.

In 1978, Tolkovsky and Levitzki articulated the notion of the collision coupling for GPCR signaling from studies on β -adrenergic receptor (β AR)-mediated adenylyl cyclase (AC) activation in turkey erythrocytes (Tolkovsky & Levitzki, 1978a). That same year, they reported that collision coupling could not explain results obtained using the same system on activation of AC by adenosine receptors. Instead, they observed a first-order process of AC activation, that could be better explained by the existence of pre-coupled GPCR-AC complexes (Braun & Levitzki, 1979; Tolkovsky & Levitzki, 1978b). Collision coupling was consistent with the 'fluid mosaic' model of the plasma membrane proposed by Singer and Nicholson in the early 1970s, which described the lipid bilayer as an isotropic milieu, allowing membrane-embedded proteins to diffuse and interact with each other by random collision (Singer & Nicolson, 1972). Initial support for collision coupling in GPCR signaling came from early studies on rhodopsin and its G protein partner transducin (Gt) in rod photoreceptor cells and on purified β AR, G proteins and AC reconstituted in phospholipid vesicles (Levitzki & Klein, 2002; Neubig, 1994).

However, the cellular models used in these influential studies were not generally representative of the common cellular environment of GPCRs and their interacting membrane signaling molecules. The outer segment disks of photoreceptors have a unique lipid composition that provides rhodopsin and Gt with much greater lateral mobility than that of most mammalian membrane proteins. Similarly, receptors and G proteins reconstituted in lipid vesicles are likely to be relatively mobile without the constraints on lateral motion imposed in cellular plasma membranes, such as interactions with the cytoskeleton and membrane nanodomains with different protein and lipid compositions, such as 'lipid rafts' (see below). Furthermore, the collision coupling model is less compatible with the existence, in the same cell, of a large variety of different GPCRs, heterotrimeric G proteins and PM-effectors (Fig. 1). In this context, the binding of an agonist must promote a series of specific and sequential intermolecular interactions between the activated GPCR, one or more heterotrimeric G proteins comprised of particular $G\alpha$, $G\beta$ and $G\gamma$ subunit combinations and, in many cases, multiple PM-effectors, including AC and phospholipase C (PLC) subtypes and a number of GPCR-modulated ion channels (Cabrera-Vera et al., 2002).

Here, we establish the concept of a GPCR-effector macromolecular membrane assembly or GEMMA. A GEMMA is defined as a pre-assembled signaling complex composed of particular combinations of GPCRs, G proteins, effectors and other associated transmembrane (TM)

proteins localized to the plasma membrane. Further, GEMMAs possess emergent functional and pharmacological characteristics making them potentially unique drug targets. Inherent in the definition of GEMMA is that the interactions between different components are membrane-delimited, which means that all the components are either intrinsic membrane or membrane-associated proteins, with no diffusible cytosolic intermediates between core components.

We must distinguish between the three primary components of membrane-delimited GPCR-mediated signaling pathways (GPCRs, G proteins and PM-effectors), from additional components that modulate or scaffold these core components. Among others, additional elements include G protein-coupled receptor kinases (GRKs) and arrestins, which are primarily recruited to the plasma membrane upon GPCR activation and may not directly interact with PM-effectors (Gurevich & Gurevich, 2019; Gurevich, Tesmer, Mushegian, & Gurevich, 2012; Homan, Glukhova, & Tesmer, 2013).

This notion revises the classical model of sequential ligand-induced association-dissociation of GPCR, G protein subunits and PM-effector, and posits that ligands can induce rearrangements within elements of these pre-assembled macromolecular complexes. Metaphorically, we can imagine that GPCRs, G proteins and PM-effectors act as 'clock gears', instead of 'billiard balls', as described by collision coupling. It can nevertheless be surmised that both alternative modes of interactions coexist in the same cell, particularly for GPCR-G protein interactions.

Within the GEMMA concept, it is necessary to address the impact of GPCR oligomerization. GEMMAs can include a number of identical or different GPCRs as well as orphan GPCRs, truncated GPCRs and other associated TM proteins, to generate unique macromolecular complexes with distinct functional and pharmacological properties. The homomeric nature of class C GPCRs is not a matter of dispute. Their obligate dimeric nature depends on an inter-protomer disulfide bridge formed via a conserved cysteine in their characteristic large binding domain (Ellaithy, Gonzalez-Maeso, Logothetis, & Levitz, 2020). The cryo-EM structures of several metabotropic glutamate receptor (mGluR) homodimers have been recently reported. These include the inactive and active conformations of mGlu₅R and mGlu₂R homodimers (Du et al., 2021; Koehl et al., 2019; Seven et al., 2021) (Fig. 2A), the active conformation of mGlu₂R and mGlu₄R homodimers coupled to a heterotrimeric Gi protein (Lin et al., 2021) and the inactive conformation of the mGlu₇R homomer and the mGlu₂R-mGlu₇R heteromer (Du et al., 2021). Whether class A (rhodopsin-like family) and family B (secretin receptor family) GPCR dimers are also constitutively formed has been a more contentious issue, even though the crystal structures of several GPCRs revealed homo-oligomers, including the chemokine CXCR₄R (Wu et al., 2010), the μ -opioid receptor (MOR) (Manglik et al., 2012) and the β_1 -adrenoceptor (β_1 AR) (Huang, Chen, Zhang, & Huang, 2013) (Fig. 2B–D).

Apart from the crystallographic evidence, during the last two decades, numerous studies using different technical approaches have provided strong support to the increasingly accepted notion that GPCR homomers and heteromers constitute primary functional GPCR




Fig. 1. G protein-coupled receptor signaling. (A) GPCRs are illustrated with the structure of retinal-bound rhodopsin in white (Palczewski et al., 2000). In humans, there are 16 G α subunits (classified as G α_s , G α_i , G α_q and G α_{12}) 5 G β and 12 G γ (G γ 1-13, but no G γ 6) subunits. Active GPCRs interact with G $\alpha\beta\gamma$ proteins, illustrated with the structure of GDP-bound G α_i 1 β 1 γ 2 in brown/yellow (Wall et al., 1995), G protein-coupled receptor kinases (GRKs), with 7 mammalian isoforms, illustrated with the structure of ATP-bound GRK1 in light orange (Singh, Wang, Maeda, Palczewski, & Tesmer, 2008), and arrestins, with 4 isoforms, illustrated with the structure of arrestin1 in red (Granzin et al., 1998). (B) Agonist binding triggers the movement of TM6 (in blue), opening an intracellular cavity for G protein arrangement, illustrated with the crystal structure of the agonist- β 2AR-Gs complex (Rasmussen et al., 2011). G α_q activates PLC, with 13 mammalian isozymes, illustrated with the crystal structure of PLC- β 3 bound to G α_q (Waldo et al., 2010), which catalyzes the hydrolysis of PIP₂ to the Ca²⁺-mobilizing second messenger inositol 1,4,5-trisphosphate (IP₃) and the PKC-activating second messenger diacylglycerol (DAG). G α_s stimulates and G α_i inhibits the activity of AC, with 9 mammalian isoforms, illustrated with the cryo-EM structure of AC9 bound to G α_s (Qi et al., 2019). G $\beta\gamma$ subunits increase the activity of Kir3 channels, illustrated with the crystal structure of the Kir3.2-G $\beta\gamma$ (Whorton and MacKinnon, 2013). Desensitization is mediated by GRK-mediated GPCR phosphorylation and subsequent binding of arrestin, illustrated with the cryo-EM structure of neurotensin NT₁ receptor in complex with arrestin2 (Huang et al., 2020). G γ isoprenylated group is shown in red.

signaling units (for reviews, see Ferré et al., 2014; Ferré, Ciruela, Casadó, & Pardo, 2020; Gomes et al., 2016; Sleno & Hébert, 2018; Gaitonde & González-Maeso, 2017; Bourque, Jones-Tabah, Devost, Clarke, & Hébert, 2020). GPCR oligomerization significantly broadens the allosteric mechanisms inherent in GPCRs, imposing conformational alterations and constraints of the individual GPCR units (Ferré et al., 2014, 2020). This gives the possibility of developing GEMMAs as targets for drug discovery, with the rationale of the existence of different GEMMAs with common components but with different oligomeric partners localized in different types of cells or cellular environments. Screening campaigns should look for molecules with the ability to promote distinct states of specific GEMMAs, using single agents or combinations of molecules with selectivity for their individual components.

2. Evidence for pre-assembly and ligand-induced rearrangement of specific GPCR, G proteins and effector complexes

2.1. Pre-assemblies of specific G α , G β and G γ subunits

Since Gilman's classic conceptualization (Gilman, 1987) it was generally assumed that upon binding to the ligand (a GPCR agonist), GPCRs undergo conformational changes activating heterotrimeric G proteins by promoting GTP binding to G α subunits and triggering the release of G $\beta\gamma$ subunits. When dissociated, both α and $\beta\gamma$ subunits modulate the activities of effector molecules directly responsible for generating cellular responses (Gilman, 1987). However, as we discuss, an alternative model suggests that G protein heterotrimers can be

found in more stable complexes that rearrange rather than dissociate upon receptor activation.

In many cases, coupling between receptors and G proteins is pleiotropic such that a single receptor can activate more than one G protein heterotrimer (Flock et al., 2017; Okashah et al., 2019), which allows initiation of multiple signal transduction pathways. However, signaling *in vivo* most often results in regulation of selective effectors with high fidelity, suggesting that the cell is endowed with different modalities to allow segregation of specific connections between different GPCRs and their effectors. In humans, there are at least 16 different isoforms of G α subunits, grouped into families- G α_s (G α_s , G α_{olf} and G α_{sXL}), G α_i (G α_i 1, G α_i 2, G α_i 3, G α_{o1} , G α_{o2} , G α_z , G α_t and G α_{gust}), G α_q (G α_q , G α_{11} , G α_{14} and $\alpha_{15/16}$) and α_{12} (α_{12} and α_{13}); and 5 different isoforms of G β (β 1-5) and 12 G γ (G γ 1-13, but no G γ 6) subunits (Downes & Gautam, 1999). The combinatorial association of different G protein subunits potentially provides the bandwidth necessary to generate a broad range of signals independently transduced from different GPCRs to different effectors transmitted by G proteins (Robishaw & Berlot, 2004).

Several seminal studies have provided evidence that specific combinations of G α , G β and G γ subunits are necessary to generate signal transduction from a given GPCR to a specific effector (Cabrera-Vera et al., 2002; Dupré, Robitaille, Rebois, & Hébert, 2009; Gudermann, Kalkbrenner, & Schultz, 1996; Rebois & Hébert, 2003). Studies in the early nineties using antisense oligonucleotides demonstrated that, even in the same cells, different combinations of specific G α , G β and G γ subunits allowed different receptors to regulate the same effector (Kleuss, Schultz, & Wittig, 1994). Somatostatin, muscarinic M₄




Fig. 2. Molecular structures of GPCR homomers. (A) Cryo-EM structures of the full length mGlu₅R in the apo inactive/open and active/closed conformations (Koehl et al., 2019). Class C GPCRs are obligate cysteine-linked homomers and each protomer is formed by a large extracellular 'Venus fly-trap' domain that binds orthosteric ligands and promotes an inter-protomer disulfide bridge formed via a conserved cysteine, a cysteine rich domain and a seven TM domain (7TM). The helix interface between protomers of the Venus fly-trap domains is represented by cylinders. Activation leads to compacting of the mGlu₅R homomer, bringing the cysteine rich domains into proximity and enabling the 7TM domains to approximate to initiate signaling. The inactive state shows a TM5-TM5 orientation but with substantial separation between the 7TM domains. Activation leads to a 20° rotation of each 7TM domain, resulting in a close contact TM6-TM6 homodimeric interface. Both protomers are colored in light blue and gray. (B) CXC₄R structure (Wu et al., 2010) reveals a homomer with an interface including TM5 and TM6. (C) MOR structure (Manglik et al., 2012) shows receptor protomers associated in pairs through two different interfaces. One interface is via TM1, TM2 and helix 8 (Hx8) (blue/white and light/dark green protomers) and the other interface comprises TM5 and TM6 (white/green protomers). (D) β₁AR structure (Huang et al., 2013) also displays two homomeric interfaces. One interface also involves TM1, TM2 and Hx8 (blue/white and light/dark green protomers), as with MOR. In contrast, the other interface engages residues from TM4, TM5 and intracellular loop 2 (ICL2) (white/green protomers).

and galanin receptors inhibited voltage gated Ca²⁺ channels by specifically coupling to Gα₂β₁γ₃, Gα₁β₃γ₄ and Gα₁β₂γ₂, respectively (Kalkbrenner et al., 1995; Kleuss, Scherübl, Hescheler, Schultz, & Wittig, 1993). The same studies showed that galanin receptors selectively used the same combination of G protein subunits, Gα₁β₂γ₂, to inhibit voltage gated Ca²⁺ channels in a different cell type (Kalkbrenner et al., 1995).

However, there is also experimental evidence indicating that one GPCR couples to different combinations of G protein subunits in different cells. Particularly significant is the example of the α_{2A}-adrenoceptor (α_{2A}AR) in the mouse brain, which, without considering the Gα subunit of the heterotrimer involved, was found to associate selectively to Gβ₂, Gγ₂ and Gγ₃ in noradrenergic cells, but to Gβ₄ and Gγ₁₂ in non-noradrenergic cells (Yim et al., 2019). Another remarkable example for the need of specific Gαβγ subunit compositions comes from genetic studies, which demonstrate that, specifically in the striatum, the signaling of adenosine A_{2A} receptors (A_{2A}R) and dopamine D₁ receptor (D₁R) with AC, more specifically the AC5 isoform, is dependent on specific coupling to Gα₁β₃γ₇ (Schwindinger et al., 2003; Schwindinger et al., 2010; Hervé, 2011; Xie et al., 2015). The involvement of specific assemblies of G protein subunits from a potential large number of combinations as components of specific complexes of GPCRs, G proteins and PM-effectors

does not seem to depend on a specific preferential affinity between those subunits, but on their synchronized ER synthesis, allowing pre-assembly and delivery to the plasma membrane.

Several studies indicate that Gαβγ heterotrimers pre-assemble before delivery to the plasma membrane, although there remains controversy over the secretory pathway involved. One study suggested that fully processed Gβγ subunits form heterotrimers with Gα on the cytosolic face of the Golgi apparatus (Michaelson, Ahearn, Bergo, Young, & Philips, 2002). However, another study favored Golgi-independent trafficking of the Gαβγ heterotrimer, where formation was prevented by a dominant mutant of the small GTPase Sar1 (Takida & Wedegaertner, 2004), responsible for the assembly of coated vesicles at the endoplasmic reticulum (ER) membrane (Mizuno-Yamasaki, Rivera-Molina, & Novick, 2012). In any case, assembly of the heterotrimer precedes acylation of the Gα subunit, which is necessary for delivery of the heterotrimer to the plasma membrane (Dupré et al., 2009; Marrari, Crouthamel, Irannejad, & Wedegaertner, 2007). The evidence for pre-assembly of the full heterotrimeric G protein in its journey to the plasma membrane indicates that the different specific Gαβγ subunit combinations are not selected from a pool of subunits localized at the plasma membrane and sorted by random collision or another so far unidentified mechanism following receptor activation.

2.2. Stable interactions between specific GPCRs, G proteins and PM-effectors

Biophysical techniques have been widely used to evaluate the possible existence of intermolecular interactions between specific GPCRs, G proteins and PM-effectors in living cells (Bouvier, 2001; Milligan, 2001; Guo et al., 2017; El Khamlichi et al., 2019; Soave, Briddon, Hill, and Stoddart (2020)). In most of these techniques, two chromophores that can change their biophysical properties when they are in contact, or in very close proximity, are separately fused to the two putative interacting proteins and transfected to mammalian cells in culture. These techniques include bioluminescence and Förster or fluorescence resonance energy transfer (BRET and FRET), time-resolved FRET (TR-FRET), bimolecular bioluminescence or fluorescence complementation (BiBC or BiFC), and fluorescence recovery after photobleaching (FRAP). A series of studies using combinations of these biophysical techniques with biochemical techniques (co-immunoprecipitation and pull-down strategies) has provided strong support for the existence of stable interactions between GPCRs and G proteins (Galés et al., 2005, 2006; Nobles, Benians, & Tinker, 2005; Dupré et al., 2006; Ayoub, Trinquet, Pflieger, & Pin, 2010; Qin, Dong, Wu, & Lambert, 2011; Damian et al., 2015; Andressen et al., 2018) and between G proteins and PM-effectors (Rebois et al., 2006; Riven, Iwanir, & Reuveny, 2006; Sadana, Dascal, & Dessauer, 2009; Yuan, Sato, Lanier, & Smrcka, 2007). Further evidence has also accumulated for the existence of direct interactions between GPCRs and PM-effectors, including L-type and N-type voltage-dependent Ca^{2+} channels (Beedle et al., 2004; Rebois et al., 2006), inwardly rectifying potassium Kir3 channels (David et al., 2006; Lavine et al., 2002) and ACs (Dupré, Baragli, Rebois, Ethier, & Hébert, 2007; Lavine et al., 2002; Navarro et al., 2018). These studies strongly support the existence of specific assemblies of GPCRs, G proteins and effectors localized in plasma membrane-delimited signaling complexes. Their specific make up raises the notion that components of such complexes could be pre-assembled before delivery to the plasma membrane. It could be demonstrated that dominant negative small GTPase isoforms promote retention of GPCR-G protein, G protein-PM-effector and GPCR-PM-effector assemblies. For instance, prevention of anterograde trafficking to the plasma membrane using mutants of the coat-recruitment small GTPase Sar1 or the also small GTPase Rab1, which guides transport vesicles to the plasma membrane (Mizuno-Yamasaki et al., 2012), reduced $\beta_2\text{AR}$ plasma membrane localization but not interactions between $\beta_2\text{AR}$ and $\text{G}\beta_1/\text{G}\gamma_2$ subunits, AC2 and $\beta_2\text{AR}$ or AC2 and β_1 or $\text{G}\gamma_2$ fused to BRET chromophores (Dupré et al., 2006; Dupré et al., 2007), indirectly indicating the pre-assembly of $\beta_2\text{AR}$ - $\beta_1\gamma_2$ -AC2 complexes.

Several recent findings suggest plausible scenarios of GPCR-G protein interactions that are different from the fully engaged and agonist-promoted interactions in which helix 5 of the Ras-GTPase domain of the $\text{G}\alpha$ protein interacts with the intracellular cavity of the GPCR formed by the agonist-promoted outward movement of TM6 (Weis & Kobilka, 2018). Whereas initial X-ray and cryo-EM structures of GPCR-G protein complexes were captured in their nucleotide-free state, recent structures based on time-resolved structural mass spectrometry techniques begin to observe alternative complexes and conformational transitions, all indicating the plasticity of interacting interfaces and dynamic nature of the GPCR-G protein complex (Du et al., 2019; Kato et al., 2019; Liu et al., 2019). Such techniques are expected to further reveal the molecular nature of GPCR-G protein pre-assemblies in the near future.

2.3. Association-dissociation of GPCRs and G proteins or ligand-induced rearrangement of GPCR-G protein complexes?

Biophysical techniques have also allowed the analysis of ligand-induced changes in the interactions between GEMMA components. The insertion of BRET chromophores in the $\alpha_{2A}\text{AR}$ and in different positions of the $\text{G}\alpha_{i1}$, β_1 and γ_2 subunits was used to establish that during the early stages of receptor activation, G protein activation is

associated with a relative movement between the $\text{G}\alpha$ and $\text{G}\beta\gamma$ subunits that does not involve their complete dissociation (Galés et al., 2006). More precisely, these BRET experiments showed a ligand-induced separation of the $\text{G}\alpha$ subunit helical domain versus the Ras-GTPase domain coupled to the $\text{G}\beta\gamma$ subunit (Galés et al., 2006). The relatively large separation of the helical and Ras-GTPase domains of the $\text{G}\alpha$ subunit, which tightly sandwich the nucleotide in all nucleotide-bound G protein crystal structures, was later on confirmed by double electron-electron resonance spectroscopy (Van Eps et al., 2011), a crystal structure of a GPCR-G protein complex (Rasmussen et al., 2011), and deuterium-exchange and electron microscopy data (Chung et al., 2011; Westfield et al., 2011).

Subsequent studies confirmed the ability of different GPCRs to transit from inactive to active conformations by a rearrangement of a pre-assembled receptor-G protein complex, such as a study on monomeric ghrelin GHS_{1a} receptor (GHS_{1a}R) reconstituted in nanodiscs, using TR-FRET and normal mode (NM) analysis (Damian et al., 2015). On the other hand, a more recent study using single-molecule tracking (SMT) with total internal reflection fluorescence (TIRF) obtained results indicating the existence of a predominant population of short-lived (1 second) complexes of $\alpha_{2A}\text{AR}$ and $\text{G}\alpha_{i1}$ labelled with two different organic fluorophores and expressed at low densities in mammalian transfected cells, in the absence of ligands (Sungkaworn et al., 2017). The authors proposed a model dependent on collision coupling, where agonists specifically regulate the kinetics of GPCR-G protein interactions by increasing their association rate (Sungkaworn et al., 2017).

Thus, the debate about a pre-assembly and rearrangement versus collision coupling and association-dissociation of GPCRs and G proteins remains open. Nevertheless, both mechanisms seem to operate and be dependent on protein density, the specific GPCR-G protein pair, and the cellular environment. With BRET and TR-FRET techniques, the GPCR protease-activated PAR_1 receptor was shown to specifically form assemblies with $\text{G}\alpha_{i1}$ but not with $\text{G}\alpha_{i2}$ (Ayoub et al., 2010). Nevertheless, $\text{G}\alpha_{i2}$ was slowly recruited upon receptor activation, indicating the possibility of the two modes of interaction of the same receptor, pre-assembly and collision coupling, depending on the G protein partner (Ayoub et al., 2010). Similar conclusions were obtained from a study using FRAP and BRET techniques examining pre-assembly and agonist-induced rearrangement of the M_3R and $\text{G}\alpha_{q1}$ (Qin et al., 2011). A six-amino-acid polybasic sequence distal to helix 8 was necessary for pre-assembly with Gq. Moreover, the polybasic epitope was also necessary for an efficient M_3R -mediated Gq activation, showing that GPCR-G protein pre-assembly significantly increased the rate of Gq-dependent signaling (Qin et al., 2011). Results obtained with FRAP and FRET techniques also showed a differential ability of 5-HT₇R and 5-HT₄R to preassemble with Gs proteins (Andressen et al., 2018). The results support a model where pre-associated 5-HT₇R- $\text{G}\alpha_{s1}\beta_1\gamma_2$ complexes undergo agonist-induced rearrangement involving a rapid movement of $\text{G}\alpha$ relative to the receptor, followed by a slower dissociation of $\text{G}\beta\gamma$ subunits from both the receptor and the $\text{G}\alpha$ subunit. In contrast, 5-HT₄R displayed properties more consistent with collision coupling (Andressen et al., 2018). These studies (Andressen et al., 2018; Ayoub et al., 2010; Qin et al., 2011) therefore indicated the possibility of the coexistence of two alternative signaling modes in the same cell, with pre-assembly providing a means to regulate more specific signaling events as compared to the collision-coupling mode.

2.4. Association-dissociation of G proteins and PM-effectors or ligand-induced rearrangement of G protein-PM-effector assemblies?

A substantial number of studies provided evidence for pre-assembly (inactive state) and rearrangement (active state) of G proteins and PM-effectors, mainly dependent on interactions with $\text{G}\beta\gamma$ subunits. $\text{G}\beta\gamma$ subunits were initially thought to passively facilitate termination of intracellular information transfer by binding to the $\text{G}\alpha$ subunit and

preventing its further spontaneous activation in the absence of receptor activation. This dogma was challenged when G $\beta\gamma$ was shown to directly activate a Kir3 channel in cardiac atrial cells (Logothetis, Kurachi, Galper, Neer, & Clapham, 1987). G $\beta\gamma$ subunits are now known to modulate different isoforms of ACs, PLCs, K⁺ and Ca²⁺ channels and many other PM-effectors (Dupré et al., 2009; Khan et al., 2013; Khan, Sung, & Hébert, 2016; Smrcka & Fisher, 2019). The G $\beta\gamma$ structure includes the seven β -sheet WD40 repeat architecture of the G β subunit (the seven “blades” of the G β subunit “propeller”) and a “hotspot” surface where the turns between blades intersect. The G $\beta\gamma$ hotspot is required for interactions with numerous effectors and it is concealed in the inactive state by its binding to the Ras-GTPase domain of the G α subunit. Upon ligand-induced activation, the hotspot is exposed and interacts with available effectors (Cabrera-Vera et al., 2002; Davis, Bonacci, Sprang, & Smrcka, 2005). Distinct but overlapping motifs of the hotspot area are involved in the interactions with the G α subunit and the different effectors (Cabrera-Vera et al., 2002; Davis et al., 2005; Mirshahi, Robillard, Zhang, Hébert, & Logothetis, 2002; Smrcka & Fisher, 2019). The sites required for agonist-induced activation of effectors by the G $\beta\gamma$ (or by the G α) subunits are likely to be inaccessible in the GDP-bound, inactive, G $\alpha\beta\gamma$ assembly, thus necessitating certain degree of dissociation of the assembly for activation of the effectors.

In addition to their key modulatory role in effector function, cumulative evidence supports an additional key role of G $\beta\gamma$ subunits in the pre-assembly of G proteins to PM-effectors in their inactive state. This has been addressed *in vitro* by demonstrating direct interactions between G $\beta\gamma$ subunits with the PM-effector when forming part of the GDP-bound G $\alpha\beta\gamma$ assembly or without involving the hotspot area. Initial evidence was obtained from pull-down experiments with purified bovine G $\beta\gamma$ subunits and the cytosolic NT and CT of the Kir3.1 subunit of Kir3 channels (Huang, Slesinger, Casey, Jan, & Jan, 1995). Kir3 channels are prototypic PM-effectors of G $\beta\gamma$ subunits and are activated by direct interactions with G $\beta\gamma$ that follow GPCR-dependent activation of Gi/o proteins (Dascal & Kahanovitch, 2015; Logothetis et al., 1987, 2015). Kir3 channels are homo- or hetero-tetrameric and can be composed of four distinct subunits, Kir3.1–4. The basic architecture of the Kir3 channel subunits consists of two TM helices with the cytosolic NT and CT and an extracellular loop which folds back to form the pore-lining ion selectivity filter. Those initial pulldown experiments showed that G $\beta\gamma$ subunits bind to both cytosolic domains and that a GDP-bound G $\alpha\beta\gamma$ assembly specifically binds to the NT of the Kir3.1 subunit, involving contacts mediated by both G $\beta\gamma$ and G α subunits (Huang et al., 1995). The results also indicated that both cytoplasmic regions of the Kir3 channel subunit are involved in the activation of the Kir3 channel (Huang et al., 1995).

Subsequent studies, using different biochemical and biophysical techniques, provided additional evidence for the involvement of the G β subunit in the interaction with Kir3 channels, both in its active and inactive states. Key residues that constituted significant contact points with Kir3 channels and were not concealed by the non-activated G α subunit were found to be involved in G $\beta\gamma$ -Kir3 channel interactions in the absence of agonist stimulation (Mirshahi et al., 2002). Experiments with internal reflection fluorescence (TIRF) microscopy combined with FRET in cells transfected with mGlu₂R, Kir3.1 or Kir3.4 fused at various positions of their cytosolic NT and CT with a FRET chromophore and G $\beta 1$ fused to the other chromophore, indicated an interaction between the G $\beta\gamma$ subunits and the Kir3 channel in the resting state and a change in FRET in the presence of the endogenous ligand glutamate (Riven et al., 2006). The results suggested that receptor activation should promote an orientation switch of the G $\beta\gamma$ dimer, without affecting the position of G α relative to the channel, which would allow the interaction of the G $\beta\gamma$ with the channel at a separate, independent site to promote opening (Riven et al., 2006).

Experiments with BRET in a cell line stably expressing β_2 AR allowed further analysis of interactions between G $\beta\gamma$ and Kir3 channels and another PM-effector, AC2, in the absence and presence of ligands,

again indicating both pre-assembly and agonist-induced rearrangement (Rebois et al., 2006). The same study provided evidence for pre-assembly and ligand-induced rearrangement of G proteins and Kir3 channels before trafficking to the plasma membrane, by fusing Kir3.1 and G $\gamma 2$ to BRET chromophores and measuring BRET changes in response to membrane-permeable and non-membrane agonists in the presence and absence of Kir3.4. In the absence of Kir3.4, the $\beta 1\gamma 2$ -Kir3.1 complex did not traffic to the plasma membrane and changes in BRET were only evident using membrane-permeable agonists (Rebois et al., 2006). These results therefore indicated an intracellular localization of functional β_2 AR as part of a GEMMA.

The subsequent publication of the crystal structure of the Kir3 channel with four Kir3.2 subunits in complex with G $\beta 1\gamma 2$ and PIP₂ confirmed the inferences made from many of the previous biochemical and biophysical studies (Glaaser & Slesinger, 2015). The biologically relevant complex consists of one channel tetramer, four G $\beta\gamma$ subunits, four PIP₂ molecules and four Na⁺ ions bound to regulatory sites (Whorton & MacKinnon, 2013). G $\beta\gamma$ subunits interact directly with the cytosolic CT and they are oriented such that the CT of the G γ subunit, which contains a covalent geranylgeranyl group, points directly to the membrane layer as if to function as an anchor (Whorton & MacKinnon, 2013) (Fig. 1B). Thus, the Kir3.2-G $\beta 1\gamma 2$ crystal structure is compatible with a physiological membrane-delimited G $\beta\gamma$ activation of Kir3. We are now waiting for structures of the Kir3 channel in complex with a GPCR and its heterotrimeric Gi protein in both inactive and active states. In the meantime, the studies described here provide evidence for pre-assembly and rearrangement of G $\alpha\beta\gamma$ -Kir3 complexes in response to agonist. Both G α and G $\beta\gamma$ subunits appear to be directly associated with Kir3 under both ligand-free inactive and agonist-induced active states. Agonist-induced rearrangement of the complex suggests separation of the G $\beta\gamma$ subunits from G α subunit, but *within a metastable complex*, with G $\beta\gamma$ subunits moving away from the initial Kir3 NT- and CT-bound G $\alpha\beta\gamma$ assembly.

The existence of G proteins and AC as a stable complex was first proposed by Levitzki based upon co-purification of AC and G proteins from turkey erythrocyte membranes independent of the activation state of the G proteins (Bar-Sinai, Marbach, Shorr, & Levitzki, 1992; Levitzki, 1988; Levitzki & Klein, 2002). A role of G $\beta\gamma$ subunits in the pre-assembly and scaffolding of ACs has also been supported by *in vitro* and *in cellulo* experiments (see below). Further, we also know that G $\beta\gamma$ subunits can directly interact with AC facilitating or inhibiting its function depending on the AC isoform. This modulatory role of G $\beta\gamma$ is also conditional on G α s-mediated activation. ACs are generally classified into four different categories based on their regulatory properties. All isoforms of TM ACs are stimulated by GTP-bound G α s. Group I includes G $\beta\gamma$ -inhibited and Ca²⁺-stimulated AC1, AC3 and AC8; group II includes G $\beta\gamma$ -stimulated AC2, AC4 and AC7; group III includes G αi and Ca²⁺-inhibited and G $\beta\gamma$ -stimulated AC5 and AC6; and group IV includes AC9, which is not regulated by G αi or Ca²⁺ and was initially believed to be forskolin-insensitive (Baldwin, Li, Brand, Watts, & Dessauer, 2019; Dessauer et al., 2017; Qi, Sorrentino, Medalia, & Korkhov, 2019). The topology of all ACs includes a long NT, two membrane-spanning domains, M1 and M2, each with six TM domains and two large cytoplasmic catalytic domains, C1 and C2. The C1 and C2 domains are homologous and dimerize to form the catalytic core at their interface, where ATP is converted to cAMP. When a GTP-bound G α s subunit binds to C2, it increases the affinity between C1 and C2, promoting catalysis, while for group III, G αi subunits bind to C1 and have the opposite effect (Baldwin et al., 2019; Dessauer et al., 2017; Sadana & Dessauer, 2009).

FRET experiments using truncated AC5 provided compelling evidence for a significant role of the NT of AC5 in scaffolding inactive, GDP-bound G $\alpha\beta\gamma$ (Sadana et al., 2009). It was also shown that AC5 NT interacts with its catalytic domains to enhance G α s- or forskolin-stimulated AC5 activity. These results support a model of G $\alpha\beta\gamma$ -AC5 interactions where the AC5 NT brings the inactive heterotrimeric G

protein and AC5 catalytic core in close proximity for efficient GPCR activation (Sadana et al., 2009). A subsequent study showed that $G\alpha\beta\gamma$ binds to the NT of most AC isoforms (Brand, Sadana, Malik, Smrcka, & Dessauer, 2015). Mutational analysis indicated that the $G\beta\gamma$ hotspot does not interact with the AC5 NT scaffolding side, although it remains necessary for conditional $G\beta\gamma$ -mediated stimulation of AC5. On the other hand, the $G\beta\gamma$ hotspot was required for both stimulating AC6 and interacting with its NT domain, indicating that $G\beta\gamma$ regulation of AC involves multiple binding events, and that the AC NT plays unique isotype-specific roles (Brand et al., 2015).

The recently reported structure of the active state (GTP γ S-bound) of AC9 in complex with $G\alpha_s$, using cryo-electron microscopy (cryo-EM) and single-molecule analysis, revealed for the first time the twelve TM domains of a TM AC isoform, as well as a helical domain that spans between the membrane and the catalytic domains of AC9 (Qi et al., 2019) (Fig. 1B). The helical domain, in fact, corresponds to a 40-residue-long cytosolic extension of TM6 and TM12 (Qi et al., 2019). This implies the necessity of dissociation of the $G\alpha$ subunit from the pre-assembled GPCR- $G\alpha\beta\gamma$ complex allowing the concealed domain of the Ras-GTPase domain of the $G\alpha$ subunit to interact with the corresponding AC catalytic domain. Similarly to Kir3 channel- $G\alpha\beta\gamma$ assembly, this suggests that ligand-induced rearrangements of the components of such complexes are associated with a separation of the $G\alpha$ subunit from the $G\beta\gamma$ subunits, but *within the framework of the assembly*, guided by the modifications in the interactions between the $G\alpha$ and $G\beta\gamma$ subunits with the AC NT.

3. Compartmentalization of signaling molecules at the plasma membrane by membrane nanodomains and scaffold proteins

3.1. Membrane nanodomains

Apart from pre-assembly of receptor-based signaling complexes, to achieve fidelity and maintain the efficiency of signaling across the plasma membrane, the cell uses two additional complementary mechanisms that promote co-localization or association of specific components of distinct GPCR signaling complexes: i) membrane nanodomains, such as lipid rafts and ii) scaffold proteins. Lipid rafts are defined as heterogeneous, dynamic, cholesterol- and sphingolipid-enriched membrane nanodomains (10–200 nm), which have the potential to form microscopic domains (>300 nm) upon clustering induced by protein-protein and protein-lipid interactions (Pike, 2006). Enriched hydrophobic components and phospholipids that contain saturated fatty acyl chains increase lipid packing and order and subsequent decreased fluidity (Ahmed, Brown, & London, 1997; Pike, 2003; Sezgin, Levental, Mayor, & Eggeling, 2017). Certain structural proteins enriched in lipid rafts serve as scaffolds or anchors for other proteins, including caveolins, glycosylphosphatidylinositol (GPI)-linked proteins and cortical actin, which also play an active part in the maintenance and remodeling of lipid rafts (Pike, 2003; Sezgin et al., 2017).

Since lipid rafts and other membrane nanodomains are not resolvable on a conventional optical microscope, super-resolution optical microscopy approaches such as photoactivated localization microscopy (PALM), stimulated emission depletion (STED) or stochastic optical reconstruction microscopy (STORM) have played a significant role in the visualization of lipid-mediated membrane protein clustering (for reviews, see Curthoys et al., 2015; Sezgin et al., 2017). For more dynamic measurements, the addition of SMT analysis has allowed to evaluate the diffusion of membrane molecules and relate it to models of heterogeneous organization of the membrane (Curthoys et al., 2015; Sezgin et al., 2017). Those include the membrane “hot spots” of interacting α_{2A} AR and $G\alpha_i1$ described by Sungkaworn et al. (2017).

The presence within lipid rafts of a variety of membrane proteins involved in cell signaling led to the consensus that membrane nanodomains play an important role in facilitating rapid and specific signal transduction events (Simons & Toomre, 2000; Smart et al.,

1999). Many GPCRs and their cognate signaling partners, including G protein subunits and AC isoforms, are enriched in lipid rafts (Ostrom, Violin, Coleman, & Insel, 2000). In the simplest scenario, lipid rafts could serve to co-localize specific GPCRs, G proteins and PM-effectors. Alternatively, rafts could contain a much larger array of signaling molecules activated when a receptor or other required molecule is recruited into the raft (Pike, 2003). There is also evidence that the targeting of GPCRs and their proximal signaling partners to lipid rafts can be cell specific (Ostrom et al., 2000; Ostrom et al., 2002). Finally, there is evidence indicating that not all lipid rafts are equivalent and may have distinctly different protein and/or lipid components that coexist in cells (Pike, 2003). However, there is no evidence for mechanisms to isolate specific GPCRs, G protein subunits and PM-effectors by segregating them into distinct lipid rafts or other membrane nanodomains and, given the large number of different signaling molecules that they can harbor, membrane nanodomains cannot be the primary means for generating signaling specificity.

3.2. Plasma membrane scaffold proteins and A kinase anchoring proteins

The most extensively studied plasma membrane scaffold proteins are those localized in and around the postsynaptic density of the glutamatergic synapse. PSD-95/Discs large/Zona occludens-1 (PDZ) domain-containing proteins (PDZ proteins) are the most abundant and PDZ domains represent the most common protein-protein interaction domain (Dunn & Ferguson, 2015). PDZ proteins directly bind to the PDZ-binding ligand motifs found in the CT region of many adhesion molecules and receptors and often contain additional modular interacting domains, which can bind to each other and to various signaling proteins encompassing both sides of the synapse (Dunn & Ferguson, 2015; Funke, Dakoji, & Bredt, 2005; Sheng & Sala, 2001; Zheng, Seibold, Horak, & Petralia, 2011). PSD-95 family PDZ domains bind to the CT of several GPCRs, including β_1 AR and β_2 AR (Dunn & Ferguson, 2015; He et al., 2006; Hu et al., 2000; Joiner et al., 2010; Li, Nooh, & Bahouth, 2013; Valentine & Haggie, 2011; Xiang, Devic, & Kobilka, 2002) and serotonin 5-HT $_{2A}$ and 5-HT $_{2C}$ receptors (Xia, Gray, Compton-Toth, & Roth, 2003). The PDZ domain of another PDZ protein family, Shank, interacts indirectly with GPCRs, including group I mGluRs through the additional scaffold protein Homer, which binds to the Shank proline-rich domain (Fig. 3A) (Bécamel et al., 2004; Dunn & Ferguson, 2015; Zheng et al., 2011). At the postsynaptic density, a key scaffolding link is determined by indirect interactions between PSD-95 and Shank proteins established by guanylate kinase-associated protein (GKAP), which refers to a family of four scaffold proteins which connect the guanylate kinase domain of PSD-95 with Shank PDZ domain (Shin et al., 2012) (Fig. 3A). Apart from specific interactions with GPCRs, direct interactions have been reported between PDZ proteins and the G protein subunit $G\gamma13$, particularly expressed in taste and olfactory cells (Li, Benard, & Margolske, 2006; Liu et al., 2012) and between Shank and the CT of the central pore subunit of the L-type voltage-dependent Cav1.3 Ca $^{2+}$ channels ($\alpha1.3$) (Olson et al., 2005). However, to our knowledge, there is no evidence to date for other direct interactions between PSD-95 and Shank families of PDZ proteins and G protein subunits or other PM-effectors.

A-kinase anchoring proteins (AKAPs) are a family of structural diverse proteins which share a common motif that binds to the regulatory subunits of the cyclic AMP-dependent protein kinase A (PKA) (Wong & Scott, 2004). More than 50 AKAPs have been identified which can be found in most cellular organelles, as well as the plasma membrane. The scaffolding of unique signaling elements by different types of AKAPs provides a framework for integration and modulation of distinct AC-cAMP-PKA-mediated cell signals (Dessauer, 2009). For instance, the plasma membrane AKAP5 (also known as AKAP79/150) targets not only PKA, but also PKC and protein phosphatase 2B (PP2B), to the inner face of the plasma membrane (Patriarchi, Buonarati, & Hell, 2018; Scott, Dessauer, & Taskén, 2013). In addition, AKAP5 binds to the intracellular




Fig. 3. Scaffolding of GPCRs, ligand-gated ion channels and PM-effectors mediated by PDZ proteins and AKAPs. (A) Complexes of ionotropic glutamate AMPAR and group I mGluRs (mGlu₁R or mGlu₅R) established by a chain of interactions of the scaffold proteins PSD-95, GKAP, Shank and Homer. Homer directly interacts with the CT of mGlu₁R or mGlu₅R, while AMPAR indirectly binds to PSD-95 by oligomerization with the TM AMPAR regulatory protein stargazin (Str). (B) Complexes of β_2 AR, AC and L-type voltage-dependent Cav1.2 Ca^{2+} channels mediated by AKAP5. (C) Complexes of β_2 AR, AC and AMPAR mediated by AKAP5 and PSD-95. By bringing together β_2 AR, AC and PKA, AKAP5-mediated complexes provide the frame for an efficient β_2 AR-dependent, PKA-mediated phosphorylation of the α 1.2 subunit of Cav1.2 channels (B) and the GluA1 subunit of AMPAR (C).

domains of several membrane-bound signaling molecules, including β_1 AR and β_2 AR (Fraser et al., 2000; Gardner, Tavalin, Goehring, Scott, & Bahouth, 2006; Li et al., 2013; Valentine & Haggie, 2011) and the PM-effectors AC5 and AC6 (Bauman et al., 2006; Efendiev et al., 2010) and L-type voltage-dependent Cav1.2 Ca^{2+} channels (Hall et al., 2007; Oliveria, Dell'Acqua, & Sather, 2007; Patriarchi et al., 2018). In addition, PSD-95 family proteins can interact through their SH3 and GK domains with a broad surface in the central region of AKAP5. PDZ proteins and AKAPs can therefore combine their interactions with different molecules to scaffold distinct macromolecular signaling complexes that include GPCRs. In this way, AKAP5 can promote the interaction of PKA with its plasma membrane-localized phosphorylation targets, either directly, such as for β_1 AR and β_2 AR (Fraser et al., 2000; Li et al., 2013; Valentine & Haggie, 2011), AC5 and AC6 (Bauman et al., 2006; Zhang et al., 2013) and the α 1.2 subunit of Cav1.2 channels (Oliveria et al., 2007), or via binding to PSD-95 family proteins, such as for the GluA1 subunit of ionotropic glutamate AMPA receptors (AMPA) (Colledge et al., 2000; Diering, Gustina, & Haganir, 2014; Joiner et al., 2010).

Interestingly, the β_2 AR has been shown to be involved in AKAP5-dependent PKA-mediated phosphorylation of both the α 1.2 subunit of Cav1.2 channels (Patriarchi et al., 2018) and the GluA1 subunit of AMPAR (Joiner et al., 2010); both events occur in dendritic spines of excitatory synapses (Patriarchi et al., 2018). Thus, at the postsynaptic density, there is evidence for the existence of different AKAP5-scaffolded macromolecular β_2 AR-AC-AKAP5 complexes that include either Cav1.2 channels or AMPAR (Figs. 3B and 3C). In both complexes, the β_2 AR does not seem to be directly connected to AKAP5. In the β_2 AR-AC-Cav1.2-AKAP5 complex, β_2 AR oligomerizes with Cav1.2 (Davare et al., 2001), which directly binds to AKAP5 (Hall et al., 2007; Oliveria et al., 2007) (Fig. 3B). In the β_2 AR-AC-AMPA-AKAP5 complex, β_2 AR

binds directly and GluA1 binds indirectly to PSD-95, through its oligomerization with the TM AMPAR regulatory protein stargazin; PSD-95 then couples to AKAP5 bringing AC5 and PKA to form the larger complex (Joiner et al., 2010; Zhang et al., 2013) (Fig. 3C). The AMPA-stargazin complex can also indirectly interact with group I mGluRs by means of a PSD-95-GKAP-Shank-Homer link (Dunn & Ferguson, 2015; Tu et al., 1999; Zheng et al., 2011) (Fig. 3A). In summary, there is evidence for specificity in the interactions of structurally similar GPCRs for some PDZ domains (He et al., 2006; Li et al., 2013; Valentine & Haggie, 2011) and different AKAPs (Fan, Shumay, Wang, & Malbon, 2001; Gardner et al., 2006; Valentine & Haggie, 2011). However, those involve relatively few GPCRs, which likely does not account for the multiplicity of specific combinations of GPCRs, G protein subunits and PM-effectors expressed in different cells or even the same cell. In summary, although complementary, neither membrane nanodomains nor scaffold proteins and AKAPs can generate the specificity of combinations of GPCRs, G protein subunits and PM-effectors provided by GEMMAs.

4. GPCR oligomerization

4.1. Homomers and heteromers as common functional GPCR units

Initial evidence for GPCR homomerization came from radioligand binding studies, with the demonstration of ligand binding cooperativity. This phenomenon has been recognized for many years and reproduced in many experimental preparations, including experiments with membrane extracts from multiple native tissues, mammalian transfected cells and other artificial systems, including GPCRs reconstituted in detergent micelles, liposomes (phospholipid vesicles with or without different proportions of cholesterol and integral membrane proteins) or

nanodiscs (phospholipid bilayer preparations held together by membrane scaffold proteins) (reviewed in Ferré et al., 2014, 2020). Additional evidence for GPCR homomerization came with pioneering studies with BRET and FRET techniques (reviewed in Bouvier, 2001; Milligan, 2001; Guo et al., 2017; El Khamlichi et al., 2019; Soave et al., 2020). This was followed by a large number of studies using these and other biophysical techniques in living cells, such as FRAP (Dorsch, Klotz, Engelhardt, Lohse, & Bünemann, 2009), TR-FRET (Cottet et al., 2012) and with single molecule-based methods, with the analysis of single fluorescence-labeled receptor molecules by fluorescence autocorrelation and cross-correlation spectroscopy (FCS and FCCS; Herrick-Davis, Grinde, Cowan, & Mazurkiewicz, 2013; Teichmann et al., 2014) and SMT analysis by TIRF (Calebiro et al., 2013; Scarselli et al., 2016). Single molecule-based methods in transfected cells also provided evidence for a differential degree of oligomerization and stoichiometry of GPCR oligomers versus monomers, depending on the type and density of GPCR (Calebiro et al., 2013; Scarselli et al., 2016; Teichmann et al., 2014). GPCR type-dependent extent of oligomerization was also recently reproduced using fluorescence-microscopy-based techniques in liposome preparations (Walsh et al., 2018).

Biophysical techniques have also been fundamental in the study of GPCR heteromers (Bourque et al., 2020; Ferré et al., 2014, 2020; Gaitonde & González-Maeso, 2017; Gomes et al., 2016; Sleno & Hébert, 2018). Furthermore, several studies also using biophysical techniques, dominant negative mutant GTPases and other complementary strategies, strongly supported the pre-assembly of not only family C (Margeta-Mitrovic, Jan, & Jan, 2000), but also family A GPCR homomers and heteromers in the ER (Décaillot, Rozenfeld, Gupta, & Devi, 2008; Dupré et al., 2006; Herrick-Davis, Weaver, Grinde, & Mazurkiewicz, 2006; Salahpour et al., 2004).

The application of techniques that selectively disrupt specific oligomeric interfaces have not only provided the means to establish the quaternary structure of GPCR oligomers, but also their emergent functional and pharmacological properties, as compared with parent monomers. Those techniques include mutating key residues, using chimeras, cysteine cross-linking techniques, or using synthetic peptides with the amino acid sequence of putative interacting domains, for instance of specific TM domains (TM peptides) (Ferré et al., 2014). Disrupting TM peptides were initially introduced to investigate the TM interface of the β_2 AR homodimer in co-immunoprecipitation experiments, suggesting the involvement of TM6 (Hébert et al., 1996). More recently, the TM-peptide strategy has been applied to BiFC experiments to identify the interfaces of several GPCR homodimers and heteromers (Guitart et al., 2014; Guitart et al., 2019; Köfalvi et al., 2020; Navarro et al., 2018; Rivera-Oliver et al., 2019). With this approach, TM6 was also found to be involved in the interface of adenosine A_{2A} receptor ($A_{2A}R$) and dopamine D_2R (D_2R) homodimers (Navarro et al., 2018). The involvement of TM6 in D_2R homomerization was further substantiated by selective ability of peptides derived from TM6 of D_2R to completely counteract the decrease in binding of a bivalent ligand (see below) with selective picomolar affinity for the D_2R homodimer (Pulido et al., 2018). TM6 has also been shown to be part of the homomeric interface of high-resolution crystallographic structures of several GPCRs, including CXC_4R and MOR (Manglik et al., 2012; Wu et al., 2010) (Fig. 2B and C). In addition, using the TM-peptide-BiFC approach, TM6 was found to be involved in several interfaces of GPCR heteromers, including adenosine A_1R -dopamine D_1R , dopamine D_1R - D_3R and adenosine $A_{2A}R$ -cannabinoid CB_1R heteromers (Guitart et al., 2014; Guitart et al., 2019; Köfalvi et al., 2020; Rivera-Oliver et al., 2019).

The fact that rearrangement of TM6 constitutes a critical ligand-induced conformational change that determines G protein activation and modulation of ligand affinity (Dupré et al., 2007), provides a framework for understanding allosteric communication through protomers of GPCR homomers or heteromers with a TM6 interface (Ferré et al., 2020; Navarro et al., 2018). Nevertheless, as reviewed elsewhere (Ferré et al., 2014), TM6 is not always involved in GPCR homomer or heteromer

interfaces (see for instance the alternative interfaces obtained with the β_1AR and MOR crystal structures, shown in Fig. 2C and D). In some cases, the homomeric interface can change in the presence of an agonist, as shown for mGluRs. Recent results obtained with the cryo-EM structures of the full-length mGlu $_5R$ shows a TM5-TM5 orientation in its inactive state, but with substantial separation between the seven TM domains. Activation of the receptor leads to a 20° rotation of each seven TM domain which results in a close contact TM6-TM6 homodimeric interface (Fig. 2A) (Koehl et al., 2019). Nevertheless, the disulfide bridge is not the only determinant of homodimerization in the inactive state of mGlu $_2R$, which shows extensive interactions along the whole length of TM4 (Du et al., 2021). Agonist binding to the mGlu $_2R$ is then associated with a TM interface change mainly contributed by TM6 (Du et al., 2021). Finally, the same GPCR can utilize different TMs in their heteromeric interactions with different GPCRs and it can also display different homomeric interfaces when forming heteromers with different GPCRs (heteromers of homodimers; see below and Köfalvi et al., 2020).

4.2. Allosteric interactions of GPCR heteromers

There is now a long list of putative GPCR receptor heteromers (Gaitonde & González-Maeso, 2017; Gomes et al., 2016), discussion of which is beyond the scope of this review. Here, we will only summarize findings for the $A_{2A}R$ - D_2R heteromer, a paradigmatic example that demonstrates the different types of allosteric interactions in the context of a GPCR heteromer which constitutes the core of a prototype GEMMA. Allosteric interactions constitute a common property of GPCR heteromers (Ferré et al., 2009, 2014, 2020). Importantly, these allosteric properties provide a framework for translational significance, for their potential role as therapeutic targets.

Allosterism is currently defined as “the process by which the interaction of a chemical or protein at one location on a protein or *macromolecular complex* (the allosteric site) influences the binding or function of the same or another chemical or protein at a topographically distinct site” (Smith & Milligan, 2010). Here we suggest three types of allosterism that can be identified in GPCR heteromers: type I allosterism, or allosterism between ligands, which entails the ability of an orthosteric ligand of one protomer in the GPCR heteromer to modify the affinity or efficacy of an orthosteric ligand of the other molecularly distinct protomer (Fig. 4A); type II allosterism, or ligand-independent allosterism, where heteromerization *per se* modifies the properties of a specific orthosteric ligand for one of the protomers of the GPCR heteromer, independent of ligand binding to the other molecularly distinct protomer (Fig. 4B); and type III allosterism, or allosterism through the effector, where the effector (PM-effector of a GEMMA) acts as an interface for the interactions between orthosteric ligands of the molecularly distinct protomers of a GPCR heteromer (Fig. 4C). Type I and III allosterisms also include the possibility of transactivation or transinhibition, by which an orthosteric ligand of one of the protomers leads to an increase or decrease in the intrinsic activity of the other molecularly different protomer.

The most described type I allosteric interaction in GPCR heteromers involves an agonist or an antagonist of one of the protomers that inhibit the affinity or efficacy of an agonist of the other molecularly distinct protomer. Such interactions are usually referred as ‘negative crosstalk’ or ‘cross-antagonism’, respectively, and their molecular mechanisms are beginning to be understood (Ferré et al., 2014, 2020) (Fig. 5). Since the first report in 1991 (Ferré, von Euler, Johansson, Fredholm, & Fuxe, 1991), $A_{2A}R$ ligands have been repeatedly reported to antagonistically modulate the binding properties of D_2R ligands in membrane preparations from transfected mammalian cells and native tissues (reviewed in Ferré et al., 2018). Demonstration of the dependence of these ligand interactions on the integrity of the $A_{2A}R$ - D_2R heteromer (with disruptive synthetic peptides or mutations of key residues of the heteromeric interface) (Bonaventura et al., 2015;




Fig. 4. Allostery in GPCR heteromers. (A) Type I allostery, in which a ligand (red spheres) binding to one protomer (green cylinders) in the GPCR heteromer can modify the affinity (ligand binding depicted by the red arrow) or the efficacy (receptor activation depicted by the wide orange arrow) of another ligand (orange spheres) binding to the partner receptor (gray cylinders) via their TM helices (see Fig. 5). (B) Type II allostery, or ligand-independent allostery, in which the affinity or efficacy of a ligand (orange spheres) for a GPCR (gray cylinders) can be modified just by heteromerization with a molecularly different GPCR (green cylinders). (C) Lateral view (left) and extracellular view (right) of a computer model of a GEMMA including two heterotetramers composed of two different GPCR homomers (one represented by white cylinders and grey surfaces and the other represented by green cylinders and surfaces), AC (light blue cylinders), Gs (brown for G α s and yellow for G β γ) and Gi (red for G α i and yellow for G β γ). This type of GEMMA, based on the A_{2A}R-D₂R-AC5 GEMMA described in Navarro et al. (2018), can explain the ability of a Gi-coupled GPCR to counteract AC activation mediated by a Gs-coupled GPCR (type III allostery). The proposed contact between the CT domain of the receptor and the G β subunit in the inactive state (Tsai et al., 2019) is shown by a colored line. To facilitate visualization of all protomers of the GPCR oligomers, one of the protomers is represented by cylinders and the other protomers are represented as colored surfaces (with different shades of the same color for the same GPCR). Also for the purpose of simplification, only both possible positions of the G α s and G α i subunits are shown: in their G β γ -associated and receptor-bound state (without surface) and in their G β γ -dissociated and AC-bound state (with surface). The agonist-induced dissociation of G α s and G α i from their respective G β γ takes place within the framework of the GEMMA. As explained in the text, the active state of this type of GEMMA, where both G α s and G α i subunits are bound to the respective catalytic domains of AC, is associated with a rearrangement of G β γ subunits and a change in the interfaces of the heterotetramers with AC.

Borroto-Escuela et al., 2010; Borroto-Escuela et al., 2010) implied a type I allostery in the A_{2A}R-D₂R heteromer. This allostery determines the therapeutic effects of A_{2A}R antagonists in Parkinson's disease, which target A_{2A}R-D₂R heteromers localized in striatopallidal neurons (for recent review, see Ferré et al., 2018). Significantly, the selective A_{2A}R antagonist istradefylline, in combination with L-DOPA, is the first non-dopaminergic drug approved by FDA for the treatment of Parkinson's Disease (Chen & Cunha, 2020).

Type II allostery has been described in the A_{2A}R-D₂R heteromer involving a specific A_{2A}R antagonist (Fig. 4B). When comparing the binding properties of several selective orthosteric A_{2A}R antagonists (including istradefylline) in mammalian transfected cells, only one compound, SCH442416, showed a selective low affinity for A_{2A}R when co-expressed with D₂R, as compared with A_{2A}R when expressed alone or co-expressed with A₁R (Orrú et al., 2011). The specific behavior of SCH442416 in the A_{2A}R-D₂R heteromer could also be demonstrated in the mouse striatum, but not in mice with conditional genetic deletion

of striatal D₂R (Ferré et al., 2018). These results represented proof of concept for the possibility of using GPCR heteromers as targets for specific ligands. Thus, it could be demonstrated that SCH442416 is a preferential presynaptic striatal A_{2A}R antagonist, because of the preferential pre-synaptic localization of A₁R-A_{2A}R heteromers in striatal glutamatergic terminals, as compared with the post-synaptic striatal localization of A_{2A}R-D₂R heteromers in striatopallidal neurons (Ferré et al., 2018). The preferential striatal presynaptic effect of SCH442416 was suggested to determine its ability to reduce cannabinoid self-administration in monkeys and its possible application to cannabinoid use disorder (Justinová, Redhi, Goldberg, & Ferré, 2014).

Apart from the ability of A_{2A}R ligands to allosterically modulate D₂R ligands, a reciprocal canonical Gs-Gi protein-dependent antagonistic interaction was discovered early on, where D₂R agonists, by activating Gi-coupled D₂R, could oppose Gs-coupled, A_{2A}R-mediated activation of AC-cAMP-PKA signaling (Kull et al., 1999). The balance between antagonistic allosteric and Gs-Gi-AC interactions determines the




Fig. 5. Mechanisms of type I allosterism in GPCR oligomers heteromers. (A) Cross-section through a prototypical class A GPCR, highlighting the agonist (white sticks), the amino acids of the orthosteric site (delineated by a white line) involved in ligand binding (orange), the conserved PIF motif (yellow), and the NPxxY and DRY motifs and Y, (yellow) that transmit the signal from the PIF motif (transmission switch) to the G protein site (Weis & Kobilka, 2018). (B) Position of these amino acids involved in ligand binding (orange spheres) and signal transmission (yellow) in models of GPCR dimers via TM5 and TM6 (top) and TM4 and TM5 and intracellular loop 2 (bottom). Clearly, a ligand-bound (type I allosterism) or ligand-free (type II allosterism) or orphan GPCR (green cylinders) can modify the affinity (orange spheres) or efficacy (yellow spheres) of the partner receptor (gray) via the protein-protein interface (blue mesh). (C) GPCRs are dynamic proteins that adopt, in a ligand-free form, a number of conformations (shades in gray) that not only involve the extracellular and intracellular sites (Weis & Kobilka, 2018), but also a potential TM oligomeric interface. An inverse agonist (red polygon) binding to one of the protomers of a GPCR heteromer triggers a high surface complementarity between TM5 and TM6 of the two molecularly different protomers, via the four-helix bundle (dashed red circle) (Manglik et al., 2012), which blocks the opening of the intracellular cavity for G protein binding at the other protomer (cross-antagonism) (Viñals et al., 2015). An agonist (green polygon) binding to one of the protomers triggers the outward movement of TM 6 (see arrows), opening the intracellular cavity for G protein binding (Rasmussen et al., 2011), whereas the TM5 and TM6 heteromeric interface impedes simultaneous agonist-induced movement of both TM6 (negative crosstalk) due to a steric clash (red cross) (Guinart et al., 2020).

final output of $A_{2A}R$ - D_2R heteromer-dependent signaling in the striatopallidal neuron (Ferré et al., 2018). Importantly, the Gs-Gi-AC interaction depends on the integrity of the $A_{2A}R$ - D_2R heteromer, since the Gi-dependent regulation of Gs-mediated AC activation could also be disrupted by TM peptides that disrupt the heteromer (Navarro et al., 2018). This discovery represents a significant shift in our understanding of crosstalk between GPCR signaling pathways, since such interactions at the effector level were previously understood as related to independent changes in second messenger levels (Zoli et al., 1993). GPCR heteromer-dependent Gs-Gi-AC interactions have been demonstrated for several other GPCR heteromers, including A_1R - D_1R , D_1R - D_3R and $A_{2A}R$ - CB_1R heteromers (Guitart et al., 2019; Köfalvi et al., 2020; Rivera-Oliver et al., 2019). In addition to GPCR heteromerization, the Gs-Gi-AC interaction suggests simultaneous respective interaction of $G_{\alpha s}$ and $G_{\alpha i}$ subunits with the C2 and C1 catalytic domains of the same molecule of AC, more specifically with group III ACs, the AC5 and AC6 isoforms. This depends on a pseudo-symmetrical arrangement of the C1 and C2 domains (Dessauer, Tesmer, Sprang, & Gilman, 1998). Another mechanism for antagonistic interactions between Gs- and Gi-coupled receptors can also occur with group I ACs, AC1, AC3 and AC8, involving interactions with $G\beta\gamma$ rather than the $G_{\alpha i}$ subunits of an activated Gi protein (Steiner, Saya, Schallmach, Simonds, & Vogel, 2006; Taussig, Quarmby, & Gilman, 1993).

The need for heteromerization of a Gs-coupled and a Gi-coupled GPCR, for oligomerization of those GPCRs with AC, and for the simultaneous binding of Gs and Gi to the C2 and C1 catalytic domains of AC, strongly suggests that a canonical antagonistic Gs-Gi-AC interaction should most commonly occur within GEMMAs that include a GPCR heteromer, Gs and Gi proteins and the PM-effector AC. Following the definition of allosterism, we can consider the antagonistic Gs-Gi-AC

interaction as an allosteric interaction of a GPCR heteromer, as part of a macromolecular complex that includes a PM-effector, which we labeled type III allosterism (Fig. 4C).

4.3. GPCR heterotetramers

The relatively large size of the heterotrimeric G protein makes simultaneous binding of more than one G protein to a GPCR unlikely. Therefore, the pentameric structure consisting of a GPCR homodimer and one heterotrimeric G protein is likely the primary functional unit, as initially supported by studies with the detergent-solubilized leukotriene B_4 receptor BLT1 (BLT₁R) (Banères & Parello, 2003). The minimal functional quaternary structure of a GPCR heteromer able to sustain a Gs-Gi-AC interaction would then be tetrameric, including Gs- and Gi-coupled homomers, or a Gs-Gi-coupled heterotetramer, as previously hypothesized (Ferré, 2015). With computational analysis of information provided from TM-peptide-BiFC experiments and several reported crystallographic structures of GPCRs alone or bound to G protein partners, it was possible to infer an optimal quaternary structure of the $A_{2A}R$ - D_2R heterotetramer: a linear structure with the two internal protomers involved in the heteromer interaction (via symmetrical TM4-TM5/TM5-TM4 interfaces) and the two external protomers of each homodimer (with a TM6/TM6 interface) coupled to their respective G proteins (Navarro et al., 2018) (Fig. 6A). Since AC5 and Golf are the major striatal AC and Gs family isoforms (Schwindinger et al., 2003, 2010; Hervé, 2011; Xie et al., 2015), evidence for $G_{\alpha\beta\gamma}$ -AC5 pre-assembly (see above) suggests that $G_{\alpha(olf)\beta\gamma}$ can indirectly pre-assemble AC5 with the $A_{2A}R$ - D_2R heterotetramer, forming a $A_{2A}R$ - D_2R -AC5 GEMMA, providing an example of an allosteric machine that integrates types I, II and III allosteric mechanisms (Fig. 6A).

Following the definition of GEMMA, the same study (Navarro et al., 2018) also provides, to our knowledge, the first non-equivocal demonstration of a ligand-dependent rearrangement of a complex including a GPCR or a GPCR oligomer and PM-effector. Using TM-peptide-BiFC experiments it could be demonstrated that specific TM domains of the $A_{2A}R$ and the D_2R directly interact with the TM domains of AC5, and that those interfaces changed upon the binding of agonists (Navarro et al., 2018). It could also be shown that disruption of the GPCRs-AC5 interfaces also disrupted the Gs-Gi-AC type III allosteric interaction within the $A_{2A}R$ - D_2R -AC5 GEMMA (Navarro et al., 2018). The quaternary structure of the $A_{2A}R$ - D_2R -AC5 GEMMA obtained by computer modeling suggested that two molecules of AC5 would be needed to interact simultaneously with $A_{2A}R$ and D_2R protomers in a single heterotetramer, while two heterotetramers are needed to allow the simultaneous interaction of $A_{2A}R$ and D_2R with a single AC5 molecule, to allow full Gs-Gi-AC interactions (Fig. 6A). According to this model, the $A_{2A}R$ - D_2R heterotetramer is the more rigid part of the complex and the model also predicts the possible extension of this basic unit into a higher-order GEMMA (Navarro et al., 2018) (Fig. 6A).

The mGlu₂R-5-HT_{2A}R heteromer is another example of a well-studied GPCR heteromer with evidence of both type I and II allosteric

mechanisms. It is localized in cortical glutamatergic pyramidal neurons and represents an important target for the hallucinogenic effects of 5-HT_{2A}R agonists and the antipsychotic effect of both 5-HT_{2A}R antagonists/inverse agonists and mGlu₂R agonists (Baki et al., 2016; Fribourg et al., 2011; González-Maeso et al., 2008; Moreno et al., 2012; Moreno et al., 2016). Its quaternary structure also seems to be tetrameric, constituted of mGlu₂R and 5-HT_{2A}R homodimers respectively coupled to Gi and Gq, with a clear involvement of TM4 in the heteromeric interface (Moreno et al., 2016; Shah, Toneatti, Gaitonde, Shin, & González-Maeso, 2020). In transfected mammalian cells expressing mGlu₂R-5-HT_{2A}R heteromers, a high-efficacy orthosteric mGlu₂R agonist promoted Gq protein-mediated increases in intracellular Ca²⁺ concentration in the absence of 5-HT_{2A}R ligands (Moreno et al., 2016). By using different mGlu₂R-mGlu₃R chimeras and mutant receptors unable to couple to their respective G proteins, it was shown that this pharmacological response involved mGlu₂R agonist-induced transactivation of 5-HT_{2A}R-Gq-mediated signaling in the mGlu₂R-5-HT_{2A}R heteromer (Moreno et al., 2016). It was also shown that transactivation required not only obligatory homomerization of mGlu₂R and the Gq protein coupling to the 5-HT_{2A}R, but also G α i coupling to the mGlu₂R (Moreno et al., 2016).




Fig. 6. The $A_{2A}R$ - D_2R heterotetramer-AC5 and mGlu₂R-5-HT_{2A}R-PLC-Kir3 GEMMAs. (A) The $A_{2A}R$ - D_2R heterotetramer-AC5 GEMMA, constituted by $A_{2A}R$ homomers coupled to Gs, D_2R homomers coupled to Gi and AC5, can be considered as a prototype of GEMMA that integrates canonical antagonistic Gs-Gi-AC interactions. (B) The mGlu₂R-5-HT_{2A}R-PLC-Kir3 GEMMA, constituted by mGlu₂R homomers coupled to Gi, 5-HT_{2A}R homomers coupled to Gq, PLC and Kir3, can be considered as a prototype of GEMMA that integrates canonical antagonistic Gi-Gq-Kir3 interactions.

Kir3 channels were also identified in transfected mammalian cells as a common PM-effector for both Gi and Gq protein-mediated signaling of the mGlu₂R-5-HT_{2A}R heteromer (Baki et al., 2016). As mentioned above, Kir3 channels are activated by direct interactions with Gβγ subunits following activation of Gi-coupled receptors (Dascal & Kahanovitch, 2015; Logothetis et al., 1987; Logothetis et al., 2015; Riven et al., 2006). However, stimulation of Kir3 channels is critically dependent on phosphatidylinositol 4,5-bisphosphate (PIP₂) (Huang, Feng, & Hilgemann, 1998; Sui, Petit-Jacques, & Logothetis, 1998). A canonical effect of Gq activation is PLC activation, which leads to PIP₂ hydrolysis and opposes Gβγ-mediated activation (Logothetis et al., 2015). When added to evidence for pre-assembly of Gαβγ with Kir3 (see above) and with PLC (Srncka & Fisher, 2019; Yuan et al., 2007), we suggest the existence of a mGlu₂R-5-HT_{2A}R-PLC-Kir3 GEMMA (Fig. 6B). This GEMMA would represent an allosteric machine that controls neuronal excitability by integrating antagonistic effects of serotonin and glutamate on Kir3 activity. As with the A_{2A}R-D₂R-AC5 GEMMA, it might be predicted that the mGlu₂R-5-HT_{2A}R-PLC-Kir3 GEMMA could be the basic unit of a higher-order GEMMA (Fig. 6B).

4.4. Oligomerization of GPCRs with orphan GPCRs, truncated GPCRs and other TM proteins

From the 800 GPCR genes identified, more than 100 GPCRs remain orphans, including 86 family A GPCRs (Davenport et al., 2013; Ngo et al., 2016). There is evidence indicating that some GPCRs will not be deorphanized because they may not have a natural ligand. Rather, they can oligomerize with other GPCRs and change their functional and pharmacological properties via allosteric interactions. This concept was originally suggested from studies with melatonin receptors and the orphan receptor GPR50 (Levoye, Dam, Ayoub, Guillaume, & Jockers, 2006). The melatonin receptor subfamily consists of MT₁R and MT₂R, with melatonin as their natural ligand, and the orphan receptor GPR50, which does not bind to melatonin or any other known endogenous ligand to date (Cecon, Oishi, & Jockers, 2018). MT₁R and MT₂R signal preferentially through Gi/o protein coupling, while GPR50 does not couple to G proteins and has a characteristic very long CT (~300 amino acids) (Oishi, Cecon, & Jockers, 2018). Several studies support the ability of the melatonin receptor family to oligomerize, including heteromerization of GPR50 with MT₁R and MT₂R (Levoye et al., 2006; Oishi et al., 2018) (Fig. 7A). Experiments using BRET in transfected mammalian cells suggested that MT₁R and MT₂R form constitutive homomers as well as MT₁R-MT₂R heteromers (Ayoub et al., 2002). In transfected cells the propensity for homomer and heteromer formation differs between both receptor subtypes, with a lower propensity of MT₂R to form homomers as compared with the MT₁R and a higher propensity of forming MT₁R-MT₂R heteromers, suggesting that the MT₂R preferentially exists as a heteromeric complex with MT₁R (Ayoub, Levoye, Delagrèze, & Jockers, 2004). In fact, by inducing expression of tagged MT₁R and MT₂R in genetically modified mice, it was possible to demonstrate the existence of MT₁R-MT₂R complexes in the retina, where they mediate the effect of melatonin on light sensitivity of rod photoreceptors (Baba et al., 2013). Significantly, in transfected mammalian cells, MT₁R-MT₂R heteromers seem to preferentially couple to Gq and promote Gq-PLC-PKC signaling, a primary signaling pathway involved in the functional effects of melatonin in rod photoreceptors (Baba et al., 2013) (Fig. 7A). GPR50 promotes a decreased ability of melatonin to bind and signal through the MT₁R, which depends on the ability of the long GPR50 CT to alter the pre-coupling or rearrangement of G proteins to MT₁R in the MT₁R-GPR50 heteromer (Levoye, Dam, Ayoub, Guillaume, Couturier, et al., 2006; Oishi et al., 2018) (Fig. 7A).

Some GPCRs exist in different splice variants, which can be expressed by the same cell and, therefore potentially oligomerize leading to distinct functional outcomes. Some of these variants represent truncated versions of receptors, missing several TM domains, often

resulting in their inability to bind to endogenous ligands *per se* (Wise, 2012). Truncated GPCR isoforms may be expressed at the plasma membrane and constitute components of GEMMAs. Those include D₃nf and GHS_{1b}, truncated splice variants of dopamine D₃ receptor (D₃R) and ghrelin GHS_{1a} receptor GHS_{1a}R, respectively. Both isoforms lack TM6 and TM7 domains, do not bind ligands or promote signaling themselves, but oligomerize with their respective functional isoforms (Chow et al., 2012; Elmhurst, Xie, O'Dowd, & George, 2000; Karpa, Lin, Kabbani, & Levenson, 2000; Leung et al., 2007; Mary et al., 2013; Navarro et al., 2016). Although under some experimental conditions both isoforms seem able to promote intracellular localization of their functional GPCR partners, there is evidence for the ability of D₃R-D₃nf and GHS_{1a}R-GHS_{1b} oligomers to localize to the plasma membrane, where the truncated partner exerts ligand-independent negative allosteric modulation (type II allosterism) of D₃R and GHS_{1a}R ligands (Elmhurst et al., 2000; Mary et al., 2013; Navarro et al., 2016). In addition, BRET and signaling experiments in transfected mammalian cells indicate that GHS_{1a}R-GHS_{1b}R can be part of a GEMMA that includes D₁R, in which ghrelin, instead of promoting its preferred GHS_{1a}R-dependent Gq-PLC signaling, activates a Gs-AC signaling, which is opposed by D₁R antagonists (Navarro et al., 2016) (Fig. 7B).

Receptor activity-modifying proteins (RAMPs) are an example of membrane-spanning accessory proteins that can alter the function of GPCRs. A small family of three proteins (RAMP1, RAMP2 and RAMP3) has substantial capacity for introducing functional diversity by directly interacting with GPCRs (Hay & Pioszak, 2016; Serafin, Harris, Nielsen, Mackie, & Caron, 2020). Structurally, RAMPs comprise a single TM domain with a long extracellular NT (90–100 amino acids) and a short intracellular CT (9 amino acids). The interactions of RAMPs with the family B GPCR calcitonin-like receptor (CLR) and the calcitonin receptor (CTR) are the most extensively studied and provide a picture of the broad influence that a single TM-domain protein can have when interacting with a GPCR (Fig. 7C). RAMPs are required to chaperone CLR to the cell surface, where RAMP-CLR complexes act as receptors for the peptide hormones calcitonin gene-related peptide (CGRP), adrenomedullin (AM) or adrenomedullin 2/intermedin (AM2), depending on the subtype of RAMP co-expressed (McLatchie et al., 1998; Serafin et al., 2020). Thus, the RAMP subtype determines which endogenous ligands preferentially bind to CLR and CLR-RAMP1, CLR-RAMP2 and CLR-RAMP3 constitute CGRP, AM and AM2 receptors, respectively (McLatchie et al., 1998; Poyner et al., 2002; Serafin et al., 2020) (Fig. 7C). CTR is the closest relative to CLR and it can reach the cell surface in the absence of RAMP, but CTR can drive RAMP translocation to the cell surface and CTR pharmacology is altered in the presence of RAMP, which increases the affinity of the endocrine hormone amylin (AMY), such that three amylin receptor subtypes (AMY_{1–3} receptors) result from each respective RAMP-CTR complex (Poyner et al., 2002).

5. Targeting specific GEMMAs

5.1. Specific GPCR ligands for individual GEMMAs

The main properties of GPCR oligomers (allosterism), orphan receptors, truncated GPCRs and other membrane-localized GPCR-associated proteins (such as RAMPs) can potentially provide significant functional and pharmacological properties to the GEMMAs that include them. The remarkable varied influence of TM RAMPs in modifying the properties of ligand binding to CLR and CTR, provides a clear demonstration for exploiting GEMMAs as targets for drug discovery. Namely, that different components of GEMMAs directly interacting with a particular GPCR can potentially modify the effects of specific ligands for this GPCR. Another reason is that localization of the same GPCR with additional components in different cells likely creates a new and unique GEMMA that can be targeted in a way distinct from the same receptor in a different complex. This raises the possibility of obtaining ligands with selectivity for the GPCRs in distinct GEMMAs which could mediate desired




Fig. 7. Oligomerization of GPCRs with orphan GPCRs, truncated GPCRs and RAMPs. (A) Homomers and heteromers of MT₁R, MT₂R and GPR50. MT₂R preferentially exists as a heteromeric complex with MT₁R and MT₁R-MT₂R heteromerization drives a change in the preference of G protein subtype coupling, from Gi to Gq. The long CT of GPR50 significantly alters MT₁R-Gi protein signaling in the MT₁R-GPR50 heteromer (see text). (B) GHS1aR forms homomers and oligomerizes with its non-functional truncated isoform GHS1bR to form homomers and heteromers. One of the properties of the GHS_{1a}R-GHS_{1b}R oligomers is the facilitation of an additional interaction with the D₁R, leading to a change in the preference of G protein subtype coupling of GHS_{1a}R, from Gq to Gs. (C) The single TM domain proteins RAMP1, RAMP2 and RAMP3 associate with CLR and form the Gs protein-coupled receptors CGRPR, AM₁R and AM₂R, respectively, determining the ligands potentially binding to CLR.

therapeutic effects of the putative ligand, while avoiding targeting GEMMAs that mediate the adverse side effects.

Type II allosterism of GPCR heteromers and the changes it brings to pharmacological properties is noted with the example of A_{2A}R heteromers discussed above. A significant clinically translational type II allosterism of GPCR heteromers for substance use disorders has been generated by the study of MOR heteromers. Since the pioneering studies from Lakshmi Devi's research group, MOR oligomerization has been the focus of a significant amount of experimental work, providing substantial evidence for MOR- δ opioid receptor (DOR) heteromerization and their role in the analgesic effects of opioids (Fujita, Gomes, & Devi, 2015). Although the initially proposed localization of MOR-DOR heteromers in the nociceptive sensory neurons of the dorsal root ganglia and dorsal horn of spinal cord was challenged (Scherrer et al., 2009), subsequent studies provided strong confirmatory evidence (Tiwari et al., 2020; Wang et al., 2010; Yekkirala et al., 2012). More recently, a major population of MOR localized in the mesencephalon, in the ventral tegmental area, has been shown to form heteromers with galanin Gal₁ receptors (Gal₁R) and to mediate the ability of opioids to activate the dopaminergic system and therefore abuse liability (Cai et al., 2019; Moreno et al., 2017). These studies support the rationale of selectively targeting MOR-DOR heteromers in the search for new effective analgesic drugs devoid of addictive properties.

Different approaches have been shown to be effective pre-clinically, including high-throughput screening of small-molecule libraries in cells expressing MOR-DOR heteromers, leading to the discovery of CYM51010, which showed a significant increase in its potency and

efficacy to promote G protein activation mediated by the MOR in heteromers with DOR (Gomes et al., 2013). CYM51010 exhibited a potent antinociceptive activity with reduced tolerance potential compared to morphine in mice (Gomes et al., 2013) and, more recently, CYM51010 was also found effective in a rat model of neuropathic pain (Tiwari et al., 2020). Because of structural similarities of CYM51010 with carfentanyl, structure-activity relationship (SAR) analysis was used to develop several carfentanyl derivatives (Faouzi et al., 2020). One of the derivatives, MP135, showed even higher relative efficacy at the MOR-DOR heteromer versus MOR alone, as compared with CYM51010 (Faouzi et al., 2020). As expected, in rodents, it produced pronounced analgesic effects, but still showed rewarding properties and was readily self-administered (Faouzi et al., 2020). Although not tested yet, MP135 might also exhibit significant potency and efficacy at the MOR-Gal₁R heteromer, which determines the dopaminergic/euphoric effects of opioids (Cai et al., 2019).

The effect of morphine, methadone and fentanyl, representatives of the three principal opioid structures, has been separately analyzed for both MOR-DOR and MOR-Gal₁R heteromers (Cai et al., 2019; Yekkirala et al., 2012; Yekkirala, Kalyuzhny, & Portoghese, 2010). The research group of Philip Portoghese found evidence for a significantly higher efficacy of the three opioids for MOR-DOR versus MOR alone (Yekkirala et al., 2010; Yekkirala et al., 2012). In a more recent study, for the first time, differential pharmacodynamic effects were observed for methadone versus morphine or fentanyl, a significant decrease in potency which depended on heteromerization of MOR with Gal₁R (Cai et al., 2019). As predicted from the predominant role of MOR-Gal₁R heteromers in the modulation of the dopaminergic system,

methadone was a much weaker activator of the dopaminergic system as compared with morphine and fentanyl and, it could be demonstrated clinically that it was also much weaker at producing euphoric effects (Cai et al., 2019). Elucidation of the molecular mechanisms of this type II allosterism in the MOR-Gal₁R heteromer should facilitate discovery of methadone-like compounds, with strong analgesic and reduced addictive properties. An additional challenge is to determine specific components of the MOR-containing GEMMAs responsible for the other major unwanted effects of opioids, such as respiratory depression and constipation.

5.2. Simultaneous targeting of different components of specific GEMMAs

Another approach to obtain opioid analgesia with low abuse liability would be using effective MOR-DOR ligands while selectively blocking the unwanted MOR signaling via the MOR-Gal₁R heteromer. This could be achieved by administering Gal₁R ligands, based on recently described type I allosterism in the MOR-Gal₁R heteromer, by which those ligands significantly reduce the affinity and efficacy of MOR agonists (Cai et al., 2019). Gal₁R are also localized in the spinal cord, where previous studies indicate their activation produces analgesic effects synergistic to those of opioids (Hua et al., 2004). Since the only available selective and potent Gal₁R ligands are galanin-derived peptides (Freimann, Kurrikoff, & Langel, 2015), the discovery of small molecules with significant brain penetrability targeting Gal₁R could provide an important new approach for the treatment of pain. Gal₁R agonists could be co-administered with lower therapeutic doses of MOR agonists, promoting analgesia and significantly reducing side effects, particularly the dopaminergic effects mediated by MOR-Gal₁R heteromers.

Type I allosterism in GPCR heteromers could then be used as a therapeutic strategy to target specific GEMMAs, either by specifically increasing the therapeutic effects or decreasing the unwanted effects of GPCR ligands. This is illustrated by the increase in the therapeutic index of L-DOPA by A_{2A}R antagonists in Parkinson's Disease (see above), which decreases the effective dose of L-DOPA required and, therefore, both short- and long-term side effects (Chen & Cunha, 2020).

A further step in drug discovery would be to obtain bivalent compounds that can simultaneously bind both protomers in the GPCR heteromer, defined as single chemical entities composed of two pharmacophore units covalently linked by an appropriate spacer. These ligands are designed to interact simultaneously within a GPCR homo- or heterooligomer, to enhance affinity and subtype selectivity and, in case of a GPCR heteromer, promote type I allosteric modulation (Daniels et al., 2005; Pulido et al., 2018; Soriano et al., 2009). Although it could be argued that the pharmacologic effects of bivalent ligands are related to the separate binding of each pharmacophore to two different GPCR oligomers, true simultaneous bivalent binding to the same GPCR oligomer can be demonstrated by dependence on the integrity of the GPCR oligomer. For instance, in a recent study, the dramatic increase of affinity of a D₂R ligand (from nanomolar to picomolar affinity) obtained when two molecules of the same ligand were attached using an appropriate spacer length was reversed upon disruption of the D₂R homomeric interface (see above) (Pulido et al., 2018).

Portoghese's group pioneered studies with bivalent ligands and opioid receptor heteromers. Their studies revealed additional implications with potential therapeutic importance. Thus, MOR-DOR bivalent ligands with specific length spacers and a MOR agonist and a DOR antagonist as pharmacophores produced analgesia without the development of tolerance (Daniels et al., 2005). This effect was attributed to the bivalent ligand-induced stabilization ("bridging") of the MOR-DOR heteromer with concomitant blockade of DOR signaling (Daniels et al., 2005), which facilitates internalization of the MOR-DOR heteromer and, therefore, subsequent development of tolerance to MOR agonist-mediated analgesia (Gomes et al., 2013).

Finally, apart from simultaneously targeting two different GPCRs, there are also possibilities of simultaneously targeting a GPCR protomer

and another component of a specific GEMMA. This is best exemplified by the recent discovery of the specific activator of Kir3.1-Kir3.2 channels, GAT1508 (Xu et al., 2020). Kir3 channels are widely expressed in the brain and in the heart, although the localization of the Kir3.1-Kir3.4 channel is predominantly in the heart, whereas Kir3.1-Kir3.2 channels are more restricted to the central nervous system (CNS) (Dascal & Kahanovitch, 2015). GAT1508 represents the first specific activator of Kir3.1-Kir3.2 channels, making them CNS-selective. The mechanism of GAT1508-induced channel activation is related to strengthening channel-PIP₂ interactions (Xu et al., 2020). Importantly, at lower concentrations that are insufficient to produce significant activation of Kir3.1-Kir3.2 channels, GAT1508 significantly potentiated the ligand-induced activating effect of an associated GPCR (such as baclofen-induced activation of the GABA_B receptor), suggesting synergism in the allosteric effects of GAT1508 and Gβγ subunits in fostering channel-PIP₂ interactions (Xu et al., 2020). This provides further selectivity of the effects of GAT1508 in the brain and evidence was provided by a specific effect of the baclofen-GAT1508 in the basolateral amygdala, with possible translational implications for post-traumatic stress disorder (Xu et al., 2020).

5.3. G protein subtype-dependent functional selectivity

The properties of a GPCR ligand can also depend on specific pre-assembled G protein subtypes, as recently shown in BRET-based experiments. This also extends the notion of functional selectivity, where a selective ligand regulates a subset G protein-dependent or independent signaling to biased G protein subtype-dependent signaling (Violin & Lefkowitz, 2007). For instance, the effect of several D₁R ligands were compared in their ability to modify BRET in cells transfected with D₁R and Gαs or Gαolf fused to BRET chromophores. The D₁R agonist dihydroxidine (DHX) was found to be a full D₁R agonist with Gαs but a weak partial agonist with Gαolf (Yano et al., 2018). It was then expected that DHX would be more effective in the cerebral cortex, where Gαs is the predominant isoform, than in the striatum, where Gαolf predominates. In fact, with electrophysiological and behavioral experiments, it was demonstrated that DHX had greater efficacy in cortical versus striatal neurons. It was then suggested that DHX or other functionally Gαs subtype-functionally selective D₁R agonists could be used as pro-cognitive drugs with reduced extrapyramidal side effects (Yano et al., 2018).

The same phenomenon had been previously observed for other GPCRs and G protein subtypes, although it was not conceptualized as functional selectivity. For instance, the selective CB₁R agonist WIN55,212-2 facilitated Gq-mediated signaling, while other classes of CB₁R agonists (including Δ⁹-tetrahydrocannabinol and the endocannabinoids 2-arachidonoylglycerol and methanandamide) preferentially signaled through Gi/o proteins (Lauckner, Hille, & Mackie, 2005). Another example is the selective ability of the A₁R agonist 5'-N-cyclopentyl-carboxyamidoadenosine, as compared with a series of adenosine analogues, to facilitate Gs-mediated signaling, although A₁R preferentially signals through Gi/o proteins (Cordeaux, Ijzerman, & Hill, 2004). In those studies, the results were interpreted as dependent on a ligand-induced stabilization of a GPCR conformation that favored preferential coupling with a specific G protein subtype. This interpretation would be in line with collision-coupling. The GEMMAs concept provides another interpretation, the ability of ligands to differentially activate GPCRs pre-assembled with specific G protein subtypes.

6. Concluding remarks and future directions

The idea that GPCRs can serve as signaling platforms for assembly of macromolecular complexes of G proteins and other related signaling proteins to provide highly efficient and spatially restricted signaling events, with no requirement for G protein subunit dissociation and lateral diffusion within the membrane, has been recurrently considered

over the years (Neubig, 1994; Levitzki & Klein, 2002; Rebois & Hébert, 2003; Dupré et al., 2006, 2009; Hepler, 2014; Navarro et al., 2018; Sleno and Hébert, 2019). The GEMMA concept restricts the components of these macromolecular assemblies to GPCRs, G proteins, PM-effectors and other associated transmembrane proteins putatively pre-assembled prior to receptor activation by agonists, although we cannot ignore the idea that these core components of transmembrane signaling complexes will interact with additional cytosolic elements that will be recruited by the assembly upon GPCR activation, including GRKs and arrestins.

The GEMMA concept represents a divergent mechanism from collision coupling and both signaling modes are likely used by the cell, depending on the GPCRs and effectors involved and their need for providing more efficient and restricted signaling (GEMMA) or less efficient but more amplified signaling (collision-coupling mode). There is also evidence suggesting that these two poles are part of a spectrum of signaling mechanisms. Mixed models have been proposed where some of the components of the macromolecular membrane complex would still dissociate (Levitzki, 1986; Neubig, 1994). Such models can still explain the catalytic behavior of GPCRs observed in some cell systems, such as the amplification of signaling seen when one receptor activates numerous AC molecules, a phenomenon that, in fact, led to the formulation of the collision-coupling model (Tolkovsky & Levitzki, 1978a). On the other hand, amplification of signaling may not be expected with a ligand-induced rearrangement without dissociation of the GEMMA components. Nevertheless, as discussed, agonist-induced rearrangement of the G protein subunits of GEMMAs including the PM-effectors Kir3 or AC might necessarily imply limited dissociation between G protein subunits within the framework of the macromolecular assembly.

Many questions about GEMMAs remain to be answered, such as whether they can form higher-order macromolecular structures, as here suggested for the A_{2A}R-D₂R-AC5 GEMMA and the mGlu₂R-5-HT_{2A}R-PLC-Kir3 GEMMA. In that case, it will be important to resolve which basic units are pre-assembled in the ER and what determines and modulates the assembly and disassembly of the final macromolecular complex. It will also be important to establish the fate of GEMMAs and whether they can internalize as intact signaling complexes. In fact, internalization of GEMMAs would provide a very plausible framework for the now well established endosomal signaling mediated by internalized GPCRs (Calebiro, Nikolaev, Persani, & Lohse, 2010; Ferrandon et al., 2009; von Zastrow & Sorkin, 2021), although current evidence suggests that β₂AR and AC may traffic independently and potentially reform in endosomes (Lazar et al., 2020). Furthermore, it will be important to determine what is the role of arrestins and other cytosolic proteins that interact with GPCRs, G proteins and PM-effectors in the modulation of internalization and both G protein-dependent and independent signaling of GEMMAs.

Apart from their relevance as functional signaling units, the GEMMA concept will be important for further exploration of GPCR pharmacology. The localization in specific cellular environments, together with the unique properties of each GEMMA, determined by the large variety of potential components, should promote new directions in the search for new therapeutic agents. Screens should focus on molecules with the ability to promote changes in the function of a specific GEMMA, using single agents or combination of molecules with selectivity for the different components in particular GEMMAs.

Our primary recommendation is the further discovery and characterization of GEMMAs in their native environments with the identification of their primary components. This can only be achieved by combining the same approaches used for identification of GPCR oligomers with proteomic, biochemical or antibody-based techniques, such as proximity ligation or AlphaLISA assays (see, for instance, Moreno-Delgado et al., 2020; Valle-León et al., 2021). Such studies should be performed in parallel with reconstitution of the putative GEMMA in an artificial system, which should allow determination of interaction

sites and interfaces between its putative components, using, for instance, the TM-peptide strategy. This should allow the establishment of unique pharmacological properties of the GEMMA that could be used for its identification in native tissue and potentially targeted for drug discovery.

We should also aim at elucidating the precise molecular structure of GEMMAs in the presence and absence of ligands that stabilize active and inactive states, which should already be possible with cryo-EM techniques. Dynamic transitions between active and inactive states could be pursued by molecular dynamics simulations. The elucidation of the molecular structures of GEMMAs and their dynamic interrelationships should provide a deeper understanding of the intricate mechanisms of membrane-delimited GPCR-mediated cell signaling and an evaluation of their potential as therapeutic targets.

Declaration of Competing Interest

J.G.-M. has a sponsored research contract with Neuristic Pharma. The remaining authors declare that there are no conflicts of interest.

Acknowledgements

This article was written as the result of a roundtable meeting of the coauthors at the University of Barcelona in February 2020, organized by the Esteve Foundation and the Institute of Neurosciences of the University of Barcelona. S.F. is supported with the intramural funds of the National Institute on Drug Abuse. C.W.D. is supported by the National Institute of General Medicine (GM60419). J.G.-M. is supported by the National Institute of Mental Health (R01MH084894 and R01MH111940). D.E.L. is supported by the National Institute of Heart, Lung and Blood Institute (HL09549-24). T.E.H. holds the Canadian Pacific Chair in Biotechnology. F.C. and L.P. are supported by FEDER-EU/“Ministerio de Ciencia, Innovación y Universidades-Agencia Estatal de Investigación” (PID2020-118511RB-I00). R.J. is supported by “Agence Nationale de la Recherche” (ANR-19-CE16-0025-01), “Fondation de la Recherche Médicale” (FRM DEQ20130326503), “Institut National de la Santé et de la Recherche Médicale” and “Centre National de la Recherche Scientifique”.

References

- Ahmed, S. N., Brown, D. A., & London, E. (1997). On the origin of sphingolipid/cholesterol-rich detergent-insoluble cell membranes: Physiological concentrations of cholesterol and sphingolipid induce formation of a detergent-insoluble, liquid-ordered lipid phase in model membranes. *Biochemistry* 36(36), 10944–10953.
- Andressen, K. W., Ulsund, A. H., Krobert, K. A., Lohse, M. J., Bünemann, M., & Levy, F. O. (2018). Related GPCRs couple differently to G_s: Preassociation between G protein and 5-HT₇ serotonin receptor reveals movement of Gα_s upon receptor activation. *FASEB Journal* 32(2), 1059–1069.
- Ayoub, M. A., Couturier, C., Lucas-Meunier, E., Angers, S., Fossier, P., Bouvier, M., et al. (2002). Monitoring of ligand-independent dimerization and ligand-induced conformational changes of melatonin receptors in living cells by bioluminescence resonance energy transfer. *The Journal of Biological Chemistry* 277(24), 21522–21528.
- Ayoub, M. A., Levoe, A., Delagrèze, P., & Jockers, R. (2004). Preferential formation of MT₁/MT₂ melatonin receptor heterodimers with distinct ligand interaction properties compared with MT₂ homodimers. *Molecular Pharmacology* 66(2), 312–321.
- Ayoub, M. A., Trinquet, E., Pflieger, K. D., & Pin, J. P. (2010). Differential association modes of the thrombin receptor PAR1 with Galphai1, Galphai2, and beta-arrestin 1. *FASEB Journal* 24(9), 3522–3535.
- Baba, K., Benleulmi-Chaachoua, A., Journé, A. S., Kamal, M., Guillaume, J. L., Dussaud, S., et al. (2013). Heteromeric MT₁/MT₂ melatonin receptors modulate photoreceptor function. *Science Signaling* 6(296), Article ra89.
- Baki, L., Fribourg, M., Younkin, J., Eltit, J. M., Moreno, J. L., Park, G., et al. (2016). Cross-signaling in metabotropic glutamate 2 and serotonin 2A receptor heteromers in mammalian cells. *Pflugers Archiv* 468(5), 775–793.
- Baldwin, T. A., Li, Y., Brand, C. S., Watts, V. J., & Dessauer, C. W. (2019). Insights into the regulatory properties of human adenylyl cyclase type 9. *Molecular Pharmacology* 95(4), 349–360.
- Banères, J. L., & Parelló, J. (2003). Structure-based analysis of GPCR function: evidence for a novel pentameric assembly between the dimeric leukotriene B₄ receptor BLT1 and the G-protein. *Journal of Molecular Biology* 329(4), 815–829.
- Bar-Sinai, A., Marbach, I., Shorr, R. G., & Levitzki, A. (1992). The GppNHP-activated adenylyl cyclase complex from turkey erythrocyte membranes can be isolated with its beta gamma subunits. *European Journal of Biochemistry* 207(2), 703–708.

- Bauman, A. L., Soughayer, J., Nguyen, B. T., Willoughby, D., Carnegie, G. K., Wong, W., et al. (2006). Dynamic regulation of cAMP synthesis through anchored PKA-adenylyl cyclase V/VI complexes. *Molecular Cell* 23(6), 925–931.
- Bécamel, C., Gavarini, S., Channir, B., Alonso, G., Galéotti, N., Dumuis, A., et al. (2004). The serotonin 5-HT_{2A} and 5-HT_{2C} receptors interact with specific sets of PDZ proteins. *The Journal of Biological Chemistry* 279(19), 20257–20266.
- Beedle, A. M., McRory, J. E., Poirot, O., Doering, C. J., Altier, C., & Barrere, C. et al. (2004). Agonist-independent modulation of N-type calcium channels by ORL1 receptors. *Nature Neuroscience* 7(2), 118–125.
- Bonaventura, J., Navarro, G., Casadó-Anguera, V., Azdad, K., Rea, W., Moreno, E., et al. (2015). Allosteric interactions between agonists and antagonists within the adenosine A_{2A} receptor-dopamine D₂ receptor heterotetramer. *Proceedings of the National Academy of Sciences of the United States of America* 112(27), E3609–E3618.
- Boroto-Escuela, D. O., Marcellino, D., Narvaez, M., Flajolet, M., Heintz, N., & Agnati, A. et al. (2010). A serine point mutation in the adenosine A_{2A}R C-terminal tail reduces receptor heteromerization and allosteric modulation of the dopamine D₂R. *Biochemical and Biophysical Research Communications* 394(1), 222–227.
- Boroto-Escuela, D. O., Romero-Fernandez, W., Tarakanov, A. O., Gómez-Soler, M., Corrales, F., & Marcellino, D. et al. (2010). Characterization of the A_{2A}R-D₂R interface: focus on the role of the C-terminal tail and the transmembrane helices. *Biochemical and Biophysical Research Communications* 402(4), 801–807.
- Bourque, K., Jones-Tabah, J., Devost, D., Clarke, P., & Hébert, T. E. (2020). Exploring functional consequences of GPCR oligomerization requires a different lens. *Progress in Molecular Biology and Translational Science* 169, 181–211.
- Bouvier, M. (2001). Oligomerization of G-protein-coupled transmitter receptors. *Nature reviews*. *Neuroscience* 2(4), 274–286.
- Brand, C. S., Sadana, R., Malik, S., Smrcka, A. V., & Dessauer, C. W. (2015). Adenylyl cyclase 5 regulation by G $\beta\gamma$ involves isoform-specific use of multiple interaction sites. *Molecular Pharmacology* 88(4), 758–767.
- Braun, S., & Levitzki, A. (1979). Adenosine receptor permanently coupled to turkey erythrocyte adenylyl cyclase. *Biochemistry* 18(10), 2134–2138.
- Cabrera-Vera, T. M., Thomas, T. O., Vanhauwe, J., Depree, K. M., Graber, S. G., & Hamm, H. E. (2002). Dissecting receptor-G protein specificity using G alpha chimeras. *Methods in Enzymology* 344, 69–81.
- Cai, N. S., Quiroz, C., Bonaventura, J., Bonifazi, A., Cole, T. O., Purks, J., et al. (2019). Opioid-galanin receptor heteromers mediate the dopaminergic effects of opioids. *The Journal of Clinical Investigation* 129(7), 2730–2744.
- Calebiro, D., Nikolaev, V. O., Persani, L., & Lohse, M. J. (2010). Signaling by internalized G-protein-coupled receptors. *Trends in Pharmacological Sciences* 31(5), 221–228.
- Calebiro, D., Rieken, F., Wagner, J., Sungkaworn, T., Zabel, U., Borzi, A., et al. (2013). Single-molecule analysis of fluorescently labeled G-protein-coupled receptors reveals complexes with distinct dynamics and organization. *Proceedings of the National Academy of Sciences of the United States of America* 110(2), 743–748.
- Cecon, E., Oishi, A., & Jockers, R. (2018). Melatonin receptors: molecular pharmacology and signalling in the context of system bias. *British Journal of Pharmacology* 175(16), 3263–3280.
- Chen, J. F., & Cunha, R. A. (2020). The belated US FDA approval of the adenosine A_{2A} receptor antagonist istradefylline for treatment of Parkinson's disease. *Purinergic Signalling* 16(2), 167–174.
- Chow, K. B., Sun, J., Chu, K. M., Tai Cheung, W., Cheng, C. H., et al. (2012). The truncated ghrelin receptor polypeptide (GHS-R1b) is localized in the endoplasmic reticulum where it forms heterodimers with ghrelin receptors (GHS-R1a) to attenuate their cell surface expression. *Molecular and Cellular Endocrinology* 348(1), 247–254.
- Chung, K. Y., Rasmussen, S. G., Liu, T., Li, S., DeVree, B. T., Chae, P. S., et al. (2011). Conformational changes in the G protein Gs induced by the β_2 adrenergic receptor. *Nature* 477(7366), 611–615.
- Colledge, M., Dean, R. A., Scott, G. K., Langeberg, L. K., Haganir, R. L., & Scott, J. D. (2000). Targeting of PKA to glutamate receptors through a MAGUK-AKAP complex. *Neuron* 27(1), 107–119.
- Cordeaux, Y., Ijzerman, A. P., & Hill, S. J. (2004). Coupling of the human A₁ adenosine receptor to different heterotrimeric G proteins: Evidence for agonist-specific G protein activation. *British Journal of Pharmacology* 143(6), 705–714.
- Cottet, M., Faklaris, O., Maurel, D., Scholler, P., Doumazane, E., Trinquet, E., et al. (2012). BRET and Time-resolved FRET strategy to study GPCR oligomerization: From cell lines toward native tissues. *Frontiers in Endocrinology* 3, 92.
- Curthoys, N. M., Parent, M., Mlodzianoski, M., Nelson, A. J., Lillieholm, J., Butler, M. B., et al. (2015). Dances with membranes: Breakthroughs from super-resolution imaging. *Current Topics in Membranes* 75, 59–123.
- Damian, M., Mary, S., Maingot, M., M'Kadmi, C., Gagne, D., Leyris, J. P., et al. (2015). Ghrelin receptor conformational dynamics regulate the transition from a preassembled to an active receptor:Gq complex. *Proceedings of the National Academy of Sciences of the United States of America* 112(5), 1601–1606.
- Daniels, D. J., Lenard, N. R., Etienne, C. L., Law, P. Y., Roerig, S. C., & Portoghesi, P. S. (2005). Opioid-induced tolerance and dependence in mice is modulated by the distance between pharmacophores in a bivalent ligand series. *Proceedings of the National Academy of Sciences of the United States of America* 102(52), 19208–19213.
- Dascal, N., & Kahanovitch, U. (2015). The roles of G $\beta\gamma$ and G α in gating and regulation of GIRK channels. *International Review of Neurobiology* 123, 27–85.
- Davare, M. A., Avdonin, V., Hall, D. D., Peden, E. M., Burette, A., Weinberg, R. J., et al. (2001). A β_2 adrenergic receptor signaling complex assembled with the Ca²⁺ channel Cav1.2. *Science* 293(5527), 98–101.
- Davenport, A. P., Alexander, S. P., Sharnan, J. L., Pawson, A. J., Benson, H. E., Monaghan, A. E., et al. (2013). International union of basic and clinical pharmacology. LXXXVIII. G protein-coupled receptor list: Recommendations for new pairings with cognate ligands. *Pharmacological Reviews* 65(3), 967–986.
- David, M., Richer, M., Mamarbachi, A. M., Villeneuve, L. R., Dupré, D. J., & Hébert, T. E. (2006). Interactions between GABA-B1 receptors and Kir 3 inwardly rectifying potassium channels. *Cellular Signalling* 18(12), 2172–2181.
- Davis, T. L., Bonacci, T. M., Sprang, S. R., & Smrcka, A. V. (2005). Structural and molecular characterization of a preferred protein interaction surface on G protein beta gamma subunits. *Biochemistry* 44(31), 10593–10604.
- Décailott, F. M., Rozenfeld, R., Gupta, A., & Devi, L. A. (2008). Cell surface targeting of mu-delta opioid receptor heterodimers by RTP4. *Proceedings of the National Academy of Sciences of the United States of America* 105(41), 16045–16050.
- Dessauer, C. W. (2009). Adenylyl cyclase–A-kinase anchoring protein complexes: The next dimension in cAMP signaling. *Molecular Pharmacology* 76(5), 935–941.
- Dessauer, C. W., Tesmer, J. J., Sprang, S. R., & Gilman, A. G. (1998). Identification of a Gialpha binding site on type V adenylyl cyclase. *The Journal of Biological Chemistry* 273(40), 25831–25839.
- Dessauer, C. W., Watts, V. J., Ostrom, R. S., Conti, M., Dove, S., & Seifert, R. (2017). International union of basic and clinical pharmacology. CI. structures and small molecule modulators of mammalian adenylyl cyclases. *Pharmacological Reviews* 69(2), 93–139.
- Diering, G. H., Gustina, A. S., & Haganir, R. L. (2014). PKA-GluA1 coupling via AKAP5 controls AMPA receptor phosphorylation and cell-surface targeting during bidirectional homeostatic plasticity. *Neuron* 84(4), 790–805.
- Dorsch, S., Klotz, K. N., Engelhardt, S., Lohse, M. J., & Binemann, M. (2009). Analysis of receptor oligomerization by FRAP microscopy. *Nature Methods* 6(3), 225–230.
- Downes, G. B., & Gautam, N. (1999). The G protein subunit gene families. *Genomics* 62(3), 544–552.
- Du, J., Wang, D., Fan, H., Xu, C., Tai, L., Lin, S., et al. (2021). Structures of human mGlu2 and mGlu7 homo- and heterodimers. *Nature* 594(7864), 589–593.
- Du, Y., Duc, N. M., Rasmussen, S., Hilger, D., Kubiak, X., Wang, L., et al. (2019). Assembly of a GPCR-G protein complex. *Cell* 177(5), 1232–1242 (e11).
- Dunn, H. A., & Ferguson, S. S. (2015). PDZ protein regulation of G protein-coupled receptor trafficking and signaling pathways. *Molecular Pharmacology* 88(4), 624–639.
- Dupré, D. J., Baragli, A., Rebois, R. V., Ethier, N., & Hébert, T. E. (2007). Signalling complexes associated with adenylyl cyclase II are assembled during their biosynthesis. *Cellular Signalling* 19(3), 481–489.
- Dupré, D. J., Robitaille, M., Ethier, N., Villeneuve, L. R., Mamarbachi, A. M., & Hébert, T. E. (2006). Seven transmembrane receptor core signaling complexes are assembled prior to plasma membrane trafficking. *The Journal of Biological Chemistry* 281(45), 34561–34573.
- Dupré, D. J., Robitaille, M., Rebois, R. V., & Hébert, T. E. (2009). The role of Gbetagamma subunits in the organization, assembly, and function of GPCR signaling complexes. *Annual Review of Pharmacology and Toxicology* 49, 31–56.
- Efendiev, R., Samelson, B. K., Nguyen, B. T., Phatarpekar, P. V., Baameur, F., Scott, J. D., et al. (2010). AKAP79 interacts with multiple adenylyl cyclase (AC) isoforms and scaffolds AC5 and -6 to alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionate (AMPA) receptors. *The Journal of Biological Chemistry* 285(19), 14450–14458.
- El Khamlichi, C., Reverchon-Assadi, F., Hervouet-Coste, N., Blot, L., Reiter, E., & Morisset-Lopez, S. (2019). Bioluminescence resonance energy transfer as a method to study protein-protein interactions: Application to G protein coupled receptor biology. *Molecules* 24(3), 537.
- Ellaithy, A., Gonzalez-Maeso, J., Logothetis, D. A., & Levitz, J. (2020). Structural and biophysical mechanisms of class C G protein-coupled receptor function. *Trends in Biochemical Sciences* 45(12), 1049–1064.
- Elmhurst, J. L., Xie, Z., O'Dowd, B. F., & George, S. R. (2000). The splice variant D3nf reduces ligand binding to the D3 dopamine receptor: Evidence for heterooligomerization. *Molecular Brain Research* 80(1), 63–74.
- Fan, G., Shumay, E., Wang, H., & Malbon, C. C. (2001). The scaffold protein gravin (cAMP-dependent protein kinase-anchoring protein 250) binds the beta 2-adrenergic receptor via the receptor cytoplasmic Arg-329 to Leu-413 domain and provides a mobile scaffold during desensitization. *The Journal of Biological Chemistry* 276(26), 24005–24014.
- Faouzi, A., Uprety, R., Gomes, I., Massaly, N., Keresztes, A. I., Le Rouzic, V., et al. (2020). Synthesis and pharmacology of a novel μ -6 opioid receptor heteromer-selective agonist based on the carfentanyl template. *Journal of Medicinal Chemistry* 63(22), 13618–13637.
- Ferrandon, S., Feinstein, T. N., Castro, M., Wang, B., Bouley, R., Potts, J. T., et al. (2009). Sustained cyclic AMP production by parathyroid hormone receptor endocytosis. *Nature Chemical Biology* 5(10), 734–742.
- Ferré, S. (2015). The GPCR heterotetramer: challenging classical pharmacology. *Trends in Pharmacological Sciences* 36(3), 145–152.
- Ferré, S., Baler, R., Bouvier, M., Caron, M. G., Devi, L. A., Durrour, T., et al. (2009). Building a new conceptual framework for receptor heteromers. *Nature Chemical Biology* 5(3), 131–134.
- Ferré, S., Bonaventura, J., Zhu, W., Hatcher-Solis, C., Taura, J., Quiroz, C., et al. (2018). Essential control of the function of the striatopallidal neuron by pre-coupled complexes of adenosine A_{2A}-dopamine D₂ receptor heterotetramers and adenylyl cyclase. *Frontiers in Pharmacology* 9, 243.
- Ferré, S., Casadó, V., Devi, L. A., Filizola, M., Jockers, R., Lohse, M. J., et al. (2014). G protein-coupled receptor oligomerization revisited: Functional and pharmacological perspectives. *Pharmacological Reviews* 66(2), 413–434.
- Ferré, S., Ciruela, F., Casadó, V., & Pardo, L. (2020). Oligomerization of G protein-coupled receptors: Still doubted? *Progress in Molecular Biology and Translational Science* 169, 297–321.
- Ferré, S., von Euler, G., Johansson, B., Fredholm, B. B., & Fuxe, K. (1991). Stimulation of high-affinity adenosine A₂ receptors decreases the affinity of dopamine D₂ receptors in rat striatal membranes. *Proceedings of the National Academy of Sciences of the United States of America* 88(16), 7238–7241.

- Flock, T., Hauser, A. S., Lund, N., Gloriam, D. E., Balaji, S., & Babu, M. M. (2017). Selectivity determinants of GPCR-G-protein binding. *Nature* 545(7654), 317–322.
- Fraser, I. D., Cong, M., Kim, J., Rollins, E. N., Daaka, Y., Lefkowitz, R. J., et al. (2000). Assembly of an A kinase-anchoring protein-beta(2)-adrenergic receptor complex facilitates receptor phosphorylation and signaling. *Current Biology* 10(7), 409–412.
- Freimann, K., Kurrikoff, K., & Langel, Ü. (2015). Galanin receptors as a potential target for neurological disease. *Expert Opinion on Therapeutic Targets* 19(12), 1665–1676.
- Fribourg, M., Moreno, J. L., Holloway, T., Provasi, D., Baki, L., Mahajan, R., et al. (2011). Decoding the signaling of a GPCR heteromeric complex reveals a unifying mechanism of action of antipsychotic drugs. *Cell* 147(5), 1011–1023.
- Fujita, W., Gomes, I., & Devi, L. A. (2015). Heteromers of μ - δ opioid receptors: New pharmacology and novel therapeutic possibilities. *British Journal of Pharmacology* 172(2), 375–387.
- Funke, L., Dakoji, S., & Bredt, D. S. (2005). Membrane-associated guanylate kinases regulate adhesion and plasticity at cell junctions. *Annual Review of Biochemistry* 74, 219–245.
- Gaitonde, S. A., & González-Maeso, J. (2017). Contribution of heteromerization to G protein-coupled receptor function. *Current Opinion in Pharmacology* 32, 23–31.
- Galés, C., Rebois, R. V., Hogue, M., Trieu, P., Breit, A., Hébert, T. E., et al. (2005). Real-time monitoring of receptor and G-protein interactions in living cells. *Nature Methods* 2(3), 177–184.
- Galés, C., Van Durm, J. J., Schaak, S., Pontier, S., Percherancier, Y., Audet, M., et al. (2006). Probing the activation-promoted structural rearrangements in preassembled receptor-G protein complexes. *Nature Structural & Molecular Biology* 13(9), 778–786.
- Gardner, L. A., Tavalin, S. J., Goehring, A. S., Scott, J. D., & Bahouth, S. W. (2006). AKAP79-mediated targeting of the cyclic AMP-dependent protein kinase to the beta1-adrenergic receptor promotes recycling and functional resensitization of the receptor. *The Journal of Biological Chemistry* 281(44), 33537–33553.
- Gilman, A. G. (1987). G proteins: transducers of receptor-generated signals. *Annual Review of Biochemistry* 56, 615–649.
- Glaaser, I. W., & Slesinger, P. A. (2015). Structural insights into GIRK channel function. *International Review of Neurobiology* 123, 117–160.
- Gomes, I., Ayoub, M. A., Fujita, W., Jaeger, W. C., Pflieger, K. D., & Devi, L. A. (2016). G protein-coupled receptor heteromers. *Annual Review of Pharmacology and Toxicology* 56, 403–425.
- Gomes, I., Fujita, W., Gupta, A., Saldanha, S. A., Negri, A., Pinello, C. E., et al. (2013). Identification of a μ - δ opioid receptor heteromer-biased agonist with antinociceptive activity. *Proceedings of the National Academy of Sciences of the United States of America* 110(29), 12072–12077.
- González-Maeso, J., Ang, R. L., Yuen, T., Chan, P., Weisstaub, N. V., López-Giménez, J. F., et al. (2008). Identification of a serotonin/glutamate receptor complex implicated in psychosis. *Nature* 452(7183), 93–97.
- Granzin, J., Wilden, U., Choe, H. W., Labahn, J., Krafft, B., & Büldt, G. (1998). X-ray crystal structure of arrestin from bovine rod outer segments. *Nature* 391(6670), 918–921.
- Gudermann, T., Kalkbrenner, F., & Schultz, G. (1996). Diversity and selectivity of receptor-G protein interaction. *Annual Review of Pharmacology and Toxicology* 36, 429–459.
- Guinard, D., Moreno, E., Galindo, L., Cuenca-Royo, A., Barrera-Conde, M., Pérez, E. J., et al. (2020). Altered signaling in CB1R-5-HT2AR heteromers in olfactory neuroepithelium cells of schizophrenia patients is modulated by cannabis use. *Schizophrenia Bulletin* 46(6), 1547–1557.
- Guitart, X., Moreno, E., Rea, W., Sánchez-Soto, M., Cai, N. S., Quiroz, C., et al. (2019). Biased G protein-independent signaling of dopamine D₁-D₃ receptor heteromers in the nucleus accumbens. *Molecular Neurobiology* 56(10), 6756–6769.
- Guitart, X., Navarro, G., Moreno, E., Yano, H., Cai, N. S., Sánchez-Soto, M., et al. (2014). Functional selectivity of allosteric interactions within G protein-coupled receptor oligomers: the dopamine D1-D3 receptor heterotetramer. *Molecular Pharmacology* 86(4), 417–429.
- Guo, H., An, S., Ward, R., Yang, Y., Liu, Y., Guo, X. X., et al. (2017). Methods used to study the oligomeric structure of G-protein-coupled receptors. *Bioscience Reports* 37(2) (BSR20160547).
- Gurevich, E. V., Tesmer, J. J., Mushegian, A., & Gurevich, V. V. (2012). G protein-coupled receptor kinases: More than just kinases and not only for GPCRs. *Pharmacology & Therapeutics* 133(1), 40–69.
- Gurevich, V. V., & Gurevich, E. V. (2019). The structural basis of the arrestin binding to GPCRs. *Molecular and Cellular Endocrinology* 484, 34–41.
- Hall, D. D., Davare, M. A., Shi, M., Allen, M. L., Weisenhaus, M., McKnight, G. S., et al. (2007). Critical role of cAMP-dependent protein kinase anchoring to the L-type calcium channel Cav1.2 via A-kinase anchor protein 150 in neurons. *Biochemistry* 46(6), 1635–1646.
- Hay, D. L., & Pioszak, A. A. (2016). Receptor activity-modifying proteins (RAMPs): New insights and roles. *Annual Review of Pharmacology and Toxicology* 56, 469–487.
- He, J., Bellini, M., Inuzuka, H., Xu, J., Xiong, Y., Yang, X., et al. (2006). Proteomic analysis of beta1-adrenergic receptor interactions with PDZ scaffold proteins. *The Journal of Biological Chemistry* 281(5), 2820–2827.
- Hébert, T. E., Moffett, S., Morello, J. P., Loisel, T. P., Bichet, D. G., Barret, C., et al. (1996). A peptide derived from a beta2-adrenergic receptor transmembrane domain inhibits both receptor dimerization and activation. *The Journal of Biological Chemistry* 271(27), 16384–16392.
- Hepler, J. R. (2014). G protein coupled receptor signaling complexes in live cells. *Cellular Logistics* 4, Article e29392.
- Herrick-Davis, K., Grinde, E., Cowan, A., & Mazurkiewicz, J. E. (2013). Fluorescence correlation spectroscopy analysis of serotonin, adrenergic, muscarinic, and dopamine receptor dimerization: The oligomer number puzzle. *Molecular Pharmacology* 84(4), 630–642.
- Herrick-Davis, K., Weaver, B. A., Grinde, E., & Mazurkiewicz, J. E. (2006). Serotonin 5-HT2C receptor homodimer biogenesis in the endoplasmic reticulum: Real-time visualization with confocal fluorescence resonance energy transfer. *The Journal of Biological Chemistry* 281(37), 27109–27116.
- Hervé, D. (2011). Identification of a specific assembly of the g protein golf as a critical and regulated module of dopamine and adenosine-activated cAMP pathways in the striatum. *Frontiers in Neuroanatomy* 5, 48.
- Homan, K. T., Glukhova, A., & Tesmer, J. J. (2013). Regulation of G protein-coupled receptor kinases by phospholipids. *Current Medicinal Chemistry* 20(1), 39–46.
- Hu, L. A., Tang, Y., Miller, W. E., Cong, M., Lau, A. G., Lefkowitz, R. J., et al. (2000). beta 1-adrenergic receptor association with PSD-95. Inhibition of receptor internalization and facilitation of beta 1-adrenergic receptor interaction with N-methyl-D-aspartate receptors. *The Journal of Biological Chemistry* 275(49), 38659–38666.
- Hua, X. Y., Hayes, C. S., Hofer, A., Fitzsimmons, B., Kilk, K., Langel, U., et al. (2004). Galanin acts at GalR1 receptors in spinal antinociception: Synergy with morphine and AP-5. *The Journal of Pharmacology and Experimental Therapeutics* 308(2), 574–582.
- Huang, C. L., Feng, S., & Hilgemann, D. W. (1998). Direct activation of inward rectifier potassium channels by PIP2 and its stabilization by Gbetagamma. *Nature* 391(6669), 803–806.
- Huang, C. L., Slesinger, P. A., Casey, P. J., Jan, Y. N., & Jan, L. Y. (1995). Evidence that direct binding of G beta gamma to the GIRK1 G protein-gated inwardly rectifying K+ channel is important for channel activation. *Neuron* 15(5), 1133–1143.
- Huang, J., Chen, S., Zhang, J. J., & Huang, X. Y. (2013). Crystal structure of oligomeric β 1-adrenergic G protein-coupled receptors in ligand-free basal state. *Nature Structural & Molecular Biology* 20(4), 419–425.
- Huang, W., Masureel, M., Qu, Q., Janetzko, J., Inoue, A., Kato, H. E., et al. (2020). Structure of the neurotensin receptor 1 in complex with β -arrestin 1. *Nature* 579(7798), 303–308.
- Joiner, M. L., Lisé, M. F., Yuen, E. Y., Kam, A. Y., Zhang, M., Hall, D. D., et al. (2010). Assembly of a beta2-adrenergic receptor-GluR1 signalling complex for localized cAMP signaling. *The EMBO Journal* 29(2), 482–495.
- Justinová, Z., Redhi, G. H., Goldberg, S. R., & Ferré, S. (2014). Differential effects of presynaptic versus postsynaptic adenosine A2A receptor blockade on Δ 9-tetrahydrocannabinol (THC) self-administration in squirrel monkeys. *The Journal of Neuroscience* 34(19), 6480–6484.
- Kalkbrenner, F., Degtiar, V. E., Schenker, M., Brendel, S., Zobel, A., Heschler, J., et al. (1995). Subunit composition of G(o) proteins functionally coupling galanin receptors to voltage-gated calcium channels. *The EMBO Journal* 14(19), 4728–4737.
- Karpa, K. D., Lin, R., Kabbani, N., & Levenson, R. (2000). The dopamine D3 receptor interacts with itself and the truncated D3 splice variant d3nf: D3-D3nf interaction causes mislocalization of D3 receptors. *Molecular Pharmacology* 58(4), 677–683.
- Kato, H. E., Zhang, Y., Hu, H., Suomivuori, C. M., Kadji, F., Aoki, J., et al. (2019). Conformational transitions of a neurotensin receptor 1-G₁ complex. *Nature* 572(7767), 80–85.
- Khan, S. M., Sleno, R., Gora, S., Zylbergold, P., Laverdure, J. P., Labbé, J. C., et al. (2013). The expanding roles of G β γ subunits in G protein-coupled receptor signaling and drug action. *Pharmacological Reviews* 65(2), 545–577.
- Khan, S. M., Sung, J. Y., & Hébert, T. E. (2016). G β γ subunits-Different spaces, different faces. *Pharmacological Research* 111, 434–441.
- Kleuss, C., Scherübel, H., Hescheler, J., Schultz, G., & Wittig, B. (1993). Selectivity in signal transduction determined by gamma subunits of heterotrimeric G proteins. *Science* 259(5096), 832–834.
- Kleuss, C., Schultz, G., & Wittig, B. (1994). Microinjection of antisense oligonucleotides to assess G-protein subunit function. *Methods in Enzymology* 237, 345–355.
- Koehl, A., Hu, H., Feng, D., Sun, B., Zhang, Y., Robertson, M. J., et al. (2019). Structural insights into the activation of metabotropic glutamate receptors. *Nature* 566(7742), 79–84.
- Köfalvi, A., Moreno, E., Cordero, A., Cai, N. S., Fernández-Dueñas, V., Ferreira, S. G., et al. (2020). Control of glutamate release by complexes of adenosine and cannabinoid receptors. *BMC Biology* 18(1), 9.
- Kull, B., Ferré, S., Arslan, G., Svenningsson, P., Fuxe, K., Owman, C., et al. (1999). Reciprocal interactions between adenosine A2A and dopamine D2 receptors in Chinese hamster ovary cells co-transfected with the two receptors. *Biochemical Pharmacology* 58(6), 1035–1045.
- Lauckner, J. E., Hille, B., & Mackie, K. (2005). The cannabinoid agonist WIN55,212-2 increases intracellular calcium via CB1 receptor coupling to Gq/11 G proteins. *Proceedings of the National Academy of Sciences of the United States of America* 102(52), 19144–19149.
- Lavine, N., Ethier, N., Oak, J. N., Pei, L., Liu, F., Trieu, P., et al. (2002). G protein-coupled receptors form stable complexes with inwardly rectifying potassium channels and adenylyl cyclase. *The Journal of Biological Chemistry* 277(48), 46010–46019.
- Lazar, A. M., Irannejad, R., Baldwin, T. A., Sundaram, A. B., Gutkind, J. S., Inoue, A., et al. (2020). G protein-regulated endocytic trafficking of adenylyl cyclase type 9. *eLife* 9, Article e58039.
- Leung, P. K., Chow, K. B., Lau, P. N., Chu, K. M., Chan, C. B., Cheng, C. H., et al. (2007). The truncated ghrelin receptor polypeptide (GHS-R1b) acts as a dominant-negative mutant of the ghrelin receptor. *Cellular Signalling* 19(5), 1011–1022.
- Levitzki, A. (1986). Beta-adrenergic receptors and their mode of coupling to adenylyl cyclase. *Physiological Reviews* 66(3), 819–854.
- Levitzki, A. (1988). From epinephrine to cyclic AMP. *Science (New York, N.Y.)* 241(4867), 800–806.
- Levitzki, A., & Klein, S. (2002). G-protein subunit dissociation is not an integral part of G-protein action. *ChemBiochem* 3(9), 815–818.
- Levoe, A., Dam, J., Ayoub, M. A., Guillaume, J. L., Couturier, C., Delagrèze, P., et al. (2006). The orphan GPR50 receptor specifically inhibits MT1 melatonin receptor function through heterodimerization. *The EMBO Journal* 25(13), 3012–3023.
- Levoe, A., Dam, J., Ayoub, M. A., Guillaume, J. L., & Jockers, R. (2006). Do orphan G-protein-coupled receptors have ligand-independent functions? New insights from receptor heterodimers. *EMBO Reports* 7(11), 1094–1098.

- Li, X., Nooh, M. M., & Bahouth, S. W. (2013). Role of AKAP79/150 protein in β 1-adrenergic receptor trafficking and signaling in mammalian cells. *The Journal of Biological Chemistry* 288(47), 33797–33812.
- Li, Z., Benard, O., & Margolskee, R. F. (2006). Ggamma13 interacts with PDZ domain-containing proteins. *The Journal of Biological Chemistry* 281(16), 11066–11073.
- Lin, S., Han, S., Cai, X., Tan, Q., Zhou, K., Wang, D., et al. (2021). Structures of G_i-bound metabotropic glutamate receptors mGlu2 and mGlu4. *Nature* 594(7864), 583–588.
- Liu, X., Xu, X., Hilger, D., Aschauer, P., Tiemann, J., Du, Y., et al. (2019). Structural insights into the process of GPCR-G protein complex formation. *Cell* 177(5), 1243–1251 (e12).
- Liu, Z., Fenech, C., Cadiou, H., Grall, S., Tili, E., Laugerette, F., et al. (2012). Identification of new binding partners of the chemosensory signaling protein G γ 13 expressed in taste and olfactory sensory cells. *Frontiers in Cellular Neuroscience* 6, 26.
- Logothetis, D. E., Kurachi, Y., Galper, J., Neer, E. J., & Clapham, D. E. (1987). The beta gamma subunits of GTP-binding proteins activate the muscarinic K⁺ channel in heart. *Nature* 325(6102), 321–326.
- Logothetis, D. E., Petrou, V. I., Zhang, M., Mahajan, R., Meng, X. Y., Adney, S. K., et al. (2015). Phosphoinositide control of membrane protein function: a frontier led by studies on ion channels. *Annual Review of Physiology* 77, 81–104.
- Manglik, A., Kruse, A. C., Kobilka, T. S., Thian, F. S., Mathiesen, J. M., Sunahara, R. K., et al. (2012). Crystal structure of the μ -opioid receptor bound to a morphinan antagonist. *Nature* 485(7398), 321–326.
- Margeta-Mitrovic, M., Jan, Y. N., & Jan, L. Y. (2000). A trafficking checkpoint controls GABA (B) receptor heterodimerization. *Neuron* 27(1), 97–106.
- Marrari, Y., Crouthamel, M., Irannejad, R., & Wedegaertner, P. B. (2007). Assembly and trafficking of heterotrimeric G proteins. *Biochemistry* 46(26), 7665–7677.
- Mary, S., Fehrentz, J. A., Damian, M., Gaibelet, G., Orcel, H., Verdié, P., et al. (2013). Heterodimerization with its splice variant blocks the ghrelin receptor 1a in a non-signaling conformation: a study with a purified heterodimer assembled into lipid discs. *The Journal of Biological Chemistry* 288(34), 24656–24665.
- McLatchie, L. M., Fraser, N. J., Main, M. J., Wise, A., Brown, J., Thompson, N., et al. (1998). RAMPs regulate the transport and ligand specificity of the calcitonin-receptor-like receptor. *Nature* 393(6683), 333–339.
- Michaelson, D., Ahearn, I., Bergo, M., Young, S., & Philips, M. (2002). Membrane trafficking of heterotrimeric G proteins via the endoplasmic reticulum and Golgi. *Molecular Biology of the Cell* 13(9), 3294–3302.
- Milligan, G. (2001). Oligomerisation of G-protein-coupled receptors. *Journal of Cell Science* 114(Pt 7), 1265–1271.
- Mirshahi, T., Robillard, L., Zhang, H., Hébert, T. E., & Logothetis, D. E. (2002). Gbeta residues that do not interact with Galpha underlie agonist-independent activity of K⁺ channels. *The Journal of Biological Chemistry* 277(9), 7348–7355.
- Mizuno-Yamasaki, E., Rivera-Molina, F., & Novick, P. (2012). GTPase networks in membrane traffic. *Annual Review of Biochemistry* 81, 637–659.
- Moreno, E., Quiroz, C., Rea, W., Cai, N. S., Mallol, J., Cortés, A., et al. (2017). Functional μ -opioid-galanin receptor heteromers in the ventral tegmental area. *The Journal of Neuroscience* 37(5), 1176–1186.
- Moreno, J. L., Miranda-Azpiazu, P., García-Bea, A., Younkin, J., Cui, M., Kozlenkov, A., et al. (2016). Allosteric signaling through an mGlu2 and 5-HT_{2A} heteromeric receptor complex and its potential contribution to schizophrenia. *Science Signaling* 9(410), Article ra5.
- Moreno, J. L., Muguza, C., Umali, A., Mortillo, S., Holloway, T., Pilar-Cuellar, F., et al. (2012). Identification of three residues essential for 5-hydroxytryptamine 2A-metabotropic glutamate 2 (5-HT_{2A}-mGlu2) receptor heteromerization and its psychoactive behavioral function. *The Journal of Biological Chemistry* 287(53), 44301–44319.
- Moreno-Delgado, D., Puigdel·l·lvol, M., Moreno, E., Rodríguez-Ruiz, M., Botta, J., Gasperini, P., et al. (2020). Modulation of dopamine D₁ receptors via histamine H₃ receptors is a novel therapeutic target for Huntington's disease. *eLife* 9, Article e51093.
- Navarro, G., Aguinaga, D., Angelats, E., Medrano, M., Moreno, E., Mallol, J., et al. (2016). A significant role of the truncated ghrelin receptor GHS-R1b in ghrelin-induced signaling in neurons. *The Journal of Biological Chemistry* 291(25), 13048–13062.
- Navarro, G., Cordero, A., Casadó-Anguera, V., Moreno, E., Cai, N. S., Cortés, A., et al. (2018). Evidence for functional pre-coupled complexes of receptor heteromers and adenylyl cyclase. *Nature Communications* 9(1), 1242.
- Neubig, R. R. (1994). Membrane organization in G-protein mechanisms. *FASEB Journal* 8(12), 939–946.
- Ngo, T., Kufareva, I., Coleman, J., Graham, R. M., Abagyan, R., & Smith, N. J. (2016). Identifying ligands at orphan GPCRs: Current status using structure-based approaches. *British Journal of Pharmacology* 173(20), 2934–2951.
- Nobles, M., Benians, A., & Tinker, A. (2005). Heterotrimeric G proteins precouple with G protein-coupled receptors in living cells. *Proceedings of the National Academy of Sciences of the United States of America* 102(51), 18706–18711.
- Oishi, A., Cecon, E., & Jockers, R. (2018). Melatonin receptor signaling: Impact of receptor oligomerization on receptor function. *International Review of Cell and Molecular Biology* 338, 59–77.
- Okashah, N., Wan, Q., Ghosh, S., Sandhu, M., Inoue, A., Vaidehi, N., et al. (2019). Variable G protein determinants of GPCR coupling selectivity. *Proceedings of the National Academy of Sciences of the United States of America* 116(24), 12054–12059.
- Oliveria, S. F., Dell'Acqua, M. L., & Sather, W. A. (2007). AKAP79/150 anchoring of calcineurin controls neuronal L-type Ca²⁺ channel activity and nuclear signaling. *Neuron* 55(2), 261–275.
- Olson, P. A., Tkatch, T., Hernandez-Lopez, S., Ulrich, S., Ilijic, E., Mugnaini, E., et al. (2005). G-protein-coupled receptor modulation of striatal CaV1.3 L-type Ca²⁺ channels is dependent on a Shank-binding domain. *The Journal of Neuroscience* 25(5), 1050–1062.
- Orrú, M., Bakešová, J., Brugarolas, M., Quiroz, C., Beaumont, V., Goldberg, S. R., et al. (2011). Striatal pre- and postsynaptic profile of adenosine A(2A) receptor antagonists. *PLoS One* 6(1), Article e16088.
- Ostrom, R. S., Liu, X., Head, B. P., Gregorian, C., Seasholtz, T. M., & Insel, P. A. (2002). Localization of adenylyl cyclase isoforms and G protein-coupled receptors in vascular smooth muscle cells: Expression in caveolin-rich and noncaveolin domains. *Molecular Pharmacology* 62(5), 983–992.
- Ostrom, R. S., Violin, J. D., Coleman, S., & Insel, P. A. (2000). Selective enhancement of beta-adrenergic receptor signaling by overexpression of adenylyl cyclase type 6: Colocalization of receptor and adenylyl cyclase in caveolae of cardiac myocytes. *Molecular Pharmacology* 57(5), 1075–1079.
- Palczewski, K., Kumasaka, T., Hori, T., Behnke, C. A., Motoshima, H., Fox, B. A., et al. (2000). Crystal structure of rhodopsin: A G protein-coupled receptor. *Science* 289(5480), 739–745.
- Patriarchi, T., Buonarati, O. R., & Hell, J. W. (2018). Postsynaptic localization and regulation of AMPA receptors and Cav1.2 by β 2 adrenergic receptor/PKA and Ca²⁺/CaMKII signaling. *The EMBO Journal* 37(20), Article e99771.
- Pike, L. J. (2003). Lipid rafts: Bringing order to chaos. *Journal of Lipid Research* 44(4), 655–667.
- Pike, L. J. (2006). Rafts defined: A report on the keystone symposium on lipid rafts and cell function. *Journal of Lipid Research* 47(7), 1597–1598.
- Poyner, D. R., Sexton, P. M., Marshall, I., Smith, D. M., Quirion, R., Born, W., et al. (2002). International Union of Pharmacology. XXXII. The mammalian calcitonin gene-related peptides, adrenomedullin, amylin, and calcitonin receptors. *Pharmacological Reviews* 54(2), 233–246.
- Pulido, D., Casadó-Anguera, V., Pérez-Benito, L., Moreno, E., Cordero, A., López, L., et al. (2018). Design of a true bivalent ligand with picomolar binding affinity for a G protein-coupled receptor homodimer. *Journal of Medicinal Chemistry* 61(20), 9335–9346.
- Qi, C., Sorrentino, S., Medalia, O., & Korkhov, V. M. (2019). The structure of a membrane adenylyl cyclase bound to an activated stimulatory G protein. *Science* 364(6438), 389–394.
- Qin, K., Dong, C., Wu, G., & Lambert, N. A. (2011). Inactive-state preassembly of G(q)-coupled receptors and G(q) heterotrimers. *Nature Chemical Biology* 7(10), 740–747.
- Rasmussen, S. G., DeVree, B. T., Zou, Y., Kruse, A. C., Chung, K. Y., Kobilka, T. S., et al. (2011). Crystal structure of the β 2 adrenergic receptor-Gs protein complex. *Nature* 477(7366), 549–555.
- Rebois, R. V., & Hébert, T. E. (2003). Protein complexes involved in heptahelical receptor-mediated signal transduction. *Receptors & Channels* 9(3), 169–194.
- Rebois, R. V., Robitaille, M., Galés, C., Dupré, D. J., Baragli, A., Trieu, P., et al. (2006). Heterotrimeric G proteins form stable complexes with adenylyl cyclase and Kir3.1 channels in living cells. *Journal of Cell Science* 119(Pt 13), 2807–2818.
- Riven, I., Iwanir, S., & Reuveny, E. (2006). GIRK channel activation involves a local rearrangement of a preformed G protein channel complex. *Neuron* 51(5), 561–573.
- Rivera-Oliver, M., Moreno, E., Álvarez-Bagnarol, Y., Ayala-Santiago, C., Cruz-Reyes, N., Molina-Castro, G. C., et al. (2019). Adenosine A₁-dopamine D₁ receptor heteromers control the excitability of the spinal motoneuron. *Molecular Neurobiology* 56(2), 797–811.
- Robishaw, J. D., & Berlot, C. H. (2004). Translating G protein subunit diversity into functional specificity. *Current Opinion in Cell Biology* 16(2), 206–209.
- Sadana, R., Dascal, N., & Dessauer, C. W. (2009). N terminus of type 5 adenylyl cyclase scaffolds Gs heterotrimer. *Molecular Pharmacology* 76(6), 1256–1264.
- Sadana, R., & Dessauer, C. W. (2009). Physiological roles for G protein-regulated adenylyl cyclase isoforms: insights from knockout and overexpression studies. *Neuro-Signals* 17(1), 5–22.
- Salahpour, A., Angers, S., Mercier, J. F., Lagacé, M., Marullo, S., & Bouvier, M. (2004). Homodimerization of the beta2-adrenergic receptor as a prerequisite for cell surface targeting. *The Journal of Biological Chemistry* 279(32), 33390–33397.
- Scarselli, M., Annibale, P., McCormick, P. J., Kolachalam, S., Aringhieri, S., Radenovic, A., et al. (2016). Revealing G-protein-coupled receptor oligomerization at the single-molecule level through a nanoscopic lens: Methods, dynamics and biological function. *The FEBS Journal* 283(7), 1197–1217.
- Scherer, G., Imamachi, N., Cao, Y. Q., Contet, C., Mennicken, F., O'Donnell, D., et al. (2009). Dissociation of the opioid receptor mechanisms that control mechanical and heat pain. *Cell* 137(6), 1148–1159.
- Schwindinger, W. F., Betz, K. S., Giger, K. E., Sabol, A., Bronson, S. K., & Robishaw, J. D. (2003). Loss of G protein gamma 7 alters behavior and reduces striatal alpha(olf) level and cAMP production. *The Journal of Biological Chemistry* 278(8), 6575–6579.
- Schwindinger, W. F., Mihalcik, L. J., Giger, K. E., Betz, K. S., Stauffer, A. M., Linden, J., Herve, D., & Robishaw, J. D. (2010). Adenosine A_{2A} receptor signaling and golf assembly show a specific requirement for the gamma7 subtype in the striatum. *The Journal of Biological Chemistry* 285(39), 29787–29796.
- Scott, J. D., Dessauer, C. W., & Taskán, K. (2013). Creating order from chaos: Cellular regulation by kinase anchoring. *Annual Review of Pharmacology and Toxicology* 53, 187–210.
- Serafin, D. S., Harris, N. R., Nielsen, N. R., Mackie, D. I., & Caron, K. M. (2020). Dawn of a New RAMPPage. *Trends in Pharmacological Sciences* 41(4), 249–265.
- Seven, A. B., Barros-Álvarez, X., de Lapeyrière, M., Papanerghi-Scott, M. M., Robertson, M. J., Zhang, C., ... Skiniotis, G. (2021). *Nature* 595(7867), 450–454.
- Sezgin, E., Levental, I., Mayor, S., & Eggeling, C. (2017). The mystery of membrane organization: Composition, regulation and roles of lipid rafts. *Molecular Cell Biology* 18(6), 361–374.
- Shah, U. H., Toneatti, R., Gaitonde, S. A., Shin, J. M., & González-Maeso, J. (2020). Site-specific incorporation of genetically encoded photo-crosslinkers locates the heteromeric interface of a GPCR complex in living cells. *Cell Chemical Biology* 27(10), 1308–1317 (e4).

- Sheng, M., & Sala, C. (2001). PDZ domains and the organization of supramolecular complexes. *Annual Review of Neuroscience* 24, 1–29.
- Shin, S. M., Zhang, N., Hansen, J., Gerges, N. Z., Pak, D. T., Sheng, M., et al. (2012). GKAP orchestrates activity-dependent postsynaptic protein remodeling and homeostatic scaling. *Nature Neuroscience* 15(12), 1655–1666.
- Simons, K., & Toomre, D. (2000). Lipid rafts and signal transduction. *Molecular Cell Biology* 20(1), 31–39.
- Singer, S. J., & Nicolson, G. L. (1972). The fluid mosaic model of the structure of cell membranes. *Science* 175(4023), 720–731.
- Singh, P., Wang, B., Maeda, T., Palczewski, K., & Tesmer, J. J. (2008). Structures of rhodopsin kinase in different ligand states reveal key elements involved in G protein-coupled receptor kinase activation. *The Journal of Biological Chemistry* 283(20), 14053–14062.
- Sleno, R., & Hébert, T. E. (2018). The dynamics of GPCR oligomerization and their functional consequences. *International Review of Cell and Molecular Biology* 338, 141–171.
- Sleno, R., & Hébert, T. E. (2019). Shaky ground - The nature of metastable GPCR signalling complexes. *Neuropharmacology* 152, 4–14.
- Smart, E. J., Graf, G. A., McNiven, M. A., Sessa, W. C., Engelman, J. A., Scherer, P. E., et al. (1999). Caveolins, liquid-ordered domains, and signal transduction. *Molecular and Cellular Biology* 19(11), 7289–7304.
- Smith, N. J., & Milligan, G. (2010). Allostery at G protein-coupled receptor homo- and heteromers: Uncharted pharmacological landscapes. *Pharmacological Reviews* 62(4), 701–725.
- Smrcka, A. V., & Fisher, I. (2019). G-protein $\beta\gamma$ subunits as multi-functional scaffolds and transducers in G-protein-coupled receptor signaling. *Cellular and Molecular Life Sciences* 76(22), 4447–4459.
- Soave, M., Briddon, S. J., Hill, S. J., & Stoddart, L. A. (2020). Fluorescent ligands: Bringing light to emerging GPCR paradigms. *British Journal of Pharmacology* 177(5), 978–991.
- Soriano, A., Ventura, R., Molero, A., Hoen, R., Casadó, V., Cortés, A., et al. (2009). Adenosine A2A receptor-antagonist/dopamine D2 receptor-agonist bivalent ligands as pharmacological tools to detect A2A-D2 receptor heteromers. *Journal of Medicinal Chemistry* 52(18), 5590–5602.
- Steiner, D., Saya, D., Schallmach, E., Simonds, W. F., & Vogel, Z. (2006). Adenylyl cyclase type-VIII activity is regulated by G(betagamma) subunits. *Cellular Signalling* 18(1), 62–68.
- Sui, J. L., Petit-Jacques, J., & Logothetis, D. E. (1998). Activation of the atrial KACH channel by the betagamma subunits of G proteins or intracellular Na⁺ ions depends on the presence of phosphatidylinositol phosphates. *Proceedings of the National Academy of Sciences of the United States of America* 95(3), 1307–1312.
- Sungkaworn, T., Jobin, M. L., Burnecki, K., Weron, A., Lohse, M. J., & Calebiro, D. (2017). Single-molecule imaging reveals receptor-G protein interactions at cell surface hot spots. *Nature* 550(7677), 543–547.
- Takida, S., & Wedegaertner, P. B. (2004). Exocytic pathway-independent plasma membrane targeting of heterotrimeric G proteins. *FEBS Letters* 567(2–3), 209–213.
- Taussig, R., Quarmany, L. M., & Gilman, A. G. (1993). Regulation of purified type I and type II adenylyl cyclases by G protein beta gamma subunits. *The Journal of Biological Chemistry* 268(1), 9–12.
- Teichmann, A., Gibert, A., Lampe, A., Grzesik, P., Rutz, C., Ferkert, J., et al. (2014). The specific monomer/dimer equilibrium of the corticotropin-releasing factor receptor type 1 is established in the endoplasmic reticulum. *The Journal of Biological Chemistry* 289(35), 24250–24262.
- Tiwari, V., He, S. Q., Huang, Q., Liang, L., Yang, F., Chen, Z., et al. (2020). Activation of μ - δ opioid receptor heteromers inhibits neuropathic pain behavior in rodents. *Pain* 161(4), 842–855.
- Tolkovsky, A. M., & Levitzki, A. (1978a). Mode of coupling between the beta-adrenergic receptor and adenylyl cyclase in turkey erythrocytes. *Biochemistry* 17(18), 3795.
- Tolkovsky, A. M., & Levitzki, A. (1978b). Coupling of a single adenylyl cyclase to two receptors: adenosine and catecholamine. *Biochemistry* 17(18), 3811–3817.
- Tsai, C. J., Marino, J., Adaixo, R., Pamula, F., Muehle, J., Maeda, S., et al. (2019). Cryo-EM structure of the rhodopsin-G α_i - $\beta\gamma$ complex reveals binding of the rhodopsin C-terminal tail to the $\beta\gamma$ subunit. *eLife* 8, Article e46041.
- Tu, J. C., Xiao, B., Naisbitt, S., Yuan, J. P., Petralia, R. S., Brakeman, P., et al. (1999). Coupling of mGluR/Homer and PSD-95 complexes by the Shank family of postsynaptic density proteins. *Neuron* 23(3), 583–592.
- Valentine, C. D., & Haggie, P. M. (2011). Confinement of $\beta(1)$ - and $\beta(2)$ -adrenergic receptors in the plasma membrane of cardiomyocyte-like H9c2 cells is mediated by selective interactions with PDZ domain and A-kinase anchoring proteins but not caveolae. *Molecular Biology of the Cell* 22(16), 2970–2982.
- Valle-León, M., Callado, L. F., Aso, E., Cajiá-Manrique, M. M., Sahlholm, K., López-Cano, M., et al. (2021). Decreased striatal adenosine A_{2A}-dopamine D₂ receptor heteromerization in schizophrenia. *Neuropsychopharmacology* 46(3), 665–672.
- Van Eps, N., Preininger, A. M., Alexander, N., Kaya, A. I., Meier, S., Meiler, J., et al. (2011). Interaction of a G protein with an activated receptor opens the interdomain interface in the alpha subunit. *Proceedings of the National Academy of Sciences of the United States of America* 108(23), 9420–9424.
- Viñals, X., Moreno, E., Lanfumey, L., Cordero, A., Pastor, A., de La Torre, R., et al. (2015). Cognitive impairment induced by Delta9-tetrahydrocannabinol occurs through heteromers between cannabinoid CB1 and serotonin 5-HT_{2A} receptors. *PLoS Biology* 13(7), Article e1002194.
- Violin, J. D., & Lefkowitz, R. J. (2007). Beta-arrestin-biased ligands at seven-transmembrane receptors. *Trends in Pharmacological Sciences* 28(8), 416–422.
- Waldo, G. L., Ricks, T. K., Hicks, S. N., Cheever, M. L., Kawano, T., Tsuboi, K., et al. (2010). Kinetic scaffolding mediated by a phospholipase C-beta and Gq signaling complex. *Science* 330(6006), 974–980.
- Wall, M. A., Coleman, D. E., Lee, E., Iñiguez-Lluhi, J. A., Posner, B. A., Gilman, A. G., et al. (1995). The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2. *Cell* 83(6), 1047–1058.
- Walsh, S. M., Mathiasen, S., Christensen, S. M., Fay, J. F., King, C., Provasi, D., et al. (2018). Single proteoliposome high-content analysis reveals differences in the homooligomerization of GPCRs. *Biophysical Journal* 115(2), 300–312.
- Wang, H. B., Zhao, B., Zhong, Y. Q., Li, K. C., Li, Z. Y., Wang, Q., et al. (2010). Coexpression of delta- and mu-opioid receptors in nociceptive sensory neurons. *Proceedings of the National Academy of Sciences of the United States of America* 107(29), 13117–13122.
- Weis, W. I., & Kobilka, B. K. (2018). The molecular basis of G protein-coupled receptor activation. *Annual Review of Biochemistry* 87, 897–919.
- Westfield, G. H., Rasmussen, S. G., Su, M., Dutta, S., DeVree, B. T., Chung, K. Y., et al. (2011). Structural flexibility of the G alpha s alpha-helical domain in the beta2-adrenoceptor Gs complex. *Proceedings of the National Academy of Sciences of the United States of America* 108(38), 16086–16091.
- Whorton, M. R., & MacKinnon, R. (2013). X-ray structure of the mammalian GIRK2- $\beta\gamma$ G-protein complex. *Nature* 498(7453), 190–197.
- Wise, H. (2012). The roles played by highly truncated splice variants of G protein-coupled receptors. *Journal of Molecular Signaling* 7(1), 13.
- Wong, W., & Scott, J. D. (2004). AKAP signalling complexes: focal points in space and time. *Molecular Cell Biology* 5(12), 959–970.
- Wu, B., Chien, E. Y., Mol, C. D., Fenalti, G., Liu, W., Katritch, V., et al. (2010). Structures of the CXCR4 chemokine GPCR with small-molecule and cyclic peptide antagonists. *Science* 330(6007), 1066–1071.
- Xia, Z., Gray, J. A., Compton-Toth, B. A., & Roth, B. L. (2003). A direct interaction of PSD-95 with 5-HT_{2A} serotonin receptors regulates receptor trafficking and signal transduction. *The Journal of Biological Chemistry* 278(24), 21901–21908.
- Xiang, Y., Devic, E., & Kobilka, B. (2002). The PDZ binding motif of the beta 1 adrenergic receptor modulates receptor trafficking and signaling in cardiac myocytes. *The Journal of Biological Chemistry* 277(37), 33783–33790.
- Xie, K., Masuho, I., Shih, C. C., Cao, Y., Sasaki, K., Lai, C. W., et al. (2015). Stable G protein-effector complexes in striatal neurons: Mechanism of assembly and role in neurotransmitter signaling. *eLife* 4, Article e10451.
- Xu, Y., Cantwell, L., Molosh, A. I., Plant, L. D., Gazdars, D., Fitz, S. D., et al. (2020). The small molecule GAT1508 activates brain-specific GIRK1/2 channel heteromers and facilitates conditioned fear extinction in rodents. *The Journal of Biological Chemistry* 295(11), 3614–3634.
- Yano, H., Cai, N. S., Xu, M., Verma, R. K., Rea, W., Hoffman, A. F., et al. (2018). Gs- versus Golf-dependent functional selectivity mediated by the dopamine D₁ receptor. *Nature Communications* 9(1), 486.
- Yekkirala, A. S., Banks, M. L., Lunzer, M. M., Negus, S. S., Rice, K. C., & Portoghesi, P. S. (2012). Clinically employed opioid analgesics produce antinociception via μ - δ opioid receptor heteromers in Rhesus monkeys. *ACS Chemical Neuroscience* 3(9), 720–727.
- Yekkirala, A. S., Kalyuzhny, A. E., & Portoghesi, P. S. (2010). Standard opioid agonists activate heteromeric opioid receptors: Evidence for morphine and [d-Ala(2)-MePhe(4)-Glyol(5)]enkephalin as selective μ - δ agonists. *ACS Chemical Neuroscience* 1(2), 146–154.
- Yim, Y. Y., Betke, K. M., McDonald, W. H., Gilsbach, R., Chen, Y., Hyde, K., et al. (2019). The in vivo specificity of synaptic G β and G γ subunits to the α_{2a} adrenergic receptor at CNS synapses. *Scientific Reports* 9(1), 1718.
- Yuan, C., Sato, M., Lanier, S. M., & Smrcka, A. V. (2007). Signaling by a non-dissociated complex of G protein $\beta\gamma$ and α subunits stimulated by a receptor-independent activator of G protein signaling, AGS8. *The Journal of Biological Chemistry* 282(27), 19938–19947.
- von Zastrow, M., & Sorkin, A. (2021). Mechanisms for regulating and organizing receptor signaling by endocytosis. *Annual Review of Biochemistry*. <https://doi.org/10.1146/annurev-biochem-081820-092427>.
- Zhang, M., Patriarchi, T., Stein, I. S., Qian, H., Matt, L., Nguyen, M., et al. (2013). Adenylyl cyclase anchoring by a kinase anchor protein AKAP5 (AKAP79/150) is important for postsynaptic β -adrenergic signaling. *The Journal of Biological Chemistry* 288(24), 17918–17931.
- Zheng, C. Y., Seabold, G. K., Horak, M., & Petralia, R. S. (2011). MAGUKs, synaptic development, and synaptic plasticity. *The Neuroscientist* 17(5), 493–512.
- Zoli, M., Agnati, L. F., Hedlund, P. B., Li, X. M., Ferré, S., & Fuxe, K. (1993). Receptor-receptor interactions as an integrative mechanism in nerve cells. *Molecular Neurobiology* 7(3–4), 293–334.