



## MESSENGER observations of planetary ion characteristics within Kelvin-Helmholtz vortices

Sae Aizawa, Jim Raines, Naoki Terada, Dominique Delcourt, Nicolas Andre

### ► To cite this version:

Sae Aizawa, Jim Raines, Naoki Terada, Dominique Delcourt, Nicolas Andre. MESSENGER observations of planetary ion characteristics within Kelvin-Helmholtz vortices. EPSC-DPS Joint Meeting 2019, Sep 2019, Genève, Switzerland. hal-03563830

**HAL Id: hal-03563830**

**<https://hal.science/hal-03563830>**

Submitted on 10 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **MESSENGER observations of planetary ion characteristics within Kelvin-Helmholtz vortices**

**Sae Aizawa** (1,3), Jim M. Raines (2), Naoki Terada (3), Dominique Delcourt (4), and Nicolas Andre (1)  
(1) IRAP, Toulouse, France (aizawas@pat.gp.tohoku.ac.jp), (2) Department of Climate and Space Sciences and Engineering, The University of Michigan, Ann Arbor, USA, (3) Graduate School of Science, Tohoku University, Sendai, Japan, (4) LPC2E, Orleans, France

### **Abstract**

The MErcury Surface, Space ENvironment, GEochemistry, and Ranging (MESSENGER) spacecraft regularly observed the magnetospheric flanks of Mercury during its orbital phase, 2011-2015. Data from the magnetometer (MAG) and Fast Imaging Plasma Spectrometer (FIPS) allow us to investigate the statistical properties of planetary ions in the presence of Kelvin-Helmholtz (KH) vortices near the duskside magnetopause. We collect data from orbits with clear signatures of KH waves, as well as consecutive orbits that do not have KH signatures, and we compare energy characteristics between KH and non-KH events. Although low planetary counts in FIPS data make the comparison of these characteristics difficult, we find that : (1) large counts of planetary ions are observed in the presence of KH waves, (2) differences in Na<sup>+</sup> energy spectra are only seen inside the magnetosphere, where they show a decelerating signature for KH events for ions with energies above 2.0 keV/e. These results suggest that planetary ions can stagnate within KH vortices and that electric field structures related to KH waves can decelerate planetary ions originating from the magnetotail region.