



Generation of acoustic tones in round jets at a Mach number of 0.9 impinging on a plate with and without a hole

Mathieu Varé, Christophe Bogey

► To cite this version:

Mathieu Varé, Christophe Bogey. Generation of acoustic tones in round jets at a Mach number of 0.9 impinging on a plate with and without a hole. *Journal of Fluid Mechanics*, 2022, 936, 10.1017/jfm.2022.47 . hal-03566628

HAL Id: hal-03566628

<https://hal.science/hal-03566628>

Submitted on 23 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Generation of acoustic tones in round jets at a Mach number of 0.9 impinging on a plate with and without a hole

Mathieu Varé[†] and Christophe Bogey

Univ Lyon, École Centrale de Lyon, INSA Lyon, Université Claude Bernard Lyon I, CNRS, Laboratoire de Mécanique des Fluides et d'Acoustique, UMR 5509, F-69134 Ecully, France

(Received xx; revised xx; accepted xx)

The generation of acoustic tones in four round jets at a Mach number of 0.9 impinging on a plate at a distance $L = 6r_0$ of the nozzle exit, where r_0 is the nozzle radius, has been investigated by large-eddy simulation. Three plates are perforated by a hole of diameter $h = 2r_0, 3r_0$ and $4.4r_0$, centered on the jet axis, whereas the fourth plate has no hole, in order to study the effects of the hole on the jet flow and acoustic fields. In all cases, acoustic tones emerge in the jet near field, upstream but also downstream of the plate for the perforated plates. Their frequencies are similar for all jets and close to those predicted for aeroacoustic feedback loops establishing between the nozzle and the plate, involving flow disturbances convected downstream and waves propagating upstream at the ambient speed of sound. Their levels, however, decrease with the hole diameter, by a few dB for $h \leq 3r_0$ but approximately by 40 dB for $h = 4.4r_0$. The features of the feedback loops are identified by computing two-dimensional space-time correlations and frequency-wavenumber spectra of the pressure fluctuations in the jet mixing layers. These loops are found to be closed by free-stream upstream-propagating guided jet waves, produced by the impingement of vortical structures on the plate for $h \leq 3r_0$ and by the scattering of the jet aerodynamic pressure fluctuations by the hole edges for $h = 4.4r_0$. Finally, an acoustic analogy is used to show that the contributions of the pressure fluctuations on the plate to the radiated noise are dominant.

1. Introduction

It is well known that intense acoustic tones are generated by the impingement of jets on a plate. Such tones were observed experimentally for high subsonic jets by many researchers, including Powell (1953), Neuwerth (1974), Preisser (1979), Ho & Nasseir (1981) and Nasseir & Ho (1982). In these studies, a staging phenomenon of the main tone frequency was measured when the nozzle-to-plate distance increases, which has led Powell (1953) to explain the tone generation by a feedback mechanism between the turbulent structures convected downstream from the nozzle to the plate and the acoustic waves propagating upstream from the plate to the nozzle. Similar feedback loops were noticed for supersonic impinging jets as well. For ideally-expanded supersonic jets, the establishment of these loops has been studied experimentally by Norum (1991) and numerically by Gojon *et al.* (2016) and Bogey & Gojon (2017). For underexpanded jets, the loops have been visualised in the experimental works of Henderson *et al.* (2005), Risborg & Soria (2009), Buchmann *et al.* (2011) and Mitchell *et al.* (2012) and in the simulations of Dauptain *et al.* (2012) and Gojon & Bogey (2017a). The upstream waves closing the feedback loop have also been analyzed in several studies. In their aeroacoustic feedback model, Ho & Nasseir (1981) assumed that they are free-stream acoustic waves propagating outside of the jet. However, they were later identified as guided

[†] Email address for correspondence: mathieu.vare@ec-lyon.fr

jet waves propagating mostly inside the jet, as suggested in the theoretical work of Tam & Ahuja (1990) for round jets and Tam & Norum (1992) for planar jets. These guided jet waves are defined by dispersion relations and classified in modes depending on their radial and azimuthal structures. In particular, their properties allow us to explain the frequencies and the axisymmetric or helical nature of the feedback tones, as highlighted in Gojon *et al.* (2016), Gojon & Bogey (2017a), Bogey & Gojon (2017) and Jaunet *et al.* (2019) for subsonic and supersonic impinging jets. They are involved in other resonance phenomena, for examples in screech generation mechanism, as studied by Gojon & Bogey (2017b), Edgington-Mitchell *et al.* (2018), Mancinelli *et al.* (2019), Edgington-Mitchell (2019), or in jet-plate interactions, as investigated by Jordan *et al.* (2018) and Tam & Chandramouli (2020). The guided jet waves also play a role in the generation of acoustic tones near the nozzle (Towne *et al.* 2017; Brès *et al.* 2018; Bogey 2021b) and in the upstream acoustic far field (Bogey 2021b) of free jets. Using a vortex-sheet model, Towne *et al.* (2017) have shown that within restricted frequency bands, both downstream and upstream-propagating guided jet waves can exist for Mach numbers between 0.82 and 1 and can interact with each other. The near-nozzle tones in both experiments and simulations are located in these bands. In a recent study, Bogey (2021a) investigated numerically the generation of these tones for jets at Mach numbers varying between 0.5 and 2. The tone properties, in terms of frequency, intensity, azimuthal structure, width and prominence, were detailed. They were shown not to strongly vary around Mach numbers of 0.82 and 1. In particular, over the whole Mach number range, the peak frequencies lie within the frequency bands of the free-stream upstream-propagating guided jet waves and the peak levels follow the typical scaling laws of jet noise. Very recently, Bogey (2021b) has shown using data from both experiments and simulations that the near-nozzle tones propagate in the far field in the upstream direction.

Jets impinging on a plate with a hole have been the subject of fewer studies than jets impinging on a flat plate. For subsonic jets, tones similar to those obtained for a non-perforated plate were first observed by Sondhauss (1854) and Rayleigh (1945) and they were referred to as hole tones. In the same way as without a hole, they are generated by a feedback mechanism between the hole edges and the nozzle. Indeed, the same staging phenomenon of the tone frequency as the nozzle-to-plate distance increases was revealed by several experiments, such as those of Chanaud & Powell (1965) and Vinoth & Rathakrishnan (2011). Hole tone generation has been studied over a wide range of Mach numbers. For instance, Langthjem & Nakano (2005) and Matsuura & Nakano (2012) simulated a jet at a Mach number of 0.03 impinging on a plate with a hole located one nozzle diameter from the jet exit. Meganathan & Vakili (2006) explored experimentally the effects of the jet velocity on the hole tone frequencies for Mach numbers varying from 0.2 up to 0.8. They noticed frequencies staging with the jet velocity, suggesting again a feedback mechanism. High subsonic and supersonic jets were considered in the experimental works of Umeda *et al.* (1988) and Umeda & Ishii (1993). Umeda *et al.* (1988) notably visualised the feedback loop between the hole edges and the nozzle for jets at Mach numbers of 0.94 and 1.54 using shadowgraphy. They also noted a tonal radiation at the feedback frequency downstream of the plate. More recently, jets impinging on a plate with a hole have been studied numerically for a Mach number of 3.7 by Kawai *et al.* (2007) and Tsutsumi *et al.* (2015) and for a Mach number of 3.1 by Troyes *et al.* (2019) and Varé & Bogey (2021), with the aim of modelling a rocket launch, during which the supersonic gas jets exhausted by the engines impinge on the launch area and are canalized in a trench dug under the rocket. No feedback loop was found for these high Mach numbers. Concerning the effects of the hole diameter on the feedback loop, Vinoth & Rathakrishnan (2011) compared the flow and pressure fields of two jets at a Mach number of 0.8 impinging on a flat plate and on a plate with a hole of same diameter as the nozzle. The two jets generate tones at similar frequencies, suggesting little effects of the

hole on the feedback mechanism. In the same way, for a Mach number of 0.94, Umeda *et al.* (1988) remarked no significant changes in the frequencies of the tones for hole diameters varying from 1.8 up to 2.4 nozzle diameters. Despite the preceding studies, several questions remain about the influence of the size of the hole on the feedback properties. In particular, the effects of the hole diameter on the tone intensities need to be carefully described. As the hole diameter increases, these intensities are expected to decrease, due to weaker interactions between the jet turbulent structures and the plate. To the best of the authors knowledge, this hypothesis has not been confirmed yet. It is also unclear whether the broadband acoustic levels depend on the hole diameter. Furthermore, it is interesting to study a jet passing through a perforated plate with no impingement of the jet turbulent structures on the plate. For such a jet, the interactions between the flow and the plate are weaker than for the cases with a small hole, which could lead to a less pronounced aeroacoustic resonance, and hence differences in the sound generation mechanisms. Finally, the tonal radiation downstream of the plate pointed out by Umeda *et al.* (1988) has not been examined thoroughly. The noise sources generating this radiation, which may be related to the feedback mechanism, have not been clearly identified. In the present work, the generation of hole tones is investigated by performing the large-eddy simulations of four subsonic round jets impinging on a plate with and without a hole. The jets are at a Mach number of 0.9 and a Reynolds number of 10^5 and are initially highly disturbed. They impinge on a plate located at a distance $L = 6r_0$ from the nozzle exit, where r_0 is the jet radius. Three of the plates have a hole of diameters $h = 2r_0$, $3r_0$ and $4.4r_0$, whereas the fourth plate has no hole. The nozzle-to-plate distance is the same as that in the experiments of Umeda *et al.* (1988), in which the establishment of a feedback loop between the nozzle and the plate was observed. The hole diameters are chosen to study noise generation mechanisms for different interactions between the jet and the plate. Indeed, for $h = 2r_0$ and $3r_0$, the jet impinges on the hole edges whereas for $h = 4.4r_0$, it passes through the hole without being distorted. The first objective of this study is to investigate the influence of the hole and its size on the tone generation. For that purpose, the characteristics of the jet flow and acoustic fields will be detailed, and compared with those obtained for a free jet with the same exit conditions (Bogey (2018, 2021a)). In particular, pressure spectra will be computed in order to highlight the tonal frequencies. The contributions of the first two azimuthal modes will also be examined in order to identify the oscillation modes of the jets. The second aim of this work is to study the upstream part of the feedback loops establishing between the nozzle and the plate, which can be related, given previous works (Tam & Ahuja (1990); Gojon *et al.* (2016); Bogey & Gojon (2017)), to the free-stream upstream-travelling guided jet waves. The production of these waves at the plate, their upstream propagation, as well as their role in generating new instability waves in the jet mixing layers when hitting the nozzle lips, will be revealed by calculating two-dimensional space-time correlations and frequency-wavenumber spectra of the pressure fluctuations in the jet shear layers. Finally, the nature of the noise sources will be investigated by separating the sound components linked to the flow turbulence and those related to the forces exerted by the flow on the plate. To this end, Curle's analogy will be employed to evaluate the contributions of the variations of the pressure on the plate to the pressure field.

The paper is organized as follows. The jet parameters and numerical methods used in the LES are documented in section 2. In section 3, the results of the simulations are presented. In particular, vorticity and pressure snapshots, mean and turbulent flow fields and pressure spectra are shown. The properties of the tones emerging in the spectra are analyzed and compared with the results of aeroacoustic feedback models. Two dimensional space-time correlations and frequency-wavenumber spectra of the pressure fluctuations in the mixing layer are provided. An acoustic analogy is then used to study the nature of the noise sources. Finally, concluding remarks are given in section 4.




Figure 1: Nozzle-exit radial profiles of (a) mean axial velocity $\langle u_z \rangle$ and (b) axial turbulence intensity $\langle u'_z u'_z \rangle^{1/2} / u_j$: — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - - free jet.

2. Parameters

2.1. Jet parameters

The jets computed in this work have a Mach number $M = u_j / c_0$ of 0.9 and a Reynolds number $Re_D = u_j D / \nu$ of 10^5 , where u_j is the jet velocity, D is the nozzle diameter and ν is the air kinematic viscosity. They exhaust from a cylindrical nozzle of radius r_0 , at ambient temperature and pressure $T_0 = 293$ K and $p_0 = 10^5$ Pa. They impinge on a plate located at $L = 6r_0$ downstream of the nozzle exit, with a width of $e = 0.4r_0$. In one case, the plate has no hole whereas for three other ones it has a hole of diameter $h = 2r_0, 3r_0$, and $4.4r_0$. The four cases are referred to as jetnohole, jetsmallhole, jetmediumhole and jetlargehole respectively. The nozzle-to-plate distance, the plate width and the hole diameter $h = 4.4r_0$ are the same as in the experiments of Umeda *et al.* (1988). At the nozzle inlet, a Blasius laminar boundary-layer profile with a boundary-layer thickness of $\delta = 0.15r_0$ is imposed for the velocity. Vortical disturbances non-correlated in the azimuthal direction are added in the boundary layer at $z = -r_0$ to create velocity fluctuations at the nozzle exit, following a procedure described in Bogey *et al.* (2011b). Their amplitude is the same as in the free jet at $M = 0.9$ in Bogey (2018, 2021a) to reach a peak turbulence intensity of 9% at the nozzle exit. The profiles of mean and root-mean-square axial velocities thus obtained at the nozzle exit are represented in figure 1. In figure 1(a), the mean velocity profiles are very similar for the four jets and are close to the Blasius laminar boundary-layer profile imposed at the inlet. As for the turbulent intensities in figure 1(b), they reach a peak of 9% for the four jets, as in the corresponding free jet.

2.2. Numerical and computational parameters

The numerical framework of this study is similar to that of recent simulations of free jets (Bogey (2021a)) and supersonic impinging jets (Gojon *et al.* (2016); Bogey & Gojon (2017)). The unsteady compressible Navier-Stokes equations are solved in cylindrical coordinates (r, θ, z) using an OpenMP based in-house solver. The time integration is performed using a six-stage Runge-Kutta algorithm (Berland *et al.* (2007)) and the spatial derivatives are evaluated with an eleven-point low-dissipation and low-dispersion finite-difference scheme (Bogey & Bailly (2004)). At the end of each time step, a selective filtering is applied in order to remove grid-to-grid oscillations (Berland *et al.* (2007)). This filter enables to dissipate kinetic turbulent energy near the grid cut-off frequency, as a subgrid-scale model (Bogey & Bailly 2006; Fauconnier *et al.* 2013; Kremer & Bogey 2015). Solid and adiabatic wall conditions are imposed to the nozzle and plate walls. The radiation boundary conditions of Tam & Dong (1996) are implemented at the lateral and radial boundaries of the computational domain.

	N_r	N_θ	N_z	$N_r \times N_\theta \times N_z$
jetnohole	559	1024	940	5.4×10^8
jetsmallhole, jetmediumhole, jetlargehole	559	1024	2430	1.4×10^9

Table 1: Mesh parameters: numbers of points N_r , N_θ and N_z in the radial, azimuthal and axial directions, and total number of points.

In these zones, they are combined with sponge zones using grid stretching and Laplacian filtering to avoid significant numerical reflections (Bogey & Bailly (2002)). The singularity at $r = 0$ is treated by applying the method of Mohseni & Colonius (2000). The first point close to the axis is thus located at $r = \Delta r/2$, where Δr is the radial mesh size near the jet axis. The effective azimuthal resolution near the origin of the polar coordinates is reduced down to $2\pi/16$ in order to increase the admissible time step of the simulation (Bogey *et al.* (2011a)).

The mesh grids used for the four simulations contain between 540 millions and 1.4 billions of points, as reported in table 1. They extend out to $r = 15r_0$ in the radial direction and down to $z = 6r_0$ and $z = 40r_0$ in the axial direction for the flat and perforated plates, respectively. The radial mesh spacing, represented in figure 2(a), is equal to $\Delta r = 0.014r_0$ on the jet centerline and decreases down to $\Delta r = 0.0036r_0$ at $r = r_0$ in the shear layers. It then increases to reach a maximum value of $\Delta r_{max} = 0.075r_0$ for $r > 6.2r_0$, which allows a Strouhal number $St = fD/u_j$ of 5.9 to be obtained for an acoustic wave with five points per wavelength. In the azimuthal direction, the grid is uniform and $N_\theta = 1024$ points are used. The axial mesh spacing Δz , presented in figure 2(b), is minimal and is equal to $\Delta z = 0.0072r_0$ at the nozzle exit. It increases and reaches $\Delta z = 0.012r_0$ between $z = 2r_0$ and $z = 4r_0$. Farther downstream, the axial mesh spacing is reduced down to $\Delta z = 0.0072r_0$ near the plate at $z = 6r_0$, as at the nozzle exit. Downstream of the plate, it raises up to $\Delta z = 0.05r_0$ at $z = 40r_0$. The extremum values of the mesh spacings and the stretching rates in both axial and radial directions are the same as in the study of Bogey (2018), where a grid convergence study was performed for the free jet with the same ejection conditions. The simulation time after the transient period is equal to $500r_0/u_j$ for all jets. During the simulations, density, velocities and pressure along the jet axis at $r = 0$, along the lip line at $r = r_0$, on the surface at $r = 15r_0$, on the hole edges at $r = h/2$, and at $z = -2r_0$, $z = 0$, on the faces of the plate at $z = L$ and $z = L + e$, and at $z = 40r_0$ are recorded at a sampling frequency corresponding to a Strouhal number of $St = 12.8$. Density, velocity components and pressure at the azimuthal angles $\theta = 0, 90, 180$ and 270 degrees are also saved at a frequency twice smaller than for the signals at constant r and z . In addition, the azimuthal Fourier coefficients of the density, pressure and velocity fields are computed up to the mode $n_\theta = 4$ for $0 \leq r \leq 15r_0$ and $0 \leq z \leq 40r_0$. The spectra presented in the next sections are calculated from these recordings and they are averaged in the azimuthal direction, when possible.

3. Results

3.1. Snapshots

Snapshots of the vorticity norm obtained down to $z = 10r_0$ for the impinging jets and the free jet are represented in figure 3. Additional videos movie1 and movie2 are also provided for jetnohole and jetmediumhole, respectively. For all jets, the mixing layers are highly disturbed near the nozzle exit. Farther downstream, they thicken with the axial distance because of the development of large-scale vortical structures. For jetnohole, jetsmallhole and




Figure 2: Variations of (a) radial and (b) axial mesh spacings; — positions of the upstream and downstream faces of the plate.




Figure 3: Snapshots in the (z, r) plane of the vorticity norm for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole, (d) jetlargehole and (e) the free jet. The color scales range from 0 up to $15u_j/r_0$, from white to red.

jetmediumhole, in figures 3(a-c), these structures impinge on the plate. A wall jet is then created, as the entire jet flow, or only a part of it, is diverted in the radial direction. For the perforated plates, indeed, a significant part of the jet flow passes through the hole. The flow fields near the impingement region are dominated by large distorted structures, which are particularly visible between $z = 4r_0$ and $z = 6r_0$. Downstream of the plate, coherent structures can also be seen in the flow. This is the case for instance around $z = 8r_0$ in figure 3(c). For jetlargehole, in figure 3(d), the mixing layers do not appear significantly affected by the plate. On the contrary, their development in the axial direction seems very similar to that for the free jet in figure 3(e).

Snapshots of the vorticity norm and of the pressure fluctuations obtained in a whole (z, r) section are given in figure 4. In figures 4(b-d), downstream of the plates with a hole, the shear layers develop down to the end of the potential cores, as for the free jet in figure 4(e). In the pressure fields, for jetnohole, jetsmallhole and jetmediumhole, strong acoustic waves are generated by the impingement of the jet turbulent structures on the plate. Periodically separated wave fronts are observed, indicating a tonal radiation upstream of the plate, but also downstream for jetsmallhole and jetmediumhole. The amplitudes of the waves are of the order of $0.01p_0$, which is much higher than the amplitude obtained in the pressure field of the free jet, typically of $0.001p_0$. In the upstream direction, the waves are slightly stronger for jetsmallhole than for jetmediumhole, but also, more surprisingly, for jetsmallhole than




Figure 4: Snapshots in the (z, r) plane of the vorticity norm in the flow and of the fluctuating pressure outside for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole, (d) jetlargehole and (e) the free jet. The color scales range from 0 up to $7u_j/r_0$ for the vorticity norm, from white to red, and between (a-c) $\pm 0.01p_0$, (d) $\pm 0.0025p_0$ and (e) $\pm 0.001p_0$ for the pressure, from blue to red.

for jetnohole. Moreover, for jetsmallhole and jetmediumhole, the amplitude of the pressure waves appear to be stronger in the upstream region than in the radial direction, which is not the case for jetnohole. Therefore, the acoustic radiations are visibly more directive in the two first cases with a perforated plate than for the no-hole case. For jetlargehole, in figure 4(d), the acoustic waves show a less organized pattern. An acoustic radiation originating from the hole can however be detected upstream and downstream of the plate. A phase jump of the pressure waves is noted across the plate, producing an acoustic pattern similar to that emitted by installed jets (Nogueira *et al.* 2017). The sound radiation seems to be caused by the scattering of the jet aerodynamic pressure fluctuations by the hole edges. The pressure waves thus generated are 4 times weaker than those produced by the other impinging jets. Despite this, they are approximately twice as strong as the mixing-noise acoustic components radiated by the free jet in figure 4(e), namely low-frequency components in the downstream direction and high-frequency components for large radiation angles. For the latter jet, the sound radiation is very directive in the downstream direction, contrary to those of the jets impinging on a perforated plate.

3.2. Flow field

The mean density fields of the jets are represented in figure 5. In all cases, density is lowest in the shear layers. Furthermore, except for jetlargehole and the free jet, compression cells are visible inside the jets. For jetnohole, in figure 5(a), three compression cells are found. The first one is located near $z = 2r_0$, and the second one is between $z = 3r_0$ and $z = 4r_0$. The third, and most intense one, corresponds to the impingement area, and extends from $z = 4.5r_0$ down to the plate in the axial direction, and up to $r = 1.8r_0$ on the plate in the radial direction. Similar compression cells have been found using Schlieren imaging in the experiments of Neuwerth (1974) for a jet at a Mach number of 0.9 impinging on a plate




Figure 5: Mean density fields in the (z, r) plane for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole, (d) jetlargehole and (e) the free jet. The color scale ranges from $0.95\rho_0$ up to $1.05\rho_0$, from blue to red.

located 6 nozzle radii from the nozzle exit. For jetsmallhole, in figure 5(b), the mean density field displays three compression zones upstream of the impingement zone. Density in these zones is stronger than that for jetnohole, which causes a more pronounced radial expansion of the jet between $z = 3r_0$ and $z = 4.5r_0$. Density is highest near the hole in the impingement area, extending only between $z = 5.7r_0$ and $z = 6.4r_0$ and up to $r = 1.4r_0$ in the radial direction in this case. Downstream of the plate, density in the potential core is constant and equal to the ambient density. The mean density field for jetmediumhole, in figure 5(c), looks like that of jetsmallhole. Density in the cells is however slightly lower, and the compression zone near the plate is smaller, and does not reach the hole edges in that case. Downstream of the plate, weak compression cells can be seen down to the end of the potential core. Finally, for jetlargehole, in figure 5(d), the mean density field resembles that of the free jet shown in figure 5(e). In particular, contrary to the other impinging jets, density in the jet potential core is close to the ambient density both upstream and downstream of the plate.

The axial variations of the jet mass flow rate m , normalized by its value $m_0 = \pi r_0^2 \rho_0 u_j$ at the nozzle exit, upstream of the plate are represented in figure 6(a). For jetnohole, the jet mass flow rate first increases, reaches $m = 1.4m_0$ at $z = 5.1r_0$, and then falls to zero on the plate. For jetsmallhole and jetmediumhole, it grows up to $m = 1.38m_0$ at $z = 5.1r_0$ in the first case and $m = 1.39m_0$ at $z = 5.5r_0$ in the second case, and then decreases down to non-zero values on the plate, due to the passing of a part of the jet flow through the hole. For these two jets, the mass flow rate is higher than that for jetnohole between $z = 2r_0$ and $z = 5r_0$. The hole in the plate therefore leads to a stronger entrainment of the fluid surrounding the jet. Finally, for jetlargehole and the free jet, the mass flow rates are very close and increases roughly linearly, in agreement with the experiments of Ricou & Spalding (1961). They are lower than those for the other jets.

The mass flow rates of the wall jets for jetnohole, jetsmallhole and jetmediumhole are shown in figure 6(b). For jetnohole, the mass flow rate increases near the impingement area, reaches a value of $1.4m_0$ at $r = 1.8r_0$, and then does not vary much down to $r = 3r_0$. Farther from the axis, it grows nearly linearly, as the wall jet approaches self similarity. This




Figure 6: Variations of (a) the jet mass flow rate m in the axial direction and (b) the mass flow rate of the wall jet m_{WJ} in the radial direction for — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - - the free jet (Bogey (2018)).

self-similarity is typical of wall jets, refer to the work of Poreh *et al.* (1967), Launder & Rodi (1983) or Van Hout *et al.* (2018), for instance. For jetsmallhole, the mass flow rate of the wall jet also grows continuously with the radial distance, but is lower than for jetnohole, which can be explained by the fact that, due to the hole in the plate, only a fraction of the jet flow is diverted in the wall jet. For jetmediumhole, the mass flow rate is still reduced compared with jetsmallhole, because of the larger hole diameter.

The profiles of centerline mean axial velocity in the impinging jets between the nozzle and the plate are presented in figure 7(a). The centerline velocity decreases down to zero at $z = 6r_0$ on the plate for jetnohole, but remains close to u_j for the three other impinging jets. Small oscillations are however visible between $z = 2r_0$ and $z = 6r_0$ for jetsmallhole and jetmediumhole. They can be linked to the compression cells observed in the density fields in these cases. Such oscillations do not appear for jetlargehole.

The variations of the jet shear-layer momentum thickness δ_θ are represented in figure 7(b). For jetlargehole, the shear-layer spreading is the same as for the free jet. The shear layers of the three other jets for which the mixing-layer turbulent structures impinge, fully or partially, on the plate, develop more rapidly. More precisely, they are thinner between $z = 0$ and $z = 4.5r_0$, and much thicker for $z \geq 4.5r_0$, for jetnohole than for the two other jets. The difference near the plate can be attributed to the stronger wall jet in the case with a non-perforated plate.

The profiles of maximum radial velocity in the wall jets are plotted in figure 7(c). For jetnohole, the maximum velocity increases in the impingement region, where the jet flow is diverted into the radial direction, up to $0.87u_j$ at $r = 2r_0$. For $r \geq 2r_0$, the wall jet then spreads cylindrically, causing a decrease of the velocity as r^{-1} . For jetsmallhole and jetmediumhole, the velocities also first grow in the impingement zone and then decay farther from this zone. They are, however, lower as the hole diameter is larger, leading to peak values only of $0.48u_j$ for jetsmallhole and $0.11u_j$ for jetmediumhole.

The rms values of axial velocity fluctuations estimated at $r = r_0$ along the nozzle-lip line are displayed in figure 8(a) between $z = 0$ and $z = 6r_0$. For the impinging jets, they are very similar to those in the free jet down to $z = 2r_0$. For jetnohole, they fall down to zero on the plate. For jetsmallhole and jetmediumhole, they also decrease between $z = 2r_0$ and $z = 4r_0$, but increase farther downstream. Finally, for jetlargehole, they grow continuously and are slightly higher than those for the free jet.

Finally, the maximum values of radial turbulence intensity in the wall jets are depicted in figure 8(b). For jetnohole, the turbulent intensity shows a local maximum at $r = r_0$. It then increases up to $0.2u_j$ at $r = 4r_0$, and decreases farther from the jet axis. For jetsmallhole and jetmediumhole, the turbulent levels are lower, and reach maximum values of $0.15u_j$ at




Figure 7: Variations of (a) mean axial velocity $\langle u_z \rangle / u_j$ at $r = 0$, and (b) shear-layer momentum thickness δ_θ / r_0 in the axial direction and of (c) the maximum mean radial velocity $\langle u_r \rangle$ in the wall jet: — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - free jet (Bogey (2018)).




Figure 8: Variations of (a) axial turbulence intensity $\langle u'_z u'_z \rangle^{1/2} / u_j$ at $r = r_0$ and (b) maximal radial turbulence intensity $\langle u'_r u'_r \rangle^{1/2} / u_j$ in the wall jet: — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - free jet.

$r = 2r_0$ and of $0.10u_j$ on the hole edges at $r = 1.5r_0$, respectively. The wall jets are therefore weaker in these two cases, as expected.

The spectra of the radial velocity fluctuations computed near the nozzle lip, at $r = r_0$ and $z = 0.4r_0$, for the impinging and the free jets are represented in figure 9 as a function of the Strouhal number. The contributions of the two first azimuthal modes are also shown. For all jets, a broadband hump is observed for both modes around a Strouhal number of $St = 1.7$, which is close to the frequency $St_\theta = f\delta_\theta/u_j = 0.016$ predicted for the Kelvin-Helmholtz instability waves using linear stability analysis (Michalke 1984). For the impinging jets, in addition, tones also emerge in the spectra, at similar frequencies but with levels depending on the presence and diameter of a hole in the plate. The tones are strong for jetnohole, jetsmallhole and jetmediumhole, but much weaker for jetlargehole. In the first case, in figures 9(a-c), very intense tones are found for the axisymmetric mode at $St = 0.41$ for jetnohole and $St = 0.40$ for the two other jets, and at their harmonic frequencies. A small peak also appears for $n_\theta = 1$ around $St = 0.7$. For both modes, the frequencies of the dominant tones are lower than those expected for the most-amplified Kelvin-Helmholtz instability waves. These results suggest that the tones are due to aeroacoustic feedback loops establishing between the nozzle and the plate, *i.e.* that the jet shear layers are forced by the acoustic waves propagating in the upstream direction closing the loops. These feedback loops and upstream-travelling waves will be investigated later. For jetlargehole, in figure 9(d), a peak can also be seen at $St = 0.40$ for $n_\theta = 0$. However, the peak has a much smaller amplitude than the dominant tones for the other impinging jets, and does not appear in the spectrum of the full velocity signal, indicating a weaker aeroacoustic resonance in this case.




Figure 9: Power spectral densities of radial velocity fluctuations u'_r obtained at $r = r_0$ and $z = 0.4r_0$ for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole, (d) jetlargehole and (e) the free jet; — full signal, — $n_\theta = 0$ and — $n_\theta = 1$, - - $St_\theta = f\delta_\theta/u_j = 0.016$.

The spectra of radial velocity fluctuations computed farther downstream in the jet shear layer, at $r = r_0$ and $z = r_0, 5r_0, 7r_0$ and $12r_0$, are shown in figure 10. At $z = r_0$ downstream of the nozzle, in figure 10(a), the spectra are very similar to those obtained at $z = 0.4r_0$. Tones are found at $St \approx 0.40$ and harmonic frequencies for jetnohole, jetsmallhole and jetmediumhole, but not for jetlargehole. At $z = 5r_0$, just upstream of the plate, in figure 10(b), tones appear in the spectra, only for the first three impinging jets as previously. Compared with the tones at $z = r_0$, they have the same frequencies, but they emerge more sharply. Downstream of the plate, in figures 10(c,d), peaks are still found in the velocity spectra for jetsmallhole and jetmediumhole, but not for jetlargehole. Close to the plate, at $z = 7r_0$, in figure 10(c), tones emerge strongly at $St = 0.4$ and $St = 0.8$ for jetsmallhole and at $St = 0.4$ for jetmediumhole whereas farther downstream, at $z = 12r_0$, in figure 10(d), only a weak tone is observed at $St = 0.4$ for these two jets. Therefore, the presence of aeroacoustic feedback loops upstream of the plate affects the jet flow development downstream of the plate. This is particularly the case in the vicinity of the plate, which explains the formation of large coherent structures in this region in the vorticity fields of figure 3(b,c).

3.3. Pressure field

3.3.1. Overall Sound Pressure Levels

The isocontours of the overall sound pressure levels obtained in the (z, r) section for the five jets are represented in figure 11(a-e). Upstream of the plate, for all impinging jets, the isolines form a lobe oriented in the upstream direction and enclosing the plate center, suggesting a link with the sound waves created in the impingement area. For jetnohole, in figure 11(a), a second lobe can be seen close to the plate. It extends in the radial direction and may be due to noise components radiated in the wall jet, which is most developed for this jet. Downstream of the plate, for jetsmallhole and jetmediumhole, in figures 11(b,c), the isolines appear as a lobe centered on the hole, highlighting an acoustic radiation from this region. For jetlargehole, in figure 11(d), their shape is different than that for the two





Figure 10: Power spectral densities of radial velocity fluctuations u'_r obtained at $r = r_0$ and (a) $z = r_0$, (b) $z = 5r_0$, (c) $z = 7r_0$ and (d) $z = 12r_0$ for — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - - the free jet.






Figure 12: Sound pressure levels (SPL) at (a) $z = 0$ and $r = 1.5r_0$ and (b) $z = 12r_0$ and $r = 15r_0$ as a function of the Strouhal number $St = fD/u_j$; — jetnohole, — jetsmallhole, — jetmediumhole, — jetlargehole and - - - free jet.

3.3.2. Pressure spectra

The pressure spectra obtained for the four impinging jets near the nozzle and downstream of the plate, when possible, are plotted in figure 12 as a function of the Strouhal number, together with the spectra for the free jet.

Near the nozzle, at $z = 0$ and $r = 1.5r_0$, in figure 12(a), several intense tones emerge for jetnohole, jetsmallhole and jetmediumhole at very similar frequencies regardless of the presence of a hole in the plate and its size. The dominant tone is located at a Strouhal number of $St = 0.41$ for jetnohole, and of $St = 0.40$ for jetsmallhole and jetmediumhole. In the three cases, it is 25 to 30 dB greater than the broadband noise level. Secondary tones, not harmonics of the dominant tones, appear at $St = 0.69$ for jetnohole, $St = 0.64$ for jetsmallhole and $St = 0.60$ for jetmediumhole. For jetlargehole, a tonal peak is also found at the Strouhal number of 0.40. However, the acoustic levels are much weaker than for the other impinging jets, and are reduced by 20 to 55 dB compared with jetnohole. They are even close to the levels estimated for the free jet for Strouhal numbers higher than 0.7, and only 10 to 15 dB higher for lower frequencies. Tonal peaks can also be noted in the spectra for the free jet, at frequencies comparable to those of the tones for the impinging jets, namely $St = 0.41, 0.70, 1.0, 1.30$ and 1.63 (Bogey (2021a)).

For the impinging jets, the tones can be assumed to be generated by feedback mechanisms between the nozzle and the plate (Powell (1953)), consisting of two steps. During the first step, coherent structures are convected in the shear layer to the plate, where their impingement produces acoustic waves. During the second step, these waves propagate upstream to the nozzle, where they excite the shear layer instability waves, creating new coherent structures and closing the feedback loop. A model was proposed by Ho & Nosseir (1981) to predict the frequencies of such a loop. It was built by considering the feedback period as the sum of two characteristic times. The first one is the time of convection of the flow structures from the nozzle to the plate and the second one is the time of propagation of the acoustic waves in the upstream direction at the ambient speed of sound c_0 . The feedback frequency f can thus be predicted by

$$f = \frac{Nu_c}{L(1 + M_c)} \quad (3.1)$$

where N is an integer, u_c is the mean convection velocity and $M_c = u_c/c_0$ is the convection Mach number. The integer N represents the order of the feedback mode and corresponds to the number of coherent structures between the nozzle and the plate. The velocity u_c , computed for all the jets in this study, is close to the classical approximation of $(2/3) \times u_j$. The feedback frequencies given by the model for $N = 3$ and $N = 5$ using this value for the

convection velocity are equal to $St = 0.42$ and 0.69 , respectively. The first value is in good agreement with the Strouhal numbers of 0.40 and 0.41 of the dominant tones emerging in the spectra. The second frequency value is close or equal to the Strouhal numbers of the tones at $St = 0.60, 0.64$ and 0.69 in the spectra of jetnohole, jetsmallhole and jetmediumhole. These results support the establishment of feedback loops at the tone frequencies. For the free jet, the tonal peaks in the spectra are due to the presence of guided jet waves, as documented by Towne *et al.* (2017) and Schmidt *et al.* (2017) and discussed very recently by Bogey (2021a). These waves will be examined in section 3.4.2. Interestingly, the feedback tones for the impinging jets and the peaks for the free jets emerge at very similar frequencies. The feedback frequencies therefore appear to be mainly determined by the properties of the upstream-propagating guided jet waves. Regarding the tones intensities, the feedback loops produce strong tones for hole diameters $h \leq 3r_0$ and much weaker tones for $h = 4.4r_0$, when the flow structures do not impinge on a solid surface.

The pressure spectra computed at $z = 12r_0$ and $r = 15r_0$ downstream of the plate are represented in figure 12(b). For jetsmallhole and jetmediumhole, intense tones emerge at $St = 0.40$ and harmonic frequencies, corresponding to the tone frequencies obtained upstream of the plate. For jetlargehole, no tones are found, but a hump emerging by about 10 dB of the broadband noise is observed for $St \leq 0.7$. This low-frequency hump resembles that noticed for installed jets by Lawrence *et al.* (2011), supporting the idea that the scattering of the jet pressure by the plate strongly contributes to the noise radiated by jetlargehole. Regarding the sound levels, they are reduced by 5 to 30 dB with respect to the previous impinging jets. Compared with those for the free jet, they are 10 dB stronger for $St \leq 0.7$ and similar for $St \geq 0.7$.

For the four impinging jets, the contributions of the first two azimuthal modes to the pressure spectra at $z = 0$ and $r = 1.5r_0$ are represented in figure 13 as a function of the Strouhal number. They are compared with the results obtained for the free jet, plotted in dashed lines. In figure 13(a), for jetnohole, the tones at $St = 0.41, 0.82$ and 1.20 are related to the axisymmetric mode, whereas the tone at $St = 0.68$ is associated with the first helical mode. This result is in agreement with the experimental work of Panickar & Raman (2007), revealing the coexistence of an axisymmetric mode and an helical instability mode for impinging jets at a Mach number higher than 0.89 . The peaks at $St = 0.41$ and $St = 0.70$ for the free jet are also linked to the modes $n_\theta = 0$ and $n_\theta = 1$, respectively, as for jetnohole. Therefore, the impingement of the jet on the plate does not change the azimuthal structure of the jet flow oscillations at a given frequency. In figures 13(b) and 13(c), the spectra for jetsmallhole and jetmediumhole are very similar to those for jetnohole in figure 13(a). The dominant tones at $St = 0.40$ and higher harmonics emerge for $n_\theta = 0$ whereas the tones at $St = 0.65$ for jetsmallhole and $St = 0.60$ for jetmediumhole occur for $n_\theta = 1$. Thus, the hole in the plate does not affect the axisymmetric or helical nature of the tones. Finally, for jetlargehole, in figure 13(d), the tonal peak around $St = 0.40$ emerges for $n_\theta = 0$, as for the other jets.

The Strouhal numbers of the tones emerging at $z = 0$ and $r = 1.5r_0$ for $n_\theta = 0$ and 1 are represented in figure 14 as a function of the hole diameter h/r_0 , with $h/r_0 = \infty$ corresponding to the free jet case. The frequencies predicted for aeroacoustic feedback loops by the model of Ho & Nosseir (1981) are also drawn using grey lines. For all impinging jets, four tones are found at similar Strouhal numbers, namely $St = 0.40$ and 1.20 for $n_\theta = 0$ in figure 14(a) and around $St = 0.7$ and 1.3 for $n_\theta = 1$ in figure 14(b), near the lines obtained using $N = 3, 5, 7$ and 9 in the feedback loop model. The tonal frequencies for $n_\theta = 0$ do not vary with the hole diameter, whereas those for $n_\theta = 1$ change slightly, for unclear reasons. It can be noted that for the free jet, the near-nozzle peaks, attributed to the upstream-propagating guided jet modes (Towne *et al.* (2017); Brès *et al.* (2018); Bogey (2021a)), also appear at the




Figure 13: Sound pressure levels (SPL) at $z = 0$ and $r = 1.5r_0$ for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole and (d) jetlargehole for the azimuthal modes $n_\theta = 0$ and $n_\theta = 1$; SPL for $n_\theta = 0$ and $n_\theta = 1$ for the free jet.




Figure 14: Strouhal numbers of the tones as a function of the hole diameter for (a) $n_\theta = 0$ and (b) $n_\theta = 1$, — frequencies obtained from equation (3.1) for increasing values of N .

four frequencies mentioned above. These results further support a link between the feedback loops and these modes. Finally, for $n_\theta = 0$, additional tones are observed, namely two tones at $St = 0.80$ and 1.20 for $h \leq 3r_0$, which are the harmonics of the first tone at $St = 0.40$ and a tone at $St = 0.60$ for $h = 4.4r_0$.

The levels of the dominant tones at $z = 0$ are represented in figure 15 as a function of the hole diameter for the modes $n_\theta = 0$ and $n_\theta = 1$. Two radial positions, $r = 1.5r_0$ and $15r_0$, are considered, and only the tones emerging by more than 3 dB from the broadband levels are displayed. For $n_\theta = 0$ in figure 15(a), near the nozzle, the level of the dominant tone is strong for $h \leq 3r_0$, reaching a peak of 180 dB for $h = 2r_0$. The amplitudes of the tone for jetsmallhole and jetmediumhole are slightly higher than for jetnohole. For $h = 4.4r_0$, the tone level is drastically reduced by about 40 dB. This can be explained by the fact that for $h \leq 3r_0$ the upstream sound waves are created by the impingement of the jet turbulent structures on the plate, whereas for $h = 4.4r_0$, they are produced by the scattering of the jet pressure fluctuations by the plate. In the latter case, the tone level is still 15 dB higher than




Figure 15: Dominant tone levels at $z = 0$ and $\circ r = 1.5r_0$ and $\triangle r = 15r_0$ as a function of the hole diameter for (a) $n_\theta = 0$ and (b) $n_\theta = 1$.

that for the free jet. Farther away from the jet axis, at $r = 15r_0$, the tone levels are highest for jetnozzle and decrease monotonously with the hole diameter, contrary to the levels of the near-nozzle tones. This is expected given the different directivities of the jets highlighted in figure 4. For $n_\theta = 1$, in figure 15(b), the level of the dominant near-nozzle tone also decreases continuously with the hole diameter. It is reduced by 5 dB between $h = 0$ and $1.5r_0$, then by 15 dB for $h = 4.4r_0$, tending towards the level for the free jet.

3.4. Investigation of the feedback loop

3.4.1. Visualisation of the feedback loop

In order to visualise the feedback loop, two-dimensional spatial correlations of the jet pressure fields are computed in a section (z, r) . The pressure fluctuations p' at a reference point (z_1, r_1) at time t are correlated with the fluctuations of pressure in the section (z, r) at time $t + \delta t$, defining the dimensionless coefficient \mathcal{R}

$$\mathcal{R}(r, z, \delta t) = \frac{\langle p'(r_1, z_1, t) p'(r, z, t + \delta t) \rangle}{\langle p'^2(r_1, z_1, t) \rangle^{1/2} \langle p'^2(r, z, t) \rangle^{1/2}} \quad (3.2)$$

where δt is the time delay between the signals and $\langle . \rangle$ denotes time averaging. In this way, the structures and time variations of the waves correlated with the pressure fluctuations at the reference point are extracted. This method has notably been used to investigate noise generation in free jets at a Mach number of 0.9 in a recent work (Bogey (2019)).

In the present study, due to the presence of tones emerging by more than 20 dB in the sound spectra, mainly associated with the mode $n_\theta = 0$, the pressure signals are strongly correlated. Therefore, links between the pressure at a reference point and flow structures can not be clearly evidenced by computing correlations from the full signals or from the axisymmetric mode. However, the tones are much less emerging for the mode $n_\theta = 1$. Correlations calculated for this mode are consequently shown in what follows.

The coefficients \mathcal{R} are first evaluated for $n_\theta = 1$ for the four impinging jets for a reference point placed near the jet nozzle at $z_1 = 0$ and $r_1 = 1.5r_0$, in order to specifically examine the characteristics of the upstream acoustic radiation. Similar results are obtained for jetnozzle, jetsmallhole and jetmediumhole. The correlations for the latter jet are represented in figure 16, for time delays δt varying from $-15r_0/u_j$ to $6r_0/u_j$. They can also be seen in the supplementary video movie3. For $\delta t = -15r_0/u_j$ and $-12r_0/u_j$, in figures 16(a,b), the correlation levels are highest around the shear layers of the jet, at $z \simeq 2.5r_0$ at $\delta t = -15r_0/u_j$ and $z \simeq 4r_0$ at $\delta t = -12r_0/u_j$. They are linked to flow coherent structures convected in the jet flow direction, which constitute the downstream part of the feedback mechanism. For $\delta t = -9r_0/u_j$ and $-6r_0/u_j$, in figures 16(c,d), the correlation levels increase and they are strongest near the hole, as the flow structures impinge on the hole edges. For $\delta t = -3r_0/u_j$, in




Figure 16: Correlations \mathcal{R} of $p'(r = 1.5r_0, z = 0, t)$ with $p'(r, z, t + \delta t)$ for $n_\theta = 1$ for (a) $\delta t = -15r_0/u_j$, (b) $\delta t = -12r_0/u_j$, (c) $\delta t = -9r_0/u_j$, (d) $\delta t = -6r_0/u_j$, (e) $\delta t = -3r_0/u_j$, (f) $\delta t = 0$, (g) $\delta t = 3r_0/u_j$ and (h) $\delta t = 6r_0/u_j$ for jetmediumhole; — circle centered on $(z = L, r = 1.5r_0)$. The color scale ranges between ± 1 , from blue to red.

figure 16(e), the correlation levels are greater than 0.5, which is significantly higher than the levels for the previous time delays. A curved region of strong positive correlations, in red, is aligned with a circle centered on the hole edge, at $z = L$ and $r = 1.5r_0$. It can be related to an acoustic wave produced by the jet impingement on the plate, which forms the upstream part of the feedback loop. The correlations are significant outside but also inside of the jet. The propagation of an upstream wave travelling at the ambient sound velocity in the jet column is thus highlighted, as noticed for supersonic impinging jets by Bogey & Gojon (2017). This wave is related to the guided jet waves studied later in section 3.4.2. For $\delta t = 0$, in figure 16(f), the correlation level is equal to 1 at the point (z_1, r_1) , as expected. Moreover, as for the previous time delay, the acoustic wavefront has a circular shape, and the correlations are strong both in the jet flow and near pressure fields. For later time delays $\delta t = 3r_0/u_j$ and $6r_0/u_j$, in figures 16(g,h), the correlation levels are lower than 0.2, except for two regions. The first one is due to the upstream propagation of the acoustic waves. The second one is found around the shear layers at $z = r_0$ for $\delta t = 3r_0/u_j$ and at $z = 3r_0$ for $\delta t = 6r_0/u_j$, revealing the generation and convection of a new coherent structure in the flow, and thus the downstream part of a new feedback cycle.

The correlations \mathcal{R} obtained for jetlargehole are displayed in figure 17 and in the additional video movie4 for time delays δt varying from $-15r_0/u_j$ to $6r_0/u_j$. For $\delta t = -15r_0/u_j$, in figure 17(a), the correlations are weak, with levels lower than 0.3. They are stronger for $\delta t = -12r_0/u_j$, in figure 17(b), exhibiting a wavepacket structure with four lobes around the shear layer at $z \simeq 3r_0$. Later, the wavepacket structure is convected down to the plate, as in figure 17(c) for $\delta t = -9r_0/u_j$ for instance. It is then scattered by the hole edges for $\delta t = -9r_0/u_j$ and $-6r_0/u_j$, in figures 17(c,d). The scattering of the aerodynamic pressure fluctuations generates acoustic waves, causing high correlation levels upstream and downstream of the plate for $\delta t = -3r_0/u_j$, aligned with a circle centered on the hole edge in figure 17(e). Upstream, in particular, the waves propagate up to the reference point in figure 17(f). As for jetmediumhole, non-negligible correlation levels are also found inside the jet, indicating the propagation of an upstream wave at the sound speed in the jet potential core. This wave belongs to the guided jet modes described in section 3.4.2. Afterward, the correlation levels finally decrease. Spots of significant correlations are nevertheless found




Figure 17: Correlations \mathcal{R} of $p'(r = 1.5r_0, z = 0, t)$ with $p'(r, z, t + \delta t)$ for $n_\theta = 1$ for (a) $\delta t = -15r_0/u_j$, (b) $\delta t = -12r_0/u_j$, (c) $\delta t = -9r_0/u_j$, (d) $\delta t = -6r_0/u_j$, (e) $\delta t = -3r_0/u_j$, (f) $\delta t = 0$, (g) $\delta t = 3r_0/u_j$ and (h) $\delta t = 6r_0/u_j$ for jetlargehole, — circle centered on $(z = L, r = 2.2r_0)$. The color scale ranges between ± 1 , from blue to red.

near the shear layers at $z \simeq r_0$ for $\delta t = 3r_0/u_j$ in figure 17(g) and $z \simeq 4r_0$ for $\delta t = 6r_0/u_j$ in figure 17(h). They can be attributed to a flow coherent structure convected downstream, initiating a new feedback cycle, as mentioned above for jetmediumhole.

Two-dimensional spatial correlations are also computed for $n_\theta = 1$ for a reference point located at $z = 2L$ and $r = 15r_0$ to shed light on the sound generation downstream of the plate. The results obtained for jetmediumhole and jetlargehole are represented in figures 18 and 19, respectively, for the time delays $\delta t = -15r_0/u_j$, $-10r_0/u_j$ and $-5r_0/u_j$. For $\delta t = -15r_0/u_j$, weak correlations are found for jetmediumhole near the hole in figure 18(a). The correlations for jetlargehole in figure 19(a) are stronger and reveal a wavepacket structure just outside of the jet shear layers upstream of the plate. Later, for $\delta t = -10r_0/u_j$ and then for $\delta t = -5r_0/u_j$, curved regions of correlation levels higher than 0.6 and centered on the hole edge are observed for both jets in figures 18(b,c) and 19(b,c). Therefore, the sound waves radiated downstream of the plate are related to phenomena occurring close to the hole edge. These phenomena can be of different nature depending on the hole diameter and consist of the interactions of the hole edge with the vortical structures passing through the hole for jetmediumhole and with the jet aerodynamic pressure waves, surrounding the flow structures, for jetlargehole.

3.4.2. Guided jet modes

In order to discuss the nature of the upstream-travelling waves closing the feedback loops in the impinging jets, frequency-wavenumber spectra have been computed as recently in Bogey (2021a). They are calculated from the pressure fluctuations in the jet shear layer at $r = r_0$ between $z = 0$ and $z = 4r_0$ for jetnohole and jetsmallhole and between $z = 0$ and $z = 10r_0$ for the other jets for the two first azimuthal modes $n_\theta = 0$ and 1. The spectra obtained for jetmediumhole, jetlargehole and the free jet are represented in figure 20 as a function of k and St , for negative wavenumbers. The results for jetnohole and jetsmallhole are not shown but they look like those for jetmediumhole. The dispersion curves of the guided jet waves given by a vortex-sheet model (Tam & Ahuja (1990)) are also displayed using red lines. Finally, the tone frequencies emerging in the LES pressure spectra in section 3.3.2 are indicated by red triangles placed at $k = 0$.




Figure 18: Correlations \mathcal{R} of $p'(r = 15r_0, z = 12r_0, t)$ with $p'(r, z, t + \delta t)$ for $n_\theta = 1$ for (a) $\delta t = -15r_0/u_j$, (b) $\delta t = -10r_0/u_j$ and (c) $\delta t = -5r_0/u_j$ for jetmediumhole; — circle centered on $(z = L, r = 1.5r_0)$. The color scale ranges between ± 1 , from blue to red.




Figure 19: Correlations \mathcal{R} of $p'(r = 15r_0, z = 12r_0, t)$ with $p'(r, z, t + \delta t)$ for $n_\theta = 1$ for (a) $\delta t = -15r_0/u_j$, (b) $\delta t = -10r_0/u_j$ and (c) $\delta t = -5r_0/u_j$ for jetlargehole; — circle centered on $(z = L, r = 2.2r_0)$. The color scale ranges between ± 1 , from blue to red.

564 The dispersion curves start on the sonic line $k = -\omega/c_0$. Farther from this line, as the
 565 wavenumber increases, in absolute value, the frequency of the waves increases up to a local
 566 maximum, then it is reduced down to a local minimum and finally increases again. Three kinds
 567 of waves can be defined depending on their positions on the dispersion curves, as proposed by
 568 Towne *et al.* (2017). The waves between the sonic line and the local maximum have a negative
 569 group velocity $v_g = d\omega/dk$ and a support outside the jet. They can be referred to as upstream-
 570 travelling free-stream waves. In the other cases, the waves are entirely confined in the jet.
 571 They propagate downstream for the waves between the two local extremum frequencies on
 572 the curves and upstream otherwise. They will be denominated as downstream- and upstream-
 573 propagating duct-like waves, respectively. For jetmediumhole, in figures 20(a,d), lines of high
 574 energy extending over the whole range of wavenumbers are found at the Strouhal numbers of
 575 the tones in the LES sound spectra. Except for the tone at $St = 1.20$ for $n_\theta = 0$, they intersect
 576 the dispersion curves of the guided jet waves predicted by the vortex-sheet model, in regions
 577 where the waves are the upstream-travelling free-stream ones. Therefore, these waves are
 578 likely to close the feedback loops generating the tones. As for the tone at $St = 1.20$, it can be
 579 expected to result from nonlinear effects between the two tones at $St = 0.40$ and $St = 0.80$.
 580 For jetlargehole, in figures 20(b,e), no lines are visible at the Strouhal numbers of the tones,
 581 contrary to jetmediumhole, which is most probably due to the quite weak tones in this case
 582 where the hole diameter is the largest. Curved bands are however clearly observed near the
 583 dispersion curves of the guided jet waves for the vortex-sheet model. They are very similar




Figure 20: Frequency-wavenumber spectra of the pressure fluctuations at $r = r_0$ for (a,d) jetmediumhole, (b,e) jetlargehole and (c,f) the free jet for (top) $n_\theta = 0$ and (bottom) $n_\theta = 1$; — dispersion curves and * lower limits of the guided jet modes for a vortex-sheet model, ◀ LES tone frequencies and - - $k = -\omega/c_0$. The greyscale levels spread over 25 dB.

to those found for the free jet in figures 20(c,f). On these bands, the intensity is high near the line $k = -\omega/c_0$, decreases farther from it and is negligible for low wavenumbers. These variations of the intensity can be explained by the fact that the guided jet waves close to the line $k = -\omega/c_0$ are free-stream waves, whereas those far from the line are duct-like waves (Tam & Ahuja 1990; Towne *et al.* 2017). Concerning the LES tones, their frequencies lie in the frequency bands of the free-stream upstream-propagating modes, highlighting again the role of these waves in the feedback loops. Moreover, as seen previously in figure 14, the tone frequencies do not vary much with the hole diameter. Despite different noise generation mechanisms, namely the scattering of the jet pressure for jetlargehole and vortical structures impinging on the plate for the other jets, they seem to be mainly determined by the frequency of the free-stream upstream-propagating guided jet waves. The tone frequencies are also close to those produced by jet-flap interactions at the same Mach number (Jordan *et al.* 2018), supporting again the idea that they are weakly influenced by the nature of the interactions between the jet and the plate.

To visualise both the downstream and the upstream parts of the feedback loops, the frequency-wavenumber spectrum obtained for the tonal case jetmediumhole for $n_\theta = 0$ is presented in figure 21 for negative but also positive wavenumbers. The lines found at the tone frequencies for $k \leq 0$ extend for $k \geq 0$. In the latter region, the strongest levels are close to the line $k = \omega/u_c$ with $u_c = (2/3) \times u_j$. Therefore, the downstream component of the feedback loops consists of flow structures convected in the jet mixing layers at the velocity u_c , as expected. These structures develop at the tone frequencies, due to the forcing of the shear layers near the nozzle lips at these frequencies, evidenced in the velocity spectra of figure 9.

3.5. Noise source investigation using Curle's analogy

3.5.1. Curle's analogy

The noise radiated by flows interacting with rigid bodies can be investigated using acoustic analogies. In particular, an analogy, in which the noise generated by such flows consists of




Figure 21: Frequency-wavenumber spectrum of the pressure fluctuations at $r = r_0$ for jetmediumhole for $n_\theta = 0$; — dispersion curves and * lower limits of the guided jet modes for a vortex-sheet model, ◄ LES tone frequencies, - - - $k = -\omega/c_0$ and - - - $k = \omega/u_c$. The greyscale levels spread over 25 dB.

two components, was proposed by Curle (1955). The first component is the aerodynamic noise produced by the flow turbulence, whereas the second one, dominant at low Mach numbers, is related to the forces exerted on the solid bodies. This type of analogy allows the separation between the direct sound field, linked to the first component, and the reflected and scattered ones, associated with the second component. It has been applied to various flows. It has allowed, for example, Gloerfelt *et al.* (2003) to investigate the noise generated by a cavity flow at a Mach number of 0.7 and Gloerfelt *et al.* (2005) to study the sound radiated by the flow around a cylinder at a Mach number of 0.12. In the first work, the authors showed that the pressure far-field predicted by the analogy is close to that obtained by a DNS computation. In an older study, Curle's analogy was employed by Preisser (1979) to correlate the far-field noise with the pressure on the plate for subsonic impinging jets.

For the present impinging jets, the term of Curle's analogy associated with the variations of pressure on the solid surfaces is considered to evaluate the acoustic field upstream and downstream of the plate. The amplitude of the pressure fluctuations due to this contribution is given by

$$p'(\mathbf{x}, t) = \frac{1}{4\pi} \frac{\partial}{\partial x_i} \int_S \frac{1}{R} P_i \left(\mathbf{y}, t - \frac{R}{c_0} \right) dS(\mathbf{y}) \quad (3.3)$$

where $R = |\mathbf{x} - \mathbf{y}|$ is the distance between a source on the plate and the observer and $P_i = (p\delta_{ij} - \tau_{ij})l_j$ is the force per unit area exerted on the fluid by the plate in the direction x_i , with τ_{ij} the viscous stress tensor and l_j the coordinates of the outward normal from the fluid. Here, the pressure normal to the plate is assumed to be dominant compared with the shear stresses, as in the work of Preisser (1979). Under this assumption and by considering that the observer is in the acoustic far field, the equation (3.3) can be simplified as

$$p'(\mathbf{x}, t) = \frac{1}{4\pi c_0} \frac{\partial}{\partial t} \int_S \frac{x_i - y_i}{R^2} p \left(\mathbf{y}, t - \frac{R}{c_0} \right) l_i dS(\mathbf{y}) \quad (3.4)$$

where the time derivative is evaluated using fourth-order centered finite differences. Equation (3.4) will be used to estimate the fluctuations of pressure both upstream and downstream of the plate. It may be noted that these fluctuations can also be evaluated using an analytical Green's function designed for a perforated plate, such as in the work of Howe (1979). This approach has not been carried out in the present work.

To examine the source terms in equation (3.4), the pressure spectra obtained on the plate at $z = L$ and $r = 3r_0$ are displayed in figure 22 for the four impinging jets. The levels are




Figure 22: Pressure spectra on the plate at $z = L$ and $r = 3r_0$ for — jetnohole, — jetsmallhole, — jetmediumhole and — jetlargehole.

highest for jetnohole and they decrease with the hole diameter. For jetnohole, jetsmallhole and jetmediumhole, tones are found at a Strouhal number of 0.4 and its harmonic frequencies, while for jetlargehole, a hump emerging by about 15 dB from the broadband levels is observed for frequencies lower than $St = 0.7$. In all cases, the spectra look like those near the nozzle in figure 12, suggesting that the spectral content of the sound waves computed using the analogy will be similar to that of the LES sound fields. The resemblance between the pressure spectra on the plate and those in the acoustic field has also been noted for supersonic impinging jets in the numerical study of Gojon & Bogey (2018).

3.5.2. Acoustic field upstream of the plate

The pressure fields obtained upstream of the plate using equation (3.4) for the four impinging jets are represented in figures 23(a-d). For all jets, periodic waves originating from the plate are observed. They look like the acoustic waves propagating in the LES pressure fields, given in figures 23(e-h), as expected. In the two cases, for $r \geq 5r_0$, the levels of fluctuating pressure are close and the wavefronts are located at similar positions. In particular, the sound waves radiated in the radial direction are well reproduced using the analogy for jetsmallhole and jetmediumhole, in figures 23(b,f) and 23(c,g). Therefore, the surfacic term of Curle's analogy is able to predict an acoustic radiation consistent with the LES results, verifying the validity of the analogy for the present study. Close to the jet flow, for $r \leq 5r_0$, some discrepancies, such as a phase shift, can be seen between the pressure waves obtained from the analogy and the LES. However, the comparison between the two pressure fields is not fully relevant here due to the fact that equation (3.4) is reliable only in the far field.

The acoustic spectra computed near the nozzle at $z = 0$ and $r = 1.5r_0$ using equation (3.4) are compared with the LES spectra in figure 24. For the four impinging jets, a good agreement is found. In particular, the effects of the hole and its size are well reproduced. For jetnohole, jetsmallhole and jetmediumhole, in figures 24(a-c), the spectra obtained from the analogy and the LES exhibit tones at the same frequencies, including the dominant tone at $St = 0.40-0.41$ and its harmonics, unsurprisingly given the frequency contents of the pressure spectra on the plate. However, the tone levels do not match perfectly. For instance, the dominant tones are 4 to 10 dB stronger in the LES than using the acoustic analogy. These discrepancies may be related to the noise radiation of aerodynamic phenomena such as the distortion of the vortical structures near the plate, not taken into account using the analogy. As for the broadband levels obtained using the analogy, for jetsmallhole and jetmediumhole, they are similar to those of the LES, with less than 3 dB of differences. For jetnohole, they are close to the LES levels for $St > 1$, but they are about 5 dB lower for $St < 1$. This result suggests that in this case, the contribution of the pressure fluctuations on the plate to the broadband noise is dominant at high frequencies.




Figure 23: Pressure fields obtained upstream of the plate (top) using equation (3.4) and (bottom) from LES for (a,e) jetnohole, (b,f) jetsmallhole, (c,g) jetmediumhole and (d,h) jetlargehole. The color scales range between (a,b,c,e,f,g) $\pm 0.01p_0$, and (d,h) $\pm 0.0025p_0$, from blue to red.

For jetlargehole, in figure 24(d), the spectra from the analogy and the LES exhibit peaks at $St = 0.40$ with comparable amplitude. Around the peaks, for $St < 0.8$, the levels obtained using the analogy are close to those of the LES with discrepancies lower than 3 dB, implying that the scattering of the jet aerodynamic pressure by the perforated plate is the main noise source. For $St > 0.8$, the levels are about 15 dB lower using the analogy than in the LES, suggesting that the high-frequency noise components are related to the flow turbulence. It can also be noted that the LES spectrum contains weak peaks for $0.8 \leq St \leq 3$ due to the presence of upstream-travelling guided jet waves (Towne *et al.* (2017), Bogey (2021a)), as mentioned previously in section 3.3.2. Naturally, these peaks are missing in the spectrum determined using Curle's analogy.

As mentioned in section 3.5.1, the reflected and scattered sound fields are related to the surfacic term of Curle's analogy, whereas the direct sound field consists of the jet mixing noise and of sound components linked to the flow distortion near the plate. With the aim of evaluating more precisely these two contributions to the pressure field, the sound levels estimated at $z = 0$ using Curle's analogy for jetsmallhole and jetlargehole are shown in figure 25, together with the levels obtained by the LES and for the free jet. The results for jetnohole and jetmediumhole, similar to those for jetsmallhole, are not represented for brevity. In all cases, the sound levels from the analogy and the LES decrease with the radial distance. For jetsmallhole, in figure 25(a), close to the nozzle, the results from the LES and the analogy exhibit discrepancies. Compared with the levels in the LES, for $r \leq 3r_0$, the levels predicted by the analogy are lower, with differences up to 9 dB, whereas for $3r_0 \leq r \leq 8r_0$, they are higher, with a maximum discrepancy of 4 dB. These differences may be due to the neglect of the aerodynamic noise sources in the analogy, especially those related to the impact of the jet mixing layers on the plate. Farther from the axis, for $r \geq 8r_0$, the levels given by the analogy are close to those from the LES, with less than 2 dB of differences. The




Figure 24: Sound pressure levels obtained at $z = 0$ and $r = 1.5r_0$ — from LES and — using equation (3.4), for (a) jetnohole, (b) jetsmallhole, (c) jetmediumhole and (d) jetlargehole, as a function of the Strouhal number.




Figure 25: Overall sound pressure levels obtained at $z = 0$ — from LES and — using equation (3.4), for (a) jetsmallhole and (b) jetlargehole, - - - levels for jetfree.

present results highlight a significant contribution of the fluctuations of pressure on the plate to the noise radiated upstream. For jetlargehole, in figure 25(b), the levels obtained using the analogy are similar to those from the LES, showing less than 3 dB of differences. This good agreement suggests that the upstream-propagating sound waves are mainly produced by the scattering of the jet pressure fluctuations by the plate.

3.5.3. Acoustic field downstream of the plate

The pressure fields obtained downstream of the plate using Curle's analogy for the three jets impinging on a perforated plate are represented in figures 26(a-c). In all cases, periodic and almost circular pressure waves are found to originate from the hole in the plates. Their wavelengths and their magnitudes are similar to those of the sound waves propagating in the LES fields, shown in figure 26(d-f). Therefore, the variations of pressure on the plate by the jet flow generate the main part of the downstream acoustic radiation for $z \leq 20r_0$.




Figure 26: Pressure fields obtained downstream of the plate (top) using equation (3.4) and (bottom) from LES for (a,d) jetsmallhole, (b,e) jetmediumhole and (c,f) jetlargehole. The color scales range between (a,b,d,e) $\pm 0.01p_0$, and (c,f) $\pm 0.0025p_0$, from blue to red.

However, the shapes of the wavefronts are slightly different, especially close to the jet flow. These differences may be due to the refraction of sound waves by the flow, not predicted by the analogy.

The pressure spectra calculated at $z = 12r_0$ and $r = 15r_0$ using the analogy are plotted in figure 27. They are very similar to those obtained from the LES. For jetsmallhole and jetmediumhole, in figures 27(a,b), they display peaks at the dominant feedback frequency $St = 0.40$ and its harmonic frequencies. For the two first peaks for jetsmallhole and the first one for jetmediumhole, the levels are close to the LES levels, with differences lower than 4 dB. Furthermore, the broadband levels determined using the analogy and the LES do not differ by more than 3 dB. Therefore, the acoustic radiation downstream of the plate seems to be mainly produced by the pressure fluctuations on the plate due to the flow impingement. Discrepancies can however be noted. Indeed, for the third and fourth tones for jetsmallhole and for the second and third tones for jetmediumhole, the levels evaluated using the analogy do not agree well with the LES results. Therefore, these tones may be due to noise generation mechanisms not taken into account by Curle's analogy, such as the interactions of turbulent structures just downstream of the plate. Finally, for jetlargehole, in figure 27(c), the spectrum predicted at $z = 12r_0$ and $r = 15r_0$ using the analogy is close to that of the LES, with a very good agreement for $St < 0.6$ and discrepancies lower than 3 dB for higher frequencies. The sound waves reaching the point at $z = 12r_0$ and $r = 15r_0$ thus appear to be mainly generated by the scattering of the jet pressure waves by the hole edge, and not by the turbulent jet developing past the hole.

To quantify more accurately the contribution of the scattered sound waves to the downstream pressure field, the sound levels obtained at $r = 15r_0$ for $z \geq 10r_0$ using the analogy for jetsmallhole and jetlargehole are displayed in figure 28. They are compared with the levels for the LES and for the free jet. The results for jetmediumhole, not shown here, look like those for jetsmallhole. For jetsmallhole, in figure 28(a), the levels from the analogy do not vary much with the axial distance, and are approximately equal to 140 dB. They are close to those of the LES down to $z = 22r_0$, with less than 2 dB of discrepancies. Farther




Figure 27: Sound pressure levels obtained at $z = 12r_0$ and $r = 15r_0$ from — LES and — using equation (3.4) for (a) jetsmallhole, (b) jetmediumhole and (c) jetlargehole, as a function of the Strouhal number.




Figure 28: Overall sound pressure levels obtained downstream of the plate at $r = 15r_0$ — from LES and — using equation (3.4), for (a) jetsmallhole and (b) jetlargehole, - - - levels for jetfree.

742 downstream, the LES levels decrease down to 130 dB at $z = 40r_0$. This reduction may be due
 743 to the refraction of the sound waves by the jet flow, neglected in the analogy. For jetlargehole,
 744 in figure 28(b), the levels in the LES are almost constant and close to 122 dB, while those
 745 for the analogy are lower and decrease with the axial distance. For $z \leq 15r_0$, the analogy
 746 provides levels lower by less than 3 dB than the LES levels, which suggests that most of the
 747 noise in this region is produced by the waves scattered by the plate. Farther downstream, the
 748 discrepancy raises up to 9 dB at $z = 40r_0$. Therefore, far from the plate, the sound radiation
 749 does not seem to be substantially due to the interactions between the jet and the plate. In
 750 particular, for $z \geq 30r_0$, the LES levels are similar to those for the free jet, implying that the
 751 jet mixing noise is the main noise component for low radiation angles.

4. Conclusion

In this paper, the production of tones by a round jet at a Mach number of 0.9 impinging on a plate with and without a hole of varying diameter h has been investigated using compressible large-eddy simulations. For all plate geometries, tones emerge in the pressure spectra. Their frequencies are found to be nearly independent of the presence and of the size of the hole in the plate, in agreement with the frequencies expected for aeroacoustic feedback loops establishing between the nozzle lips and the plate, involving turbulent structures convected downstream by the jet flow and acoustic waves travelling upstream. The latter waves moreover appear to belong to the family of the free-stream upstream-propagating guided waves of the jet, allowing us to explain the azimuthal structure of the jet oscillation mode at each tone frequency. Regarding the tone intensities, they decrease when the hole in the plate is larger, weakly when the hole diameter increases from $h = 2r_0$ to $h = 3r_0$, but very strongly between $h = 3r_0$ and $h = 4.4r_0$. The reason for that is shown to be the different mechanisms producing the acoustic waves closing the feedback loops. Indeed, these waves are generated directly by the impingement of the jet flow structures on the plate for the two smallest holes, but by the scattering of the jet aerodynamic pressure fluctuations by the plate for the largest hole, leading to much weaker tones in that case. These results emphasize the importance of the nature of the interactions between the jet flow and the plate in producing strongly emerging acoustic tones. Finally, an acoustic analogy based on the variations of the pressure on the plate has also been used to study the nature of the dominant noise sources. It provides results in good agreement with the LES, showing the significant contribution of the pressure fluctuations on the plate to the pressure field. In the future, it could be interesting to examine whether the present phenomena observed for a high-subsonic jet will also be obtained for supersonic jets impinging on a perforated plate. In particular, this may not be the case for impinging rocket jets, for which no acoustic tones have been reported in the literature.

Acknowledgments

This work was financed by ArianeGroup and the DGA (Direction Générale de l'Armement) and by the IRICE IJES project RA0014963 (Installed Jet Effect Simulator, FEDER-FSE Rhône-Alpes). It was granted access to the HPC resources of PMCS2I (Pôle de Modélisation et de Calcul en Sciences de l'Ingénieur et de l'Information) of Ecole Centrale de Lyon, PSMN (Pôle Scientifique de Modélisation Numérique) of ENS de Lyon and P2CHPD (Pôle de Calcul Hautes Performances Dédiés) of Université Lyon I, members of FLMSN (Fédération Lyonnaise de Modélisation et Sciences Numériques), partner of EQUIPEX EQUIP@MESO, and to the resources of TGCC (Très Grand Centre de calcul du CEA) under the allocation 2020-2a0204 made by GENCI (Grand Equipement National de Calcul Intensif). It was performed within the framework of the Labex CeLyA of Université de Lyon, operated by the French National Research Agency (grant no. ANR-10-LABX-0060/ANR-16-IDEX-0005).

Declaration of interests

The authors report no conflict of interest.

REFERENCES

- BERLAND, J., BOGEY, C., MARSDEN, O. & BAILLY, C. 2007 High-order, low dispersive and low dissipative explicit schemes for multiple-scale and boundary problems. *Journal of Computational Physics* **224** (2), 637–662.

- BOGEY, C. 2018 Grid sensitivity of flow field and noise of high-Reynolds-number jets computed by large-eddy simulation. *International Journal of Aeroacoustics* **17** (4-5), 399–424.
- BOGEY, C. 2019 Two-dimensional features of correlations in the flow and near pressure fields of Mach number 0.9 jets. *Tech. Rep.* 2019-0806. AIAA Paper.
- BOGEY, C. 2021a Acoustic tones in the near-nozzle region of jets: characteristics and variations between mach numbers 0.5 and 2. *Journal of Fluid Mechanics* **921**.
- BOGEY, C. 2021b Tones in the acoustic far field of jets in the upstream direction. *AIAA Journal*.
- BOGEY, C. & BAILLY, C. 2002 Three-dimensional non-reflective boundary conditions for acoustic simulations: far-field formulation and validation test cases. *Acta Acustica united with Acustica* **88** (4), 463–471.
- BOGEY, C. & BAILLY, C. 2004 A family of low dispersive and low dissipative explicit schemes for flow and noise computations. *Journal of Computational Physics* **194** (1), 194–214.
- BOGEY, C. & BAILLY, C. 2006 Large-eddy simulations of transitional round jets: influence of the Reynolds number on flow development and energy dissipation. *Physics of Fluids* **18** (6), 065101.
- BOGEY, C., DE CACQUERAY, N. & BAILLY, C. 2011a Finite differences for coarse azimuthal discretization and for reduction of effective resolution near origin of cylindrical flow equations. *Journal of Computational Physics* **230** (4), 1134–1146.
- BOGEY, C. & GOJON, R. 2017 Feedback loop and upwind-propagating waves in ideally expanded supersonic impinging round jets. *Journal of Fluid Mechanics* **823**, 562–591.
- BOGEY, C., MARSDEN, O. & BAILLY, C. 2011b Large-eddy simulation of the flow and acoustic fields of a Reynolds number 10^5 subsonic jet with tripped exit boundary layers. *Physics of Fluids* **23** (3), 035104.
- BRÈS, G., JORDAN, P., JAUNET, V., LE RALLIC, M., CAVALIERI, A., TOWNE, A., LELE, S., COLONIUS, T. & SCHMIDT, O. 2018 Importance of the nozzle-exit boundary-layer state in subsonic turbulent jets. *Journal of Fluid Mechanics* **851**, 83–124.
- BUCHMANN, N.A., MITCHELL, D.M., INGVOSEN, K. M., HONNERY, D.R. & SORIA, J. 2011 High spatial resolution imaging of a supersonic underexpanded jet impinging on a flat plate. *Proceedings of the 6th Australian Conference on Laser Diagnostics in Fluid Mechanics and Combustion* **116**.
- CHANAUD, R.C. & POWELL, A. 1965 Some experiments concerning the hole and ring tone. *The Journal of the Acoustical Society of America* **37** (5), 902–911.
- CURLE, N. 1955 The influence of solid boundaries upon aerodynamic sound. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **231** (1187), 505–514.
- DAUPTAIN, A., GICQUEL, L.Y. & MOREAU, S. 2012 Large-eddy simulation of supersonic impinging jets. *AIAA Journal* **50** (7), 1560–1574.
- EDGINGTON-MITCHELL, D. 2019 Aeroacoustic resonance and self-excitation in screeching and impinging supersonic jets—a review. *International Journal of Aeroacoustics* **18** (2-3), 118–188.
- EDGINGTON-MITCHELL, D., JAUNET, V., JORDAN, P., TOWNE, A., SORIA, J. & HONNERY, D. 2018 Upstream-travelling acoustic jet modes as a closure mechanism for screech. *Journal of Fluid Mechanics* **855**.
- FAUCONNIER, D., BOGEY, C. & DICK, E. 2013 On the performance of relaxation filtering for large-eddy simulation. *J. Turbul.* **14** (1), 22–49.
- GLOERFELT, X., BAILLY, C. & JUVÉ, D. 2003 Direct computation of the noise radiated by a subsonic cavity flow and application of integral methods. *Journal of sound and vibration* **266** (1), 119–146.
- GLOERFELT, X., PÉROT, F., BAILLY, C. & JUVÉ, D. 2005 Flow-induced cylinder noise formulated as a diffraction problem for low Mach numbers. *Journal of Sound and Vibration* **287** (1-2), 129–151.
- GOJON, R. & BOGEY, C. 2017a Flow structure oscillations and tone production in underexpanded impinging round jets. *AIAA Journal* **55** (6), 1792–1805.
- GOJON, R. & BOGEY, C. 2017b Numerical study of the flow and the near acoustic fields of an underexpanded round free jet generating two screech tones. *International Journal of Aeroacoustics* **16** (7-8), 603–625.
- GOJON, R. & BOGEY, C. 2018 Flow features near plate impinged by ideally expanded and underexpanded round jets. *AIAA Journal* **56** (2), 445–457.
- GOJON, R., BOGEY, C. & MARSDEN, O. 2016 Investigation of tone generation in ideally expanded supersonic planar impinging jets using large-eddy simulation. *Journal of Fluid Mechanics* **808**, 90–115.
- HENDERSON, B., BRIDGES, J. & WERNET, M. 2005 An experimental study of the oscillatory flow structure of tone-producing supersonic impinging jets. *Journal of Fluid Mechanics* **542**, 115–137.
- HO, C.-M. & NOSSEIR, N. S. 1981 Dynamics of an impinging jet. Part 1. the feedback phenomenon. *Journal of Fluid Mechanics* **105**, 119–142.
- HOWE, M.S. 1979 On the theory of unsteady high Reynolds number flow through a circular aperture.

- 851 *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **366** (1725),
852 205–223.
- 853 JAUNET, V., MANCINELLI, M., JORDAN, P., TOWNE, A., EDGINGTON-MITCHELL, D. M., LEHNASCH, G. &
854 GIRARD, S. 2019 Dynamics of round jet impingement. *AIAA Paper 2019-2769*.
- 855 JORDAN, P., JAUNET, V., TOWNE, A., CAVALIERI, A., COLONIUS, T., SCHMIDT, O. & AGARWAL, A. 2018 Jet-flap
856 interaction tones. *Journal of Fluid Mechanics* **853**, 333–358.
- 857 KAWAI, S., TSUTSUMI, S., TAKAKI, R. & FUJII, K. 2007 Computational aeroacoustic analysis of overexpanded
858 supersonic jet impingement on a flat plate with/without hole. *ASME/JSME 2007 5th Joint Fluids
859 Engineering Conference 2007-37563* pp. 1163–1167.
- 860 KREMER, F. & BOGEY, C. 2015 Large-eddy simulation of turbulent channel flow using relaxation filtering:
861 Resolution requirement and Reynolds number effects. *Computers & Fluids* **116**, 17–28.
- 862 LANGTHJEM, M. A. & NAKANO, M. 2005 The jet hole-tone oscillation cycle subjected to acoustic excitation:
863 a numerical study based on an axisymmetric vortex method. *The Proceedings of the International
864 Conference on Jets, Wakes and Separated Flows* **2005**, 745–750.
- 865 LAUNDER, B. E. & RODI, W. 1983 The turbulent wall jet measurements and modeling. *Annual review of fluid
866 mechanics* **15** (1), 429–459.
- 867 LAWRENCE, J., AZARPEYVAND, M. & SELF, R. 2011 Interaction between a flat plate and a circular subsonic
868 jet. *AIAA Paper 2011-2745*.
- 869 MANCINELLI, M., JAUNET, V., JORDAN, P. & TOWNE, A. 2019 Screech-tone prediction using upstream-
870 travelling jet modes. *Experiments in Fluids* **60** (1), 22.
- 871 MATSUURA, K. & NAKANO, M. 2012 A throttling mechanism sustaining a hole tone feedback system at very
872 low Mach numbers. *Journal of Fluid Mechanics* **710**, 569–605.
- 873 MEGANATHAN, A. & VAKILI, A. 2006 An experimental study of acoustic and flow characteristics of hole
874 tones. *AIAA Paper 2006-1015*.
- 875 MICHALKE, A. 1984 Survey on jet instability theory. *Progress in Aerospace Sciences* **21**, 159–199.
- 876 MITCHELL, D. M., HONNERY, D. R. & SORIA, J. 2012 The visualization of the acoustic feedback loop
877 in impinging underexpanded supersonic jet flows using ultra-high frame rate schlieren. *Journal of
878 Visualization* **15** (4), 333–341.
- 879 MOHSENI, K. & COLONIUS, T. 2000 Numerical treatment of polar coordinate singularities. *Journal of
880 Computational Physics* **157** (2), 787–795.
- 881 NEUWERTH, G. 1974 Acoustic feedback of a subsonic and supersonic free jet which impinges on an obstacle.
882 *NASA Technical Translation F-15719*.
- 883 NOGUEIRA, P.A.S., CAVALIERI, A.V.G. & JORDAN, P. 2017 A model problem for sound radiation by an installed
884 jet. *Journal of Sound and Vibration* **391**, 95–115.
- 885 NORUM, T. D. 1991 Supersonic rectangular jet impingement noise experiments. *AIAA Journal* **29** (7),
886 1051–1057.
- 887 NOSSEIR, N. S. & HO, C.-M. 1982 Dynamics of an impinging jet. Part 2. the noise generation. *Journal of
888 Fluid Mechanics* **116**, 379–391.
- 889 PANICKAR, P. & RAMAN, G. 2007 Criteria for the existence of helical instabilities in subsonic impinging jets.
890 *Physics of Fluids* **19** (10), 103–106.
- 891 POREH, M., TSUEI, Y.G. & CERMAK, J. E. 1967 Investigation of a turbulent radial wall jet. *Journal of Applied
892 Mechanics* **34** (2).
- 893 POWELL, A. 1953 On edge tones and associated phenomena. *Acta Acustica United with Acustica* **3** (4),
894 233–243.
- 895 PREISSER, J. S. 1979 Fluctuating surface pressure and acoustic radiation for subsonic normal jet impingement.
896 *NASA Technical Paper 1361*.
- 897 RAYLEIGH, LORD 1945 The theory of sound, vols. I and II.
- 898 RICOU, F.P. & SPALDING, D.B. 1961 Measurements of entrainment by axisymmetrical turbulent jets. *Journal
899 of fluid mechanics* **11** (1), 21–32.
- 900 RISBORG, A & SORIA, J 2009 High-speed optical measurements of an underexpanded supersonic jet
901 impinging on an inclined plate. In *Proceedings of SPIE 7126, 28th International Congress on
902 High-Speed Imaging and Photonics*. International Society for Optics and Photonics.
- 903 SCHMIDT, O. T., TOWNE, A., COLONIUS, T., CAVALIERI, A. V. G., JORDAN, P. & BRÈS, G. A. 2017 Wavepackets
904 and trapped acoustic modes in a turbulent jet: coherent structure eduction and global stability. *Journal
905 of Fluid Mechanics* **825**, 1153–1181.
- 906 SONDHAUSS, C. 1854 Über die beim Ausströmen der Luft entstehenden Töne. *Annalen der Physik* **167** (2),
907 214–240.

- TAM, C.K.W. & AHUJA, K.K. 1990 Theoretical model of discrete tone generation by impinging jets. *Journal of Fluid Mechanics* **214**, 67–87.
- TAM, C.K.W. & CHANDRAMOULI, S. 2020 Jet-plate interaction tones relevant to over-the-wing engine mount concept. *Journal of Sound and Vibration* **486**, 115378.
- TAM, C.K.W. & DONG, Z. 1996 Radiation and outflow boundary conditions for direct computation of acoustic and flow disturbances in a non uniform mean flow. *Journal of Computational Acoustics* **4** (02), 175–201.
- TAM, C.K.W. & NORUM, T.D. 1992 Impingement tones of large aspect ratio supersonic rectangular jets. *AIAA Journal* **30** (2), 304–311.
- TOWNE, A., CAVALIERI, A. V. G., JORDAN, P., COLONIUS, T., SCHMIDT, O., JAUNET, V. & BRÈS, G. A. 2017 Acoustic resonance in the potential core of subsonic jets. *Journal of Fluid Mechanics* **825**, 1113–1152.
- TROYES, J. N., VUILLOT, F., LANGENAI, A. & LAMBARÉ, H. 2019 Coupled CFD-CAA simulation of the noise generated by a hot supersonic jet impinging on a flat plate with exhaust hole. *AIAA Paper* 2019-2752 .
- TSUTSUMI, S., TAKAKI, R., IKAIDA, H. & TERASHIMA, K. 2015 Numerical aeroacoustics analysis of a scaled solid jet impinging on flat plate with exhaust hole. *30th International Symposium on Space Technology and Science* .
- UMEDA, Y. & ISHII, R. 1993 Hole tone generation from highly choked jets. *The Journal of the Acoustical Society of America* **94** (2), 1058–1066.
- UMEDA, Y., MAEDA, H. & ISHII, R. 1988 Hole tone generated from almost choked to highly choked jets. *AIAA Journal* **26** (9), 1036–1043.
- VAN HOUT, R., RINSKY, V. & GROBMAN, Y. G. 2018 Experimental study of a round jet impinging on a flat surface: flow field and vortex characteristics in the wall jet. *International Journal of Heat and Fluid Flow* **70**, 41–58.
- VARÉ, M. & BOGEY, C. 2021 Large-eddy simulations of the flow and acoustic fields of a rocket jet impinging on a perforated plate. *AIAA Paper* 2021-2152 .
- VINOTH, B.R. & RATHAKRISHNAN, E. 2011 Effect of impinging plate geometry on the self-excitation of subsonic impinging jets. *Journal of Fluids and Structures* **27** (8), 1238–1251.