



HAL
open science

Synthesis, characterization and electrochemical behavior of new bis(fluoroalkyl) ferrocenylphosphonates and their tin tetrachloride complexes

Hanen Mechi, M.A.K. a K Sanhoury, F. Laribi, Eric Manoury, M. Mastouri, N. Raouafi, Azzedine Bousseksou, M. T Ben Dhia

► To cite this version:

Hanen Mechi, M.A.K. a K Sanhoury, F. Laribi, Eric Manoury, M. Mastouri, et al.. Synthesis, characterization and electrochemical behavior of new bis(fluoroalkyl) ferrocenylphosphonates and their tin tetrachloride complexes. *Journal of Organometallic Chemistry*, 2022, 957, pp.122178. 10.1016/j.jorganchem.2021.122178 . hal-03576457

HAL Id: hal-03576457

<https://hal.science/hal-03576457>

Submitted on 16 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis, characterization and electrochemical behaviour of new bis(fluoroalkyl) ferrocenylphosphonates and their tin tetrachloride complexes

Hanen Mechi^{a,b}, M.A.K Sanhoury^{a,c}, F. Laribi^a, E. Manoury^b, M. Mastouri^d, N. Raouafi^d, A. Bousseksou^b, M. T. Ben Dhia^a

^a *Laboratory of Structural Organic Chemistry: Synthesis and Physicochemical Studies, Department of Chemistry, Faculty of Sciences of Tunis, El Manar, University Campus I 2092, Tunis, Tunisia.*

^b *Laboratoire de Chimie de Coordination (CNRS) 205 Route de Narbonne, 31400 Toulouse, France.*

^c *Materials Chemistry Research Unit, Faculty of Sciences and Techniques, UNA, Nouakchott, Mauritania.*

^d *Lab of Analytical Chemistry, Department of Chemistry, Faculty of Sciences of Tunis, El Manar, University Campus I 2092, Tunis, Tunisia.*

Abstract

The synthesis and characterization of new fluoroalkyl and non-fluorinated long chain alkyl ferrocenylphosphonates from reaction of ferrocene with corresponding phosphorochloridates are reported. These phosphonates were used as ligands for the preparation of new tin(IV) chloride complexes. All these ferrocenylphosphonate derivatives were characterized with multinuclear (¹H, ¹³C, ¹⁹F, ³¹P and ¹¹⁹Sn) NMR, IR and UV-vis spectroscopic techniques. The electrochemical behavior of these compounds was also investigated using cyclic voltammetry; the results show that fluoroalkyl groups render the reverse one-electron transfer oxidation reaction more difficult ($\Delta E_{1/2} = 310\text{-}330$ mV) as compared to non fluorinated alkyl groups ($\Delta E_{1/2} = 215\text{-}261$ mV).

Keywords: Ferrocenylphosphonates, fluoroalkyl alcohols, cyclic voltammetry.

1. Introduction

Organophosphorus chemistry have gained considerable interest due to their widespread applications in various areas such as agriculture [1,2], biology [3,4], pharmaceuticals [5-7] and chemical synthesis [8,9]. Among these, one of the most common classes are phosphonates, which are attracting increasing interest as important structural motifs that are found in a variety of naturally occurring compounds [10-12], as anticancer [13,14], inhibitors of hepatitis C virus [15,16], anti-HIV [17] and antimalarial agents [18].

On the other hand, ferrocene and ferrocenyl derivatives are well known for their ability to undergo reversible one-electron oxidation, they have been used extensively as single-electron transfer (SET) reagents in organometallic and coordination chemistry to switch the redox state of complexes [19-22]. Ferrocene-containing organophosphorus compounds are increasingly

investigated in organic synthesis as synthetic intermediates [23,24], potential ligands, in catalysis [25] and as sensitizers in dye sensitized solar cells [26].

The incorporation of fluoroalkyl groups has been shown to modify the physical and chemical properties of molecules giving them special applications in the fields of pharmaceuticals, catalysis and materials science [27-29]. It was also shown that fluoroalkyl chains generally provide a higher thermal stability to the molecule, facilitating both hits storage and handling [30,31]. They are also used as biocompatible surfactants as emulsion stabilizers and vesicle-forming components [32,33].

In continuation of our exploration in the synthesis of variously substituted phosphoramidates [34,35], we describe, herein, the synthesis of a series of new fluoroalkyl and alkyl ferrocenylphosphonates and corresponding tin tetrachloride complexes. The products were obtained in good yields and fully characterized by multinuclear (^1H , ^{13}C , ^{19}F , ^{31}P and ^{119}Sn NMR) NMR, IR, cyclic voltammetry (CV), HRMS for phosphonates and elemental analysis for the corresponding tin(IV) complexes. The effect of fluoroalkyl and non-fluorinated long chain groups on the electrochemical behavior of these phosphonate derivatives is also investigated.

2. Experimental

All reactions were carried out under an inert atmosphere. Solvents were carefully dried by conventional methods and distilled under argon before use. All commercially available chemicals ferrocene, *t*BuLi 1.7 M solution in pentane and *t*BuOK were used as received from their suppliers. Dialkyl and bis(fluoroalkyl) chlorophosphates [36] and diethyl ferrocenylphosphonate (**3**) [37] were prepared following literature procedures.

2.1 General procedure for the preparation of ferrocenylphosphonates $\text{FcP}(\text{O})(\text{OR})_2$ (**1-6**)

The phosphonates (**1-6**) were prepared in a similar manner to the method reported by Oms *et al.* [37] with some modifications: KO*t*Bu (72 mg, 0.65 mmol) was added to a solution of ferrocene (1.0 g, 5.38 mmol) in THF (20 mL) at room temperature and stirred for 30 min. The solution was then cooled to $-78\text{ }^\circ\text{C}$ and *t*-BuLi (6.32 mL, 10.76 mmol, 1.7 M in pentane) was added dropwise. The reaction mixture was stirred at $-78\text{ }^\circ\text{C}$ for 1 h. The chlorophosphate (6.73 mmol) in THF was added to the orange solution at $-78\text{ }^\circ\text{C}$. The solution slowly warmed to room temperature and left until the reaction was complete. The resulting deep brown solution was concentrated under vacuo; water (50 mL) was added to the residue and solution extracted

with diethyl ether (3x20 mL). The combined organic fractions were dried over Na₂SO₄, filtered and evaporated *in vacuo*. The crude product was purified by flash column chromatography.

FcP(O)(OCH₂CF₃)₂ (1)

Flash chromatography Pentane to Et₂O. Yellow solid, MP = 54°C. FT-IR (ν/cm⁻¹) 3019, 2401, 1712, 1421, 1215(P=O), 1175, 1106, 1075, 984, 751, 668. RMN ¹H (400 MHz, CDCl₃) δ 4.55 (s, 5H), 4.47 – 4.37 (m, 4H), 4.34 (d, *J* = 4.3 Hz, 4H). ¹⁹F NMR (376 MHz, CDCl₃) δ -75.10 (t, *J* = 7.9 Hz). ¹³C NMR (101 MHz, CDCl₃) 122,73(qd, ¹*J*_{C-F} = 278.76 Hz, ³*J*_{C-P} = 9 Hz), 72.27 (d, *J* = 15.0 Hz, C3), 71.68 (d, *J* = 16.8 Hz, C2), 70.17 (s, C_PC), 62.65 (d, *J*¹_{P-C} = 225,23 Hz, C1), 61.68 (q (d, *J*_{P-O-C} = 3.0 Hz), P(OCH₂CF₃)₂). ³¹P NMR (162 MHz, CDCl₃) δ 30.23 (s). Anal. calcd for Anal. calcd for C₁₄H₁₃F₆FeO₃P: C, 39.10; H, 3.05 %. Found:

FcP(O)(OCH₂C₂F₅)₂ (2)

Flash chromatography Pentane to Et₂O/MeOH (9:1). Brown liquid. FT-IR (ν/cm⁻¹) 3020, 2401, 1704, 1423, 1215 (P=O), 1156, 1027, 749, 668. ¹H NMR (400 MHz, CDCl₃) δ 4.58 – 4.45 (m, 8H), 4.34 (s, 5H). ¹⁹F NMR (377 MHz, CDCl₃) δ -83.56 (s), -124.49 (s). ¹³C NMR (101 MHz, CDCl₃) δ 111,86 (tq, ¹*J*_{C-F} = 248.46 Hz, ²*J*_{C-F} = 29 Hz, CF₂), 118,35 (qt, ¹*J*_{C-F} = 286.84 Hz, ²*J*_{C-F} = 35 Hz, CF₃), 72.29 (d, *J* = 15.2 Hz, C3), 71.57 (d, *J* = 17.1 Hz, C2), 70.12 (s, C_PC), 62.29 (d, *J*¹_{P-C} = 225,23 Hz, C1), 61 (t (d, *J*_{P-O-C} = 5.0 Hz), P(OCH₂C₂F₅)₂). ³¹P NMR (162 MHz, CDCl₃) δ 30.34 (s). Anal. calcd for Anal. calcd for C₁₆H₁₃F₆FeO₃P: C, 39.25; H, 2.47 %. Found:

FcP(O)(OHex)₂ (4)

Flash chromatography hexane gradient to hexane/EtOAc 9:1. Pale brown liquid. ¹H NMR (400 MHz, CDCl₃) δ 4.50 – 4.45 (m, 2H), 4.39 – 4.34 (m, 2H), 4.27 (s, 5H), 4.10 – 3.94 (m, 4H), 1.71 – 1.58 (m, 4H), 1.41 – 1.21 (m, 12H), 0.86 (t, *J* = 6.7 Hz, 6H). ¹³C NMR (101 MHz, CDCl₃) 71.50 (d, *J* = 15.4 Hz, C3), 71.11 (d, *J* = 13.9 Hz, C2), 69.79 (s, C_PC), 66.88 (d, *J*¹_{P-C} = 216,14 Hz, C1), 65.64 (d, *J*_{P-O-C} = 6.0 Hz), 31.31 (s), 30.48 (d, *J* = 6.6 Hz), 25.24(s), 22.91 (s), 22.51 (s), 13.97 (s). ³¹P NMR (162 MHz, CDCl₃) δ 25.79 (s). Anal. calcd for C₂₂H₃₅FeO₃P: C, 60.84; H, 8.12 %. Found:

FcP(O)(OHep)₂ (5)

Flash chromatography hexane/EtOAc gradient to EtOAc/MeOH (4:1). Pale Brown liquid. ¹H NMR (400 MHz, CDCl₃) δ 4.52 (t, *J* = 4.4 Hz, 2H), 4.41 (t, *J* = 4.6 Hz, 2H), 4.32 (s, 5H), 4.25 – 3.80 (m, 5H), 1.77 – 1.64 (m, 5H), 1.49 – 1.28 (m, 10H), 0.92 (t, *J* = 7.1 Hz, 7H). ³¹P NMR (162 MHz, CDCl₃) δ 25.77 (s).

FcP(O)(OOct)₂ (6)

Flash chromatography hexane/EtOAc gradient to EtOAc/MeOH (4:1). Brown liquid. ¹H NMR (400 MHz, CDCl₃) δ 4.52 (d, *J* = 10.4 Hz, 2H), 4.40 (s, 2H), 4.31 (s, 5H), 4.18 – 4.08 (m, 5H), 1.76 – 1.58 (m, 5H), 1.30 (ddd, *J* = 18.7, 9.4, 4.8 Hz, 27H), 0.89 (t, *J* = 6.9 Hz, 7H). ³¹P NMR (162 MHz, CDCl₃) δ 25.75 (s).

2.2 General procedure for the preparation of complexes (7-10)

A solution of ferrocenyphosphonates (0.46 mmol) in dry CH₂Cl₂ (10 mL) was slowly added to SnCl₄ (0.06 g, 0.23 mmol) in CH₂Cl₂ (20 mL) and the mixture stirred under N₂ for 24 h. The volatiles were then removed *in vacuo* and the complex [SnCl₄L₂] washed with anhydrous hexane and dried *in vacuo* to give the complexes as brown oils.

[SnCl₄(1)₂] (7)

Yield: 0.20 g (80%); FT-IR (cm⁻¹) ν = 1159 (P=O), 1077 (P-O-C), 749, 668; ¹H NMR (300 MHz, CDCl₃) δ 4.93 – 4.60 (m, 8H), 4.52 (s, 5H). ³¹P NMR (121 MHz, CDCl₃) δ 28.06 (¹J_{P-C} = 223 Hz), 27.67. ¹¹⁹Sn NMR (112 MHz, CDCl₃) -714.89. Anal. calcd for C₂₈H₂₆Cl₄F₁₂Fe₂O₆P₂Sn: C, 30.86; H, 2.41 %. Found:

[SnCl₄(2)₂] (8)

Yield: 0.27 g (90%); FT-IR (cm⁻¹) ν = 1177 (P=O), 1126 (P-O-C), 754, 668; ¹H NMR (300 MHz, CDCl₃) δ 5.03 – 4.7 (m, 8H), 4.52 (s, 5H). ³¹P NMR (121 MHz, CDCl₃) δ 28.19 (¹J_{P-C} = 216.25 Hz), 25.06. ¹¹⁹Sn NMR (112 MHz, CDCl₃) -714.19, -647.97. Anal. calcd for C₃₂H₂₆Cl₄F₂₀Fe₂O₆P₂Sn: C, 29.80; H, 2.03 (%); Found:

[SnCl₄(3)₂] (9)

Yield: 0.18 g (90%); ¹H NMR (400 MHz): δ = 4.50-4.38 (m, 4H), 4.30 (s, 5H), 4.12 (m, 4H), 1.33 (t, *J* = 7.1 Hz, 6H). RMN ³¹P (121 MHz, CDCl₃) δ 21.83 (¹J_{P-C} = 232.32 Hz). NMR ¹¹⁹Sn (112 MHz, CDCl₃) -728.82.

[SnCl₄(4)₂] (10)

Yield: 0.18 g (71 %); FT-IR (cm⁻¹) ν = 1126 (P=O), 1159 (P-O-C), 738, 668; ¹H NMR (300 MHz, CDCl₃) δ 5.20 (s, 5H), 5.57-6.5 (m, 4H), 3.81-3.94 (m, 4H), 1.17-1.37 (m, 12H), 0.85-

0.87 (m, 6H). ^{31}P NMR (121 MHz, CDCl_3) δ 19.47, 16.01. ^{119}Sn NMR (112 MHz, CDCl_3) -660. Anal. calcd for $\text{C}_{44}\text{H}_{70}\text{Cl}_4\text{F}_{20}\text{Fe}_2\text{O}_6\text{P}_2\text{Sn}$: C, 48.12; H, 6.42 (%). Found:

2.3 Physical measurements

Melting points were measured using an “Electrothermal 9100 : apparatus and are uncorrected. ^1H (300 MHz), ^{13}C (75.4 MHz), ^{19}F NMR (282 MHz) and ^{31}P (121 MHz) NMR spectra were recorded with a Bruker ACIII-300 spectrometer. TMS was used as the internal standard for ^1H and ^{13}C , CFCl_3 for ^{19}F and H_3PO_4 for ^{31}P NMR spectroscopy. HRMS data were obtained from a high resolution MicromassmicrOTOF-Q II 10027 spectrometer with electron spray ionization method. Infrared spectra were recorded on a YL2000 FT-IR spectrometer using KBr pellets. Cyclic voltammograms and potential square wave voltammograms were obtained in a three-electrode cell. A GCE working electrode, a platinum wire auxiliary electrode and an Ag/AgCl reference electrode were used to record voltammograms. The electrolyte was $n\text{Bu}_4\text{NPF}_6$ (0.1 M in dichloromethane). The concentration of electro-active species was 0.001 M. Pulse amplitude of 25 mV and a scan rate of 5 mVs^{-1} were used to record the potential square wave voltammograms. UV/Vis spectra were recorded on a Unico 2802 spectrophotometer using chloroform as solvent.

3. Results and discussion

3.1. Synthesis of ferrocenylphosphonates and their complexes

The synthesis of the new ferrocenylphosphonates $\text{FcP}(\text{O})(\text{OR})_2$ (**1**, **2** and **4-6**) was performed using the $^t\text{BuLi}/^t\text{BuOK}$ system at low temperature, following a similar methodology reported by Oms *et al.* [37]. Thus, the reaction of ferrocene (Fc), using the Schlosser–Lochmann base ($^t\text{BuLi}/\text{K}^t\text{OtBu}$) in THF at -78°C , with different phosphorochloridates gave the phosphonates **1-6** in yields of 30–79% (Table 1). The isolated yields were, in general, better for fluoroalkyl derivatives. The previously known phosphonate (**3**), $\text{FcP}(\text{O})(\text{OEt})_2$ [37], was prepared in the same way and used for comparison purposes.

Table 2: Synthesis of ferrocenylphosphonate complexes, $[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OR})_2)_2]$ (**7-10**).

Complex	Yield (%)
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OCH}_2\text{CF}_3)_2)_2]$ (7)	80
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OCH}_2\text{C}_2\text{F}_5)_2)_2]$ (8)	90
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OEt})_2)_2]$ (9)	90
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OHex})_2)_2]$ (10)	71

3.2. IR and NMR Spectroscopic characterization

The strong bands observed in the infrared spectra within the range 1214-1250 cm^{-1} for ferrocenylphosphonates (**1-4**) are assigned to $\nu_{\text{P}=\text{O}}$. The P-O stretching vibrations display relatively strong absorptions in the range 1067-1075 cm^{-1} . These stretching vibration values are in good agreement with literature data for related derivatives [38,39]. Interestingly, the P=O stretching vibrations for complexes (**7-10**) are shifted to lower frequencies while the P-O vibrations are shifted to higher frequencies upon coordination with tin tetrachloride compared to that in the free ligands (**1-4**) (Table 3). This suggests that the P=O bond is weakened upon complexation, consistent with the coordination to the tin center through the P=O group. The magnitude of the coordination shift ($\Delta\nu_{\text{P}=\text{O}}$) in such complexes is in the range 38-88 cm^{-1} and would be directly proportional to the stability of the complex, which depends upon the donor ability of the parent ligand. Thus, the most pronounced coordination shift (88 cm^{-1}) is observed for non alkyl phosphonate complexes **9** and **10**, in agreement with the expected weakening effect of fluoroalkyl groups on the donor ability of fluoroalkyl phosphonates in complexes **7** and **8** (See Table 3).

Table 3 : IR data (ν/cm^{-1}) for ferrocenylphosphonate complexes (**7-10**).

Complex	$\nu(\text{P=O})$		$\nu(\text{P-O})$	
	L	Complex ($\Delta\nu_{(\text{P=O})}$)	L	Complex ($\Delta\nu_{(\text{P-O})}$)
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OCH}_2\text{CF}_3)_2)_2]$ (7)	1215	1159 (56)	1074	1177 (103)
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OCH}_2\text{C}_2\text{F}_5)_2)_2]$ (8)	1215	1177 (38)	1075	1126 (51)
$[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OHex})_2)_2]$ (10)	1214	1126 (88)	1074	1159 (85)

All ferrocenylphosphonates (**1-6**) were also characterized by their NMR (^1H , ^{13}C , ^{19}F and ^{31}P) spectra, which are all consistent with the proposed structures. Indeed, the ^{31}P NMR spectra show that the resonances of fluoroalkyl phosphonates (**1** and **2**) are shifted downfield (30 ppm) as compared to non fluorinated derivatives (**3-6**) (25 ppm) [40].

The ^{31}P NMR spectra of corresponding tin(IV) complexes (**7-10**) show a shift of the resonance signals towards the strong fields compared to those of the free ligands. Such a behaviour is explained by electron effects resulting from a modification in electron density at the phosphorus due to the coordination of the P=O oxygen atom of the ligand to the tin atom. This coordination shift is more pronounced for non fluoroalkyl phosphonate derivatives, in agreement with our above mentioned IR data. It is important to note that the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectra of complexes (**7-10**) show two types of signals: the high frequency major resonance flanked with Sn satellites is assigned to the *trans* isomer and the low frequency signal which is relatively broad due to fast expected ligand exchange is attributed to the *cis* form (Figures 1(a and c)).

In order to gain more insights into the structure of complexes (**7-10**) in solution and in addition to ^{31}P NMR, we examined these adducts using ^{119}Sn NMR spectroscopy. The corresponding ^{119}Sn NMR spectra of complexes **7** and **8** in CDCl_3 display at room temperature only one triplet feature at -715 ppm ($^2J_{\text{P-Sn}} = 227$ Hz) for **7** (Figure 1(b)) and a triplet at -714 ppm ($^2J_{\text{P-Sn}} = 222$ Hz) together with a broad signal at -648 ppm in the region of hexacoordinated tin species for **8** (Figure 1(d)). The triplet resonance observed in the ^{119}Sn NMR spectra is due to $^{119}\text{Sn}\text{-}^{31}\text{P}$ coupling in the *trans* isomers, while the large signal is ascribed to the labile *cis* isomer. This is consistent with the ^{31}P NMR spectra where signals

displaying Sn satellites with the corresponding coupling constants were observed. The results are also in fair agreement with the related bidentate ferrocenyl phosphonate tin(VI) complexes $[(Fc^{1,1'})_2SnCl_4]$ and $[(Fc^{1,2})_2SnCl_4]$ [41] in which the ^{119}Sn NMR spectra display one triplet at 707 ppm ($^2J_{P-Sn} = 215$ Hz) and -694 ppm ($^2J_{P-Sn} = 176$ Hz) for $Fc^{1,1'}$ and $Fc^{1,2}$ complexes, respectively.




Figure 1 ^{31}P NMR spectra for ferrocenyl phosphonate complexes **9** (a), **7** (c) and **8** (e); ^{119}Sn NMR spectra for **9** (b), **7** (d) and **8** (f) in $CDCl_3$ at 25 °C.

Table 4: NMR data (δ /ppm and J /Hz) for ferrocenylphosphonate ($L = \mathbf{1-4}$) and ferrocenylphosphonate complexes $[\text{SnCl}_4(L)_2]$ ($\mathbf{7-10}$).

Compound	^{31}P			$\Delta\delta^{31}\text{P}^a$		^{119}Sn ($^2J_{P-Sn}$)		<i>Trans</i> (%) ^b
	L	<i>Cis</i>	<i>Trans</i>	<i>Cis</i>	<i>Trans</i>	<i>Cis</i>	<i>Trans</i>	
$[\text{SnCl}_4(\mathbf{1})_2]$ (7)	30.23	27.67	28.06	2.56	2.17	-	-715 (227)	79
$[\text{SnCl}_4(\mathbf{2})_2]$ (8)	30.34	25.06	28.19	5.28	2.15	-648	-714 (222)	52
$[\text{SnCl}_4(\mathbf{3})_2]$ (9)	26.70	-	21.83	-	4.87	-	-729 (236)	100
$[\text{SnCl}_4(\mathbf{4})_2]$ (10)	25.79	19.47	-	6.32	-	-660	-714 (224)	-

^a $\Delta\delta^{31}\text{P} = \delta(^{31}\text{P})_L - \delta(^{31}\text{P})_{\text{comp.}}$; ^b measured from ^{31}P NMR.

The above NMR data show that the ferrocenyl phosphonate complexes exist in solution as mixtures of *trans* and *cis* isomers. Complexes **7** and **9** exist as almost a single isomer with a significant predominance of the *trans* form, whereas complex **8** and **10** exist as two isomers with an observed predominance for the *cis* isomer. The difference observed in closely related complexes **7** and **8** could be interpreted by the weakening effect of the $\text{OCH}_2\text{C}_2\text{F}_5$ groups on the donor ability of the ligand in the latter adduct.

3.3. Electrochemical behaviour

Cyclic voltammetry and potential square wave experiments were carried out in dichloromethane containing 0.1 M tetrabutylammonium hexafluorophosphate as supporting electrolyte. The cyclic voltammogram of $\text{FcP}(\text{O})(\text{OCH}_2\text{CF}_3)_2$ (**1**) (1 mM) at a scan rate of 100 mVs^{-1} is shown in Fig. 2. The half wave potentials of different compounds prepared are given in Table 5, along with the direct comparison with ferrocene. As observed before, the presence of phosphonate group directly linked to the ferrocenyl ring renders the oxidation more difficult ($\Delta E_{1/2} = 215\text{-}261$ mV) [37], consistent with the electron-withdrawing effect of the phosphonate substituent. This effect is enhanced by the presence of fluoroalkyl groups in our case for **1** and **2** and their related tin complexes **7** and **8**. For example, comparison of the electrochemical behaviour of fluoroalkyl and non-fluorinated long chain group

ferrocenylphosphonates shows that the presence of fluoroalkyl groups renders the reverse one-electron transfer oxidation reaction approximately 100 mV more difficult as compared to that of non fluorinated alkyl group ($\Delta E_{1/2} = 310\text{-}330$ mV for **1** and **2** vs. 215 mV for **4**). Complex **7** is easier to oxidize than its related ligand (**1**), whilst complex **8** shows no difference with its ligand (**2**) (Table 5). This could be explained by a more pronounced ligand dissociation associated to its exchange with *cis* isomer in the latter complex (**8**), in fair agreement with NMR data discussed above. The potential square wave voltammetry (PWSV) was used to determine the half wave potential of complex **9** (Fig. 3). Complexes **9** and **10** containing non fluorinated ligands **3** and **4** display, in general, the same electrochemical trends as those observed for related non fluorinated ferrocenyl phosphonate derivatives [36].




Figure 2 Cyclic voltammogram of ferrocenylphosphonate derivative **1** and $I = f(v^{1/2})$.




Figure 3 PSWV curve of the complex $[\text{SnCl}_4(\text{FcP}(\text{O})(\text{OEt})_2)_2]$ (**9**).

Table 5 Oxidation of ferrocenyl phosphorus derivatives by voltammetry, in DCM, 0.1 M.

Compound	E _{pa} (mV) /(Ag/AgCl)	E _{pc} (mV) /(Ag/AgCl)	ΔE ^a (mV)	E _{1/2} ^b (mV)	ΔE _{1/2} ^c (mV) /FcH
FcH	700	330	370	515	0
FcP(O)(OCH ₂ CF ₃) ₂ (1)	910	770	140	840	325
FcP(O)(OCH ₂ C ₂ F ₅) ₂ (2)	880	780	100	830	315
FcP(O)(OEt) ₂ (3) ^d	648	581	67	614	261
FcP(O)(OH _{ex}) ₂ (4)	810	650	160	730	215
[SnCl ₄ (1) ₂] (7)	875	756	119	815	300
[SnCl ₄ (2) ₂] (8)	880	780	100	830	325
[SnCl ₄ (3) ₂] (9)	-	-	-	780 ^e	265
[SnCl ₄ (4) ₂] (10)	790	690	100	740	225

^a ΔE = E_{pa}-E_{pc}; ^b ΔE_{1/2} = (1/2)(E_{pa}+E_{pc}); ^c ΔE_{1/2} = E_{1/2}/(Ag/AgCl)-E_{1/2}(FcH+/FcH)/(Ag/AgCl),

^d Values taken from [32]; ^e Determined with PSWV.

3.4. UV-vis characterization

To gain more insights into the electronic structures of phosphonates **1–4** and related tin(IV) complexes **7–10**, their UV-vis spectra were recorded in chloroform, which display bands at 580–651 nm, 436–463 nm, shoulders at 332–356 nm and strong bands at 297–331 nm (Fig. 4). This trend is in general consistent with related ferrocenyl phosphonates [24]. The low energy absorption is most probably assigned to ferrocene [42]. The absorption bands of phosphonates are slightly bathochromic shifted upon coordination to the tin center (Fig. 4)




Figure 4 UV-Vis spectra of fluoroalkyl ferrocenylphosphonates **1** (black) and **2** (green); and their tin complexes **7** (red) and **8** (blue) in CHCl_3 .

4. Conclusion

We have synthesized a series of new ferrocenylphosphonates and related tin tetrachloride complexes. These were characterized by NMR, IR, CV and UV-vis techniques. The results show that the presence of fluoroalkyl groups affects more significantly both the reactivity and electrochemical behavior of ferrocenylphosphonates and corresponding tin(IV) adducts as compared to non fluorinated counterparts. The biological activity of these ferrocenyl phosphonate derivatives as well as their coordination chemistry towards other metal cations are already under investigation.

Acknowledgements

We are grateful to the Tunisian Ministry of High Education and Scientific Research for financial support (LR99ES14) of this research.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by Tunisian Ministry of High Education and Scientific Research [grant number (LR99ES14)].

References

- [1] For an example of a phosphorus containing herbicide, see: J.E. Franz, M.K. Mao, J.A. Sikorski, Glyphosate: A Unique Global Herbicide, American Chemical Society, Washington, DC, 1997.
- [2] M.S. Shaik, M.R. Nadiveedhi, M. Gundluru, S. Poola, R. Allagadda, A. R. Chippada, S. R. Cirandur, Green synthesis of diethyl((2-iodo-4-(trifluoromethyl)phenyl)amino(aryl)methyl) phosphonates as a potent α -glucosidase inhibitors, *Synth. Commun.* 50 (2020) 1532-2432.
- [3] F.H. Westheimer, Why nature chose phosphates, *Science* 235 (1987) 1173-1178.
- [4] H. Seto, T. Kuzuyama, Bioactive natural products with carbon–phosphorus bonds and their biosynthesis, *Nat. Prod. Rep.* 16 (1999) 589-596.
- [5] For representative examples of phosphorus containing pharmaceutical agents, see: P. Kafarski, B. Lejczak, Aminophosphonic acids of potential medical importance, *Curr. Med. Chem.* 1 (2001) 301-312.
- [6] P. Kafarski, B. Lejczak, Phosphorus, Biological activity of aminophosphonic acids Sulfur, Silicon *Relat. Elem.* 63 (1991) 193.
- [7] F.A. Cotton, G. Wilkinson, C.A. Murillo, M. Bochmann, *Advanced Inorganic Chemistry*, sixth ed., John Wiley & Sons, New York, 1999.
- [8] T. Rein, T.M. Pedersen, Asymmetric wittig type reactions, *Synthesis* 5 (2002) 579-594.
- [9] N. Bricklebank, Ylides and related species, *Organophosphorus Chem.* 33 (2003) 289-320.
- [10] L.Y. Wu, M.O. Anderson, Y. Toriyabe, J. Maung, T.Y. Campbell, C. Tajon, M. Kazak, J. Moser, C.E. Berkman, The Molecular Pruning of a Phosphoramidate Peptidomimetic Inhibitor of Prostate-Specific Membrane Antigen, *Bioorg. Med. Chem.* 15 (2007) 7434-7443.

- [11] H. Chapman, M. Kernan, E. Prisbe, J. Rohloff, M. Sparacino, T. Terhorst, R. Yu, Practical synthesis, separation, and stereochemical assignment of the pmpa pro-drug gs-7340, *Nucleos. Nucleot. Nucl. Acids* 20 (2001) 621-628.
- [12] H. Chapman, M. Kernan, J. Rohloff, M. Sparacino, T. Terhorst, Purification of PMPA amidate prodrugs by SMB chromatography and x-ray crystallography of the diastereomerically pure GS-7340, *Nucleos. Nucleot. Nucl. Acids* 20 (2001) 1085-1090.
- [13] M. Lewandowska, P. Ruszkowski, D. Baraniak, A. Czarnecka, N. Kleczewska, L. Celewicz, Synthesis of 3'-azido-2',3'-dideoxy-5-fluorouridine phosphoramidates and evaluation of their anticancer activity, *Eur. J. Med. Chem.* 67 (2013) 188–195.
- [14] Z. Wang, C. Guo, W. Xie, C. Liu, C. Xiao, Z. Tan, Novel phosphoramidates with porphine and nitrogenous drug: One-pot synthesis and orientation to cancer cells, *Eur. J. Med. Chem.* 45 (2010) 890–895.
- [15] M. Donghi, B. Attenni, C. Gardelli, A.D. Marco, F. Fiore, C. Giuliano, R. Laufer, J. F. Leone, V. Pucci, M. Rowley, F. Narjes, Synthesis and evaluation of novel phosphoramidate prodrugs of 2'-methyl cytidine as inhibitors of hepatitis C virus NS5B polymerase, *Bioorg. Med. Chem. Lett.* 19 (2009) 1392–1395.
- [16] M.J. Sofia, W. Chang, P. A. Furman, R.T. Mosley, B.S. Ross, Nucleoside, nucleotide, and non-nucleoside inhibitors of hepatitis C virus NS5B RNA dependent RNA-polymerase, *J. Med. Chem.* 55 (2012) 2481–2531.
- [17] N.F. Zakirova, A.V. Shipitsyn, M.V. Jasko, M.M. Prokofjeva, V.L. Andronova, G.A. Galegov, V.S. Prassolov, S.N. Kochetkov, Phosphoramidate derivatives of acyclovir: synthesis and antiviral activity in HIV-1 and HSV-1 models in vitro, *Bioorg. Med. Chem.* 20 (2012) 5802–5809.
- [18] C. Mara, E. Dempsey, A. Bell, J.W. Barlow, Synthesis and evaluation of phosphoramidate and phosphorothioamidate analogues of amiprofos methyl as potential antimalarial agents, *Bioorg. Med. Chem. Lett.* 21 (2011) 6180–6183.
- [19] P. Stepnicka, *Ferrocenes: Ligands, Materials and Biomolecules*, Wiley, Hoboken, 2008.
- [20] R.G. Array_S, J. Adrio, J.C. Carretero, Chiral ferrocene ligands in asymmetric catalysis, *Angew. Chem.* 118 (2006) 7836.

- [21] D.R. van Staveren, N. Metzler-Nolte, Bioorganometallic chemistry of ferrocene, *Chem. Rev.* 104 (2004) 5931.
- [22] N.G. Connelly, W.E. Geiger, Chemical Redox Agents for Organometallic Chemistry, *Chem. Rev.* 96 (1996) 877.
- [23] F. Horký, I. Císarovà, Jirí Schulz, P. Stepnika, Synthesis and structural characterisation of 1'-(diphenylphosphino) ferrocene-1-phosphonic acid, its ammonium salts and Pd(II) complexes, *Journal of Organometallic Chemistry* 891 (2019) 44-53.
- [24] R. Shekurov, M. Khrizanforov, T. Gerasimova, Z. Yamaleeva, K. Ivshin, A. Lakomkina, I. Bezkishko, A. Kononov, O. Sinyashin, Y. Budnikova, O. Kataeva, V. Miluykov, Electrochemical Properties and Structure of Multi-Ferrocenyl Phosphorus Thioesters, *Molecules* 25 (2020) 939-948.
- [25] C.O. Turrin, E. Manoury, A.M. Caminade, Ferrocenyl Phosphorhydrazone Dendrimers Synthesis, and Electrochemical and Catalytic Properties, *Molecules* 25 (2020) 447-457.
- [26] T.J. Ajayi, M. Ollengo, L. le Roux, M.N. Pillay, R.J. Staples, S.M. Biros, K. Wenderich, B. Mei, W.E. van Zyl, Heterodimetallic Ferrocenyl Dithiophosphonate Complexes of Nickel(II), Zinc(II) and Cadmium(II) as Sensitizers for TiO₂-Based Dye-Sensitized Solar Cells, *Chemistry Select* 4 (2019) 7416–7424.
- [27] R.D. Chambers, *Organofluorine Chemistry: Fluorinated Alkenes and Reactive Intermediates*, Springer, New York, NY, 2010.
- [28] T. Hiyama, *Organofluorine Compounds: Chemistry and Applications*, Springer, New York, NY, 2000.
- [29] A. Thebti, I. Chniti, M.A. Sanhoury, I. Chehidi, I.H. Ouzari, A. Boudabous, Antimicrobial Activity of Highly Fluorinated Thiocarbamates and Dithiocarbamates, *Curr. Chem. Biol.* 13 (2019) 120-128.
- [30] C.M. Timperley, R.E. Arbon, S.A. Saunders, M.J. Waters, Fluorinated phosphorus compounds: Part 6. The synthesis of bis(fluoroalkyl) phosphites and bis(fluoroalkyl) phosphorohalidates, *J. Fluorine Chem.* 113 (2002) 65-78.
- [31] M. Shinmena, K. Sasaharaa, S. Nakamuraa, T. Kanbarab, T. Yajima, Synthesis of perfluoroalkylene oligo(ethylene glycol) alternative polymer via photoinduced polyaddition, *J. Fluorine Chem.* 229 (2020) 109417.

- [32] M.P. Krafft, P. vierling, J.G. Riess, Synthesis and preliminary data on the biocompatibility and emulsifying properties of perfluoroalkylated phosphoramidates as injectable surfactants, *Eur. J. Med. Chem.* 26 (1991) 545-550.
- [33] V.M. Sadtler, F. Jeanneaux, M.P. Krafft, J. Rábai, J.G. Riess, Perfluoroalkylated amphiphiles with a morpholinophosphate or a dimorpholinophosphate polar head group, *New J. Chem.* 22 (1998) 609-613.
- [34] L. Bahri, T. Barhoumi-Slimi, R Mallek, M.A.K. Sanhoury, B. Crousse, M.T. Ben Dhia, One-pot synthesis of new highly substituted allylic phosphorodiamidates, *J. Fluorine Chem.* 189 (2016) 96–101.
- [35] M. M'haiham, K. Ebeid, F. Ebnou, C.L. Carpenter-Warren, A.M.Z. Slawin, J.D. Woollins, M.T. Ben Dhia, M.A. Sanhoury, Synthesis, characterization, and structures of zinc(II) and cadmium(II) complexes with phosphoramides bearing cyclic amino groups *J. Coord. Chem.* 73 (2020) 1-14.
- [36] R.A. Bowman, J.R. Stock, J.A. Jackson, Improved synthesis of bis(2,2,2-trifluoroethyl) phosphorochloridate, *Organic Preparations and Procedures International: The New Journal for Organic Synthesis* 31 (1999) 230-231.
- [37] O. Oms, F. Maurel, F. Carre, J.L. Bideau, A. Vioux, D. Leclercq, Improved synthesis of diethyl ferrocenylphosphonate, crystal structure of $(\text{FcPO}_3\text{Et}_2)_2 \cdot \text{ZnCl}_2$, and electrochemistry of ferrocenylphosphonates, $\text{FcP}(\text{O})(\text{OR})_2$, $\text{FcCH}_2\text{P}(\text{O})(\text{OR})_2$, $1,10\text{-fc}[\text{P}(\text{O})(\text{OR})_2]_2$ and $[\text{FcP}(\text{O})(\text{OEt})_2]_2 \cdot \text{ZnCl}_2$ ($\text{Fc}=(\eta^5\text{C}_5\text{H}_5)\text{Fe}(\eta^5\text{C}_5\text{H}_4)$, $\text{fc}=(^5\text{C}_5\text{H}_4)\text{Fe}(^5\text{C}_5\text{H}_4)$, $\text{R}=\text{Et}, \text{H}$), *Journal of Organometallic Chemistry* 689 (2004) 2654–2661.
- [38] C.M. Timperley, S.N. Marriott, M.J. Waters, Fluorinated phosphorus compounds: Part 7. The reactions of bis(fluoroalkyl) phosphorochloridates with nitrogen nucleophiles, *J. Fluorine Chem.* 113 (2002) 111-122.
- [39] C.M. Timperley, M. Bird, J.F. Broderick, I. Holden, I.J. Morton, M.J. Waters, Fluorinated phosphorus compounds: Part 1. The synthesis and reactions of some fluoroalkyl phosphoryl compounds, *J. Fluorine Chem.* 104 (2000) 215-223.
- [40] S.R. Alley, W. Henderson, Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic, *J. Organomet. Chem.* 216 (2001) 637–639
- [41] M. Gawron, C. Dietz, M. Lutter, A. Duthie, V. Jouikov, K. Jurkschat, Different Complexation Behavior of P-Functionalized Ferrocene Derivatives Towards SnCl_2 , SnCl_4 and

SnPh₂Cl₂: Auto-ionization and Redox-Type Reactions, *Chemistry A European Journal* 21 (2015) 16609-16622. .

[42] T. Gerasimova, R. Shekurov, L. Gilmanova, A. Laskin, S. Katsyuba, V. Kovalenko, M. Khrizanforov, V. Milyukov, O; Sinyashin, IR and UV study of reversible water-induced structural transformations of poly (manganese 1, 10-ferrocenediyl-bis (H-phosphinate)) and poly (cobalt 1,10-ferrocenediyl-bis (H-phosphinate)), *J. Mol. Struc.* 1166 (2018) 237–242.