



**HAL**  
open science

## A neuroergonomic approach to performance estimation in a psychomotor vigilance task

Emilie Jahanpour, Xiaoqi Xu, Marcel F. Hinss, Nicolas Drougard, Raphaëlle  
N. Roy

► **To cite this version:**

Emilie Jahanpour, Xiaoqi Xu, Marcel F. Hinss, Nicolas Drougard, Raphaëlle N. Roy. A neuroergonomic approach to performance estimation in a psychomotor vigilance task. Neuroergonomics Conference 2021, Sep 2021, Munich, Germany. pp.0. hal-03576486

**HAL Id: hal-03576486**

**<https://hal.science/hal-03576486>**

Submitted on 16 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A neuroergonomic approach to performance estimation in a psychomotor vigilance task

Emilie Soheila Jahanpour<sup>\*1</sup>, Xiaoqi Xu<sup>\*1,2</sup>, Marcel Francis Hinss<sup>1,3</sup>, Nicolas Drougard<sup>1,2</sup>,  
Raphaëlle N. Roy<sup>1,2</sup>

[\*denotes equal contribution]

[<sup>1</sup>ISAE-SUPAERO, Université de Toulouse, France]

[<sup>2</sup>Artificial and Natural Intelligence Toulouse Institute (ANITI), Université de Toulouse, France]

[<sup>3</sup>Maastricht University, Netherlands]

**Keywords:** passive BCI; EEG; ECG; performance; vigilance; PVT

**Introduction:** Passive brain-computer interfaces (pBCI; tools that enable an implicit mental state estimation) have gained attention in a wide range of applications, including performance and vigilance monitoring in high-risk work settings (Lotte & Roy, 2019). Vigilance can be defined as the ability to maintain sustained attention to a stimulus for an extended period of time (Al-Shargie et al., 2019), and is influenced by the time of day and fatigue (Lim and Dinges, 2008). A vigilance decrement impacts performance over time (called time-on-task -TOT) during tedious monitoring tasks, resulting in slower reaction times or increased errors (Pattyn et al., 2008). This effect is experienced in all kinds of activities such as in aeronautics where pilots can experience a performance drop during the flight (Wiggins, 2011). Hence, vigilance and performance estimation is a crucial step towards the implementation of safer work settings. Machine learning applied to physiological measures, such as cerebral activity (via electroencephalogram -EEG- recordings), is a promising way to estimate performance. EEG's spectral activity is impacted by fluctuations in vigilance (Matousek & Petersén, 1983) and the power in both theta and alpha bands can be considered as robust biomarkers of mental fatigue (Tran et al., 2020). Numerous studies have attempted to estimate performance during a vigilance task based on EEG measures (Tian et al. 2018), or electrocardiographic (ECG) measures (e.g., heart rate -HR-, and its variability -HRV; Chua et al., 2012). To our knowledge, performance estimation during monotonous tasks has not reached a high accuracy, and pBCI pipelines could be improved. Hence, the objective of this study is to employ a comprehensive neuroergonomic approach to vigilance and performance characterization for a typical vigilance task: the Psychomotor Vigilance Task (PVT; Dinges & Powell, 1985) encompassing statistical analyses, as well as EEG-based performance classification using pre-stimulus signal.

**Methods:** Ten volunteers (3 females;  $M_{age}=25$ ,  $sd=3$ ; ethical number from Univ. Toulouse: 2021-342) performed a 10-minute PVT (i.e., 90 stimuli) from a task battery. They had to complete a fatigue questionnaire (Karolinska Sleepiness Scale; Åkerstedt & Gillberg, 1990) and their response time (RT) was measured. In addition, EEG and ECG activities were recorded using an ActiChamp system (63+1 electrodes). Data were processed in two ways: i) TOT analysis: data were split into ten 1-minute windows; and ii) Performance analysis: response-based epoching (-2:0s for EEG; -10:0s for ECG), only the 30 best and 30 worst trials were kept (labeled according to RT), and also used for performance classification. Considering statistical analyses, one-way repeated measures ANOVAs and paired samples t-tests (Student and Wilcoxon), were performed separately on each dataset. To perform TOT analysis on

RT, the signal was cut into 10 periods of 9 simultaneous stimuli and the reciprocal RT (mean  $1/RT$ ) was calculated on each period. EEG data were filtered (1-40 Hz) and ocular artifacts were automatically removed (SOBI method). The theta, alpha, and beta power were extracted for three electrode clusters (frontal, central, posterior). The Task Load Index (TLI:  $\theta_{a_{Fz}}/\alpha_{p_z}$ ) and the engagement ratio ( $\beta/[\theta+\alpha]$ ; average on all electrodes) were also calculated. ECG data were filtered (1-40 Hz) and normalized using a 1-minute eyes-open resting state period. HR and HRV (as SDNN) were then extracted. For EEG-based estimation, a dimensionality reduction method based on Laplacian was applied (Xu et al., 2021), and a minimum distance to mean with geodesic filtering classifier (FgMDM) was trained and tested using a 10-fold cross-validation procedure.

**Results:** There was no significant difference in subjective fatigue (regardless of PVT order in battery). There was a significant linear downward trend on reciprocal RT ( $p<.05$ ; other contrasts n.s.; Fig.1.A). There was a significant effect of TOT on alpha power at frontal ( $p<.05$ ,  $\eta_p^2=.26$ ) and posterior ( $p<.05$ ,  $\eta_p^2=.27$ ) sites, as well as on the engagement ratio ( $p<.05$ ,  $\eta_p^2=.31$ ; Fig.1.B). TOT also significantly impacted HR ( $p<.001$ ,  $\eta_p^2=.31$ ; Fig.1.C) and HRV ( $p<.05$ ,  $\eta_p^2=.27$ ). Regarding performance (best/worst trials), alpha power was significantly lower for the best than worst trials at frontal ( $p<.01$ ,  $r_b=-.93$ ), and posterior ( $p<.01$ ,  $r_b=-.89$ ) sites, while the TLI was higher for the best trials ( $p<.05$ ,  $d=.87$ ). In terms of performance classification, the FgMDM classifier achieved a mean accuracy of 58.2% without dimensionality reduction. By selecting the number of dimensions that gives the best accuracy for each subject, the mean accuracy with dimensionality reduction was 70.5% (mean number of dimensions = 27.3, Fig.2.A). By comparison, the principal component analysis (PCA) achieved a mean accuracy of 71.1% (mean number of dimensions = 23).

**Discussion:** The results showed that the neuroergonomic approach employed in this study enabled us to assess physiological modulations due to vigilance fluctuations, such as alpha power and engagement ratio decreased over time. However, the HR drop observed in the first minutes may be due to the initial presentation novelty (Kelsey et al., 1999). Also, analyses of the best and worst trials showed that vigilance decreased during the bad trials. However, some results were not as strong as those obtained in the literature. This may be due to either the short task duration (10 minutes) compared to the several sessions performed over several hours or days in the literature, or because our participants were not sleep deprived. Besides, these analyses were performed on a small number of participants ( $n=10$ ). Regarding performance estimation, the classification accuracy was low for several reasons. Firstly, there is little difference in terms of vigilance during the 10-minute PVT task, even though we have chosen the 30 best and worst trials. Secondly, the number of training samples is quite limited (i.e., 60 per subject). In this case, dimensionality reduction becomes essential as shown by the results. Both dimensionality reduction methods (non-supervised) significantly improve accuracy while reducing the computation time. Moreover, on average the Laplacian-based method performs better in lower dimensions than PCA (Fig.2.B). In summary, this study shows that even with a short task and a normal level of fatigue, it is possible to observe an impact of a monotonous task on behavioral and physiological measures at the group level. Yet, it remains difficult to implement an accurate performance estimation pipeline using a single short session at the individual level.




Figure 1. A. Reciprocal reaction time (mean  $1/RT$ ) of each time period corresponding to 9 stimuli, dotted line represents the linear trend; B. EEG engagement ratio ( $\beta/(\theta+\alpha)$ ) for each minute; C. Heart rate (HR) normalized over 1-minute of open eyes resting state across minutes ; Error bars represent 95% confidence interval (\*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$ )




Figure 2. A. Best classification accuracy across all dimensions for each subject; B. Mean classification accuracy across all subjects as a function of dimension. Adjusted chance level calculated from (Combrisson and Jerbi, 2015).

## References:

- Åkerstedt, T., and Gillberg, M. (1990). Subjective and Objective Sleepiness in the Active Individual. *Int. J. Neurosci.* 52, 29–37. doi:10.3109/00207459008994241.
- Al-Shargie, F., Tariq, U., Mir, H., Alawar, H., Babiloni, F., and Al-Nashash, H. (2019). Vigilance Decrement and Enhancement Techniques: A Review. *Brain Sci.* 9, 178. doi:10.3390/brainsci9080178.
- Chua, E. C.-P., Tan, W.-Q., Yeo, S.-C., Lau, P., Lee, I., Mien, I. H., et al. (2012). Heart Rate Variability Can Be Used to Estimate Sleepiness-related Decrements in Psychomotor Vigilance during Total Sleep Deprivation. *Sleep* 35, 325–334. doi:10.5665/sleep.1688.
- Combrisson, E., and Jerbi, K. (2015). Exceeding chance level by chance: The caveat of theoretical chance levels in brain signal classification and statistical assessment of decoding accuracy. *J. Neurosci. Methods* 250, 126–136. doi:10.1016/j.jneumeth.2015.01.010.
- Dinges, D. F., and Powell, J. W. (1985). Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *Behav. Res. Methods Instrum. Comput.* 17, 652–655. doi:10.3758/BF03200977.
- Kelsey, R. M., Blascovich, J., Tomaka, J., Leitten, C. L., Schneider, T. R., and Wiens, S. (1999). Cardiovascular reactivity and adaptation to recurrent psychological stress: Effects of prior task exposure. *Psychophysiology* 36, 818–831. doi:10.1111/1469-8986.3660818.
- Lim, J., and Dinges, D. F. (2008). Sleep deprivation and vigilant attention. *Ann. N. Y. Acad. Sci.* 1129, 305–322. doi:10.1196/annals.1417.002.
- Lotte, F., and Roy, R. N. (2019). “Chapter 7 - Brain-Computer Interface Contributions to Neuroergonomics,” in *Neuroergonomics*, eds. H. Ayaz and F. Dehais (Academic Press), 43–48. doi:10.1016/B978-0-12-811926-6.00007-5.
- Matousek, M., and Petersén, I. (1983). A method for assessing alertness fluctuations from EEG spectra. *Electroencephalogr. Clin. Neurophysiol.* 55, 108–113. doi:10.1016/0013-4694(83)90154-2.
- Pattyn, N., Neyt, X., Henderickx, D., and Soetens, E. (2008). Psychophysiological investigation of vigilance decrement: Boredom or cognitive fatigue? *Physiol. Behav.* 93, 369–378. doi:10.1016/j.physbeh.2007.09.016.
- Tian, S., Wang, Y., Dong, G., Pei, W., and Chen, H. (2018). Mental Fatigue Estimation Using EEG in a Vigilance Task and Resting States. in *2018 40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, 1980–1983. doi:10.1109/EMBC.2018.8512666.
- Tran, Y., Craig, A., Craig, R., Chai, R., and Nguyen, H. (2020). The influence of mental fatigue on brain activity: Evidence from a systematic review with meta-analyses. *Psychophysiology* 57, e13554. doi:10.1111/psyp.13554.
- Wiggins, M. W. (2011). Vigilance decrement during a simulated general aviation flight. *Appl. Cogn. Psychol.* 25, 229–235. doi:10.1002/acp.1668.
- Xu, X., Drougard, N., and Roy, R. N. (2021). Dimensionality Reduction via the Laplace-Beltrami Operator: Application to EEG-based BCI. in *2021 10th International IEEE/EMBS Conference on Neural Engineering (NER)*, 347–350. doi:10.1109/NER49283.2021.9441404.