- 1 Influence of exposure assessment methods on associations between long-term exposure to
- 2 outdoor fine particulate matter and risk of cancer in the French cohort Gazel
- 3 Authors: Emeline Lequy^{1,2*}, Mohammad Javad Zare Sakhvidi³, Danielle Vienneau^{4,5}, Kees de
- 4 Hoogh^{4,5}, Jie Chen⁶, Jean-François Dupuy⁷, Valérie Garès⁷, Emilie Burte³, Olivier Bouaziz⁸,
- 5 Alain Le Tertre⁹, Vérène Wagner⁹, Ole Hertel¹⁰, Jesper Heile Christensen¹⁰, Sergey Zhivin¹,
- 6 Jack Siemiatycki², Marcel Goldberg¹, Marie Zins¹, Bénédicte Jacquemin³
- 7 1: Unité "Cohortes en Population" UMS 011 Inserm/Université de Paris/Université Paris
- 8 Saclay/UVSQ, Villejuif, France
- 9 2: Centre de recherche du centre hospitalier de l'université de Montréal, Québec, Canada
- 10 3: Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail)
- 11 UMR_S 1085, F-35000 Rennes, France
- 12 4: Swiss Tropical and Public Health Institute, Basel, Switzerland.
- 13 5: University of Basel, Basel, Switzerland.
- 14 6: Utrecht University, Utrecht, the Netherlands
- 15 7: Univ Rennes, INSA, CNRS, IRMAR UMR 6625, F-35000 Rennes, France
- 16 8 : MAP5, UMR CNRS 8145, Université de Paris, Paris, France
- 17 9: Santé publique France, Saint-Maurice, France
- 18 10: Dep. Env. Science, Aarhus University, Frederiksborgvej 399, 4000 Roskilde, Denmark
- 19
- 20
- 21
- 22 *Address correspondence to:
- 23 Emeline Lequy, UMS 011, Hôpital Paul Brousse, 16 avenue Paul Vaillant Couturier, 94807
 24 VILLEJUIF CEDEX, France. Email: <u>e.lequy@gmail.com</u>. Phone number: +33 (0)1 77 74 74 23.
- 25
- 26

27 Declaration of competing financial interests:

28 The authors declare they have no actual or potential competing financial interests.

29 Table of Content

31	Supplementary Methods	. 3
32	Lag period	.3
33	Specificities of the Cox model	. 3
34	Hazard ratios of the two-piece linear model	. 3
35	Supplementary Figures	.4
36	Supplementary Tables	10
37		

39 Supplementary Methods

- 40 Lag period
- 41 We implemented a 10-year lag period, since 10 years is the period estimated for a cancer to
- 42 develop, except hematopoietic cancers with a small frequency in our dataset.
- 43

44 Specificities of the Cox model

- 45 We organized the data in a counting process style, with one entry per available year per
- 46 participant until incidence/censoring. In our dataset, PM2.5 concentrations overall
- 47 decreased with time, while participants aged and cancer incidence risk increased. For these
- 48 reasons, using annual exposure variables may lead to spurious associations. Besides, we were
- 49 interested in the association of cancer incidence and long-term PM2.5 exposure, available at
- 50 an annual time step. The extended Cox model with time-dependent variables estimates the
- 51 instantaneous risk with the exposure at the date of the event: using only the annual levels of
- 52 exposure may not estimate the association between incident cancer and long-term exposure.
- 53 We addressed these problems by using cumulative exposures for each participant from
- 54 baseline to incidence or censoring, as previously used (Leguy et al., 2021), and by adjusting
- 55 for calendar time, dichotomized with a cut-off in 2007.
- 56 Hazard ratios of the two-piece linear model
- 57 We provided two types of hazard ratios (HR) and their 95% confidence intervals: the first HR
- 58 will be that provided directly by the model and corresponds to the slope of the first part of the
- 59 curve (per one IQR increase of cumulative exposure); the second will approximate the value
- 60 of the plateau using the coefficients of the Cox model to calculate the HR and 95% CI at the
- 61 80th percentile of cumulative exposure ($_{380 \mu g}/m^3$ for the LUR, $_{215 \mu g}/m^3$ for Gazel-Air).

62 Supplementary Figures



- 64 Figure S1: residential places of Gazel participants (gray dots) in mainland
- 65 France in 1989 (n=20,430).

66

63



68 Figure S2: flowchart of the selection of Gazel participants included in the

- 69 analyses pertaining to all-site and lung cancer.
- 70



72

73





76 **iteration number for the imputed data**. Smokpac: smoking pack-years; fdep2009:

deprivation index (put back as nonimputed values in the final database); smokpas: passive

- 78 smoking at work or at home; alcoclass: alcohol consumption; family1: marital status; vegfr:
- 79 fruit and vegetable intake; ses: socioeconomic status; educ1: education; density2: urban
- 80 classification (put back as nonimputed values in the final database).



83 Figure S4: Associations between cumulative PM2.5 and all-site incident cancer (left, with 293,188 person-years and 3,711 incident cancer cases) and lung 84 incident cancer (right, with 254,135 person-years and 349 incident cancer cases) 85 in the Gazel cohort, expressed as hazard ratio (HR) with the lowest exposure as 86 87 the reference for each exposure assessment method. Cumulative exposure to PM2.5 88 was estimated by the LUR or the Gazel-Air exposure assessments in separate Cox models 89 with attained age as underlying time-scale and time-dependent variables, adjusted for sex, 90 cumulative smoking pack-years, passive smoking, alcohol use, BMI, education, 91 socioeconomic status, family status, fruit and vegetable consumption, occupational exposure 92 to lung carcinogens, age at inclusion and calendar time. The x-axis represents the cumulative 93 exposure for both exposure assessment methods, the LUR at the top of the axis and Gazel-Air at the 94 bottom.





97 Figure S5: Bland-Altman graphs depicting the differences (y-axis) between the

98 LUR and the Gazel-Air exposure values for Gazel participant in function of the

99 mean exposure (x-axis) between the two exposure assessment methods, whether

100 by European NUTS-1 region (combining all the years) or by year (combining all

101 **the regions).** From top-left, left panel: NUTS1 from one to eight; from top-left, right panel:

102 follow-up year from 0 (1989) to 19 (2008).





105 **Figure S6: exposure-response relationships between PM2.5 and all-site cancer**

- 106 risk in the Gazel cohort using a spline function for 293,188 person-years (3,711
- 107 cases). Main analysis, sensitivity analyses, and effect modification analyses.
- 108 Hazard ratios and confidence intervals with the lowest exposure as the reference for each

- 109 exposure assessment method were estimated by separate Cox model with attained age as
- 110 underlying time-scale and time-dependent variables, adjusted for sex, cumulative smoking
- 111 pack-years, passive smoking, alcohol use, BMI, education, socioeconomic status, family
- 112 status, vegetable and fruit consumption, occupational exposure to lung carcinogens, age at
- 113 inclusion and calendar time. Exposures were lagged 10 years. Participants were excluded
- 114 from the analysis if they were diagnosed with cancer before 1999. The x-axis represents the
- 115 cumulative exposure for both exposure assessment methods, the LUR at the top of the axis and Gazel-
- 116 Air at the bottom.

117 Supplementary Tables

- 118 Table S1: associations between long-term exposure to PM2.5 and first-occurring
- 119 all-site incident cancer in the Gazel cohort using two exposure models using log-
- 120 transformed cumulative exposure (Hazard Ratios 95% Confidence Interval).
- 121 Hazard ratios correspond to the increased risk of cancer for 1-unit increase of the natural log-
- 122 transformed cumulative exposure to PM2.5.

		LUR		Gazel-Air	
		HR	95% CI	HR	95% CI
Main		1.21	1.07-1.37	1.19	1.07-1.33
Sensitivity analyses					
	Further adjusted for area-level				
	deprivation	1.24	1.10-1.40	1.22	1.09-1.36
	Using address-level geocodes	1.28	0.98-1.66	1.26	1.00-1.58
	Complete-case analysis	1.15	0.93-1.51	1.21	1.02-1.45
	Missing data as category	1.20	1.04-1.38	1.17	1.04-1.32
	Missing data as median/mode	1.22	1.08-1.38	1.21	1.09-1.35
Effect modificiation					
By sex	Women	1.25	0.97-1.61	1.33	1.04-1.69
	Men	1.20	1.04-1.38	1.16	1.02-1.31
By smoking status	Neversmoker	1.24	1.01-1.53	1.28	1.06-1.54
	Eversmoker	1.20	1.03-1.40	1.15	1.00-1.32
By distance to the road:	<500m	1.27	1.01-1.59	1.21	0.99-1.48
	>500m	1.21	1.04-1.41	1.21	1.06-1.38
By urban classification:	Urban (in the Paris region)	1.24	0.84-1.82	1.26	0.83-1.92
	Urban (out of the Paris region)	1.20	0.82-1.74	1.01	0.74-1.40
	Semi-urban	1.17	0.89-1.53	1.21	0.96-1.53
	Rural	1.12	0.85-1.49	1.20	0.94-1.53

Extended Cox model with attained age as time axis, and with time-varying exposure to cumulative exposure to PM2.5, adjusted for sex (included with a strata function), age at enrollment, calendar time (time-varying, dichotomized at year 2007), cumulative pack-years (time-varying), passive smoking, alcohol intake (time-varying), socioeconomic status, marital status (time-varying), body mass index (time-varying), occupational exposure to lung carcinogens, and consumption of fruit and vegetable (time-varying).

123

125 Table S2: stratified associations between long-term exposure to PM2.5 and first-

126 occurring all-site incident cancer in the Gazel cohort using two exposure models.

- 127 Hazard Ratios (HR) and their 95% Confidence Interval (CI) correspond to the increased risk
- 128 of cancer below ("slope", per one IQR increase of cumulative exposure) and above ("plateau")
- 129 the 65th percentile of exposure.

			LUR				Gazel-A	Air		
	Exposure		<p65* (<="" td=""><td>slope)</td><td>>p65 (p</td><td>lateau)</td><td><p65 (s)<="" td=""><td>lope)</td><td>>p65 (p</td><td>lateau)</td></p65></td></p65*>	slope)	>p65 (p	lateau)	<p65 (s)<="" td=""><td>lope)</td><td>>p65 (p</td><td>lateau)</td></p65>	lope)	>p65 (p	lateau)
Effect		Cases	HR	95% CI	HR**	CI**	HR	95% CI	HR**	CI**
modificiation										
By sex	Women	755	1.51	0.98-2.30	1.72	0.73-4.07	1.60	1.06-2.41	1.88	0.83-4.04
	Men	2956	1.53	1.21-1.94	1.56	1.18-2.06	1.39	1.13-1.72	1.39	1.12-1.72
By smoking	Neversmoker	878	1.37	0.96-1.96	1.30	0.72-2.36	1.36	0.98-1.89	1.36	0.84-2.22
status	Eversmoker	2833	1.62	1.25-2.11	1.77	1.27-2.46	1.48	1.17-1.86	1.53	1.19-1.98
By distance	<500m	1,244	1.69	1.15-2.48	2.10	1.04-4.26	1.60	1.13-2.26	1.86	1.06-3.26
to the road	>500m	2,467	1.47	1.14-1.89	1.47	1.06-2.06	1.37	1.09-1.72	1.39	1.07-1.80
By urban	Urban (in the	423	1.48	0.77-2.84	1.51	0.34-6.80	1.43	0.67-3.06	1.29	0.20-8.49
classification	Parisregion)									
	Urban (out of the	503	1.61	0.85-3.06	1.64	0.25-10.80	1.12	0.63-2.00	0.91	0.20-4.09
	Paris region)									
	Semi-urban	878	1.27	0.80-2.01	1.01	0.34-3.06	1.49	0.99-2.24	1.44	0.59-3.52
	Rural	722	1.26	0.78-2.03	1.08	0.25-4.61	1.26	0.84-1.89	1.18	0.34-4.09

Extended Cox model with attained age as time axis, and with time-varying exposure to cumulative exposure to PM2.5, adjusted for sex (included with a strata function), age at enrollment, calendar time (time-varying, dichotomized at year 2007), cumulative pack-years (time-varying), passive smoking, alcohol intake (time-varying), socioeconomic status, marital status (time-varying), body mass index (time-varying), occupational exposure to lung carcinogens, and consumption of fruit and vegetable (time-varying). *p65 is the 65th percentile of cumulative exposure and corresponds to 315 μ g/m³ and to 185 μ g/m³ for the LUR and the Gazel-Air exposure assessments, respectively. IQR= 216 μ g/m³ for the LUR, 127 μ g/m³ for Gazel-Air.**: the HR and CI were calculated for the 80th percentile (380 μ g/m³ for the LUR, 215 μ g/m³ for Gazel-Air).

130

132 Table S1: associations between all-site incident cancer and all co-variables in the

122	Gazal cohort as Hazard Ratio (HP) and 95% Confidence	Intorvals (05%CI)
133	Gazer conort as nazaru kauo (FIR) and 95% Confidence	Intervals (95%CI).

Variable	HR	95% CI
Smoking (for 5 cumulative pack-years)	1.04	1.03-1.05
Passive smoking - no	1 (reference)	
Yes	1.01	0.94-1.09
Alcohol use - light drinker	1 (reference)	
Abstinent	1.04	0.91-1.20
Moderate drinker	1.05	0.96-1.15
Heavy drinker	1.23	1.11-1.37
Unclear pattern	1.02	0.90-1.15
Socio-economic status - low	1 (reference)	
Intermediate	0.93	0.85-1.02
High	0.95	0.84-1.08
Family status - single	1 (reference)	
Not single	0.99	0.89-1.11
Body Mass Index (for 1 unit)	1.01	1.00-1.02
Vegetable & fruit consumption - every day or almost	1 (reference)	
Never or less than once a week	1.72	1.18-2.52
Once or twice a week	1.11	0.95-1.31
More than twice a week, not everyday	1.09	0.99-1.20
Education - 9-11 years	1 (reference)	
12-13 years	0.90	0.78-1.03
14-15 years	1.00	0.86-1.16
Other secondary diploma	0.93	0.81-1.06
Other diploma	0.92	0.74-1.15
Occupational exposure* - none	1 (reference)	
One	0.95	0.84-1.07
Two	1.00	0.89-1.11
Three or more	0.97	0.88-1.06
Age at inclusion (for one year)	1.02	1.00-1.04
Calendar year - before 2007	1 (reference)	
After 2007	1.22	1.07-1.39

*: to nine selected lung carcinogens.

Extended Cox model with attained age as time axis, and with time-varying exposure to cumulative exposure to PM_{2.5}, adjusted for sex (included with a strata function), age at enrollment, calendar time (time-varying, dichotomized at year 2007), cumulative pack-years (time-varying), passive smoking, alcohol intake (time-varying), socioeconomic status, marital status (time-varying), body mass index (time-varying), occupational exposure to lung carcinogens, and consumption of fruit and vegetable (time-varying). Results obtained with 3,711 cases of all-site cancer and 293,188 person-years.

134