



**HAL**  
open science

## Revisiting the Gas-Phase Rate Coefficient of the $\text{OH} + \text{SO}_2 + \text{M} \rightarrow \text{HSO}_3 + \text{M}$ Reaction

Max R McGillen, H Elothmani, Yangang Ren, Li Zhou, Mahmoud Idir, Souad El Hajjaji, Véronique Daële, A. Mellouki

► **To cite this version:**

Max R McGillen, H Elothmani, Yangang Ren, Li Zhou, Mahmoud Idir, et al.. Revisiting the Gas-Phase Rate Coefficient of the  $\text{OH} + \text{SO}_2 + \text{M} \rightarrow \text{HSO}_3 + \text{M}$  Reaction. The 11th International Conference on Chemical Kinetics, Jun 2019, Orleans, France. hal-03580140

**HAL Id: hal-03580140**

**<https://hal.science/hal-03580140>**

Submitted on 2 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Revisiting the gas-phase rate coefficient of the $\text{OH} + \text{SO}_2 + \text{M} \rightarrow \text{HSO}_3 + \text{M}$ reaction

M.R. McGillen<sup>1,2,\*</sup>, H. Elothmani<sup>1</sup>, Y. Ren<sup>1</sup>, L. Zhou<sup>1</sup>, M. Idir<sup>1</sup>, V. Daële<sup>1</sup>, A.R. Ravishankara<sup>1,2,3,4</sup>  
and A. Mellouki<sup>1</sup>

<sup>1</sup>CNRS-INSIS, Institut de Combustion, Aérodynamique, Réactivité et Environnement 1C, Avenue de la recherche scientifique, 45071 Orléans cedex 2, France

<sup>2</sup>Le Studium Loire Valley Institute for Advanced Studies, Orléans, France.

<sup>3</sup>Departments of Chemistry and Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA

<sup>4</sup>NOAA Earth System Research Laboratory (ESRL) Chemical Sciences Division, 325 Broadway, Boulder, CO 80305, USA

## Keywords

*Hydroxyl, Sulfur dioxide, Kinetics, Pressure dependence, Temperature dependence.*

Oxidation of  $\text{SO}_2$  to sulfuric acid impacts acid precipitation and aerosol nucleation in Earth's atmosphere in remote and polluted environments. This oxidation can take place in both the liquid and gas phase. Only the gas phase oxidation is expected to lead to new particles because of the clustering reactions of  $\text{H}_2\text{SO}_4$ . This aerosol nucleation has a major effect upon air quality and the Earth's radiative balance, and is of crucial importance to the chemistry of the atmosphere.

The rate limiting step in this process is the reaction of OH radicals with  $\text{SO}_2$  to form  $\text{HSO}_3$ . The pressure- and temperature-dependent reaction of  $\text{OH} + \text{SO}_2$  has been studied many times previously – since its importance was first recognized in the 1970s. Notwithstanding, some of the most recent literature has cast doubt on much of this data, especially under conditions that are relevant to atmospheric chemistry.

Here, we present measurements of the rate coefficient using the pulsed laser photolysis–laser induced fluorescence technique as a function of temperature (249–373 K) and of pressure in helium, argon and nitrogen bath gases (30–600 Torr). In addition, relative rate measurements using a chamber at 760 Torr ( $\text{N}_2$ ,  $\text{O}_2$  and air) were also performed to corroborate our absolute observations. By utilizing these new data, together with the available literature data, an updated pressure- and temperature-dependent parameterization will be provided. This allows the atmospheric impact of this reaction to be constrained with a new level of certainty.

\*Corresponding author: e-mail address