



HAL
open science

Projective classes as images of accessible functors

Friedrich Wehrung

► **To cite this version:**

| Friedrich Wehrung. Projective classes as images of accessible functors. 2022. hal-03580184v1

HAL Id: hal-03580184

<https://hal.science/hal-03580184v1>

Preprint submitted on 18 Feb 2022 (v1), last revised 2 May 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROJECTIVE CLASSES AS IMAGES OF ACCESSIBLE FUNCTORS

FRIEDRICH WEHRUNG

ABSTRACT. We are dealing with projective classes (in short PC) over first-order vocabularies with no restrictions on the (possibly infinite) arities of relation or operation symbols. We verify that $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$ for any infinite cardinal λ , and that if λ is singular, then $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{PC}(\mathcal{L}_{\infty\lambda^+})$. If λ is regular, then a class of structures over a λ -ary vocabulary is $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable iff it is the image of a λ -continuous functor on a λ -accessible category. It follows that many PC classes of structures, previously known not to be closed under elementary equivalence over any $\mathcal{L}_{\infty\lambda}$, are not co-PC over $\mathcal{L}_{\infty\infty}$. Those classes arise from diverse contexts including convex ℓ -subgroup lattices of lattice-ordered groups, ideal lattices of rings, nonstable K_0 -theory of rings, coordinatization of sectionally complemented modular lattices, and real spectra of commutative unital rings. For example, the class of posets of finitely generated two-sided ideals of all unital rings is PC but not co-PC over $\mathcal{L}_{\infty\infty}$. We also provide a negative solution to a problem, raised in 2011 by Gillibert and the author, asking for an extension of the condensate construction to commutative diagrams indexed by arbitrary bounded posets.

CONTENTS

1. Introduction	2
1.1. Intractability results	2
1.2. PC and RPC classes	3
1.3. Back to GIS and images of accessible functors	4
2. Basic concepts	5
2.1. Sets, ordinals, cardinals, posets	5
2.2. Categories and functors	5
2.3. Infinitary languages	6
3. Projective and relatively projective classes	7
4. A first case of conjugacy: partial λ -ary truncating	8
5. An $\mathcal{L}_{\infty\lambda}$ version of the Skolem normal form	12
6. Another case of conjugacy: from RPC to PC	16
7. RPC classes as images of continuous functors	22
8. The case of singular cardinals	26
9. Back-and-forth systems, anti-elementarity, and Tuuri's Interpolation Theorem	30
9.1. Karttunen's back-and-forth systems	30

Date: February 18, 2022.

2010 Mathematics Subject Classification. 03C75, 18C35, 03C40, 06D35, 16D70, 16E50, 14P10.

Key words and phrases. First-order; vocabulary; infinitary language; infinitely deep language; projective; accessible; category; continuous; functor; colimit; lattice; ℓ -group; ring; real spectrum; condensate.

9.2. The infinitely deep languages $\mathcal{M}_{\kappa\lambda}$	32
10. Application to intractability results	33
10.1. Posets of convex ℓ -subgroups in ℓ -groups	33
10.2. Posets of ideals in rings	35
10.3. Nonstable K_0 -theory of rings	35
10.4. Coordinatizability of sectionally complemented modular lattices	35
10.5. Real spectra of commutative unital rings	37
11. Beyond co-PC: Ploščica's Condition and ℓ -spectra	38
12. A finite bounded poset for which condensates do not work	40
References	43

1. INTRODUCTION

1.1. Intractability results. There are numerous results stating that a given class \mathcal{C} , of structures over a given vocabulary (with possibly infinite arities) \mathbf{v} , is “intractable” in the sense that \mathcal{C} is not closed under $\mathcal{L}_{\infty\lambda}$ -elementary equivalence for any infinite cardinal λ (due to a classical result of Chang [4, Proposition 6], this is equivalent to saying that \mathcal{C} is not the class of models of any class of $\mathcal{L}_{\infty\lambda}$ -sentences). Many of those results take advantage of \mathcal{C} being the image of a functor Φ defined on a category \mathcal{S} , with Φ taking values in all structures over a given vocabulary and both \mathcal{S} and Φ “well-behaved” in some sense. Good behavior is here expressed by stating that the category \mathcal{S} and the functor Φ are both *accessible* as defined in Makkai and Paré [20] (see Subsection 2.2 for definitions). Examples, originating in various references including Goodearl [11], Jónsson [17], Keimel [19], Mundici [23], investigated in Mellor and Tressl [22], Wehrung [33, 37], include the following:

- (1) \mathcal{S} is a large enough accessible category of lattice-ordered groups (in short *ℓ -groups*) and Φ sends every object G of \mathcal{S} to its lattice $\text{Cs}_c G$ of finitely generated convex ℓ -subgroups (resp., its semilattice $\text{Id}_c G$ of finitely generated ℓ -ideals);
- (2) \mathcal{S} is a large enough accessible category of rings and Φ sends every object R of \mathcal{S} to its semilattice $\text{Id}_c R$ of finitely generated two-sided ideals;
- (3) \mathcal{S} is a large enough accessible category of unital rings and Φ sends every object R of \mathcal{S} to the monoid $V(R)$ of all isomorphism classes of finitely generated projective right R -modules;
- (4) \mathcal{S} is a large enough accessible category of von Neumann regular rings, possibly with additional structure, and Φ sends every object R of \mathcal{S} to the lattice $\mathbb{L}(R)$ of all its principal right ideals;
- (5) \mathcal{S} is the category of all commutative unital rings and Φ sends every object A of \mathcal{S} to the Stone dual of its real spectrum.

The meaning of “large enough” varies from one context to the other. The corresponding results, described in Section 10, all state a level of intractability that takes the following general form, involving the concept of a *PC-definable class* which we shall discuss in more detail shortly (cf. Subsection 1.2).

General Intractability Scheme (GIS). *Classes \mathcal{C}_0 and \mathcal{C}_1 of structures are constructed such that \mathcal{C}_0 is $\text{PC}(\mathcal{L}_{\infty\infty})$ -definable, $\mathcal{C}_0 \subseteq \mathcal{C}_1$, and there is no $\text{co-PC}(\mathcal{L}_{\infty\infty})$ -definable class \mathcal{C} such that $\mathcal{C}_0 \subseteq \mathcal{C} \subseteq \mathcal{C}_1$.*

1.2. PC and RPC classes. We are dealing with classes of structures definable over infinitary languages, with possible infinite arities. Standard such classes are provided by σ -complete Boolean algebras or Banach algebras. A less standard such class is provided by Tarski's *cardinal algebras* [30].

For a logic \mathcal{L} (a formal definition of that concept will not be needed here), recall that a class \mathcal{C} of \mathbf{v} -structures is

- *projective over \mathcal{L}* (in short $\text{PC}(\mathcal{L})$) if it is the class of \mathbf{v} -reducts of all models of some \mathcal{L} -sentence over a larger vocabulary \mathbf{w} ;
- *relatively projective over \mathcal{L}* (in short $\text{RPC}(\mathcal{L})$) if it is the class of relativizations to \mathbf{U} of \mathbf{v} -reducts of all models of some \mathcal{L} -sentence over a larger vocabulary \mathbf{w} containing a unary predicate symbol \mathbf{U} .

(For formal definitions, see Definition 3.1.) A class is *co-projective* (in abbreviation co-PC) if its complement is projective. A typical example of a co-PC, non PC (over $\mathcal{L}_{\infty\infty}$) class of structures is the class of all complete Boolean algebras (cf. Cole and Dickmann [5, Theorem 3]). At that point a remark is in order.

Remark 1.1. The equation $\text{PC}(\mathcal{L}_{\infty\omega}) = \text{RPC}(\mathcal{L}_{\infty\omega})$, stated in Oikkonen [24, Corollary 1.3], is extended in Oikkonen [25, Remark 1.1], with a sketch of proof, to the logics $\mathcal{L}_{\kappa\lambda}$, where κ and λ are regular cardinals such that $\alpha^\beta < \kappa$ whenever $\alpha < \kappa$ and $\beta < \lambda$. However, the given argument, stated there for relational vocabularies, does not apply to vocabularies containing infinitary operation symbols *a priori*:

- the possible presence of many operation symbols, even within a single $\mathcal{L}_{\kappa\lambda}$ -sentence¹, runs counter to the possibility of a suitable Löwenheim-Skolem type Theorem as required in the argument of [25, Remark 1.1];
- the possible presence of operation symbols with large arity prevents, in Definition 3.1, the \mathbf{v} -closedness of $\mathbf{U}^{\mathbf{N}}$ from being expressible by an $\mathcal{L}_{\infty\lambda}$ -sentence.

We will clarify the interaction between those concepts, in light of accessible categories, by establishing, in full generality, the following results:

- $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$, for any infinite cardinal λ (cf. Corollary 8.7).
- $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{PC}(\mathcal{L}_{\infty\lambda^+})$, for any singular cardinal λ (cf. Theorem 8.6).
- For any regular cardinal λ , every $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable class of structures, over a vocabulary \mathbf{v} in which all operations have arity smaller than λ , is the image of a faithful² λ -continuous functor from a λ -accessible category to \mathbf{v} -structures (cf. Theorem 6.9). The assumption that all operations in \mathbf{v} have arity smaller than λ cannot be dispensed with (cf. Example 5.6).
- For any regular cardinal λ , any λ -accessible category \mathcal{S} , and any λ -continuous functor Φ from \mathcal{S} to all structures over a given λ -ary vocabulary \mathbf{v} , the image of Φ is $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable (cf. Theorem 7.1). The assumption that \mathbf{v} be λ -ary cannot be dispensed with (cf. Example 8.8).

In the course of establishing those results, we will observe that *every $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable class of structures, over a vocabulary \mathbf{v} , is the class of \mathbf{v} -reducts of models of a sentence \mathbf{E} over a vocabulary \mathbf{w} containing \mathbf{v} such that $\mathbf{w} \setminus \mathbf{v}$ is λ -ary; moreover,*

¹Consider the sentence $(\forall x)(x \neq f(\mathbf{g}_\xi(x) \mid \xi < \theta))$ for large θ .

²Throughout this work it seems that faithfulness, of the representing functor, marks the advantage of PC over RPC; see, in particular, Theorem 6.9.

\mathcal{E} can be taken a conjunction of universal $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentences (cf. Theorem 4.11 and Proposition 5.4).³

1.3. Back to GIS and images of accessible functors. In the context of Subsection 1.1(2) above, our General Intractability Scheme GIS yields that the class of posets of finitely generated ideals of unital rings (resp., unital von Neumann regular rings) is not $\text{co-PC}(\mathcal{L}_{\infty\infty})$ -definable (cf. Theorem 10.3). More generally, in GIS, all interesting instances of \mathcal{C} and \mathcal{C}_0 turn out to be images of accessible functors, thus $\text{PC}(\mathcal{L}_{\infty\infty})$ -definable (thus lending some optimality to the scheme). The proof of GIS then runs as follows. If \mathcal{C} were $\text{co-PC}(\mathcal{L}_{\infty\infty})$ -definable, then Tuuri’s Interpolation Theorem from [32] would imply the existence of a sentence, in the infinitely deep language $\mathcal{M}_{\infty\infty}$, whose class of models is intermediate between \mathcal{C}_0 and \mathcal{C} , thus between \mathcal{C}_0 and \mathcal{C}_1 . By a result of Karttunen [18], that class of models is closed under strong back-and-forth equivalence \simeq_λ (cf. Definition 9.1) for large enough λ . This would in turn contradict earlier results by the author [37], employing an extension of the concept of “condensate” introduced in Gillibert and Wehrung [10], and entailing the existence of \simeq_λ -equivalent models, one in \mathcal{C}_0 and the other in the complement of \mathcal{C}_1 .

In the specific context of ℓ -ideal lattices of ℓ -groups, we extend, in Section 11, this method to an *a priori* non co-PC -condition, that we call *Ploščica’s Condition*. In particular, we verify, under the Generalized Continuum Hypothesis GCH, that Ploščica’s Condition, together with all currently known $\mathcal{L}_{\infty\infty}$ -sentences satisfied by all lattices of finitely generated ℓ -ideals in Abelian ℓ -groups, is still not sufficient to characterize the latter objects (we get a counterexample of cardinality \aleph_4).

The construction, in Tůma and Wehrung [31], of a diagram \mathcal{D}_\boxtimes of finite Boolean $(\vee, 0)$ -semilattices that cannot be lifted by any diagram of congruence lattices of (among others) majority algebras⁴, raised the hope for a negative solution to the well known open question asking whether every distributive algebraic lattice is the congruence lattice of a majority algebra. That hope was also supported by the condensate construction from Gillibert and Wehrung [10], introduced as a means to turn diagram counterexamples, *indexed by lattices*, to object counterexamples. However, the diagram \mathcal{D}_\boxtimes above is indexed by a variant of the six-element bounded poset P_0 represented on the left hand side of Figure 12.2; *that poset is not a lattice*. This was the original motivation for [10, Problem 1], which asked for an extension of [10, Theorem 3.4.2] (called there CLL) to arbitrary bounded posets. In Section 12 we use our “image of a functor” approach to solve that problem in the negative. This solution involves the six-element bounded poset P_0 represented on the left hand side of Figure 12.2. Roughly speaking, it consists of a faithful, finitely accessible functor Φ , surjective on objects, from a (finitary) variety \mathcal{V} of algebras to the category \mathbf{DSLat}^0 of all distributive $(\vee, 0)$ -semilattices with $(\vee, 0)$ -homomorphisms, with a P_0 -indexed commutative diagram in \mathbf{DSLat}^0 that cannot be lifted, in a strong sense, with respect to Φ .

³The latter result would not follow from some hypothetical infinitary translation of Skolem normal form reduction *a priori*: for one thing, even the less demanding prenex normal form reduction may fail in $\mathcal{L}_{\infty\lambda}$ (cf. Dickmann [6, Fact 1.1.1]).

⁴A *majority algebra* is a set endowed with a ternary operation m satisfying the equations $m(x, x, y) = m(x, y, x) = m(y, x, x) = x$.

2. BASIC CONCEPTS

2.1. Sets, ordinals, cardinals, posets. Like for many works involving proper classes, a natural axiomatic environment for this work is Bernays-Gödel set theory (cf. Jech [14, p. 70]) with the axiom of choice for sets. The cardinality of a set X will be denoted by $\text{card } X$. For a cardinal κ , a set X , or, more generally, a structure with universe X , is κ -small if $\text{card } X < \kappa$; we denote by $[X]^{<\kappa}$ the set of all κ -small subsets of X . We denote set exponentiation by $(X, Y) \mapsto {}^X Y$ and cardinal exponentiation by $(\alpha, \beta) \mapsto \alpha^\beta$. For cardinals α and κ , we set $\alpha^\kappa \stackrel{\text{def}}{=} \sup\{\alpha^\gamma \mid \gamma < \kappa\}$. We denote by $\text{dom } f$ and $\text{rng } f$ the domain and the range of a function f , respectively. Disjoint unions will be denoted in the form $X \sqcup Y$.

For every element a in a poset P , we shall denote by $P \downarrow a$, or $\downarrow a$ if P is understood, the set $\{x \in P \mid x \leq a\}$. A subset X of P is a *lower subset* of P if $P \downarrow x \subseteq X$ whenever $x \in X$. If, in addition, $P = (P, \vee, 0)$ is a join-semilattice with least element 0 (in abbreviation a $(\vee, 0)$ -semilattice), a nonempty lower subset I of P is an *ideal* if it closed under finite joins. Then I induces a join-congruence \equiv_I on P , by letting $x \equiv_I y$ hold if there exists $z \in I$ such that $x \vee z = y \vee z$. The quotient $(\vee, 0)$ -semilattice is then denoted by P/I . A poset (P, \leq) is a *tree* if it has a least element and every principal ideal $P \downarrow a$, where $a \in P$, is well-ordered.

2.2. Categories and functors. For more details we refer the reader to Adámek and Rosický [1].

Every category \mathcal{S} will be identified with its class of arrows (aka morphisms), and we denote by $\text{Ob } \mathcal{S}$ the class of objects (aka identities) of \mathcal{S} . If $\varphi: A \rightarrow B$ is a morphism in \mathcal{S} , we write $A = \mathbf{d}(\varphi)$ and $B = \mathbf{r}(\varphi)$. Commutative diagrams, indexed by a poset P , will be denoted in the form $\vec{S} = (S_{p_1}, \sigma_{p_1, p_2} \mid p_1 \leq p_2 \text{ in } P)$ (where each $\sigma_{p_1, p_2}: S_{p_1} \rightarrow S_{p_2}$). A cocone above \vec{S} will then be denoted in the form $(S, \sigma_p \mid p \in P)$, with the usual commutation relations (e.g., $\sigma_{p_1} = \sigma_{p_2} \circ \sigma_{p_1, p_2}$ whenever $p_1 \leq p_2$ in P) assumed. Colimit cocones will be written in the form

$$(S, \sigma_p \mid p \in P) = \varinjlim (S_{p_1}, \sigma_{p_1, p_2} \mid p_1 \leq p_2 \text{ in } P) \quad \text{within } \mathcal{S}$$

(this notation defines $(S, \sigma_p \mid p \in P)$ up to isomorphism).

Let λ be a regular cardinal. A subcategory \mathcal{S}^\dagger of a category \mathcal{S} is λ -directed colimit-dense if every object of \mathcal{S} is the colimit, within \mathcal{S} , of a λ -directed commutative diagram of \mathcal{S}^\dagger . The category \mathcal{S} is λ -accessible if it has all λ -directed colimits and it has a λ -directed colimit-dense full small subcategory of λ -presentable objects; then we denote by $\mathbf{Pres}_\lambda \mathcal{S}$ any set of representatives, relatively to isomorphy, for the λ -presentable objects of \mathcal{S} , and we may take $\mathcal{S}^\dagger = \mathbf{Pres}_\lambda \mathcal{S}$. If, in addition, \mathcal{S} is cocomplete, we say that \mathcal{S} is *locally λ -presentable*. A functor $\Phi: \mathcal{S} \rightarrow \mathcal{T}$ is λ -continuous (resp., λ -accessible) if it preserves λ -directed colimits (resp., it is λ -continuous and \mathcal{S} and \mathcal{T} are both λ -accessible categories). We will omit the prefix “ λ ” if it is existentially quantified, and replace it by “finitely” in case $\lambda = \omega$; for instance, “accessible” means “ λ -accessible for some λ ” and “finitely accessible” means “ ω -accessible”.

Slightly deviating from the terminology in use in Gillibert and Wehrung [10] and Wehrung [37], we define, for any functor $\Phi: \mathcal{A} \rightarrow \mathcal{B}$,

- the *range* of Φ , $\text{rng } \Phi \stackrel{\text{def}}{=} \{\Phi(A) \mid A \in \text{Ob } \mathcal{A}\}$,
- the *image* of Φ , $\text{im } \Phi = \Phi \mathcal{A} \stackrel{\text{def}}{=} \{B \in \text{Ob } \mathcal{B} \mid (\exists A \in \text{Ob } \mathcal{A})(B \cong \Phi(A))\}$.

We denote by **Set** the category of all sets with maps.

2.3. Infinitary languages. We are dealing with first-order structures over vocabularies with possibly infinite arities (cf. Adámek and Rosický [1, Ch. 5]). A vocabulary \mathbf{v} consists of a set of relation and operation symbols, with which it is occasionally identified, together with the arity map $\mathbf{s} \mapsto \text{ar}(\mathbf{s})$, usually, but not always (exceptions will occur in Sections 4 and 6) with values in ordinals (nonzero on relation symbols). We denote by \mathbf{Var} the set of all variables, by $\#\mathbf{v}$ the number of symbols in \mathbf{v} , and by \mathbf{v}_{cst} , \mathbf{v}_{ope} , and \mathbf{v}_{rel} the sub-vocabularies of \mathbf{v} consisting of the constant symbols, operation symbols, and relation symbols, respectively. Further, we set $\overline{\text{ar}}(\mathbf{s}) \stackrel{\text{def}}{=} \text{card var}(\mathbf{s})$ for every symbol \mathbf{s} of \mathbf{v} . For a cardinal λ , the vocabulary \mathbf{v} is λ -ary if $\overline{\text{ar}}(\mathbf{s}) < \lambda$ for every symbol \mathbf{s} of \mathbf{v} . Syntactical objects such as formulas or symbols of variable, relation, operation will usually⁵ be denoted in sans serif fonts (e.g., \mathbf{E} , \mathbf{x} , \mathbf{R} , \mathbf{f}), while their semantic counterparts, if defined, will be denoted in math fonts (e.g. x , R , f). We shall denote by $\text{var}(\mathbf{s})$ (resp., $\text{fvar}(\mathbf{s})$) the set of all variables (resp., free variables) occurring in a term or formula (resp., formula) \mathbf{s} .

We denote by ∞ an extra object set on top of all cardinal numbers. Whenever $\lambda \leq \kappa \leq \infty$, the infinitary language $\mathcal{L}_{\kappa\lambda}(\mathbf{v})$ is the class (or, in case $\kappa < \infty$, the set) of all formulas obtained by closing the atomic formulas of \mathbf{v} under negations (denoted in the form $\neg\mathbf{E}$), disjunctions of less than κ formulas with less than λ free variables altogether (denoted in the form $\bigvee_{i \in I} \mathbf{E}_i$), and existential quantifiers over λ -small sets of variables (denoted in the form $(\exists \mathbf{X})\mathbf{E}$). *Preformulas* are defined similarly, except that now any disjunction of preformulas is a preformula (without any restriction on the free variables). Conjunctions and universal quantifiers are defined as usual, *via* $\bigwedge = \neg \bigvee \neg$ and $\forall = \neg \exists \neg$, respectively. This choice of \neg , \bigvee , and \exists as primitive connectives affects the definition of a subformula, as for example $\exists \mathbf{x} \neg \mathbf{E}(\mathbf{x})$ is a subformula of $\forall \mathbf{x} \mathbf{E}(\mathbf{x})$. This is of direct importance in the definition of the sentences $\widehat{\mathbf{E}}$ introduced in Notation 5.1. Indexed sets of variables will be denoted in the form $\vec{x}_{(I)}$ where $\vec{x} = (x_i \mid i \in I)$ is a one-to-one map from I into the variable symbols. Quantifiers, if applied to indexed sets of variables, will be denoted in the form $\mathbf{Q} \vec{x}_{(I)}$ (or just $\mathbf{Q}\mathbf{x}$ if I is a singleton), where \mathbf{Q} is either \exists or \forall .

The universe M of a \mathbf{v} -structure $\mathbf{M} = (M, \dots)$ will be denoted by $|\mathbf{M}|$. For a sub-vocabulary \mathbf{u} of \mathbf{v} , we denote by $\mathbf{M}|_{\mathbf{u}}$ the \mathbf{u} -reduct of \mathbf{M} . The interpretation of a symbol \mathbf{s} of \mathbf{v} in \mathbf{M} will be denoted by $\mathbf{s}^{\mathbf{M}}$. Homomorphisms, embeddings, and isomorphisms of \mathbf{v} -structures are defined the usual way; for example, a \mathbf{v} -homomorphism $\varphi: \mathbf{M} \rightarrow \mathbf{N}$ is an embedding iff it is one-to-one and $\vec{x} \in \mathbf{R}^{\mathbf{M}} \Leftrightarrow \varphi \vec{x} \in \mathbf{R}^{\mathbf{N}}$ whenever $\mathbf{R} \in \mathbf{v}_{\text{rel}}$ and $\vec{x} \in \text{ar}(\mathbf{R})\mathbf{M}$.

We denote by $\mathbf{Str}(\mathbf{v})$ the category whose objects are the \mathbf{v} -structures and whose arrows are the \mathbf{v} -homomorphisms. The category $\mathbf{Str}(\mathbf{v})$ is locally presentable (cf. Adámek and Rosický [1, Theorem 5.30]). We will often need the description of λ -directed colimits, in $\mathbf{Str}(\mathbf{v})$, given by the following lemma.

Lemma 2.1 (Folklore). *Let λ be a regular cardinal and let \mathbf{v} be a vocabulary with \mathbf{v}_{ope} λ -ary. Let P be a λ -directed poset and let $\vec{\mathbf{M}} = (\mathbf{M}_p, \varphi_{p,q} \mid p \leq q \text{ in } P)$*

⁵With the exception of the unary predicate symbol \mathbf{U} involved in the definition of a relatively projective class in Definition 3.1.

be a P -indexed commutative diagram in $\mathbf{Str}(\mathbf{v})$. Then a cocone $(M, \varphi_p \mid p \in P)$ above \vec{M} is the colimit of \vec{M} iff

- (i) $M = \bigcup \{\varphi_p[M_p] \mid p \in P\}$;
- (ii) for all $p \in P$ and all $x, y \in M_p$, $\varphi_p(x) = \varphi_p(y)$ implies that there exists $q \geq p$ such that $\varphi_{p,q}(x) = \varphi_{p,q}(y)$;
- (iii) $\mathbf{R}^M = \bigcup \{\varphi_p \mathbf{R}^{M_p} \mid p \in P\}$ whenever $R \in \mathbf{v}_{\text{rel}}$. If $\overline{\text{ar}}(\mathbf{R}) < \lambda$, this amounts to verifying that for all $p \in P$ and $(x_\xi \mid \xi \in \text{ar}(\mathbf{R})) \in {}^{\text{ar}(\mathbf{R})}M_p$,

$$(\varphi_p(x_\xi) \mid \xi \in \text{ar}(\mathbf{R})) \in \mathbf{R}^M \implies (\exists q \geq p)((\varphi_{p,q}(x_\xi) \mid \xi \in \text{ar}(\mathbf{R})) \in \mathbf{R}^{M_q}).$$

3. PROJECTIVE AND RELATIVELY PROJECTIVE CLASSES

The following encompasses the classical definitions of PC and RPC classes (cf. Ebbinghaus [7]) stated for vocabularies without restrictions on arities.

Definition 3.1. Let \mathcal{L} be a logic of the form either $\mathcal{L}_{\infty\lambda}$, for an infinite cardinal λ , or $\mathcal{L}_{\infty\infty}$, and let \mathbf{v} be a vocabulary. A class \mathcal{C} of \mathbf{v} -structures is

- *abstract* if every isomorphic copy of a member of \mathcal{C} belongs to \mathcal{C} ;
- *small-abstract* if it is the class of isomorphic copies of all members of a set of \mathbf{v} -structures;
- *projective over \mathcal{L}* , in short $\text{PC}(\mathcal{L})$ -*definable* or just $\text{PC}(\mathcal{L})$, if there are a vocabulary \mathbf{w} extending \mathbf{v} and a sentence E from $\mathcal{L}(\mathbf{w})$ such that

$$\mathcal{C} = \{N \upharpoonright \mathbf{v} \mid N \in \text{Mod}_{\mathbf{w}}(E)\}; \quad (3.1)$$

- *relatively projective over \mathcal{L}* , in short $\text{RPC}(\mathcal{L})$ -*definable* or just $\text{RPC}(\mathcal{L})$, if there are a vocabulary \mathbf{w} extending \mathbf{v} , a unary predicate symbol U in \mathbf{w} , and a sentence E of $\mathcal{L}(\mathbf{w})$ such that

$$\mathcal{C} = \{U^N \upharpoonright \mathbf{v} \mid N \in \text{Mod}_{\mathbf{w}}(E), U^N \text{ is } \mathbf{v}\text{-closed}\} \quad (3.2)$$

(where “ \mathbf{v} -closed” means closed, within N , under all operations from \mathbf{v}_{ope} . Note that the closedness of U^N under \mathbf{v} cannot be expressed by an $\mathcal{L}_{\infty\lambda}$ -sentence unless \mathbf{v}_{ope} is λ -ary; in the latter case, that closedness statement can of course be incorporated into E);

- *co-PC*(\mathcal{L}) if its complement $\mathbf{Str}(\mathbf{v}) \setminus \mathcal{C}$ is $\text{PC}(\mathcal{L})$.

As discussed in Ebbinghaus [7], both concepts of PC and RPC can be defined for most logics. However, as illustrated in Oikkonen [25], the resulting concepts are sensitive to the restrictions put on the vocabularies \mathbf{v} and \mathbf{w} . That possible ambiguity is addressed in Definition 3.1 by removing all the possible restrictions on \mathbf{v} and \mathbf{w} . In particular, even in case $\mathcal{L} = \mathcal{L}_{\infty\lambda}$, \mathbf{v} may be λ -ary whereas \mathbf{w} is not. The resulting possible ambiguity in the definitions of PC and RPC will be addressed in Theorem 4.11: *if \mathbf{v} is λ -ary, then we may take \mathbf{w} λ -ary as well.*

Of course, every PC class is also RPC. It is well known that the converse fails for various logics, including classical first-order logic $\mathcal{L}_{\omega\omega}$; see for example Oikkonen [24]. We will see (cf. Corollary 8.7) that the equality $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$ holds in full generality, over any vocabulary.

The well known fact that the isomorphism class of any model is $\text{PC}(\mathcal{L}_{\infty\omega})$ is stated in Oikkonen [24, page 260]. We include that observation in the following lemma, along with proofs, for convenience.

Lemma 3.2. *The following statements holds, for any vocabulary \mathbf{v} :*

- (1) *Let λ be an infinite cardinal and let $(\mathcal{C}_i \mid i \in I)$ be a collection, indexed by a set I , of classes of \mathbf{v} -structures. If each \mathcal{C}_i is $\text{PC}(\mathcal{L}_{\infty\lambda})$, then so are $\bigcup_{i \in I} \mathcal{C}_i$ and $\bigcap_{i \in I} \mathcal{C}_i$.*
- (2) *Every small-abstract class of \mathbf{v} -structures is both $\text{PC}(\mathcal{L}_{\infty\omega})$ (via a single universal $\mathcal{L}_{\infty\omega}$ -sentence) and $\text{co-PC}(\mathcal{L}_{\infty\omega})$.*

Proof. *Ad (1).* We may assume that there are vocabularies $\mathbf{v}_i \supseteq \mathbf{v}$, pairwise intersecting in \mathbf{v} , with $\mathcal{L}_{\infty\lambda}(\mathbf{v}_i)$ -sentences \mathbf{E}_i such that each $\mathcal{C}_i = \{\mathbf{N} \upharpoonright_{\mathbf{v}} \mid \mathbf{N} \in \text{Mod}_{\mathbf{v}_i}(\mathbf{E}_i)\}$. Setting $\mathbf{v}^* \stackrel{\text{def}}{=} \bigcup_{i \in I} \mathbf{v}_i$, $\bigcup_{i \in I} \mathcal{C}_i$ is the class of \mathbf{v} -reducts of all models of the \mathbf{v}^* -sentence $\bigvee_{i \in I} \mathbf{E}_i$ and $\bigcap_{i \in I} \mathcal{C}_i$ is the class of \mathbf{v} -reducts of all models of $\bigwedge_{i \in I} \mathbf{E}_i$.

Ad (2). By (1) (and its proof) it suffices to settle the case of the isomorphism class \mathcal{C} of a single \mathbf{v} -structure \mathbf{M} . Letting \mathbf{C} be a set of constant symbols, outside \mathbf{v} , indexing \mathbf{M} , \mathcal{C} is the class of \mathbf{v} -reducts of the universal $\mathcal{L}_{\infty\omega}(\mathbf{v} \cup \mathbf{C})$ -sentence obtained as the conjunction of the diagram of \mathbf{M} (each element of \mathbf{M} being indexed by the corresponding element of \mathbf{C}) and the universal $\mathcal{L}_{\infty\omega}(\mathbf{C})$ -sentence stating that every element belongs to \mathbf{C} . Thus \mathcal{C} is $\text{PC}(\mathcal{L}_{\infty\omega})$ -definable.

Denote by \leq a binary relation symbol. There are $\mathcal{L}_{\infty\omega}(\leq)$ -sentences \mathbf{E}_α , ranging over all ordinals α , such that a structure (P, \leq) satisfies \mathbf{E}_α iff \leq is a linear order and (α, \leq) embeds into (P, \leq) as a lower subset (cf. Dickmann [6, p. 319]). It follows that for every cardinal κ , the class of all \mathbf{v} -structures of cardinality at least κ is the class of all \mathbf{v} -reducts of the $\mathcal{L}_{\infty\omega}(\mathbf{v} \sqcup \{\leq\})$ -sentence \mathbf{E}_κ ; whence it is $\text{PC}(\mathcal{L}_{\infty\omega})$.

Letting $\kappa > \text{card } \mathbf{M}$, the complement of \mathcal{C} is the union of the class of all \mathbf{v} -structures of cardinality at least κ (which is $\text{PC}(\mathcal{L}_{\infty\omega})$ as observed in the paragraph above) and the small-abstract class of all κ -small structures outside \mathcal{C} (which is $\text{PC}(\mathcal{L}_{\infty\omega})$ by the above); whence it is $\text{PC}(\mathcal{L}_{\infty\omega})$. Now apply (1). \square

4. A FIRST CASE OF CONJUGACY: PARTIAL λ -ARY TRUNCATING

For an infinite cardinal λ and a vocabulary \mathbf{v} , any $\text{PC}(\mathcal{L}_{\infty\lambda})$ class of \mathbf{v} -structures is the class of \mathbf{v} -reducts of all models of some $\mathcal{L}_{\infty\lambda}$ -sentence in a larger vocabulary \mathbf{w} (cf. Definition 3.1). No restriction is set on the arities of the symbols in either \mathbf{v} or \mathbf{w} *a priori*. We shall verify in this section that \mathbf{w} may be taken in such a way that $\mathbf{w} \setminus \mathbf{v}$ is λ -ary; a similar result holds for RPC (cf. Theorem 4.11).

Our basic idea is quite simple: since all atomic formulas of $\mathcal{L}_{\infty\lambda}$ have less than λ variables, we may replace every \mathbf{w} -term \mathbf{t} by a term $\tilde{\mathbf{t}}$ in a new λ -ary vocabulary $\tilde{\mathbf{w}}$; and similarly, every atomic formula in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ by an atomic formula in $\tilde{\mathbf{w}}$. This translation process should not affect symbols from \mathbf{v} . It usually introduces more symbols than the original \mathbf{w} has. In order to be able to translate back $\tilde{\mathbf{w}}$ -structures to \mathbf{w} -structures, a ‘‘coherence sentence’’ \mathbf{G} needs to be introduced (cf. Notation 4.4), that needs to be satisfied by all $\tilde{\mathbf{w}}$ -structures in question.

Notation 4.1. For a vocabulary \mathbf{w} and an infinite cardinal λ , we shall denote by $\mathbf{T}_\lambda(\mathbf{w})$ (resp., $\mathbf{A}_\lambda(\mathbf{w})$) the set of all terms (resp., atomic formulas) of \mathbf{w} on less than λ variables. Hence, $\mathbf{A}_\lambda(\mathbf{w})$ is the set of all atomic formulas in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$.

We shall now describe an inductive procedure that, given vocabularies $\mathbf{v} \subseteq \mathbf{w}$ and an infinite cardinal λ , produces a vocabulary $\tilde{\mathbf{w}} = \text{col}_\lambda(\mathbf{w}, \mathbf{v})$, containing $\mathbf{v} \cup \mathbf{w}_{\text{cst}}$ and collapsing $\mathbf{w} \setminus (\mathbf{v} \cup \mathbf{w}_{\text{cst}})$ to a new λ -ary vocabulary, usually with more symbols than \mathbf{w} . The new symbols will be denoted in the form \mathbf{s} or \mathbf{E} ; their arities will be sets of variables (as opposed to mere ordinals).

Definition 4.2. We set $\tilde{s} \stackrel{\text{def}}{=} s$ whenever s is either a variable or a constant symbol from \mathbf{w} . In case $s = f(s_\xi \mid \xi \in \text{ar}(f)) \in T_\lambda(\mathbf{w})$, we define a term \tilde{s} , in a possibly larger vocabulary, by

$$\tilde{s} \stackrel{\text{def}}{=} \begin{cases} f(\tilde{s}_\xi \mid \xi \in \text{ar}(f)), & \text{if } f \in \mathbf{v}, \\ \dot{s}(u \mid u \in \text{var}(s)) & \text{(usually written as } \dot{s}), \text{ otherwise,} \end{cases} \quad (4.1)$$

where, in the second case (i.e., $f \notin \mathbf{v}$), \dot{s} is a new operation symbol with arity $\text{var}(s)$.

Let $E \in A_\lambda(\mathbf{w})$. If E has the form $s = t$, for $s, t \in T_\lambda(\mathbf{w})$, then we define \tilde{E} as $\tilde{s} = \tilde{t}$. If E does not have that form (we will then say that E is *relational*), then $E = R(s_\xi \mid \xi \in \text{ar}(R))$, for $R \in \mathbf{w}_{\text{rel}}$ and terms s_ξ of \mathbf{w} with less than λ variables altogether; we then define an atomic formula \tilde{E} , with the same variables as E but in a possibly larger vocabulary, by

$$\tilde{E} \stackrel{\text{def}}{=} \begin{cases} R(\tilde{s}_\xi \mid \xi \in \text{ar}(R)), & \text{if } R \in \mathbf{v}, \\ \dot{E}(u \mid u \in \text{var}(E)) & \text{(usually written as } \dot{E}), \text{ otherwise,} \end{cases} \quad (4.2)$$

where, in the second case (i.e., $R \notin \mathbf{v}$), \dot{E} is a new relation symbol with arity $\text{var}(E)$.

The common vocabulary $\tilde{\mathbf{w}} = \text{col}_\lambda(\mathbf{w}, \mathbf{v})$ for all those syntactical objects is

$$\mathbf{v} \cup \{\dot{s} \mid s \text{ as in the second case of (4.1)}\} \cup \{\dot{E} \mid E \text{ as in the second case of (4.2)}\}.$$

In particular, $\mathbf{w} \cap \tilde{\mathbf{w}} = \mathbf{v} \cup \mathbf{w}_{\text{cst}}$.

Lemma 4.3. *In the context above, $\text{var}(\tilde{s}) = \text{var}(s)$ and $\text{var}(\tilde{E}) = \text{var}(E)$. Moreover, $\tilde{\mathbf{w}} \setminus \mathbf{v}$ is λ -ary.*

Notation 4.4 (Coherence sentence). Denote by \mathbf{G} the conjunction of all sentences of the following form:

$$\left(\forall_{(\text{var}(s))} \vec{x} \right) \left(\forall_{(\text{var}(t))} \vec{y} \right) \left(\bigwedge_{\xi \in \text{ar}(f)} (\tilde{s}_\xi(\vec{x}) = \tilde{t}_\xi(\vec{y})) \Rightarrow \tilde{s}(\vec{x}) = \tilde{t}(\vec{y}) \right), \quad (4.3)$$

ranging over all pairs (s, t) of terms in $T_\lambda(\mathbf{w})$ of the form $s = f(s_\xi \mid \xi \in \text{ar}(f))$ and $t = f(t_\xi \mid \xi \in \text{ar}(f))$ (i.e., with the same underlying f), and

$$\left(\forall_{(\text{var}(E))} \vec{x} \right) \left(\forall_{(\text{var}(F))} \vec{y} \right) \left[\left(\tilde{E}(\vec{x}) \wedge \bigwedge_{\xi \in \text{ar}(f)} (\tilde{s}_\xi(\vec{x}) = \tilde{t}_\xi(\vec{y})) \right) \Rightarrow \tilde{F}(\vec{y}) \right], \quad (4.4)$$

ranging over all pairs (E, F) of atomic formulas in $A_\lambda(\mathbf{w})$ of the form $E = R(s_\xi \mid \xi \in \text{ar}(R))$ and $F = R(t_\xi \mid \xi \in \text{ar}(R))$ (i.e., with the same underlying R). By construction, \mathbf{G} is an $\mathcal{L}_{\infty\lambda}(\tilde{\mathbf{w}})$ -sentence.

Note that in the above, dummy variables are allowed for: for example, $\tilde{s}_\xi(\vec{x})$ should be $\tilde{s}_\xi(\vec{x} \upharpoonright_{\text{var}(s_\xi)})$, and so on. To avoid clutter, we keep the former notation.

We shall now establish a correspondence between \mathbf{w} -structures and $\tilde{\mathbf{w}}$ -structures.

Definition 4.5. Let \mathbf{M} be a \mathbf{w} -structure and let \mathbf{N} be a $\tilde{\mathbf{w}}$ -structure. We say that (\mathbf{M}, \mathbf{N}) is a *conjugate pair*, or, equivalently, that \mathbf{M} is *left conjugate to* \mathbf{N} , respectively that \mathbf{N} is *right conjugate to* \mathbf{M} , if the following statements hold:

- (1) $\mathbf{M} \upharpoonright_{\mathbf{v} \cup \mathbf{w}_{\text{cst}}} = \mathbf{N} \upharpoonright_{\mathbf{v} \cup \mathbf{w}_{\text{cst}}}$;
- (2) For every term $s = f(s_\xi \mid \xi \in \text{ar}(f)) \in T_\lambda(\mathbf{w})$ with $f \notin \mathbf{v}$,

$$(\forall \vec{a} \in {}^{\text{var}(s)}M) (\dot{s}^{\mathbf{N}}(\vec{a}) = s^{\mathbf{M}}(\vec{a})). \quad (4.5)$$

(3) For every atomic formula $E = R(s_\xi \mid \xi \in \text{ar}(R)) \in A_\lambda(\mathbf{w})$ with $R \notin \mathbf{v}$,

$$(\forall \vec{a} \in \text{var}^{(E)}M)(\vec{a} \in \dot{E}^N \iff \mathbf{M} \models E(\vec{a})). \quad (4.6)$$

Lemma 4.6. *The following statements hold, for any conjugate pair (\mathbf{M}, \mathbf{N}) :*

- (1) *The equation $s^{\mathbf{M}}(\vec{a}) = \tilde{s}^{\mathbf{N}}(\vec{a})$ holds, for all $s \in T_\lambda(\mathbf{w})$ and all $\vec{a} \in \text{var}^{(s)}M$.*
- (2) *The equivalence $\mathbf{N} \models \tilde{E}(\vec{a}) \iff \mathbf{M} \models E(\vec{a})$ holds, for all $E \in A_\lambda(\mathbf{w})$ and all $\vec{a} \in \text{var}^{(E)}M$.*
- (3) *$\mathbf{N} \models G$ (cf. Notation 4.4).*

Proof. *Ad (1).* We argue by induction on the term s . The result is obvious if s is either a constant (cf. Definition 4.5(1)) or a variable. Let $s = f(s_\xi \mid \xi \in \text{ar}(f))$. If $f \in \mathbf{v}$, then $f^{\mathbf{M}} = f^{\mathbf{N}}$ (cf. Definition 4.5(1)) so

$$\begin{aligned} \tilde{s}^{\mathbf{N}}(\vec{a}) &= f^{\mathbf{M}}(\tilde{s}_\xi^{\mathbf{N}}(\vec{a}) \mid \xi \in \text{ar}(f)) && \text{(cf. (4.1))} \\ &= f^{\mathbf{M}}(s_\xi^{\mathbf{M}}(\vec{a}) \mid \xi \in \text{ar}(f)) && \text{(by induction hypothesis)} \\ &= s^{\mathbf{M}}(\vec{a}). \end{aligned}$$

If $f \notin \mathbf{v}$, then a direct application of (4.5) yields the relation $\tilde{s}^{\mathbf{N}}(\vec{a}) = s^{\mathbf{M}}(\vec{a})$.

Ad (2). By (1) above, the given statement is obvious in case E has the form $s = t$. Now let $E = R(s_\xi \mid \xi \in \text{ar}(R))$. If $R \in \mathbf{v}$, then $R^{\mathbf{M}} = R^{\mathbf{N}}$ (cf. Definition 4.5(1)) so

$$\begin{aligned} \mathbf{N} \models \tilde{E}(\vec{a}) &\Leftrightarrow (\tilde{s}_\xi^{\mathbf{N}}(\vec{a}) \mid \xi \in \text{ar}(R)) \in R^{\mathbf{M}} && \text{(cf. (4.2))} \\ &\Leftrightarrow (s_\xi^{\mathbf{M}}(\vec{a}) \mid \xi \in \text{ar}(R)) \in R^{\mathbf{M}} && \text{(by (1) above)} \\ &\Leftrightarrow \mathbf{M} \models E(\vec{a}). \end{aligned}$$

If $R \notin \mathbf{v}$, then a direct application of (4.6) yields that $\mathbf{N} \models \tilde{E}(\vec{a})$ iff $\mathbf{M} \models E(\vec{a})$.

Ad (3). We start with the sentences of the form (4.3). In that context, let $\vec{a} \in \text{var}^{(s)}M$ and $\vec{b} \in \text{var}^{(t)}M$ such that $\tilde{s}_\xi^{\mathbf{N}}(\vec{a}) = \tilde{t}_\xi^{\mathbf{N}}(\vec{b})$ whenever $\xi \in \text{ar}(f)$. By (1) above, this means that $s_\xi^{\mathbf{M}}(\vec{a}) = t_\xi^{\mathbf{M}}(\vec{b})$ whenever $\xi \in \text{ar}(f)$. By applying $f^{\mathbf{M}}$, we get $s^{\mathbf{M}}(\vec{a}) = t^{\mathbf{M}}(\vec{b})$, thus, by (1) above, $\tilde{s}^{\mathbf{N}}(\vec{a}) = \tilde{t}^{\mathbf{N}}(\vec{b})$, as required.

We now deal with sentences of the form (4.4). In that context, let $\vec{a} \in \text{var}^{(E)}M$ and $\vec{b} \in \text{var}^{(F)}M$ such that $\mathbf{N} \models \tilde{E}(\vec{a})$ whereas $\tilde{s}_\xi^{\mathbf{N}}(\vec{a}) = \tilde{t}_\xi^{\mathbf{N}}(\vec{b})$ whenever $\xi \in \text{ar}(R)$. By (1) and (2) above, this means that $\mathbf{M} \models E(\vec{a})$ whereas $s_\xi^{\mathbf{M}}(\vec{a}) = t_\xi^{\mathbf{M}}(\vec{b})$ whenever $\xi \in \text{ar}(R)$. Therefore, $\mathbf{M} \models F(\vec{b})$, thus, by (2) above, $\mathbf{N} \models \tilde{F}(\vec{b})$, as required. \square

Lemma 4.7. *Every \mathbf{w} -structure is left conjugate to a unique $\tilde{\mathbf{w}}$ -structure.*

Proof. The missing interpretations $\dot{s}^{\mathbf{N}}$ and $\dot{E}^{\mathbf{N}}$ are necessarily given by (4.5) and (4.6), respectively. They define the $\tilde{\mathbf{w}}$ -structure \mathbf{N} uniquely. \square

Lemma 4.8. *Every $\tilde{\mathbf{w}}$ -structure \mathbf{N} , satisfying the statement \mathbf{G} , is right conjugate to some \mathbf{w} -structure \mathbf{M} .*

Proof. We need to define the missing interpretations in \mathbf{M} of all symbols in $\mathbf{w} \setminus (\mathbf{v} \cup \mathbf{w}_{\text{cst}})$, in such a way that all instances of (4.5) and (4.6) hold. We start with operation symbols in $\mathbf{w} \setminus (\mathbf{v} \cup \mathbf{w}_{\text{cst}})$.

Claim 1. *For every $f \in \mathbf{w}_{\text{ope}}$, there exists a map $f^{\mathbf{M}}: \text{ar}(f)M \rightarrow M$, equal to $f^{\mathbf{N}}$ in case $f \in \mathbf{v}_{\text{ope}}$, such that for every term of the form $s = f(s_\xi \mid \xi \in \text{ar}(f))$ in $T_\lambda(\mathbf{w})$ and every $\vec{a} \in \text{var}^{(s)}M$,*

$$f^{\mathbf{M}}(\tilde{s}_\xi^{\mathbf{N}}(\vec{a}) \mid \xi \in \text{ar}(f)) = \tilde{s}^{\mathbf{N}}(\vec{a}). \quad (4.7)$$

Proof of Claim. In order to prove that (4.7) correctly defines a partial function f^M , we need to verify that for all terms of the form $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ and $\mathbf{t} = f(\mathbf{t}_\xi \mid \xi \in \text{ar}(f))$ in $\mathbf{T}_\lambda(\mathbf{w})$, all $\vec{a} \in \text{var}(\mathbf{s})M$, and all $\vec{b} \in \text{var}(\mathbf{t})M$, the conjunction over all $\xi \in \text{ar}(f)$ of $\tilde{\mathbf{s}}_\xi^N(\vec{a}) = \tilde{\mathbf{t}}_\xi^N(\vec{b})$ implies $\tilde{\mathbf{s}}^N(\vec{a}) = \tilde{\mathbf{t}}^N(\vec{b})$. This follows from our assumption that N satisfies \mathbf{G} .

Now suppose that $f \in \mathbf{v}_{\text{ope}}$. Then $\tilde{\mathbf{s}} = f(\tilde{\mathbf{s}}_\xi \mid \xi \in \text{ar}(f))$ (cf. (4.1)), thus

$$\tilde{\mathbf{s}}^N(\vec{a}) = f^N(\tilde{\mathbf{s}}_\xi^N(\vec{a}) \mid \xi \in \text{ar}(f)). \quad \square \text{ Claim 1.}$$

This yields a $(\mathbf{v} \cup \mathbf{w}_{\text{ope}})$ -structure, specializing the intended M thus which we will also temporarily denote by M , satisfying all equations (4.7).

Claim 2. *The relation $\mathbf{s}^M(\vec{a}) = \tilde{\mathbf{s}}^N(\vec{a})$ holds, for every $\mathbf{s} \in \mathbf{T}_\lambda(\mathbf{w})$ and every $\vec{a} \in \text{var}(\mathbf{s})M$.*

Proof of Claim. We argue by induction on the term \mathbf{s} . The given conclusion is trivial if \mathbf{s} is either a variable or a constant. Let $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$. Then

$$\begin{aligned} \mathbf{s}^M(\vec{a}) &= f^M(\mathbf{s}_\xi^M(\vec{a}) \mid \xi \in \text{ar}(f)) \\ &= f^M(\tilde{\mathbf{s}}_\xi^N(\vec{a}) \mid \xi \in \text{ar}(f)) && \text{(by our induction hypothesis)} \\ &= \tilde{\mathbf{s}}^N(\vec{a}) && \text{(cf. (4.7)).} \end{aligned} \quad \square \text{ Claim 2.}$$

Now let $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ with $f \notin \mathbf{v}$. Using Claim 2, we get $\mathbf{s}^M(\vec{a}) = \tilde{\mathbf{s}}^N(\vec{a})$. By (4.1), $\tilde{\mathbf{s}}^N(\vec{a}) = \dot{\mathbf{s}}^N(\vec{a})$. Therefore, all statements of the form (4.5) hold.

It remains to construct interpretations R^M , for $R \in \mathbf{w}_{\text{rel}} \setminus \mathbf{v}$, in such a way that all statements of the form (4.6) hold. This means that whenever $E = R(\mathbf{s}_\xi \mid \xi \in \text{ar}(R))$ in $\mathbf{T}_\lambda(\mathbf{w})$ and $\vec{a} \in \text{var}(E)M$, the equivalence

$$(\mathbf{s}_\xi^M(\vec{a}) \mid \xi \in \text{ar}(R)) \in R^M \iff \vec{a} \in \dot{E}^N$$

holds. In order for those to yield a consistent definition of R^M , we need to verify that whenever $F = R(\mathbf{t}_\xi \mid \xi \in \text{ar}(R)) \in \mathbf{A}_\lambda(\mathbf{w})$ and $\vec{b} \in \text{var}(F)M$ such that

$$\mathbf{s}_\xi^M(\vec{a}) = \mathbf{t}_\xi^M(\vec{b}) \quad \text{whenever } \xi \in \text{ar}(R), \quad (4.8)$$

the equivalence $\vec{a} \in \dot{E}^N \iff \vec{b} \in \dot{F}^N$ — in other words (cf. (4.2)), $N \models (\tilde{E}(\vec{a}) \iff \tilde{F}(\vec{b}))$ — holds. Now from (4.8) and Claim 2 it follows that

$$\tilde{\mathbf{s}}_\xi^N(\vec{a}) = \tilde{\mathbf{t}}_\xi^N(\vec{b}) \quad \text{whenever } \xi \in \text{ar}(R). \quad (4.9)$$

Since N satisfies \mathbf{G} , it also satisfies the desired $\tilde{E}(\vec{a}) \iff \tilde{F}(\vec{b})$. \square

Notation 4.9. We extend the definition of \tilde{E} (cf. (4.2)) from all formulas in $\mathbf{A}_\lambda(\mathbf{w})$ to all formulas in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$, by setting $E = \neg F \Rightarrow \tilde{E} = \neg\tilde{F}$; $E = \bigvee_{i \in I} E_i \Rightarrow \tilde{E} = \bigvee_{i \in I} \tilde{E}_i$; $E = (\exists X)F \Rightarrow \tilde{E} = (\exists X)\tilde{F}$.

A straightforward induction argument, based on Lemma 4.6(2), then yields the following.

Lemma 4.10. *Let (M, N) be a conjugate pair. Then the equivalence $N \models \tilde{E}(\vec{a}) \iff M \models E(\vec{a})$ holds, for all $E \in \mathcal{L}_{\infty\lambda}(\mathbf{w})$ and all $\vec{a} \in \text{fvar}(E)M$.*

Theorem 4.11. *Let \mathbf{v} be a vocabulary and let λ be an infinite cardinal. Then a class \mathcal{C} of \mathbf{v} -structures is $\text{PC}(\mathcal{L}_{\infty\lambda})$ (resp., $\text{RPC}(\mathcal{L}_{\infty\lambda})$) iff there are a vocabulary \mathbf{w} containing \mathbf{v} (resp., $\mathbf{v} \cup \{\mathbf{U}\}$ for a unary predicate \mathbf{U}), such that $\mathbf{w} \setminus \mathbf{v}$ is λ -ary, together with a sentence \mathbf{E} in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$, such that $\mathcal{C} = \{\mathbf{N} \upharpoonright \mathbf{v} \mid \mathbf{N} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E})\}$ (resp., $\mathcal{C} = \{\mathbf{U}^{\mathbf{N}} \upharpoonright \mathbf{v} \mid \mathbf{N} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E}), \mathbf{U}^{\mathbf{N}} \text{ v-closed}\}$). In particular, if \mathbf{v} is λ -ary, then so is \mathbf{w} .*

Proof. We present the argument for RPC; the argument for PC is similar. By definition, there are a unary predicate symbol \mathbf{U} , a vocabulary $\mathbf{w} \supseteq \mathbf{v} \cup \{\mathbf{U}\}$, and a sentence \mathbf{E} in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ such that

$$\mathcal{C} = \{\mathbf{U}^{\mathbf{M}} \upharpoonright \mathbf{v} \mid \mathbf{M} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E}), \mathbf{U}^{\mathbf{M}} \text{ is } \mathbf{v}\text{-closed}\}$$

Set $\tilde{\mathbf{w}} \stackrel{\text{def}}{=} \text{col}_{\lambda}(\mathbf{w}, \mathbf{v} \cup \{\mathbf{U}\})$. Since $\tilde{\mathbf{w}} \setminus \mathbf{v}$ is λ -ary, it thus suffices to prove that $\mathcal{C} = \mathcal{C}'$ where

$$\mathcal{C}' \stackrel{\text{def}}{=} \{\mathbf{U}^{\mathbf{N}} \upharpoonright \mathbf{v} \mid \mathbf{N} \in \mathbf{Mod}_{\tilde{\mathbf{w}}}(\tilde{\mathbf{E}} \wedge \mathbf{G}), \mathbf{U}^{\mathbf{N}} \text{ is } \mathbf{v}\text{-closed}\}$$

Let $\mathbf{A} \in \mathcal{C}$. By definition, $\mathbf{A} = \mathbf{U}^{\mathbf{M}} \upharpoonright \mathbf{v}$ for some $\mathbf{M} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E})$ such that $\mathbf{U}^{\mathbf{M}}$ is \mathbf{v} -closed. By Lemma 4.7, \mathbf{M} has a right conjugate $\mathbf{N} \in \mathbf{Str}(\tilde{\mathbf{w}})$. By Lemmas 4.6(3) and 4.10, \mathbf{N} satisfies $\tilde{\mathbf{E}} \wedge \mathbf{G}$; whence $\mathbf{A} = \mathbf{U}^{\mathbf{M}} \upharpoonright \mathbf{v} = \mathbf{U}^{\mathbf{N}} \upharpoonright \mathbf{v} \in \mathcal{C}'$. Conversely, let $\mathbf{A} \in \mathcal{C}'$. Then $\mathbf{A} = \mathbf{U}^{\mathbf{N}} \upharpoonright \mathbf{v}$ for some $\mathbf{N} \in \mathbf{Mod}_{\tilde{\mathbf{w}}}(\tilde{\mathbf{E}} \wedge \mathbf{G})$ such that $\mathbf{U}^{\mathbf{N}}$ is \mathbf{v} -closed. By Lemma 4.8, \mathbf{N} has a left conjugate \mathbf{M} . By Lemma 4.10 and since $\mathbf{N} \models \tilde{\mathbf{E}}$, we get $\mathbf{M} \models \mathbf{E}$; whence $\mathbf{A} = \mathbf{U}^{\mathbf{N}} \upharpoonright \mathbf{v} = \mathbf{U}^{\mathbf{M}} \upharpoonright \mathbf{v} \in \mathcal{C}$. \square

5. AN $\mathcal{L}_{\infty\lambda}$ VERSION OF THE SKOLEM NORMAL FORM

In this section we shall verify that the $\text{PC}(\mathcal{L}_{\infty\lambda})$ (or $\text{RPC}(\mathcal{L}_{\infty\lambda})$) character of a given class of \mathbf{v} -structures can be witnessed by a conjunction of universal $\mathcal{L}_{\infty\lambda}$ -sentences. This will lead, under the assumption that \mathbf{v}_{ope} be λ -ary, to a representation of any $\text{RPC}(\mathcal{L}_{\infty\lambda})$ class as the image of a λ -continuous functor on a λ -accessible category (cf. Theorem 5.5).

Up to Lemma 6.7 we shall fix an infinite cardinal λ and a vocabulary \mathbf{w} .

Notation 5.1. For any formula $\mathbf{E} \in \mathcal{L}_{\infty\lambda}(\mathbf{w})$, we construct inductively a quantifier-free formula $\bar{\mathbf{E}} \in \mathcal{L}_{\infty\lambda}$, over a vocabulary containing \mathbf{w} , as follows. If \mathbf{E} is atomic, let $\bar{\mathbf{E}} \stackrel{\text{def}}{=} \mathbf{E}$. If $\mathbf{E} = \bigvee_{i \in I} \mathbf{E}_i$, let $\bar{\mathbf{E}} = \bigvee_{i \in I} \bar{\mathbf{E}}_i$. If $\mathbf{E} = \neg \mathbf{F}$, let $\bar{\mathbf{E}} = \neg \bar{\mathbf{F}}$. If $\mathbf{E}(\vec{x})$ is $(\exists \vec{y}) \mathbf{F}(\vec{x}, \vec{y})$ (call such formulas \exists -prefixed), let $\bar{\mathbf{E}}(\vec{x}) \stackrel{\text{def}}{=} \bar{\mathbf{F}}(\vec{x}, \vec{f}_{\mathbf{E}}(\vec{x}))$, where $\vec{f}_{\mathbf{E}}(\vec{x}) \stackrel{\text{def}}{=} (\mathbf{f}_{\mathbf{E},j}(\vec{x}) \mid j \in J)$ with new operation symbols $\mathbf{f}_{\mathbf{E},j}$ (meant as Skolem functions)⁶. Observe that $\bar{\mathbf{E}}$ is quantifier-free, with the same free variables as \mathbf{E} . Denote by $\mathbf{w}[\mathbf{E}]$ its vocabulary.

For any formula $\mathbf{E}(\vec{x}) = (\exists \vec{y}) \mathbf{F}(\vec{x}, \vec{y})$, denote by $\text{sk}_{\mathbf{E}}$ the following $\mathcal{L}_{\infty\lambda}$ -sentence:

$$(\forall \vec{x})(\forall \vec{y}) \left(\bar{\mathbf{F}}(\vec{x}, \vec{y}) \Rightarrow \bar{\mathbf{F}}(\vec{x}, \vec{f}_{\mathbf{E}}(\vec{x})) \right). \quad (\text{sk}_{\mathbf{E}})$$

The conjunction $\hat{\mathbf{E}}$ of the $\text{sk}_{\mathbf{F}}$, over all \exists -prefixed subformulas \mathbf{F} of \mathbf{E} , is thus a conjunction of universal sentences from $\mathcal{L}_{\infty\lambda}(\mathbf{w}[\mathbf{E}])$.

Lemma 5.2. *Every \mathbf{w} -structure \mathbf{M} expands to some $\mathbf{N} \in \mathbf{Mod}_{\mathbf{w}[\mathbf{E}]}(\hat{\mathbf{E}})$.*

⁶Strictly speaking, $I = \text{fvar}(\mathbf{E})$ and $J = \text{fvar}(\mathbf{F}) \setminus I$. For better readability, we keep the notation involving indexed sets of variables.

Proof. A standard existence argument for Skolem expansions. For every ordinal α , let us denote by \mathcal{E}_α the set of all subformulas of \mathbf{E} with quantifier rank $\leq \alpha$, by \mathcal{E}'_α the set of all \exists -prefixed members of \mathcal{E}_α , and by \mathbf{w}_α the union of \mathbf{w} with the vocabulary of all $\bar{\mathbf{F}}$ with $\mathbf{F} \in \mathcal{E}_\alpha$ (equivalently, all $\bar{\mathbf{F}}$ with $\mathbf{F} \in \mathcal{E}'_\alpha$). Denoting by θ the quantifier rank of \mathbf{E} , we construct inductively a $(\theta + 1)$ -indexed sequence $(\mathbf{N}_\alpha \mid \alpha \leq \theta)$ of structures, with $\mathbf{N}_0 = \mathbf{M}$, $\mathbf{N}_\alpha \in \mathbf{Str}(\mathbf{w}_\alpha)$, $\alpha \leq \beta \Rightarrow \mathbf{N}_\alpha = \mathbf{N}_\beta \upharpoonright_{\mathbf{w}_\alpha}$, and each $\mathbf{N}_\alpha \models \widehat{\mathbf{F}}$ whenever $\mathbf{F} \in \mathcal{E}_\alpha$. Let $\alpha > 0$ and suppose $(\mathbf{N}_\xi \mid \xi < \alpha)$ constructed. For each subformula $\mathbf{F}(\vec{x}) = (\exists \vec{y}) \mathbf{G}(\vec{x}, \vec{y})$ of \mathbf{E} with quantifier rank α and each $\vec{a} \in {}^I M$, we define $\bar{f}_F^{\mathbf{N}_\alpha}(\vec{a})$ as any $\vec{b} \in {}^J M$ such that $\bigcup_{\xi < \alpha} \mathbf{N}_\xi$ satisfies $\bar{\mathbf{G}}(\vec{a}, \vec{b})$ if such a \vec{b} exists, and any element of ${}^J M$ otherwise. By construction, $\mathbf{N}_\alpha \models \text{sk}_F$ whenever $\mathbf{F} \in \mathcal{E}'_\alpha$. Set $\mathbf{N} \stackrel{\text{def}}{=} \mathbf{N}_\theta$. \square

Lemma 5.3. *Let $\mathbf{E} \in \mathcal{L}_{\infty\lambda}(\mathbf{w})$ and let \mathbf{F} be a subformula of \mathbf{E} . Then every $\mathbf{w}[\mathbf{E}]$ -structure satisfies⁷ the implication $\widehat{\mathbf{E}} \Rightarrow (\mathbf{F} \Leftrightarrow \bar{\mathbf{F}})$.*

Proof. By induction on \mathbf{F} . The atomic case and the \bigvee, \neg steps are all straightforward (we use the observation that $\models (\widehat{\mathbf{G}} \Rightarrow \widehat{\mathbf{F}})$ whenever \mathbf{F} is a subformula of \mathbf{G}). Let $\mathbf{F}(\vec{x})$ be $(\exists \vec{y}) \mathbf{G}(\vec{x}, \vec{y})$, let $\mathbf{M} \in \mathbf{Mod}_{\mathbf{w}[\mathbf{E}]}(\widehat{\mathbf{E}})$, and let $\vec{a} \in {}^I M$.

Suppose first that $\mathbf{M} \models \bar{\mathbf{F}}(\vec{a})$. Since $\mathbf{M} \models \text{sk}_F$, $\mathbf{M} \models \bar{\mathbf{G}}(\vec{a}, \vec{b})$ where $\vec{b} \stackrel{\text{def}}{=} \bar{f}_F^{\mathbf{M}}(\vec{a})$. By induction hypothesis, $\mathbf{M} \models \mathbf{G}(\vec{a}, \vec{b})$; whence $\mathbf{M} \models \mathbf{F}(\vec{a})$.

Suppose, conversely, that $\mathbf{M} \models \mathbf{F}(\vec{a})$, that is, $\mathbf{M} \models \mathbf{G}(\vec{a}, \vec{b})$ for some \vec{b} . By induction hypothesis, $\mathbf{M} \models \bar{\mathbf{G}}(\vec{a}, \vec{b})$. Since $\mathbf{M} \models \text{sk}_F$, it follows that $\mathbf{M} \models \bar{\mathbf{G}}(\vec{a}, \bar{f}_F^{\mathbf{M}}(\vec{a}))$; that is, $\mathbf{M} \models \bar{\mathbf{F}}(\vec{a})$. \square

Proposition 5.4. *Let \mathbf{v} be a vocabulary and let λ be an infinite cardinal. Then a class \mathcal{C} of \mathbf{v} -structures is $\text{RPC}(\mathcal{L}_{\infty\lambda})$ (resp., $\text{PC}(\mathcal{L}_{\infty\lambda})$) iff there are a vocabulary \mathbf{w} containing \mathbf{v} , with $\mathbf{w} \setminus \mathbf{v}$ λ -ary and a unary predicate symbol \mathbf{U} for the RPC case, together with a conjunction \mathbf{E} of universal $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentences, such that*

$$\mathcal{C} = \{\mathbf{U}^M \upharpoonright_{\mathbf{v}} \mid \mathbf{M} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E}), \mathbf{U}^M \text{ } \mathbf{v}\text{-closed}\}, \quad (5.1)$$

(resp., $\mathcal{C} = \{\mathbf{M} \upharpoonright_{\mathbf{v}} \mid \mathbf{M} \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E})\}$).

Proof. We deal with the statement on RPC; the argument for PC is similar. By Theorem 4.11, there exists a vocabulary \mathbf{w} containing $\mathbf{v} \cup \{\mathbf{U}\}$, with $\mathbf{w} \setminus \mathbf{v}$ λ -ary, together with a sentence \mathbf{E} in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$, such that (5.1) holds. Note that $\mathbf{w}[\mathbf{E}] \setminus \mathbf{w}$ consists only of operations with λ -small arities (viz. the $f_{F,j}$). By Lemma 5.2, every \mathbf{w} -structure \mathbf{M} expands to some $\mathbf{N} \in \mathbf{Mod}_{\mathbf{w}[\mathbf{E}]}(\widehat{\mathbf{E}})$. Further, by Lemma 5.3, $\mathbf{N} \models \bar{\mathbf{E}}$ iff $\mathbf{N} \models \mathbf{E}$, equivalently $\mathbf{M} \models \mathbf{E}$. Therefore,

$$\mathcal{C} = \{\mathbf{U}^N \upharpoonright_{\mathbf{v}} \mid \mathbf{N} \in \mathbf{Mod}_{\mathbf{w}[\mathbf{E}]}(\widehat{\mathbf{E}} \wedge \bar{\mathbf{E}}), \mathbf{U}^N \text{ } \mathbf{v}\text{-closed}\}.$$

Observe that $\bar{\mathbf{E}}$ is universal, whereas $\widehat{\mathbf{E}}$ is a conjunction of universal sentences. \square

Theorem 5.5. *Let λ be a regular cardinal, let \mathbf{v} be a vocabulary with \mathbf{v}_{ope} λ -ary, and let \mathcal{C} be an $\text{RPC}(\mathcal{L}_{\infty\lambda})$ class of \mathbf{v} -structures. Then there are a λ -accessible category \mathcal{S} and a λ -continuous functor $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ such that $\mathcal{C} = \text{im } \Phi = \text{rng } \Phi$.*

⁷With the usual convention that $\models \mathbf{F}(\vec{x})$ means $\models (\forall \vec{x})\mathbf{F}(\vec{x})$.

Moreover, \mathcal{S} can be taken a full subcategory, closed under λ -directed colimits, of $\mathbf{Str}(\mathbf{w})$ with \mathbf{w} -embeddings, for some vocabulary \mathbf{w} extending \mathbf{v} with $\mathbf{w} \setminus \mathbf{v}$ λ -ary.

We will see in Theorem 6.9 that the functor Φ can be taken faithful.

Proof. By Proposition 5.4, there are a vocabulary \mathbf{w} , containing $\mathbf{v} \cup \{\mathbf{U}\}$ with $\mathbf{w} \setminus \mathbf{v}$ λ -ary, together with a conjunction \mathbf{E} of universal $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentences, such that (5.1) holds. Set $\mathcal{S} \stackrel{\text{def}}{=} \mathbf{Mod}_{\mathbf{w}}(\mathbf{E})$, with \mathbf{w} -embeddings as morphisms.

We claim that \mathcal{S} is λ -accessible. First observe that since \mathbf{E} is a conjunction of universal sentences of $\mathcal{L}_{\infty\lambda}(\mathbf{w})$, \mathbf{w}_{ope} is λ -ary, and by Lemma 2.1, \mathcal{S} has all λ -directed colimits, which are, up to isomorphism, the λ -directed unions. Since every member of \mathcal{S} is the λ -directed union of its λ -generated substructures (i.e., those generated by a λ -small subset), every λ -presentable member of \mathcal{S} is λ -generated. Let, conversely, \mathbf{A} be a λ -generated member of \mathcal{S} , let $\mathbf{M} = \bigcup_{p \in P} \mathbf{M}_p$ with P λ -directed, and suppose that \mathbf{A} is a submodel of \mathbf{M} . Since \mathbf{A} is λ -generated, $\mathbf{M} = \bigcup_{p \in P} \mathbf{M}_p$, and P is λ -directed, there exists $p \in P$ such that $\mathbf{A} \subseteq \mathbf{M}_p$. We shall prove that $\mathbf{R}^{\mathbf{A}} = \text{ar}(\mathbf{R})\mathbf{A} \cap \mathbf{R}^{\mathbf{M}_p}$ whenever $\mathbf{R} \in \mathbf{w}_{\text{rel}}$. For all $q \geq p$, \mathbf{M}_p is a submodel of \mathbf{M}_q thus $\mathbf{R}^{\mathbf{M}_p} = \text{ar}(\mathbf{R})\mathbf{M}_p \cap \mathbf{R}^{\mathbf{M}_q}$. Since $\mathbf{R}^{\mathbf{M}} = \bigcup_{q \in P} \mathbf{R}^{\mathbf{M}_q}$, it follows that $\mathbf{R}^{\mathbf{M}_p} = \text{ar}(\mathbf{R})\mathbf{M}_p \cap \mathbf{R}^{\mathbf{M}}$. Therefore, $\mathbf{R}^{\mathbf{A}} = \text{ar}(\mathbf{R})\mathbf{A} \cap \mathbf{R}^{\mathbf{M}} = \text{ar}(\mathbf{R})\mathbf{A} \cap \mathbf{R}^{\mathbf{M}_p}$, as desired. This proves that the λ -presentable members of \mathcal{S} are exactly its λ -generated members. Since those are obviously λ -directed colimit-dense in \mathcal{S} , it follows that \mathcal{S} is λ -accessible.

For any $\mathbf{N} \in \text{Ob } \mathcal{S}$, define $\Phi(\mathbf{N}) \stackrel{\text{def}}{=} \mathbf{U}^{\mathbf{N}} \downarrow_{\mathbf{v}}$. For any arrow $\alpha: \mathbf{M} \hookrightarrow \mathbf{N}$ within \mathcal{S} (so α is a \mathbf{w} -embedding), define $\Phi(\alpha)$ as the induced map from $\mathbf{U}^{\mathbf{M}}$ into $\mathbf{U}^{\mathbf{N}}$. By Lemma 2.1, Φ preserves λ -directed colimits. By (5.1), $\mathcal{C} = \text{rng } \Phi = \text{im } \Phi$. \square

The following example shows that the assumption that \mathbf{v}_{ope} be λ -ary cannot be dispensed with in the statement of Theorem 5.5.

Example 5.6 (The Idempotent). *Let λ be a regular cardinal and let the vocabulary \mathbf{u}_{λ} consist of the single operation symbol \mathbf{f} with $\text{ar}(\mathbf{f}) = \lambda$. Set $\mathcal{J}_{\lambda} \stackrel{\text{def}}{=} \mathbf{Mod}_{\mathbf{v}_{\lambda}}(\mathbf{J}_{\lambda})$ where \mathbf{J}_{λ} is the $\mathcal{L}_{\infty\lambda}(\mathbf{v}_{\lambda})$ -sentence $(\forall \mathbf{x})(\mathbf{f}(\mathbf{x}, \mathbf{x}, \dots) = \mathbf{x})$. Then \mathcal{J}_{λ} is not the image of any λ -continuous functor from any λ -accessible category to $\mathbf{Str}(\mathbf{u}_{\lambda})$.*

Proof. We shall in fact prove the following stronger statement (note the formal analogy with part of the statement of Theorem 7.1):

Let \mathcal{S} be a category, with a subcategory \mathcal{S}^{\dagger} whose objects form a set, and let $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{u}_{\lambda})$ be a functor such that

- (1) *every object S of \mathcal{S} is a colimit of a λ -directed commutative diagram \vec{S} in \mathcal{S}^{\dagger} such that $\Phi(S) = \varinjlim \Phi(\vec{S})$;*
- (2) *\mathcal{S} has all colimits from \mathcal{S}^{\dagger} indexed by the poset λ , and those colimits are preserved by Φ ;*
- (3) *$\text{im } \Phi$ is contained in \mathcal{J}_{λ} .*

Then $\text{im } \Phi$ is small-abstract.

(This indeed implies the statement of Example 5.6: take $\mathcal{S}^{\dagger} \stackrel{\text{def}}{=} \mathbf{Pres}_{\lambda} \mathcal{S}$.)

Suppose, to the contrary, that \mathcal{S} , \mathcal{S}^{\dagger} , and Φ satisfy (1)–(3) above, with $\text{im } \Phi$ not small-abstract. Let κ be an upper bound for all the cardinalities of $|\Phi(S)|$ for $S \in \text{Ob } \mathcal{S}^{\dagger}$, with $\kappa \geq \lambda$. Since $\text{im } \Phi$ is not small-abstract, there exists an object S of \mathcal{S} such that $(M, f) \stackrel{\text{def}}{=} \Phi(S) \in \text{rng } \Phi$ with $\kappa^{\lambda} < \text{card } M$. The statement that

$(M, f) \in \mathcal{J}_\lambda$ means that f is idempotent, that is, $f(x, x, \dots) = x$ whenever $x \in M$. By our assumption (1), there exists a λ -directed colimit cocone, within \mathcal{S} , of the form $(S, \sigma_p \mid p \in P) = \varinjlim (S_p, \sigma_{p,q} \mid p \leq q \text{ in } P)$, which is preserved by Φ ; whence, setting $(M_p, f_p) \stackrel{\text{def}}{=} \Phi(S_p)$, $\sigma_p \stackrel{\text{def}}{=} \Phi(\sigma_p)$, and $\sigma_{p,q} \stackrel{\text{def}}{=} \Phi(\sigma_{p,q})$, we get

$$((M, f), \sigma_p \mid p \in P) = \varinjlim ((M_p, f_p), \sigma_{p,q} \mid p \leq q \text{ in } P) \quad \text{within } \mathbf{Str}(\mathbf{u}_\lambda). \quad (5.2)$$

The set $\overline{M} \stackrel{\text{def}}{=} \bigcup_{p \in P} \sigma_p[M_p]$ is contained in M .

Claim 1. M is the closure of \overline{M} under f .

Proof of Claim. Denote by M' the closure of \overline{M} under f and by f' the domain-range restriction of f from ${}^\lambda M'$ to M' . Then each domain-range restriction $\sigma'_p: M_p \rightarrow M'$ of σ_p is an f-homomorphism from (M_p, f_p) to (M', f') , so those maps define a cocone above $((M_p, f_p), \sigma_{p,q} \mid p \leq q \text{ in } P)$, and so the universal property of the colimit yields a unique f-homomorphism $\varphi: M \rightarrow M'$ such that each $\varphi \circ \sigma_p = \sigma'_p$. Denoting by $\varepsilon: (M', f') \hookrightarrow (M, f)$ the inclusion map, it follows that each $\varepsilon \circ \varphi \circ \sigma_p = \varepsilon \circ \sigma'_p = \sigma_p$; whence $\varepsilon \circ \varphi = \text{id}_M$. Since ε is one-to-one, it is thus bijective; that is, $M = M'$. \square Claim 1.

Claim 2. $\text{card } \overline{M} > \kappa^\lambda$.

Proof of Claim. If $\text{card } \overline{M} \leq \kappa^\lambda$, then, since f has arity λ and by Claim 1, $\text{card } M \leq \kappa^\lambda$, a contradiction. \square Claim 2.

Claim 3. *There exists a strictly increasing λ -sequence $(p_\xi \mid \xi < \lambda)$ of elements in P such that the λ -sequence $(\sigma_{p_\xi}[M_{p_\xi}] \mid \xi < \lambda)$ is strictly increasing with respect to set inclusion.*

Proof of Claim. Let $\alpha < \lambda$ and suppose having performed the construction up to α , getting $(p_\xi \mid \xi < \alpha)$. Since $\text{card}(\bigcup_{\xi < \alpha} \sigma_{p_\xi}[M_{p_\xi}]) \leq \kappa \leq \kappa^\lambda$, it follows from Claim 2 that $\sigma_{p_\alpha}[M_{p_\alpha}] \not\subseteq \bigcup_{\xi < \alpha} \sigma_{p_\xi}[M_{p_\xi}]$ for some $p_\alpha \in P$. Since P is λ -directed, p_α may be taken an upper bound of $\{p_\xi \mid \xi < \alpha\}$. Then $\sigma_{p_\xi}[M_{p_\xi}] \subsetneq \sigma_{p_\alpha}[M_{p_\alpha}]$ whenever $\xi < \alpha$. \square Claim 3.

Renaming the p_ξ obtained in Claim 3, we may assume that $p_\xi = \xi$ for all $\xi < \lambda$ (so λ is a sub-poset of P). Pick $a_\xi \in \sigma_{\xi+1}[M_{\xi+1}] \setminus \sigma_\xi[M_\xi]$, for each $\xi < \lambda$. Setting $D \stackrel{\text{def}}{=} \bigcup_{\alpha < \lambda} {}^\lambda \sigma_\alpha[M_\alpha]$, it follows that $\vec{a} \stackrel{\text{def}}{=} (a_\xi \mid \xi < \lambda)$ belongs to ${}^\lambda \overline{M} \setminus D$. Moreover, for each $\alpha < \lambda$ and $\vec{x} \in {}^\lambda \sigma_\alpha[M_\alpha]$, $f \sigma_\alpha(\vec{x}) = \sigma_\alpha f_\alpha(\vec{x}) \in \overline{M}$; so $f[D] \subseteq \overline{M}$.

We can thus endow the set $N \stackrel{\text{def}}{=} \overline{M} \sqcup \{\infty\}$, for an extra element $\infty \notin N$, with the map $g: {}^\lambda N \rightarrow N$ defined by

- $g(\vec{x}) = f(\vec{x})$ whenever $\vec{x} \in D$ (in that case $g(\vec{x}) \in \overline{M}$);
- $g(\infty, \infty, \dots) \stackrel{\text{def}}{=} a_0$;
- $g(\vec{x}) = \infty$ in all other cases.

It follows from our Assumption (2) that the colimit

$$(S', \sigma'_\xi \mid \xi < \lambda) = \varinjlim (S_\xi, \sigma_{\xi,\eta} \mid \xi \leq \eta < \lambda)$$

exists in \mathcal{S} , and further, setting $(M', f') \stackrel{\text{def}}{=} \Phi(S')$ and $\sigma'_\xi \stackrel{\text{def}}{=} \Phi(\sigma'_\xi)$,

$$((M', f'), \sigma'_\xi \mid \xi < \lambda) \stackrel{\text{def}}{=} \varinjlim ((M_\xi, f_\xi), \sigma_{\xi,\eta} \mid \xi \leq \eta < \lambda). \quad (5.3)$$

Since $(M', f') = \Phi(S') \in \text{rng } \Phi \subseteq \mathcal{J}_\lambda$ (cf. (3)), the map f' is idempotent. Moreover, for each $\xi < \lambda$, the domain-range restriction $\tau_\xi: (M_\xi, f_\xi) \rightarrow (N, g)$ of σ_ξ is an f -homomorphism; so those maps define a cocone above $((M_\xi, f_\xi), \sigma_{\xi, \eta} \mid \xi \leq \eta < \lambda)$ within $\mathbf{Str}(\mathbf{u}_\lambda)$. By the universal property of the colimit (5.3), there exists a unique f -homomorphism $\psi: (M', f') \rightarrow (N, g)$ such that each $\psi \circ \sigma'_\xi = \tau_\xi$.

For each $\xi < \lambda$, pick $u_\xi \in M_{\xi+1}$ such that $a_\xi = \sigma_{\xi+1}(u_\xi)$; then $a'_\xi \stackrel{\text{def}}{=} \sigma'_{\xi+1}(u_\xi)$ belongs to M' , and so does $t \stackrel{\text{def}}{=} f'(a'_\xi \mid \xi < \lambda)$. Now each $\psi(a'_\xi) = \psi \sigma'_{\xi+1}(u_\xi) = \tau_{\xi+1}(u_\xi) = \sigma_{\xi+1}(u_\xi) = a_\xi$, whence $\psi(t) = g(a_\xi \mid \xi < \lambda) = \infty$. It follows that

$$g(\infty, \infty, \dots) = g(\psi(t), \psi(t), \dots) = \psi(f'(t, t, \dots)) = \psi(t) = \infty;$$

a contradiction since $g(\infty, \infty, \dots) = a_0$. \square

6. ANOTHER CASE OF CONJUGACY: FROM RPC TO PC

In general, a vocabulary \mathbf{v} may have operation symbols of arity λ or larger. This makes it harder to write an RPC class, of \mathbf{v} -structures, in PC form: the statement that U^N be \mathbf{v} -closed cannot *a priori* be incorporated into \mathbf{E} in (3.2). Similarly, the statement that a relation, with arity λ or greater, is the graph of a function, cannot be expressed by an $\mathcal{L}_{\infty\lambda}$ -sentence. Nonetheless, we shall see that this difficulty can be circumvented, enabling us to prove (cf. Theorem 6.8) that $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$ for any regular cardinal λ .

Until Lemma 6.7 we shall fix an infinite cardinal λ together with a unary predicate symbol U and vocabularies \mathbf{v}, \mathbf{w} such that $\mathbf{v} \cup \{U\} \subseteq \mathbf{w}$. We shall also set $T_\lambda^*(\mathbf{w}) \stackrel{\text{def}}{=} T_\lambda(\mathbf{w}) \setminus \text{Var}$, and

$$\tilde{\mathbf{w}} \stackrel{\text{def}}{=} \mathbf{v} \sqcup \{\dot{s} \mid s \in T_\lambda^*(\mathbf{w}) \setminus \mathbf{v}_{\text{cst}}\} \sqcup \{\dot{E} \mid E \in A_\lambda(\mathbf{w}) \text{ relational}\} \sqcup \{J_{s,t} \mid s, t \in T_\lambda(\mathbf{w})\},$$

where each $\text{ar}(\dot{s}) \stackrel{\text{def}}{=} \text{var}(s)$, each $\text{ar}(\dot{E}) \stackrel{\text{def}}{=} \text{var}(E)$, and each $J_{s,t}$ is a relation symbol with arity a disjoint union of $\text{var}(s)$ and $\text{var}(t)$. Partly following (4.1) and (4.2), for $s \in T_\lambda(\mathbf{w})$ and $E \in A_\lambda(\mathbf{w})$, the terms $\tilde{s} \in T_\lambda(\tilde{\mathbf{w}})$ and atomic formulas $\tilde{E} \in A_\lambda(\tilde{\mathbf{w}})$, with the same variables as s and E , respectively, are now defined as follows:

$$\tilde{s} \stackrel{\text{def}}{=} \begin{cases} s, & \text{if } s \in \mathbf{v}_{\text{cst}} \cup \text{Var}, \\ \dot{s}(u \mid u \in \text{var}(s)) & \text{(usually written as } \dot{s}), \text{ otherwise,} \end{cases} \quad (6.1)$$

$$\tilde{E} \stackrel{\text{def}}{=} \begin{cases} J_{s,t}(\vec{x}, \vec{y}), & \text{if } E \text{ has the form } s(\vec{x}) = t(\vec{y}), \\ \dot{E}(u \mid u \in \text{var}(E)) & \text{(usually written as } \dot{E}), \text{ otherwise.} \end{cases} \quad (6.2)$$

The following definition may be viewed as an adaptation, to the present context, of the “ κ -partitions” introduced in Oikkonen [24, Notation 2.1].

Definition 6.1. A \mathbf{w} -structure M is a (λ, U) -*distension* of a \mathbf{v} -structure A if $A = U^M \upharpoonright_{\mathbf{v}}$ and $M = \{s^M(\vec{a}) \mid s \in T_\lambda(\mathbf{w}) \text{ and } \vec{a} \in \text{var}(s)A\}$.

Observe that in the context of Definition 6.1, U^M is necessarily \mathbf{v} -closed in N .

Notation 6.2 (Coherence sentence). Denote by H the conjunction of all $\mathcal{L}_{\infty\lambda}(\tilde{\mathbf{w}})$ -sentences of the following form (where z, z_1, z_2 are variables, $s, s_i, t \in T_\lambda(\mathbf{w})$,

$f \in \mathbf{w}_{\text{ope}}, R \in \mathbf{w}_{\text{rel}}$:

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x}) J_{\mathbf{s}, \mathbf{s}}(\vec{x}, \vec{x}); \quad (6.3)$$

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x})(\forall_{(\text{var}(\mathbf{t}))} \vec{y})(J_{\mathbf{s}, \mathbf{t}}(\vec{x}, \vec{y}) \Rightarrow J_{\mathbf{t}, \mathbf{s}}(\vec{y}, \vec{x})); \quad (6.4)$$

$$(\forall_{(\text{var}(\mathbf{s}_1))} \vec{x}_1)(\forall_{(\text{var}(\mathbf{s}_2))} \vec{x}_2)(\forall_{(\text{var}(\mathbf{s}_3))} \vec{x}_3) \left((J_{\mathbf{s}_1, \mathbf{s}_2}(\vec{x}_1, \vec{x}_2) \wedge J_{\mathbf{s}_2, \mathbf{s}_3}(\vec{x}_2, \vec{x}_3)) \Rightarrow J_{\mathbf{s}_1, \mathbf{s}_3}(\vec{x}_1, \vec{x}_3) \right); \quad (6.5)$$

$$J_{\mathbf{c}, \mathbf{z}}(\mathbf{c}), \quad \text{whenever } \mathbf{c} \in \mathbf{v}_{\text{cst}}; \quad (6.6)$$

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x})(\forall_{(\text{var}(\mathbf{t}))} \vec{y})(J_{\mathbf{s}, \mathbf{t}}(\vec{x}, \vec{y}) \Rightarrow \tilde{\mathbf{s}}(\vec{x}) = \tilde{\mathbf{t}}(\vec{y})); \quad (6.7)$$

$$(\forall \mathbf{x}) J_{\mathbf{z}_1, \mathbf{z}_2}(\mathbf{x}, \mathbf{x}); \quad (6.8)$$

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x})(\forall_{(\text{var}(\mathbf{t}))} \vec{y}) \left(\left(\bigwedge_{\xi \in \text{ar}(f)} J_{\mathbf{s}_\xi, \mathbf{t}_\xi}(\vec{x}, \vec{y}) \right) \Rightarrow J_{\mathbf{s}, \mathbf{t}}(\vec{x}, \vec{y}) \right),$$

provided $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ and $\mathbf{t} = f(\mathbf{t}_\xi \mid \xi \in \text{ar}(f))$ in $\mathbf{T}_\lambda(\mathbf{w})$; (6.9)

(Here and elsewhere we allow for dummy variables in notations such as $J_{\mathbf{s}_\xi, \mathbf{t}_\xi}(\vec{x}, \vec{y})$.)

$$(\forall_{(\text{var}(\mathbf{E}))} \vec{x})(\forall_{(\text{var}(\mathbf{F}))} \vec{y}) \left(\left(\tilde{\mathbf{E}}(\vec{x}) \wedge \bigwedge_{\xi \in \text{ar}(\mathbf{R})} J_{\mathbf{s}_\xi, \mathbf{t}_\xi}(\vec{x}, \vec{y}) \right) \Rightarrow \tilde{\mathbf{F}}(\vec{y}) \right),$$

provided $\mathbf{E} = \mathbf{R}(\mathbf{s}_\xi \mid \xi \in \text{ar}(\mathbf{R}))$ and $\mathbf{F} = \mathbf{R}(\mathbf{t}_\xi \mid \xi \in \text{ar}(\mathbf{R}))$ in $\mathbf{A}_\lambda(\mathbf{w})$; (6.10)

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x}) \left(\left(\bigwedge_{\xi \in \text{ar}(f)} J_{\mathbf{s}_\xi, \mathbf{z}}(\vec{x}, \tilde{\mathbf{s}}_\xi(\vec{x})) \right) \Rightarrow \right.$$

$$\left. \left(\tilde{\mathbf{s}}(\vec{x}) = f(\tilde{\mathbf{s}}_\xi(\vec{x}) \mid \xi \in \text{ar}(f)) \wedge J_{\mathbf{s}, \mathbf{z}}(\vec{x}, \tilde{\mathbf{s}}(\vec{x})) \right) \right),$$

provided $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ in $\mathbf{T}_\lambda(\mathbf{w})$ with $f \in \mathbf{v}_{\text{ope}}$; (6.11)

$$(\forall_{(\text{var}(\mathbf{E}))} \vec{x}) \left(\left(\bigwedge_{\xi \in \text{ar}(\mathbf{R})} J_{\mathbf{s}_\xi, \mathbf{z}}(\vec{x}, \tilde{\mathbf{s}}_\xi(\vec{x})) \right) \Rightarrow \right.$$

$$\left. \left(\tilde{\mathbf{E}}(\vec{x}) \Leftrightarrow \mathbf{R}(\tilde{\mathbf{s}}_\xi(\vec{x}) \mid \xi \in \text{ar}(\mathbf{R})) \right) \right),$$

provided $\mathbf{E} = \mathbf{R}(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ in $\mathbf{A}_\lambda(\mathbf{w})$ with $\mathbf{R} \in \mathbf{v}_{\text{rel}}$, (6.12)

$$(\forall_{(\text{var}(\mathbf{s}))} \vec{x}) \left(\tilde{\mathbf{U}}\mathbf{s}(\vec{x}) \Leftrightarrow J_{\mathbf{s}, \mathbf{z}}(\vec{x}, \tilde{\mathbf{s}}(\vec{x})) \right). \quad (6.13)$$

Definition 6.3. Let \mathbf{M} be a \mathbf{w} -structure and let \mathbf{A}' be a $\tilde{\mathbf{w}}$ -structure. We say that $(\mathbf{M}, \mathbf{A}')$ is a *conjugate pair*, or, equivalently, that \mathbf{M} is *left conjugate to* \mathbf{A}' , respectively that \mathbf{A}' is *right conjugate to* \mathbf{M} , if the following statements hold:

- (1) $\mathbf{A}' \upharpoonright_{\mathbf{v}} = \mathbf{U}^{\mathbf{M}} \upharpoonright_{\mathbf{v}}$ (denote that \mathbf{v} -structure by \mathbf{A});
- (2) $M = \{\mathbf{s}^{\mathbf{M}}(\vec{a}) \mid \mathbf{s} \in \mathbf{T}_\lambda(\mathbf{w}) \text{ and } \vec{a} \in \text{var}(\mathbf{s})A\}$;
- (3) for all $\mathbf{s}, \mathbf{t} \in \mathbf{T}_\lambda(\mathbf{w})$, $\vec{a} \in \text{var}(\mathbf{s})A$, and $\vec{b} \in \text{var}(\mathbf{t})A$, $\mathbf{s}^{\mathbf{M}}(\vec{a}) = \mathbf{t}^{\mathbf{M}}(\vec{b})$ implies that $\tilde{\mathbf{s}}^{\mathbf{A}'}(\vec{a}) = \tilde{\mathbf{t}}^{\mathbf{A}'}(\vec{b})$;
- (4) for every relational $\mathbf{E} \in \mathbf{A}_\lambda(\mathbf{w})$, $\tilde{\mathbf{E}}^{\mathbf{A}'} = \{\vec{a} \in \text{var}(\mathbf{E})A \mid \mathbf{M} \models \mathbf{E}(\vec{a})\}$;

(5) for all $s, t \in T_\lambda(\mathbf{w})$, all $\vec{a} \in \text{var}(s)A$, and all $\vec{b} \in \text{var}(t)A$, $\mathbf{A}' \models J_{s,t}(\vec{a}, \vec{b})$ iff $\mathbf{M} \models s(\vec{a}) = t(\vec{b})$.

Lemma 6.4. *For every conjugate pair $(\mathbf{M}, \mathbf{A}')$ and $\mathbf{A} \stackrel{\text{def}}{=} \mathbf{A}' \upharpoonright_{\mathbf{v}}$, \mathbf{M} is a (λ, U) -distension of \mathbf{A} and \mathbf{A}' satisfies \mathbf{H} .*

Proof. The statement that \mathbf{M} is a (λ, U) -distension of \mathbf{A} trivially follows from items (1) and (2) of Definition 6.3. By letting t be a variable in (3), we obtain

(3') For all $s \in T_\lambda(\mathbf{w})$ and all $\vec{a} \in \text{var}(s)A$, $s^{\mathbf{M}}(\vec{a}) \in A$ implies that $\tilde{s}^{\mathbf{A}'}(\vec{a}) = s^{\mathbf{M}}(\vec{a})$.

The verification of the statements (6.3)–(6.13) in \mathbf{A}' breaks down into the following observations:

- (6.3)–(6.5): use Definition 6.3(5);
- (6.6): use $c^{\mathbf{A}} = c^{\mathbf{M}}$ whenever $c \in \mathbf{v}_{\text{cst}}$;
- (6.7): use (5) and (3) from Definition 6.3;
- (6.8): use Definition 6.3(5);
- (6.9): Definition 6.3(5);
- (6.10): use (5) and (4) from Definition 6.3;
- (6.11): suppose that $\mathbf{A}' \models J_{s_\xi, z}(\vec{a}, \tilde{s}_\xi(\vec{a}))$ for all $\xi \in \text{ar}(f)$. By (3') and Definition 6.3(5), this means that the element $b_\xi \stackrel{\text{def}}{=} s_\xi^{\mathbf{M}}(\vec{a})$ belongs to A and $b_\xi = \tilde{s}_\xi^{\mathbf{A}'}(\vec{a})$, whenever $\xi \in \text{ar}(f)$. Since $f \in \mathbf{v}_{\text{ope}}$, the element $b \stackrel{\text{def}}{=} s^{\mathbf{M}}(\vec{a}) = f^{\mathbf{M}}(b_\xi \mid \xi \in \text{ar}(f)) = f^{\mathbf{A}}(b_\xi \mid \xi \in \text{ar}(f))$ belongs to A . By (3') and Definition 6.3(5), $b = \tilde{s}^{\mathbf{A}'}(\vec{a})$ and $\mathbf{A}' \models J_{s,z}(\vec{a}, b)$.
- (6.12): a similar argument as for (6.11), using (3') and Definition 6.3(4).
- (6.13): $\mathbf{A}' \models \widetilde{U}s(\vec{a})$ iff $\mathbf{M} \models Us(\vec{a})$ iff $s^{\mathbf{M}}(\vec{a}) \in A$ (use (4) and (1) from Definition 6.3). \square

Lemma 6.5. *The following statements hold, for any \mathbf{v} -structure \mathbf{A} :*

- (1) *Every (λ, U) -distension \mathbf{M} of \mathbf{A} has a right conjugate \mathbf{A}' .*
- (2) *Every expansion \mathbf{A}' of \mathbf{A} to $\widetilde{\mathbf{w}}$, satisfying \mathbf{H} , has a left conjugate \mathbf{M} .*

Proof. *Ad (1).* We need to define the interpretations in \mathbf{A}' of all symbols in $\widetilde{\mathbf{w}} \setminus \mathbf{v}$. Pick $o \in A$.

- For every $s \in T_\lambda(\mathbf{w}) \setminus \mathbf{v}_{\text{cst}}$ and every $\vec{a} \in \text{var}(s)A$, we set

$$\dot{s}^{\mathbf{A}'}(\vec{a}) \stackrel{\text{def}}{=} \begin{cases} s^{\mathbf{M}}(\vec{a}), & \text{if } s^{\mathbf{M}}(\vec{a}) \in A; \\ o, & \text{otherwise.} \end{cases}$$

- For every relational $E \in \mathbf{A}_\lambda(\mathbf{w})$, we set

$$\dot{E}^{\mathbf{A}'} \stackrel{\text{def}}{=} \left\{ \vec{a} \in \text{ar}(E)A \mid \mathbf{M} \models E(\vec{a}) \right\}.$$

- For all $s, t \in T_\lambda(\mathbf{w})$, we set

$$J_{s,t}^{\mathbf{A}'} \stackrel{\text{def}}{=} \left\{ (\vec{a}, \vec{b}) \in \text{var}(s)A \times \text{var}(t)A \mid \mathbf{M} \models s(\vec{a}) = t(\vec{b}) \right\}.$$

The verifications of (1)–(5) in Definition 6.3 are straightforward.

Ad (2). Setting $\mathcal{M} \stackrel{\text{def}}{=} \{(s, \vec{a}) \mid s \in T_\lambda(\mathbf{w}) \text{ and } \vec{a} \in \text{var}(s)A\}$, we define a binary relation \sim on \mathcal{M} by the rule

$$(s, \vec{a}) \sim (t, \vec{b}) \quad \text{if } \mathbf{A}' \models J_{s,t}(\vec{a}, \vec{b}), \quad \text{for all } (s, \vec{a}), (t, \vec{b}) \in \mathcal{M}. \quad (6.14)$$

Since \mathbf{A}' satisfies all statements (6.3)–(6.5), \sim is an equivalence relation on \mathcal{M} . Denoting the \sim -equivalence class of (\mathbf{s}, \vec{a}) by $[\mathbf{s}, \vec{a}]$, we set

$$M \stackrel{\text{def}}{=} \mathcal{M}/\sim = \{[\mathbf{s}, \vec{a}] \mid (\mathbf{s}, \vec{a}) \in \mathcal{M}\}.$$

To avoid clutter, we will frequently extend the notation $[\mathbf{s}, \vec{a}]$ to pairs (\mathbf{s}, \vec{a}) where $\vec{a} \in {}^X A$ with $\text{var}(\mathbf{s}) \subseteq X$, by setting $[\mathbf{s}, \vec{a}] \stackrel{\text{def}}{=} [\mathbf{s}, \vec{a}]_{\text{var}(\mathbf{s})}$. We further set $\varepsilon(a) \stackrel{\text{def}}{=} [z, a]$ (for $a \in A$), which is independent of the variable z by (6.8), whenever $a \in A$. By (6.7) for \mathbf{s}, \mathbf{t} variables, ε is one-to-one. We set $\mathbf{c}^M \stackrel{\text{def}}{=} [c, \emptyset]$ whenever $c \in \mathbf{w}_{\text{cst}}$. By (6.6), $\varepsilon(\mathbf{c}^A) = \mathbf{c}^M$ whenever $c \in \mathbf{v}_{\text{cst}}$.

Claim 1. *For every $(\mathbf{s}, \vec{a}) \in \mathcal{M}$, $[\mathbf{s}, \vec{a}] \in \varepsilon A$ iff $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}, z}(\vec{a}, \tilde{\mathbf{s}}(\vec{a}))$.*

Proof of Claim. If $[\mathbf{s}, \vec{a}] \in \varepsilon A$, then there exists $b \in A$ such that $[\mathbf{s}, \vec{a}] = [z, b]$, that is, $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}, z}(\vec{a}, b)$. By (6.7) with \mathbf{t} a variable, $b = \tilde{\mathbf{s}}^{\mathbf{A}'}(\vec{a})$, whence $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}, z}(\vec{a}, \tilde{\mathbf{s}}(\vec{a}))$. The argument can be followed backwards. \square Claim 1.

Claim 2. *For every $f \in \mathbf{w}_{\text{ope}}$, there exists a map $f^M : \text{ar}(f)M \rightarrow M$ such that for all $\vec{x} = (x_\xi \mid \xi \in \text{ar}(f)) \in \text{ar}(f)M$,*

$$f^M(\vec{x}) = \begin{cases} [\mathbf{s}, \vec{a}], & \text{if } \mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f)) \in \mathbf{T}_\lambda(\mathbf{w}), \vec{a} \in \text{var}(\mathbf{s})A, \text{ each } x_\xi = [\mathbf{s}_\xi, \vec{a}], \\ \varepsilon f^A(\vec{b}), & \text{if } f \in \mathbf{v}_{\text{ope}} \text{ and } \vec{x} = \varepsilon(\vec{b}) \text{ for some } \vec{b} \in \text{ar}(f)A. \end{cases}$$

Proof of Claim. The case where $f \in \mathbf{v}_{\text{cst}}$ has already been covered above (via $\varepsilon(\mathbf{c}^A) = \mathbf{c}^M$). Suppose from now on that $f \notin \mathbf{v}_{\text{cst}}$. If $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f))$ and $\mathbf{t} = f(\mathbf{t}_\xi \mid \xi \in \text{ar}(f))$ in $\mathbf{T}_\lambda(\mathbf{w})$, with each $x_\xi = [\mathbf{s}_\xi, \vec{a}] = [\mathbf{t}_\xi, \vec{b}]$, so $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}_\xi, \mathbf{t}_\xi}(\vec{a}, \vec{b})$, then, by (6.9), $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}, \mathbf{t}}(\vec{a}, \vec{b})$, that is, $[\mathbf{s}, \vec{a}] = [\mathbf{t}, \vec{b}]$. Now if $f \in \mathbf{v}_{\text{ope}}$, $\mathbf{s} = f(\mathbf{s}_\xi \mid \xi \in \text{ar}(f)) \in \mathbf{T}_\lambda(\mathbf{w})$, and each $x_\xi = [\mathbf{s}_\xi, \vec{a}] = \varepsilon(b_\xi)$, then, by the proof of Claim 1, each $b_\xi = \tilde{\mathbf{s}}_\xi^{\mathbf{A}'}(\vec{a})$, so, setting $b \stackrel{\text{def}}{=} \tilde{\mathbf{s}}^{\mathbf{A}'}(\vec{a})$, it follows from (6.11) that $b = f^A(\vec{b})$ and $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}, z}(\vec{a}, b)$; whence $[\mathbf{s}, \vec{a}] = [z, b] = \varepsilon(b)$. \square Claim 2.

Claim 3. *Let $R \in \mathbf{w}_{\text{rel}}$, set*

$$R^* \stackrel{\text{def}}{=} \{([\mathbf{s}_\xi, \vec{a}] \mid \xi \in \text{ar}(R)) \mid E \stackrel{\text{def}}{=} R(\mathbf{s}_\xi \mid \xi \in \text{ar}(R)) \in \mathbf{T}_\lambda(\mathbf{w}), \\ \vec{a} \in \text{var}(E)A, \text{ and } \mathbf{A}' \models \tilde{E}(\vec{a})\},$$

and $R^M \stackrel{\text{def}}{=} \begin{cases} \varepsilon R^A \cup R^*, & \text{if } R \in \mathbf{v}_{\text{rel}}, \\ R^*, & \text{otherwise.} \end{cases}$. Then the following statements hold:

- (1) *If $R \in \mathbf{v}_{\text{rel}}$, then $R^M \cap \text{ar}(R)\varepsilon A = \varepsilon R^A$.*
- (2) *Whenever $E = R(\mathbf{s}_\xi \mid \xi \in \text{ar}(R)) \in \mathbf{A}_\lambda(\mathbf{w})$ and $\vec{a} \in \text{var}(E)A$, $([\mathbf{s}_\xi, \vec{a}] \mid \xi \in \text{ar}(R)) \in R^M$ iff $\mathbf{A}' \models \tilde{E}(\vec{a})$.*

Proof of Claim. Ad (1). Let $E = R(\mathbf{s}_\xi \mid \xi \in \text{ar}(R))$ in $\mathbf{T}_\lambda(\mathbf{w})$ and $\vec{a} \in \text{var}(E)A$. Moreover, let $\vec{b} = (b_\xi \mid \xi \in \text{ar}(R)) \in \text{ar}(R)A$ such that each $[\mathbf{s}_\xi, \vec{a}] = [z, b_\xi]$; that is, $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}_\xi, z}(\vec{a}, b_\xi)$. By the proof of Claim 1, each $b_\xi = \tilde{\mathbf{s}}_\xi^{\mathbf{A}'}(\vec{a})$. By (6.12), it follows that $\mathbf{A}' \models \tilde{E}(\vec{a})$ iff $\vec{b} \in R^A$, as desired.

Ad (2). Let $([\mathbf{s}_\xi, \vec{a}] \mid \xi \in \text{ar}(R)) \in R^M$. If $([\mathbf{s}_\xi, \vec{a}] \mid \xi \in \text{ar}(R)) \in R^*$ then there are $F = R(\mathbf{t}_\xi \mid \xi \in \text{ar}(R)) \in \mathbf{A}_\lambda(\mathbf{w})$ and $\vec{b} \in \text{var}(F)A$ such that $\mathbf{A}' \models \tilde{F}(\vec{b})$ and each $[\mathbf{s}_\xi, \vec{a}] = [\mathbf{t}_\xi, \vec{b}]$; that is, $\mathbf{A}' \models \mathbf{J}_{\mathbf{s}_\xi, \mathbf{t}_\xi}(\vec{a}, \vec{b})$. The desired conclusion then follows from (6.10). Suppose now that $R \in \mathbf{v}_{\text{rel}}$ and $([\mathbf{s}_\xi, \vec{a}] \mid \xi \in \text{ar}(R)) \in \varepsilon R^A$. There exists

$\vec{b} \in \mathbf{R}^{\mathbf{A}}$ such that each $[s_\xi, \vec{a}] = [z, b_\xi]$; so $\mathbf{A}' \models J_{s_\xi, z}(\vec{a}, \tilde{s}_\xi(\vec{a}))$ and $b_\xi = \tilde{s}_\xi^{\mathbf{A}'}(\vec{a})$. Since $\mathbf{A}' \models \mathbf{R}(b_\xi \mid \xi \in \text{ar}(\mathbf{R}))$, it follows from (6.12) that $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$.

Conversely, if $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$, then $([s_\xi, \vec{a}] \mid \xi \in \text{ar}(\mathbf{R})) \in R^* \subseteq \mathbf{R}^{\mathbf{M}}$. \square Claim 3.

In particular, Claims 2 and 3 ensure that the map ε is a \mathbf{v} -embedding from \mathbf{A} into the \mathbf{w} -structure \mathbf{M} thus defined.

Claim 4. $\varepsilon \mathbf{A} = \mathbf{U}^{\mathbf{M}} \upharpoonright_{\mathbf{v}}$.

Proof of Claim. Since ε is a \mathbf{v} -embedding, it suffices to verify that $[s, \vec{a}] \in \varepsilon \mathbf{A}$ iff $[s, \vec{a}] \in \mathbf{U}^{\mathbf{M}}$ whenever $(s, \vec{a}) \in \mathcal{M}$. By Claim 1, $[s, \vec{a}] \in \varepsilon \mathbf{A}$ is equivalent to $\mathbf{A}' \models J_{s, z}(\vec{a}, \tilde{s}(\vec{a}))$. The statement $[s, \vec{a}] \in \mathbf{U}^{\mathbf{M}}$ means that $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$ where $\mathbf{E} \stackrel{\text{def}}{=} \mathbf{U}(s)$. By (6.13), the two are equivalent. \square Claim 4.

Claim 5. $s^{\mathbf{M}}(\varepsilon \vec{a}) = [s, \vec{a}]$ whenever $(s, \vec{a}) \in \mathcal{M}$.

Proof of Claim. If $s = c \in \mathbf{w}_{\text{cst}}$ then the desired relation follows from the definition $c^{\mathbf{M}} = [c, \emptyset]$. For nonconstant s , argue by induction on s and apply Claim 2. \square Claim 5.

By Claims 4 and 5, items (1) and (2) of Definition 6.3 are both satisfied; that is, \mathbf{M} is a (λ, \mathbf{U}) -distension of \mathbf{A} (or, strictly speaking, of its isomorphic copy $\varepsilon \mathbf{A}$).

Let us verify Definition 6.3(3). Suppose that $s^{\mathbf{M}}(\varepsilon \vec{a}) = t^{\mathbf{M}}(\varepsilon \vec{b})$. By Claim 5, $(s, \vec{a}) \sim (t, \vec{b})$, that is, $\mathbf{A}' \models J_{s, t}(\vec{a}, \vec{b})$. Using (6.7), it follows that $\tilde{s}^{\mathbf{A}'}(\varepsilon \vec{a}) = \tilde{t}^{\mathbf{A}'}(\varepsilon \vec{b})$.

Now Definition 6.3(4). For all $\vec{a} \in \text{var}^{(\mathbf{E})} \mathbf{A}$, \mathbf{M} satisfies $\mathbf{E}(\varepsilon \vec{a})$ iff the vector $(s_\xi^{\mathbf{M}}(\varepsilon \vec{a}) \mid \xi \in \text{ar}(\mathbf{R}))$ belongs to $\mathbf{R}^{\mathbf{M}}$. Since each $s_\xi^{\mathbf{M}}(\varepsilon \vec{a}) = [s_\xi, \vec{a}]$ (cf. Claim 5) and by Claim 3, this holds iff $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$.

For Definition 6.3(5), apply Claim 5 together with (6.14). \square

Notation 6.6. Suppose that λ is regular. We extend the assignment $\mathbf{E} \mapsto \tilde{\mathbf{E}}$, originally defined on all atomic formulas of $\mathcal{L}_{\infty\lambda}(\mathbf{w})$, as follows.

- If \mathbf{E} is $\bigvee_{i \in I} \mathbf{E}_i$, then $\tilde{\mathbf{E}}$ is $\bigvee_{i \in I} \tilde{\mathbf{E}}_i$.
- If \mathbf{E} is $\neg \mathbf{F}$, then $\tilde{\mathbf{E}}$ is $\neg \tilde{\mathbf{F}}$.
- Let $\mathbf{E}(\vec{x})$ be $(\exists \vec{y}) F(\vec{x}, \vec{y})$, where $\text{card } J < \lambda$. For every $t \in {}^J \mathbf{T}_\lambda(\mathbf{w})$, setting

$\text{var}(t) \stackrel{\text{def}}{=} \bigcup_{j \in J} \text{var}(t_j)$ (note that *since* λ *is regular*, that set is λ -small), we denote by $\mathbf{F}^t(\vec{x}, \vec{u})$ the formula $F(\vec{x}, (t_j(\vec{u}) \mid j \in J))$, then define $\tilde{\mathbf{E}}(\vec{x})$ as

$$\bigvee_{t \in {}^J \mathbf{T}_\lambda(\mathbf{w})} (\exists_{(\text{var}(t))} \vec{u}) \tilde{\mathbf{F}}^t(\vec{x}, \vec{u}).$$

Lemma 6.7. *Let λ be regular, let $(\mathbf{M}, \mathbf{A}')$ be a conjugate pair, and set $\mathbf{A} \stackrel{\text{def}}{=} \mathbf{A}' \upharpoonright_{\mathbf{v}}$. Then for every formula $\mathbf{E}(\vec{x}) \in \mathcal{L}_{\infty\lambda}(\mathbf{w})$ and every $\vec{a} \in {}^I \mathbf{A}$, $\mathbf{M} \models \mathbf{E}(\vec{a})$ iff $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$.*

Proof. The case where \mathbf{E} is atomic follows from the definition of conjugacy (cf. Definition 6.3, especially items (4) and (5)). The induction steps corresponding to disjunction and negation are trivial. Now let $\mathbf{E}(\vec{x})$ be $(\exists \vec{y}) F(\vec{x}, \vec{y})$, where I

and J are both λ -small, and let $\vec{a} \in {}^I \mathbf{A}$. If $\mathbf{M} \models \mathbf{E}(\vec{a})$ then there exists $\vec{y} \in {}^J \mathbf{M}$ such that $\mathbf{M} \models F(\vec{a}, \vec{y})$. By Definition 6.3(2), for each $j \in J$ there are $t_j \in \mathbf{T}_\lambda(\mathbf{w})$

and $\vec{u}_j \in \text{var}(\mathfrak{t}_j)A$ such that $y_j = \mathfrak{t}_j^M(\vec{u}_j)$. Since λ is regular, $\sum_{j \in J} \text{card var}(\mathfrak{t}_j) < \lambda$ and we may thus assume, after a suitable reindexing, that each $y_j = \mathfrak{t}_j^M(\vec{u})$ for a single vector $\vec{u} \in \text{var}(\mathfrak{t})A$ with $\text{card var}(\mathfrak{t}) < \lambda$; whence $\mathbf{M} \models \mathbf{F}^{\mathfrak{t}}(\vec{a}, \vec{u})$. By our induction hypothesis, $\mathbf{A}' \models \tilde{\mathbf{F}}^{\mathfrak{t}}(\vec{a}, \vec{u})$; whence $\mathbf{A}' \models \tilde{\mathbf{E}}(\vec{a})$. This argument can be followed backwards, thus the desired equivalence holds. \square

Theorem 6.8. *Let λ be a regular cardinal. Then $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$.*

Proof. Let \mathbf{v} be a vocabulary and let \mathcal{C} be an $\text{RPC}(\mathcal{L}_{\infty\lambda})$ class of \mathbf{v} -structures. By Proposition 5.4, there are a vocabulary \mathbf{w} containing \mathbf{v} , with $\mathbf{w} \setminus \mathbf{v}$ λ -ary and a unary predicate symbol \mathbf{U} , together with a conjunction \mathbf{E} of universal $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentences, such that (3.2) holds.

Claim.

$$\mathcal{C} = \{U^M \upharpoonright_{\mathbf{v}} \mid M \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E}), \\ M \text{ is a } (\lambda, \mathbf{U})\text{-distension of } U^M, U^M \mathbf{v}\text{-closed}\}. \quad (6.15)$$

Proof of Claim. We prove the nontrivial containment. Let $\mathbf{A} \in \mathcal{C}$. By (3.2), $\mathbf{A} = U^N \upharpoonright_{\mathbf{v}}$ for some $N \in \mathbf{Mod}_{\mathbf{w}}(\mathbf{E})$ such that U^N is \mathbf{v} -closed. Set

$$M \stackrel{\text{def}}{=} \{s^N(\vec{a}) \mid s \in T_{\lambda}(\mathbf{w}), \vec{a} \in \text{var}(s)A\}.$$

Since some operations of \mathbf{v} may have arity λ or larger, we cannot ensure that M is \mathbf{v} -closed within N *a priori*. However, picking $o \in A$, we may define an alternate interpretation of each $f \in \mathbf{w}_{\text{ope}}$ on M by setting $f^M(x) \stackrel{\text{def}}{=} f^N(x)$ if either there are $s = f(s_{\xi} \mid \xi < \text{ar}(f)) \in T_{\lambda}(\mathbf{w})$ and $\vec{a} \in \text{var}(s)A$ with each $x_{\xi} = s_{\xi}^N(\vec{a})$, in which case $f^N(x) = s^N(\vec{a}) \in M$, or $x \in \text{ar}(f)A$, in which case $f^N(x) = f^A(x) \in A$; and $f^M(x) \stackrel{\text{def}}{=} o$ in all other cases. Setting $R^M \stackrel{\text{def}}{=} R^N$ whenever $R \in \mathbf{w}_{\text{rel}}$, M is now a (λ, \mathbf{U}) -distension of \mathbf{A} . Since $s^M(\vec{a}) = s^N(\vec{a})$ whenever $s \in T_{\lambda}(\mathbf{w})$ and $\vec{a} \in \text{var}(s)A$, and since M is a \mathbf{w}_{rel} -substructure of N , every universal $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentence satisfied by N is also satisfied by M . Since $N \models \mathbf{E}$, it follows that $M \models \mathbf{E}$. \square Claim.

Now set $\mathcal{C}' \stackrel{\text{def}}{=} \{\mathbf{A}' \upharpoonright_{\mathbf{v}} \mid \mathbf{A}' \in \mathbf{Mod}_{\tilde{\mathbf{w}}}(\tilde{\mathbf{E}} \wedge \mathbf{H})\}$. In order to complete the proof of Theorem 6.8 it suffices to verify that $\mathcal{C} = \mathcal{C}'$.

By the Claim above, every $\mathbf{A} \in \mathcal{C}$ is $U^M \upharpoonright_{\mathbf{v}}$ for some (λ, \mathbf{U}) -distension M of \mathbf{A} . By Lemma 6.5, M has a right conjugate \mathbf{A}' . By Lemma 6.7, $\mathbf{A}' \models \tilde{\mathbf{E}}$. By Lemma 6.4, $\mathbf{A}' \models \mathbf{H}$. Hence, $\mathbf{A} \in \mathcal{C}'$, thus completing the proof that $\mathcal{C} \subseteq \mathcal{C}'$.

Conversely, every $\mathbf{A} \in \mathcal{C}'$ is $\mathbf{A}' \upharpoonright_{\mathbf{v}}$ for some $\mathbf{A}' \in \mathbf{Mod}_{\tilde{\mathbf{w}}}(\tilde{\mathbf{E}} \wedge \mathbf{H})$. By Lemma 6.5, \mathbf{A}' has a left conjugate M , which is necessarily a (λ, \mathbf{U}) -distension of \mathbf{A} . By Lemma 6.7, $M \models \mathbf{E}$. Hence, $\mathbf{A} = U^M \upharpoonright_{\mathbf{v}} \in \mathcal{C}$, thus completing the proof that $\mathcal{C}' \subseteq \mathcal{C}$. \square

Now that we know that $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$, we can take into account the projectivity of the class \mathcal{C} in Theorem 5.5. We obtain an additional faithfulness information on Φ .

Theorem 6.9. *Let λ be a regular cardinal, let \mathbf{v} be a λ -ary vocabulary, and let \mathcal{C} be an $\text{RPC}(\mathcal{L}_{\infty\lambda})$ class (equivalently, a $\text{PC}(\mathcal{L}_{\infty\lambda})$ class) of \mathbf{v} -structures. Then there are a λ -accessible category \mathcal{S} and a faithful λ -continuous functor $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ such that $\mathcal{C} = \text{im } \Phi = \text{rng } \Phi$. Moreover, \mathcal{S} can be taken a full subcategory, closed*

under λ -directed colimits, of $\mathbf{Str}(\mathbf{w})$ with \mathbf{w} -embeddings, for some vocabulary \mathbf{w} extending \mathbf{v} with $\mathbf{w} \setminus \mathbf{v}$ λ -ary.

Proof. We imitate the proof of Theorem 5.5, incorporating the PC = RPC information. By definition, there are a λ -ary vocabulary \mathbf{w} containing \mathbf{v} and a sentence E of $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ such that $\mathcal{C} = \{\mathbf{N}|_{\mathbf{v}} \mid \mathbf{N} \in \mathbf{Mod}_{\mathbf{w}}(E)\}$. Define \mathcal{S} as in the proof of Theorem 5.5, and let $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$, $\mathbf{N} \mapsto \mathbf{N}|_{\mathbf{v}}$ be the forgetful functor. As in the proof of Theorem 5.5, \mathcal{S} is λ -accessible, Φ is λ -continuous, and $\mathcal{C} = \text{rng } \Phi = \text{im } \Phi$. Moreover, the functor Φ is obviously faithful. \square

7. RPC CLASSES AS IMAGES OF CONTINUOUS FUNCTORS

The main aim of this section is to provide a converse to the results of Section 5, by proving (cf. Theorem 7.1) that the image of any λ -accessible functor $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$, for a λ -accessible category \mathcal{S} and a λ -ary vocabulary \mathbf{v} , is $\text{RPC}(\mathcal{L}_{\infty\lambda})$ (thus, due to Theorem 6.8, $\text{PC}(\mathcal{L}_{\infty\lambda})$). We prove something a bit more general.

Theorem 7.1. *Let λ be a regular cardinal, let \mathbf{v} be a λ -ary vocabulary, let \mathcal{S} be a category, and let $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ be a functor. We suppose that \mathcal{S} has a λ -directed colimit-dense small subcategory \mathcal{S}^\dagger , such that all λ -directed colimits from \mathcal{S}^\dagger exist in \mathcal{S} and are preserved by Φ . Then the image of Φ is $\text{RPC}(\mathcal{L}_{\infty\lambda})$ -definable, via an extension of \mathbf{v} by a unary predicate U together with a collection of binary predicates. Hence it is also $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable.*

Proof. Our assumptions imply immediately that $\text{im } \Phi$ is the class of all \mathbf{v} -structures \mathbf{M} for which there are a commutative diagram $(S_{p_1}, \sigma_{p_1, p_2} \mid p_1 \leq p_2 \text{ in } P)$ in \mathcal{S}^\dagger , with P λ -directed, and a colimit cocone

$$(\mathbf{M}, \sigma_p \mid p \in P) = \varinjlim (\Phi(S_{p_1}), \Phi(\sigma_{p_1, p_2}) \mid p_1 \leq p_2 \text{ in } P) \quad \text{within } \mathbf{Str}(\mathbf{v}). \quad (7.1)$$

To that end, let the vocabulary \mathbf{w} consist of \mathbf{v} together with the following collection of additional relation symbols:

- a unary predicate U and a binary relation symbol \leq ;
- binary relation symbols D_φ , indexed by morphisms $\varphi \in \mathcal{S}^\dagger$;
- binary relation symbols $F_{S, u}$, for $S \in \text{Ob } \mathcal{S}^\dagger$ and $u \in |\Phi(S)|$.

Let us describe the semantic interpretation of each of those additional relation symbols, and write down the corresponding sentence in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$.

- U should be interpreted by the original model \mathbf{M} , and its complement (meant as the P in (7.1)) should be a λ -directed poset. The required $\mathcal{L}_{\infty\lambda}$ -sentences are thus:

$$U \text{ is nonempty, and closed under all operations in } \mathbf{v}; \quad (7.2)$$

$$\leq \text{ is a } \lambda\text{-directed partial ordering on the complement of } U. \quad (7.3)$$

Let us, from now on, use the abbreviations $\mathbf{P}(\mathbf{p})$, $(\forall^U \mathbf{x})\mathbf{E}$, $(\exists^U \mathbf{x})\mathbf{E}$, $(\forall^{\mathbf{P}} \mathbf{p})\mathbf{E}$, $(\exists^{\mathbf{P}} \mathbf{p})\mathbf{E}$, for $\neg U(\mathbf{p})$, $(\forall \mathbf{x})(U(\mathbf{x}) \Rightarrow \mathbf{E})$, $(\exists \mathbf{x})(U(\mathbf{x}) \wedge \mathbf{E})$, $(\forall \mathbf{p})(\neg U(\mathbf{p}) \Rightarrow \mathbf{E})$, and $(\exists \mathbf{p})(\neg U(\mathbf{p}) \wedge \mathbf{E})$ respectively.

- The relation symbols D_φ , for $\varphi \in \mathcal{S}^\dagger$, capture the labeling of the commutative diagram $(S_{p_1}, \sigma_{p_1, p_2} \mid p_1 \leq p_2 \text{ in } P)$. For $p, q \in P$, the relation $D_\varphi(p, q)$ is intended to express the statement $\varphi = \sigma_{p, q}$. The required

$\mathcal{L}_{\infty\lambda}$ -sentences are thus:

$$(\forall^P \mathbf{p}, \mathbf{q})(D_\varphi(\mathbf{p}, \mathbf{q}) \Rightarrow \mathbf{p} \leq \mathbf{q}); \quad (7.4)$$

$$(\forall^P \mathbf{p}, \mathbf{q})\left(\mathbf{p} \leq \mathbf{q} \Rightarrow \bigvee_{\varphi \in \mathcal{S}^\dagger} D_\varphi(\mathbf{p}, \mathbf{q})\right); \quad (7.5)$$

$$(\forall^P \mathbf{p}, \mathbf{q})\neg(D_{\varphi_1}(\mathbf{p}, \mathbf{q}) \wedge D_{\varphi_2}(\mathbf{p}, \mathbf{q})), \quad \text{whenever } \varphi_1 \neq \varphi_2; \quad (7.6)$$

$$(\forall^P \mathbf{p}, \mathbf{q})\left(D_\varphi(\mathbf{p}, \mathbf{q}) \Rightarrow (D_{\mathbf{d}(\varphi)}(\mathbf{p}, \mathbf{p}) \wedge D_{\mathbf{r}(\varphi)}(\mathbf{q}, \mathbf{q}))\right); \quad (7.7)$$

$$(\forall^P \mathbf{p}, \mathbf{q}, \mathbf{r})\left((D_\varphi(\mathbf{p}, \mathbf{q}) \wedge D_\psi(\mathbf{q}, \mathbf{r})) \Rightarrow D_{\psi \circ \varphi}(\mathbf{p}, \mathbf{r})\right). \quad (7.8)$$

- The relation symbols $F_{S,u}$, for $S \in \text{Ob } \mathcal{S}^\dagger$ and $u \in |\Phi(S)|$, capture the representation of \mathbf{M} as the colimit in (7.1). More precisely, the relation $F_{S,u}(p, x)$ is intended to express the conjunction of $p \in P$, $S = S_p$, $u \in |\Phi(S)|$, and $x = \sigma_p(u)$. The corresponding $\mathcal{L}_{\infty\lambda}$ -statements are then obtained by translating the conditions in Lemma 2.1:

$$(\forall^P \mathbf{p})(\forall \mathbf{x})(F_{S,u}(\mathbf{p}, \mathbf{x}) \Rightarrow U(\mathbf{x})); \quad (7.9)$$

$$(\forall^P \mathbf{p})(D_S(\mathbf{p}, \mathbf{p}) \Leftrightarrow (\exists \mathbf{x})F_{S,u}(\mathbf{p}, \mathbf{x})); \quad (7.10)$$

$$(\forall^P \mathbf{p})(\forall \mathbf{x}, \mathbf{y})\left((F_{S,u}(\mathbf{p}, \mathbf{x}) \wedge F_{S,u}(\mathbf{p}, \mathbf{y})) \Rightarrow \mathbf{x} = \mathbf{y}\right); \quad (7.11)$$

$$(\forall^P \mathbf{p}, \mathbf{q})(\forall \mathbf{x})\left((F_{\mathbf{d}(\varphi),u}(\mathbf{p}, \mathbf{x}) \wedge D_\varphi(\mathbf{p}, \mathbf{q})) \Rightarrow F_{\mathbf{r}(\varphi),\Phi(\varphi)(u)}(\mathbf{q}, \mathbf{x})\right) \quad (7.12)$$

$$(\forall^U \mathbf{x})(\exists^P \mathbf{p}) \bigvee_{S \in \text{Ob } \mathcal{S}^\dagger, u \in |\Phi(S)|} F_{S,u}(\mathbf{p}, \mathbf{x}); \quad (7.13)$$

$$(\forall^P \mathbf{p})(\forall_{(\text{ar}(\mathbf{R}))} \vec{x})\left(\bigwedge_{\beta \in \text{ar}(\mathbf{R})} F_{S,u_\beta}(\mathbf{p}, x_\beta) \Rightarrow R(x_\beta \mid \beta \in \text{ar}(\mathbf{R}))\right) \\ \text{for } \mathbf{R} \in \mathbf{v}_{\text{rel}}, \text{ provided } (u_\beta \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^{\Phi(S)}; \quad (7.14)$$

$$(\forall^P \mathbf{p})(\forall_{(\text{ar}(\mathbf{f}))} \vec{x})(\forall \mathbf{x})\left((F_{S,u}(\mathbf{p}, \mathbf{x}) \wedge \bigwedge_{\beta \in \text{ar}(\mathbf{f})} F_{S,u_\beta}(\mathbf{p}, x_\beta)) \Rightarrow \right. \\ \left. \mathbf{x} = \mathbf{f}(x_\beta \mid \beta \in \text{ar}(\mathbf{f}))\right) \\ \text{for } \mathbf{f} \in \mathbf{v}_{\text{ope}}, \text{ provided } u = \mathbf{f}^{\Phi(S)}(u_\beta \mid \beta \in \text{ar}(\mathbf{f})); \quad (7.15)$$

$$(\forall^P \mathbf{p})(\forall \mathbf{x})\left((F_{S,u_1}(\mathbf{p}, \mathbf{x}) \wedge F_{S,u_2}(\mathbf{p}, \mathbf{x})) \Rightarrow \bigvee_{\substack{\varphi: S \rightarrow T \text{ within } \mathcal{S}^\dagger, \\ \Phi(\varphi)(u_1) = \Phi(\varphi)(u_2) = u}} (\exists^P \mathbf{q})D_\varphi(\mathbf{p}, \mathbf{q})\right); \quad (7.16)$$

$$\begin{aligned}
(\forall^P \mathbf{p})(\forall_{(\text{ar}(\mathbf{R}))} \vec{x}) & \left(\left(\bigwedge_{\beta \in \text{ar}(\mathbf{R})} F_{S, u_\beta}(\mathbf{p}, \mathbf{x}_\beta) \wedge R(\mathbf{x}_\beta \mid \beta \in \text{ar}(\mathbf{R})) \right) \Rightarrow \right. \\
& \bigvee_{\substack{\varphi: S \rightarrow T \text{ within } \mathcal{S}^\dagger, \\ (\Phi(\varphi)(u_\beta) \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^{\Phi(T)}}} (\exists^P \mathbf{q}) D_\varphi(\mathbf{p}, \mathbf{q}) \Big) \\
& \text{for } \mathbf{R} \in \mathbf{v}_{\text{rel}}. \quad (7.17)
\end{aligned}$$

The conjunction \bigwedge , of all possible sentences (7.2)–(7.17), is an $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ -sentence. Therefore, in order to complete the proof that \mathcal{C} is $\text{RPC}(\mathcal{L}_{\infty\lambda})$, it suffices to verify that a \mathbf{v} -structure \mathbf{M} belongs to \mathcal{C} iff it is $\mathbf{U}^{\mathbf{N}} \downarrow_{\mathbf{v}}$ for some \mathbf{w} -structure \mathbf{N} satisfying \bigwedge .

Suppose first that $\mathbf{M} = \mathbf{U}^{\mathbf{N}} \downarrow_{\mathbf{v}}$ for some \mathbf{w} -structure \mathbf{N} satisfying \bigwedge . We shall construct a λ -directed colimit representation of the form (7.1). Since \mathbf{N} satisfies both (7.2) and (7.3), the interpretation \leq of \leq in \mathbf{N} is a λ -directed partial ordering on $P \stackrel{\text{def}}{=} N \setminus M$. Moreover, each $D_\varphi \stackrel{\text{def}}{=} D_\varphi^{\mathbf{N}}$ is contained in \leq (use (7.4)), whereas each $F_{S,u} \stackrel{\text{def}}{=} F_{S,u}^{\mathbf{N}}$ is contained in $P \times M$.

Let $p, q \in P$ with $p \leq q$. Due to (7.5) and (7.6), there are unique $\sigma_{p,q}, S_p, S_q \in \mathcal{S}^\dagger$ such that $(p, q) \in D_{\sigma_{p,q}}$, $(p, p) \in D_{S_p}$, and $(q, q) \in D_{S_q}$. Due to (7.7), both $S_p = \mathbf{d}(\sigma_{p,p})$ and $S_q = \mathbf{r}(\sigma_{q,q})$ are objects in \mathcal{S}^\dagger . Hence $\sigma_{p,q}: S_p \rightarrow S_q$. If $p = q$ then $\sigma_{p,q} = S_p$ is itself an identity in \mathcal{S}^\dagger . Whenever $p \leq q \leq r$ in P , it follows from (7.8) that $\sigma_{p,r} = \sigma_{q,r} \circ \sigma_{p,q}$. Therefore, $(S_p, \sigma_{p,q} \mid p \leq q \text{ in } P)$ is a commutative diagram.

Let $p \in P$ and set $S \stackrel{\text{def}}{=} S_p$; so $(p, p) \in D_S$. Let $u \in |\Phi(S)|$. Due to (7.10) and (7.11), there exists a unique $\sigma_p(u) \in M$ such that $(p, \sigma_p(u)) \in F_{S,u}$. This defines a map $\sigma_p: \Phi(S) \rightarrow M$.

We claim that σ_p is a \mathbf{v} -homomorphism. Let $\mathbf{R} \in \mathbf{v}_{\text{rel}}$, let $(u_\beta \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^{\Phi(S)}$, and set $x_\beta \stackrel{\text{def}}{=} \sigma_p(u_\beta)$ for each $\beta \in \text{ar}(\mathbf{R})$. Then each $(p, x_\beta) \in F_{S, u_\beta}$, thus, due to (7.14), $(x_\beta \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^M$. Likewise, let $\mathbf{f} \in \mathbf{v}_{\text{ope}}$ and let $u = \mathbf{f}^{\Phi(S)}(u_\beta \mid \beta \in \text{ar}(\mathbf{f}))$ within $\Phi(S)$. Set $x \stackrel{\text{def}}{=} \sigma_p(u)$ and $x_\beta \stackrel{\text{def}}{=} \sigma_p(u_\beta)$ for each $\beta \in \text{ar}(\mathbf{f})$. Then $(p, x) \in F_{\xi, S, u}$ whereas each $(p, x_\beta) \in F_{S, u_\beta}$. Due to (7.15), it follows that $x = \mathbf{f}^M(x_\beta \mid \beta \in \text{ar}(\mathbf{f}))$. This completes the proof of our claim that σ_p is a \mathbf{v} -homomorphism.

Let $p \leq q$ in P . Set $S \stackrel{\text{def}}{=} S_p$, $T \stackrel{\text{def}}{=} S_q$, and $\varphi \stackrel{\text{def}}{=} \sigma_{p,q}$. Let $u \in |\Phi(S)|$ and set $x \stackrel{\text{def}}{=} \sigma_p(u)$, $y \stackrel{\text{def}}{=} \sigma_q(\Phi(\varphi)(u))$. Then $(p, q) \in D_\varphi$ and $(p, x) \in F_{S,u}$. Due to (7.12), $(q, x) \in F_{T, \Phi(\varphi)(u)}$; that is, $x = y$. This completes the proof that $\sigma_p = \sigma_q \circ \Phi(\sigma_{p,q})$.

Therefore, $(\mathbf{M}, \sigma_p \mid p \in P)$ is a cocone above $(\Phi(S_p), \Phi(\sigma_{p,q}) \mid p \leq q \text{ in } P)$ within $\mathbf{Str}(\mathbf{v})$.

We shall now verify, using Lemma 2.1, that it is a colimit cocone.

Let $x \in M$. Due to (7.13), there are $p \in P$ and $u \in |\Phi(S_p)|$ such that $(p, x) \in F_{S_p, u}$; whence $x = \sigma_p(u)$. Therefore, $M = \bigcup \{ \sigma_p[\Phi(S_p)] \mid p \in P \}$.

Let $p \in P$, $S \stackrel{\text{def}}{=} S_p$, and $u_1, u_2 \in |\Phi(S)|$ such that $\sigma_p(u_1) = \sigma_p(u_2)$ — denote by x that element of M . Then (p, x) belongs to each F_{S, u_i} , thus, due to (7.16), there exists $q \in P$ such that $p \leq q$ and $\Phi(\sigma_{p,q})(u_1) = \Phi(\sigma_{p,q})(u_2)$.

Similarly, let $\mathbf{R} \in \mathbf{v}_{\text{rel}}$ and $(u_\beta \mid \beta \in \text{ar}(\mathbf{R})) \in \text{ar}(\mathbf{R})|\Phi(S)$ such that $(\sigma_p(u_\beta) \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^M$. Since each $(p, x_\beta) \in F_{S, u_\beta}$ and due to (7.17), there exists $q \in P$, with $p \leq q$, such that $(\sigma_{p,q}(u_\beta) \mid \beta \in \text{ar}(\mathbf{R})) \in \mathbf{R}^{\Phi(S_q)}$. Owing to

Lemma 2.1, this completes the proof that

$$(\Phi(S_p), \sigma_p \mid p \in P) = \varinjlim (\Phi(S_p), \Phi(\sigma_{p,q}) \mid p \leq q \text{ in } P).$$

Therefore, \mathbf{M} belongs to $\text{im } \Phi$.

Let, conversely, $\mathbf{M} \in \text{im } \Phi$, as witnessed by a λ -directed commutative diagram $(S_{p_1}, \sigma_{p_1, p_2} \mid p_1 \leq p_2 \text{ in } P)$ in \mathcal{S}^\dagger and a colimit cocone within \mathcal{S} as in (7.1). We may assume that $P \cap M = \emptyset$. Define the \mathbf{w} -structure \mathbf{N} , with universe $N \stackrel{\text{def}}{=} M \sqcup P$, as follows:

- $U^{\mathbf{N}} \stackrel{\text{def}}{=} M$;
- The binary relation symbol \leq is interpreted as the partial ordering \leq of P ;
- for every $R \in \mathbf{v}_{\text{rel}}$, $R^{\mathbf{N}} \stackrel{\text{def}}{=} R^M$;
- for every $f \in \mathbf{v}_{\text{ope}}$, $f^{\mathbf{N}}$ is any map from $\text{ar}(f)N$ to M extending f^M ;
- for every $\varphi \in \mathcal{S}^\dagger$, we set

$$D_\varphi \stackrel{\text{def}}{=} \{(p, q) \in P \times P \mid p \leq q \text{ and } \sigma_{p,q} = \varphi\};$$

- for every $S \in \text{Ob } \mathcal{S}^\dagger$ and $u \in |\Phi(S)|$, we set

$$F_{S,u} \stackrel{\text{def}}{=} \{(p, \sigma_p(u)) \mid p \in P, S = S_p\}.$$

We verify that those interpretations satisfy all instances of (7.2)–(7.17) within \mathbf{N} .

For (7.2)–(7.4) this is obvious. For (7.5) and (7.6) we must take $\varphi = \varphi_1 = \varphi_2 \stackrel{\text{def}}{=} \sigma_{p,q}$. For (7.7) we just observe that $\sigma_{p,q}$ is an arrow from $\sigma_{p,p} = S_p$ to $\sigma_{q,q} = S_q$. For (7.8), we apply the equality $\sigma_{p,r} = \sigma_{q,r} \circ \sigma_{p,q}$. For (7.9) this follows from the containment $F_{S,u} \subseteq P \times M$. For (7.10) and (7.11), just observe that each σ_p is a function defined on $|\Phi(S_p)|$. For (7.12), φ is necessarily $\sigma_{p,q}$ so we just need to apply the equality $\sigma_p = \sigma_q \circ \Phi(\sigma_{p,q})$. For (7.13) this follows from Lemma 2.1(i). The statements (7.14) and (7.15) follow from σ_p being a \mathbf{v} -homomorphism. Finally, (7.16) and (7.17) follows from items (ii) and (iii) of Lemma 2.1, respectively.

The final part of the statement of Theorem 7.1, that $\text{im } \Phi$ is $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable, now follows from Theorem 6.8. \square

Corollary 7.2. *Let λ be a regular cardinal and let \mathbf{v} be a λ -ary vocabulary. Then for any small subcategory \mathcal{T} of $\mathbf{Str}(\mathbf{v})$, the class $\mathcal{T}_{(\lambda)}$ of all colimits of λ -directed commutative diagrams from \mathcal{T} is $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable.*

Proof. View $\mathcal{T}_{(\lambda)}$ as a full subcategory of $\mathbf{Str}(\mathbf{v})$ and apply Theorem 7.1 to the inclusion functor from $\mathcal{S} \stackrel{\text{def}}{=} \mathcal{T}_{(\lambda)}$ into $\mathbf{Str}(\mathbf{v})$, with $\mathcal{S}^\dagger \stackrel{\text{def}}{=} \mathcal{T}$. \square

Note that the objects of \mathcal{T} may not be λ -presentable in $\mathbf{Str}(\mathbf{v})$. Besides, in contrast to the context of Adámek and Rosický [1, Theorem 2.26], $\mathcal{T}_{(\lambda)}$ is only a class of objects (as opposed to a category).

Examples 7.3. Consider the following small subcategories $\mathcal{S}_0, \mathcal{S}_1, \mathcal{S}_2$ of the category $\mathbf{Set}_{\neq \emptyset}$ of all nonempty sets (viewed as $\mathbf{Str}(\emptyset)$):

- \mathcal{S}_0 is the category of all finite subsets of ω with all inclusion maps;
- \mathcal{S}_1 is the category of all finite subsets of ω with all one-to-one maps;
- \mathcal{S}_2 is the category with the unique object ω and its identity map.

Then $(\mathcal{S}_0)_{(\omega)}$ is the class of all at most countable sets, $(\mathcal{S}_1)_{(\omega)}$ is the class of all sets, and $(\mathcal{S}_2)_{(\omega)}$ is the class of all infinite sets.

Corollary 7.4. *Let λ be a regular cardinal, let \mathbf{v} be a λ -ary vocabulary, let \mathcal{S} be a λ -accessible category, and let $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ be a λ -continuous functor. Then the image of Φ is $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable.*

Proof. Apply Theorem 7.1 with $\mathcal{S}^\dagger \stackrel{\text{def}}{=} \mathbf{Pres}_\lambda \mathcal{S}$. □

Bringing together Theorem 5.5 and Corollary 7.4, we obtain the following.

Theorem 7.5. *Let λ be a regular cardinal and let \mathbf{v} be a λ -ary vocabulary. For any class \mathcal{C} of \mathbf{v} -structures, the following are equivalent:*

- (i) \mathcal{C} is $\text{PC}(\mathcal{L}_{\infty\lambda})$;
- (ii) \mathcal{C} is $\text{RPC}(\mathcal{L}_{\infty\lambda})$;
- (iii) *there are a λ -accessible category \mathcal{S} and a λ -continuous functor $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ such that $\mathcal{C} = \text{im } \Phi$.*

Although, by Theorem 6.8, the condition that \mathbf{v} be λ -ary is not needed for the equivalence (i) \Leftrightarrow (ii) in Theorem 7.5, this is not the case for both implications (i) \Rightarrow (iii) and (iii) \Rightarrow (i) (cf. Examples 5.6 and 8.8, respectively).

Corollary 7.6. *Let \mathbf{v} be a vocabulary. For any class \mathcal{C} of \mathbf{v} -structures, the following are equivalent:*

- (i) \mathcal{C} is $\text{PC}(\mathcal{L}_{\infty\infty})$;
- (ii) \mathcal{C} is $\text{RPC}(\mathcal{L}_{\infty\infty})$;
- (iii) *there are an accessible category \mathcal{S} and a continuous functor $\Phi: \mathcal{S} \rightarrow \mathbf{Str}(\mathbf{v})$ such that $\mathcal{C} = \text{im } \Phi$.*

Proof. By the Uniformization Theorem for accessible categories and functors (Adámek and Rosický [1, 2.19]; see also Proposition 2.3.5 and Theorem 2.3.10 in Makkai and Paré [20]), for any accessible category \mathcal{S} and any continuous functor Φ from \mathcal{S} to some accessible category, there are arbitrarily large cardinals κ such that \mathcal{S} is a κ^+ -accessible category and Φ is a κ^+ -accessible functor. Apply Theorem 7.5. □

8. THE CASE OF SINGULAR CARDINALS

Due to the regularity requirement on the infinite cardinal λ in Lemma 6.7, the argument of the proof of Theorem 6.8 does not extend to the case where λ is singular. The latter will require a completely different approach, based on the following simple observation of cardinal arithmetic. This will lead to Theorem 8.6, which, together with Theorem 6.8, shows that $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$ for any infinite cardinal λ (regardless of whether λ is regular).

Lemma 8.1. *The following statements hold, for every infinite cardinal λ :*

- (1) *If λ is regular, then there are arbitrarily large cardinals κ such that $\kappa^\lambda < \kappa^\lambda$.*
- (2) *If λ is singular, then for every cardinal $\kappa \geq 2^\lambda$ there exists $\alpha < \lambda$ such that $\kappa^\lambda = \kappa^\alpha$; hence $\kappa^\lambda = \kappa^\lambda$.*

Proof. *Ad (1).* Let α be an ordinal and set $\kappa \stackrel{\text{def}}{=} \beth_{\alpha+\lambda}$ (we are using the standard notation for Beth numbers). Then $\kappa^\lambda = \kappa$ has cofinality λ . On the other hand (cf. Jech [14, Corollary 5.13]), $\text{cf}(\kappa^\lambda) > \lambda$; whence $\kappa^\lambda < \kappa^\lambda$.

Ad (2). The conclusion is obvious for $\kappa = 2^\lambda$ (take $\alpha = 1$) so we may assume that $\kappa > 2^\lambda$. The least cardinal μ such that $\kappa^\lambda = \mu^\lambda$ satisfies $\mu > 2^\lambda$ as well. Note

that $\mu > \lambda$, and $\bar{\mu}^\lambda < \mu$ whenever $\bar{\mu} < \mu$ (otherwise $\mu \leq \bar{\mu}^\lambda$ thus $\kappa^\lambda = \mu^\lambda \leq \bar{\mu}^\lambda$ and thus $\kappa^\lambda = \bar{\mu}^\lambda$, a contradiction).

If $\text{cf } \mu > \lambda$, then, using Jech [14, Theorem 5.20(iii,a)], $\kappa^\lambda = \mu^\lambda = \mu \leq \kappa$ thus $\kappa^\lambda = \kappa$ and we are done.

Suppose that $\text{cf } \mu \leq \lambda$. Since λ is singular, $\alpha \stackrel{\text{def}}{=} \text{cf } \mu < \lambda$. Moreover, using Jech [14, Theorem 5.20(iii,b)], $\kappa^\lambda = \mu^\lambda = \mu^\alpha \leq \kappa^\alpha$; whence $\kappa^\lambda = \kappa^\alpha$. \square

Definition 8.2. A *multiterm*, in a vocabulary \mathbf{v} , is a family $\mathbf{t} = (\mathbf{t}_j \mid j \in J)$ of terms of \mathbf{v} . If each \mathbf{t}_j has all its variables in a set I , we will write $\mathbf{t}: {}^I\mathbf{M} \rightarrow {}^J\mathbf{M}$. Moreover, for any \mathbf{v} -structure \mathbf{M} , we will denote $\mathbf{t}^{\mathbf{M}}: {}^I\mathbf{M} \rightarrow {}^J\mathbf{M}$, $x \mapsto (\mathbf{t}_j^{\mathbf{M}}(x) \mid j \in J)$, and call $\mathbf{t}^{\mathbf{M}}$ a *multiterm function* on \mathbf{M} . Multiterms $\mathbf{s}: {}^I\mathbf{M} \rightarrow {}^J\mathbf{M}$ and $\mathbf{t}: {}^J\mathbf{M} \rightarrow {}^K\mathbf{M}$ may be composed to a multiterm $\mathbf{t} \circ \mathbf{s}: {}^I\mathbf{M} \rightarrow {}^K\mathbf{M}$ in the natural way, that respects the composition of the associated multiterm functions.

Lemma 8.3. For all cardinals α and λ , with λ singular and $0 < \alpha < \lambda$, there are a λ -ary vocabulary \mathbf{b} , consisting of operations only, and multiterms $\mathbf{g}: {}^\alpha\mathbf{M} \rightarrow {}^\lambda\mathbf{M}$ and $\mathbf{h}: {}^\lambda\mathbf{M} \rightarrow {}^\alpha\mathbf{M}$ of \mathbf{b} , such that every set M with $(\text{card } M)^\alpha = (\text{card } M)^\lambda$ expands to a \mathbf{b} -structure \mathbf{M} such that \mathbf{M} satisfies both statements $(\forall \vec{x})_{(\alpha)} ((\mathbf{h} \circ \mathbf{g})(\vec{x}) = \vec{x})$ and $(\forall \vec{y})_{(\lambda)} ((\mathbf{g} \circ \mathbf{h})(\vec{y}) = \vec{y})$ (that we shall from now on abbreviate by $\mathbf{h} \circ \mathbf{g} = \text{id}$ and $\mathbf{g} \circ \mathbf{h} = \text{id}$, respectively).

Note. Setting $\kappa \stackrel{\text{def}}{=} \text{card } M$ and by virtue of the well known identity $\kappa^\lambda = (\kappa^\lambda)^{\text{cf } \lambda}$, $\kappa^\alpha = \kappa^\lambda$ is equivalent to $\kappa^\alpha = \kappa^\lambda$, that is, $\kappa^\alpha = \kappa^\beta$ whenever $\alpha \leq \beta < \lambda$. Also, without the restriction that \mathbf{b} be λ -ary, Lemma 8.3 would be trivial.

Proof. We begin with the case where $\text{cf } \lambda \leq \alpha \leq \lambda$. Let $\lambda = \sum_{i < \text{cf } \lambda} \lambda_i$ where each $\alpha \leq \lambda_i < \lambda$. Let \mathbf{b} consist of operation symbols $\mathbf{f}_{i,j}$ and $\mathbf{f}'_{j,i}$, for $i < \text{cf } \lambda$ and $j < \alpha$, where each $\text{ar}(\mathbf{f}_{i,j}) = \alpha$ and $\text{ar}(\mathbf{f}'_{j,i}) = \lambda_i$. Moreover, pick a bijection $(\text{cf } \lambda) \times \alpha \rightarrow \alpha$, $(i, k) \mapsto \langle i, k \rangle$, with inverse $\alpha \rightarrow (\text{cf } \lambda) \times \alpha$, $k \mapsto (p(k), q(k))$. We let

$$\mathbf{g}_j(\vec{x})_{(\alpha)} \stackrel{\text{def}}{=} \mathbf{f}_{i,\bar{j}}(x_{\langle i,k \rangle} \mid k < \alpha), \text{ whenever } j = \sum_{i' < i} \lambda_{i'} + \bar{j} \text{ with } \bar{j} < \lambda_i;$$

$$\mathbf{g}(\vec{x})_{(\alpha)} \stackrel{\text{def}}{=} (\mathbf{g}_j(\vec{x}) \mid j < \lambda);$$

$$\mathbf{h}_k(\vec{y})_{(\lambda)} \stackrel{\text{def}}{=} \mathbf{f}'_{q(k),p(k)}(y_{\sum_{i' < p(k)} \lambda_{i'} + \bar{j}} \mid \bar{j} < \lambda_{p(k)}), \text{ whenever } k < \alpha;$$

$$\mathbf{h}(\vec{y})_{(\lambda)} \stackrel{\text{def}}{=} (\mathbf{h}_k(\vec{y}) \mid k < \alpha).$$

Let us define interpretations $f_{i,j}$ and $f'_{j,i}$, of the symbols $\mathbf{f}_{i,j}$ and $\mathbf{f}'_{j,i}$, on any set M , with cardinality κ , such that $\kappa^\alpha = \kappa^\lambda$. For each $i < \text{cf } \lambda$, since $\kappa^\alpha = \kappa^{\lambda_i}$, there are mutually inverse bijections $f_i: {}^\alpha M \rightarrow {}^{\lambda_i} M$, $x \mapsto (f_{i,j}(x) \mid j < \lambda_i)$ and $f'_i: {}^{\lambda_i} M \rightarrow {}^\alpha M$, $y \mapsto (f'_{j,i}(y) \mid j < \alpha)$. For each $x \in {}^\alpha M$, we define $g(x)$ as the concatenation of all $f_i(x_{\langle i,j \rangle} \mid j < \alpha)$ for $i < \text{cf } \lambda$. Hence g is a bijection from ${}^\alpha M$ onto ${}^\lambda M$, and for each $x \in {}^\alpha M$, $g(x) = (g_j(x) \mid j < \alpha)$ where, for $j = \sum_{i' < i} \lambda_{i'} + \bar{j}$ with $\bar{j} < \lambda_i$, $g_j(x) = f_{i,\bar{j}}(x_{\langle i,k \rangle} \mid k < \alpha)$.

For each $x \in {}^\alpha M$ and $i < \text{cf } \lambda$, set $x_{(i)} \stackrel{\text{def}}{=} (x_{\langle i,j \rangle} \mid j < \alpha)$. Set $h = g^{-1} = (h_k \mid k < \alpha)$. Let $y \in {}^\lambda M$ and $x = h(y)$. Then for each $i < \text{cf } \lambda$, $f_i(x_{(i)}) =$

$(y_{\sum_{i' < i} \lambda_{i'} + \bar{j}} \mid \bar{j} < \lambda_i)$, thus $x_{(i)} = f'_i(y_{\sum_{i' < i} \lambda_{i'} + \bar{j}} \mid \bar{j} < \lambda_i)$. It follows that each $h_k(y) = x_k = x_{(p(k))}(q(k)) = f'_{q(k), p(k)}(y_{\sum_{i' < p(k)} \lambda_{i'} + \bar{j}} \mid \bar{j} < \lambda_{p(k)})$.

If $0 < \alpha < \text{cf } \lambda$, then $\kappa^\alpha = \kappa^\lambda$ entails $\kappa^{\text{cf } \lambda} = \kappa^\alpha = \kappa^\lambda$, thus it suffices to compose the multiterms associated to the latter equation, *via* the argument above, to a pair of multiterms giving rise to mutually inverse bijections between κ^α and $\kappa^{\text{cf } \lambda}$. \square

Let α and λ be cardinals, with λ singular and $0 < \alpha < \lambda$, and let \mathbf{v} and \mathbf{w} be vocabularies with $\mathbf{v} \subseteq \mathbf{w}$ and $\mathbf{w} \setminus \mathbf{v}$ consisting only of relations with arities ordinals smaller than or equal to λ . Let \mathbf{b} be a λ -ary vocabulary, which we may assume to be disjoint from \mathbf{w} , with terms \mathbf{g} and \mathbf{f} satisfying the conclusion of Lemma 8.3.

Denote⁸ by $\mathbf{w}|\alpha$ the vocabulary with the same symbols and arities as \mathbf{w} , except for the relation symbols $R \in \mathbf{w} \setminus \mathbf{v}$ with $\text{ar}(R) = \lambda$, each of which is replaced by a new relation symbol R_α with arity α .

To any formula E in $\mathcal{L}_{\infty\lambda^+}(\mathbf{w} \cup \mathbf{b})$ we associate the collection of its (\mathbf{g}, \mathbf{h}) -contractions (with respect to the pair (\mathbf{g}, \mathbf{h})), which are *preformulas* of $\mathcal{L}_{\infty\lambda}((\mathbf{w}|\alpha) \cup \mathbf{b})$ with the same free variables as E , by the following inductive process:

- If E is atomic, then it is its only (\mathbf{g}, \mathbf{h}) -contraction, except in case $E = R(\vec{t})$ for a relation symbol $R \in \mathbf{w} \setminus \mathbf{v}$ with arity λ , in which case the only (\mathbf{g}, \mathbf{h}) -contraction of E is $R_\alpha(\mathbf{h}(\vec{t}))$.
- If E is $\neg F$, then its (\mathbf{g}, \mathbf{h}) -contractions are the $\neg F'$ for (\mathbf{g}, \mathbf{h}) -contractions F' of F .
- If E is $\bigvee_{i \in I} E_i$, then its (\mathbf{g}, \mathbf{h}) -contractions are the $\bigvee_{i \in I} E'_i$ for (\mathbf{g}, \mathbf{h}) -contractions E'_i of E_i .
- If E is $(\exists Y)F$, where $Y \subseteq \text{fvar}(F)$ has less than λ elements, then the (\mathbf{g}, \mathbf{h}) -contractions of E are the $(\exists Y)F'$ for (\mathbf{g}, \mathbf{h}) -contractions F' of F .
- If E is $(\exists Y)F$, where $Y \subseteq \text{fvar}(F)$ has λ elements, then the (\mathbf{g}, \mathbf{h}) -contractions of E are all formulas of the form $(\exists_{(\alpha)} \vec{z})F'(y_j \mapsto g_j(\vec{z}) \mid j < \lambda)$ (with free variables of E being assigned to themselves), for one-to-one enumerations $(y_j \mid j < \lambda)$ of Y , with all $z_i \in Y$.

Although a formula usually has more than one (\mathbf{g}, \mathbf{h}) -contraction, all its (\mathbf{g}, \mathbf{h}) -contractions are logically equivalent.

Every \mathbf{w} -structure \mathbf{M} such that $(\text{card } M)^\alpha = (\text{card } M)^\lambda$ has a $(\mathbf{w} \cup \mathbf{b})$ -expansion, which we shall still denote by \mathbf{M} , such that the interpretations $g \stackrel{\text{def}}{=} \mathbf{g}^{\mathbf{M}}$ and $h \stackrel{\text{def}}{=} \mathbf{h}^{\mathbf{M}}$ are mutually inverse. The (\mathbf{g}, \mathbf{h}) -contraction of \mathbf{M} is the $((\mathbf{w}|\alpha) \cup \mathbf{b})$ -structure on M , where all symbols in $\mathbf{w} \cup \mathbf{b}$ which are not relation symbols in $\mathbf{w} \setminus \mathbf{v}$ with arity λ have the same interpretations in \mathbf{M} and \mathbf{M}' , and for every relation symbol $R \in \mathbf{w} \setminus \mathbf{v}$ with arity λ ,

$$R_\alpha^{\mathbf{M}'} \stackrel{\text{def}}{=} \{\vec{x} \in {}^\alpha M \mid \mathbf{M} \models R\mathbf{g}(\vec{x})\} = h[R^{\mathbf{M}}]. \quad (8.1)$$

Lemma 8.4. *Every $((\mathbf{w}|\alpha) \cup \mathbf{b})$ -structure \mathbf{M}' , satisfying both statements $\mathbf{h} \circ \mathbf{g} = \text{id}$ and $\mathbf{g} \circ \mathbf{h} = \text{id}$, is the (\mathbf{g}, \mathbf{h}) -contraction of a unique $(\mathbf{w} \cup \mathbf{b})$ -structure \mathbf{M} satisfying those statements.*

Proof. For any relation symbol $R \in \mathbf{w} \setminus \mathbf{v}$ with arity λ , $R^{\mathbf{M}}$ can be recovered from $R_\alpha^{\mathbf{M}'}$ *via* (8.1): namely, $R^{\mathbf{M}} = g[R_\alpha^{\mathbf{M}'}]$. \square

⁸This notation is slightly overloaded: indeed, $\mathbf{w}|\alpha$ also depends of \mathbf{v} .

Lemma 8.5. *Let M be a $(\mathbf{w} \cup \mathbf{b})$ -structure satisfying both statements $\mathbf{h} \circ \mathbf{g} = \text{id}$ and $\mathbf{g} \circ \mathbf{h} = \text{id}$, and let M' be the (\mathbf{g}, \mathbf{h}) -contraction of M . Then for every $E \in \mathcal{L}_{\infty\lambda+}(\mathbf{w} \cup \mathbf{b})$, every (\mathbf{g}, \mathbf{h}) -contraction E' of E , and every $\vec{a} \in \text{var}^{(E)}M$, $M \models E(\vec{a})$ iff $M' \models E'(\vec{a})$.*

Proof. By induction on the formula E . If E is atomic, then the result is trivial unless $E = R(\vec{t})$ where $R \in \mathbf{w} \setminus \mathbf{v}$ with arity λ and $\vec{t} = (t_\xi \mid \xi < \lambda)$ for terms t_ξ of $\mathbf{w} \cup \mathbf{b}$. Then $M' \models E'(\vec{a})$ iff $M' \models R_\alpha(\mathbf{h}(\vec{t}(\vec{a})))$, iff $M \models R(\mathbf{gh}(\vec{t}(\vec{a})))$ (because each $t_\xi^M(\vec{a}) = t_\xi^{M'}(\vec{a})$), iff $M \models R(\vec{t}(\vec{a}))$, that is, $M \models E(\vec{a})$. The induction steps corresponding to disjunction and negation are trivial.

The only nontrivial remaining step is thus the one where E is $(\exists Y)F$ where Y is a set of free variables of F with cardinality λ . Any (\mathbf{g}, \mathbf{h}) -contraction E' of E has the form $(\exists \vec{z})F'(y_j \mapsto \mathbf{g}_j(\vec{z}) \mid j < \lambda)$, for a (\mathbf{g}, \mathbf{h}) -contraction F' of F , a one-to-one enumeration $(y_j \mid j < \lambda)$ of Y , and all $z_i \in Y$. Hence $M' \models E'(\vec{a})$ iff there exists $\vec{c} \in {}^\alpha M$ such that $M' \models F'(y_j \mapsto \mathbf{g}_j(\vec{c}) \mid j < \lambda)$. Since g is surjective, $(g_j(\vec{c}) \mid j < \lambda)$ can take any value in ${}^\lambda M$, so $M' \models E'(\vec{a})$ iff $M' \models F'(\vec{a}, \vec{b})$ for some $\vec{b} \in {}^Y M$. By the induction hypothesis, this is equivalent to saying that $M \models F(\vec{a}, \vec{b})$ for some $\vec{b} \in {}^Y M$; that is, $M \models E(\vec{a})$. \square

Theorem 8.6. *Let λ be a singular cardinal. Then $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{PC}(\mathcal{L}_{\infty\lambda+}) = \text{RPC}(\mathcal{L}_{\infty\lambda+}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$.*

Proof. We know from Theorem 6.8 that $\text{PC}(\mathcal{L}_{\infty\lambda+}) = \text{RPC}(\mathcal{L}_{\infty\lambda+})$. Hence, due to the trivial containments $\text{PC}(\mathcal{L}_{\infty\lambda}) \subseteq \text{RPC}(\mathcal{L}_{\infty\lambda}) \subseteq \text{RPC}(\mathcal{L}_{\infty\lambda+})$, it suffices to prove the containment $\text{PC}(\mathcal{L}_{\infty\lambda+}) \subseteq \text{PC}(\mathcal{L}_{\infty\lambda})$.

Let \mathcal{C} be a $\text{PC}(\mathcal{L}_{\infty\lambda+})$ -class over a vocabulary \mathbf{v} . By Theorem 4.11, there are a vocabulary \mathbf{w} containing \mathbf{v} , with each symbol in $\mathbf{w} \setminus \mathbf{v}$ of arity an ordinal smaller than or equal to λ , and a sentence $E \in \mathcal{L}_{\infty\lambda+}(\mathbf{w})$ such that $\mathcal{C} = \{N \upharpoonright_{\mathbf{v}} \mid N \in \text{Mod}_{\mathbf{w}}(E)\}$. Replacing every operation in $\mathbf{w} \setminus \mathbf{v}$ by a new relation symbol, viewed as its graph, and modifying E accordingly, we may assume that $\mathbf{w} \setminus \mathbf{v}$ consists of relation symbols only.

For every nonzero cardinal $\alpha < \lambda$, let \mathbf{b}_α be a λ -ary vocabulary, with terms \mathbf{g}_α and \mathbf{h}_α , satisfying the conclusion of Lemma 8.3. Any $(\mathbf{g}_\alpha, \mathbf{h}_\alpha)$ -contraction E_α of E is a preformula in $\mathcal{L}_{\infty\lambda}((\mathbf{w} \upharpoonright_\alpha) \cup \mathbf{b}_\alpha)$ with the same free variables as E (there are none); thus E_α is a sentence in $\mathcal{L}_{\infty\lambda}((\mathbf{w} \upharpoonright_\alpha) \cup \mathbf{b}_\alpha)$. We may assume that \mathbf{w} and the \mathbf{b}_α are mutually disjoint. We may also assume that the additional relation symbols R_α are all pairwise distinct and do not belong to $\mathbf{w} \cup \bigcup_{\alpha < \lambda} \mathbf{b}_\alpha$. The disjunction E^* , over all nonzero $\alpha < \lambda$, of all statements $(\mathbf{g}_\alpha \circ \mathbf{h}_\alpha = \text{id}) \wedge (\mathbf{h}_\alpha \circ \mathbf{g}_\alpha = \text{id}) \wedge E_\alpha$, is an $\mathcal{L}_{\infty\lambda}$ -statement over $\mathbf{w}^* \stackrel{\text{def}}{=} \bigcup_{0 < \alpha < \lambda} ((\mathbf{w} \upharpoonright_\alpha) \cup \mathbf{b}_\alpha)$. Hence, $\mathcal{C}^* \stackrel{\text{def}}{=} \{N \upharpoonright_{\mathbf{v}} \mid N \in \text{Mod}_{\mathbf{w}^*}(E^*)\}$ is a $\text{PC}(\mathcal{L}_{\infty\lambda})$ -definable class. Therefore, in order to conclude the proof, it suffices, by Lemma 3.2, to verify that $\mathcal{C}^* \subseteq \mathcal{C}$ and that $\mathcal{C} \setminus \mathcal{C}^*$ is small-abstract.

Let $M \in \mathcal{C}^*$. By definition, there are a nonzero cardinal $\alpha < \lambda$ and a \mathbf{w}^* -structure N^* , satisfying $(\mathbf{g}_\alpha \circ \mathbf{h}_\alpha = \text{id}) \wedge (\mathbf{h}_\alpha \circ \mathbf{g}_\alpha = \text{id}) \wedge E_\alpha$, such that $M = N^* \upharpoonright_{\mathbf{v}}$. By Lemma 8.4, $N^* \upharpoonright_{(\mathbf{w} \upharpoonright_\alpha) \cup \mathbf{b}_\alpha}$ is the $(\mathbf{g}_\alpha, \mathbf{h}_\alpha)$ -contraction of a unique $(\mathbf{w} \cup \mathbf{b}_\alpha)$ -structure N_α . Since $N^* \models E_\alpha$, it follows from Lemma 8.5 that $N_\alpha \models E$. Therefore, $M = N^* \upharpoonright_{\mathbf{v}} = N_\alpha \upharpoonright_{\mathbf{v}}$ belongs to \mathcal{C} . This completes the proof that $\mathcal{C}^* \subseteq \mathcal{C}$.

Let $M \in \mathcal{C}$ with cardinality $\kappa \geq 2^\lambda$. Then $M = N \upharpoonright_{\mathbf{v}}$ for some $N \in \text{Mod}_{\mathbf{w}}(E)$. By Lemma 8.1, there exists a nonzero cardinal $\alpha < \lambda$ such that $\kappa^\alpha = \kappa^\lambda$. By Lemma 8.3, N expands to a $(\mathbf{w} \cup \mathbf{b}_\alpha)$ -structure N_α satisfying both statements

$h_\alpha \circ g_\alpha = \text{id}$ and $g_\alpha \circ h_\alpha = \text{id}$. The (g_α, h_α) -contraction N'_α of N_α is a $((\mathbf{w}|\alpha) \cup \mathbf{b}_\alpha)$ -structure, thus it expands in turn to a \mathbf{w}^* -structure N^* . From $N \models E$ it follows that $N_\alpha \models E$, thus, by Lemma 8.5, $N'_\alpha \models E_\alpha$; that is, $N^* \models E_\alpha$. Therefore, $N^* \models E^*$, so $M = N^* \upharpoonright_{\mathbf{v}} \in \mathcal{C}^*$. This proves that $\mathcal{C} \setminus \mathcal{C}^*$ is contained in the small-abstract class $\{M \in \mathbf{Str}(\mathbf{v}) \mid \text{card } M < 2^\lambda\}$. \square

Bringing together Theorems 6.8 and 8.6, we obtain:

Corollary 8.7. $\text{PC}(\mathcal{L}_{\infty\lambda}) = \text{RPC}(\mathcal{L}_{\infty\lambda})$, for any infinite cardinal λ .

The influence of the regularity / singularity of λ on the behavior of $\text{PC}(\mathcal{L}_{\infty\lambda})$ (thus also $\text{RPC}(\mathcal{L}_{\infty\lambda})$) can be expressed by the following example. That example also shows that the condition that \mathbf{v} be λ -ary cannot be dispensed with in the statement of Theorem 7.5.

Example 8.8 (The Empty). *Let λ be an infinite cardinal and let the vocabulary \mathbf{v}_λ consist of the single relation symbol R with $\text{ar}(R) = \lambda$. Set $\mathcal{E}_\lambda \stackrel{\text{def}}{=} \mathbf{Mod}_{\mathbf{v}_\lambda}(E_\lambda)$ where E_λ is the $\mathcal{L}_{\infty\lambda+}(\mathbf{v}_\lambda)$ -sentence $(\forall \vec{x})_{(\lambda)} \neg R(\vec{x})$. Then \mathcal{E}_λ is the range of a finitely continuous functor on $\mathbf{Set}_{\neq \emptyset}$. Moreover, \mathcal{E}_λ is $\text{PC}(\mathcal{L}_{\infty\lambda})$ iff λ is singular.*

Proof. Let $\Phi: \mathbf{Set}_{\neq \emptyset} \rightarrow \mathcal{E}_\lambda$, $M \mapsto (M, \emptyset)$, naturally extended at morphism level. Then \mathcal{E}_λ is the range of Φ . Moreover, \mathcal{E}_λ is obviously $\text{PC}(\mathcal{L}_{\infty\lambda+})$. By Theorem 8.6, if λ is singular, then \mathcal{E}_λ is also $\text{PC}(\mathcal{L}_{\infty\lambda})$.

Finally, let λ be regular and suppose that $\mathcal{E}_\lambda = \{M \upharpoonright_{\mathbf{v}_\lambda} \mid M \in \mathbf{Mod}_{\mathbf{w}}(E)\}$ for some vocabulary \mathbf{w} extending \mathbf{v}_λ and some statement $E \in \mathcal{L}_{\infty\lambda}(\mathbf{w})$. Set $\theta \stackrel{\text{def}}{=} (\text{card } \mathcal{T}_\lambda(\mathbf{w}))^\lambda$ (cf. Notation 4.1). By Lemma 8.1, there exists an infinite cardinal $\kappa \geq \theta$ such that $\kappa^\lambda < \kappa^\lambda$. Since $(\kappa, \emptyset) \in \mathcal{E}_\lambda$, there exists $M \in \mathbf{Mod}_{\mathbf{w}}(E)$ such that $(\kappa, \emptyset) = (M, R^M)$.

Let $(v_\xi \mid \xi < \lambda)$ be a λ -sequence of distinct variables, set $X_\alpha \stackrel{\text{def}}{=} \{v_\xi \mid \xi < \alpha\}$ for each $\alpha < \lambda$, and

$$D \stackrel{\text{def}}{=} \left\{ (s_\xi^M(\vec{a}) \mid \xi < \lambda) \mid \text{all } s_\xi \in \mathcal{T}_\lambda(\mathbf{w}), (\exists \alpha < \lambda) \left(\bigcup_{\xi < \lambda} \text{var}(s_\xi) \subseteq X_\alpha \text{ and } \vec{a} \in X_\alpha \right) \right\}.$$

Then $\text{card } D \leq \theta \cdot \kappa^\lambda = \kappa^\lambda$. Since $D \subseteq {}^\lambda \kappa$ and $\kappa^\lambda < \kappa^\lambda$, there exists $\vec{a} \in {}^\lambda \kappa \setminus D$. Let M' be the model obtained from M by changing R^M (which is empty) to $\{\vec{a}\}$. Since every formula in $\mathcal{L}_{\infty\lambda}(\mathbf{w})$ has less than λ free variables, and since $\{\vec{a}\} \cap D = \emptyset \cap D = \emptyset$, M and M' satisfy the same atomic $\mathcal{L}_{\infty\lambda}$ -sentences with parameters from κ , hence (as $|M| = |M'| = \kappa$) the same $\mathcal{L}_{\infty\lambda}$ -sentences. In particular, $M' \models E$, so $(\kappa, \{\vec{a}\}) = M' \upharpoonright_{\mathbf{v}_\lambda}$ belongs to \mathcal{E}_λ ; a contradiction. \square

9. BACK-AND-FORTH SYSTEMS, ANTI-ELEMENTARITY, AND TUURI'S INTERPOLATION THEOREM

9.1. Karttunen's back-and-forth systems. There are a number of works establishing *non-representability*, of a given class \mathcal{C} of \mathbf{v} -structures for some vocabulary \mathbf{v} , as the class of models of any *class* of $\mathcal{L}_{\infty\lambda}$ -sentences, for any regular cardinal λ . In those works, this kind of intractability result is always achieved by establishing, in one form or another, that \mathcal{C} is not closed under a suitable notion of back-and-forth equivalence. It turns out that the following instance of the latter, introduced

in Karttunen [18] as “partial isomorphism with weak λ -extension property”, denoted there as $\mathbf{M} \cong_w^{\lambda_e} \mathbf{N}$ and that we will denote here in the form $\mathbf{M} \triangleleft_\lambda \mathbf{N}$, is sufficiently general to encompass all those situations.

Definition 9.1. Let λ be a regular cardinal and let \mathbf{M} and \mathbf{N} be models over the same vocabulary. A λ -back-and-forth system, in short λ -BFS, between \mathbf{M} and \mathbf{N} , is a poset $(\mathcal{F}, \trianglelefteq)$, together with a function $f \mapsto \bar{f}$ with domain \mathcal{F} , such that each \bar{f} is an isomorphism from a submodel $\mathbf{d}(f)$ of \mathbf{M} onto a submodel $\mathbf{r}(f)$ of \mathbf{N} , and the following statements hold:

- (1) For all $f, g \in \mathcal{F}$, $f \trianglelefteq g$ implies that $\bar{f} \subseteq \bar{g}$ (i.e., \bar{g} extends \bar{f}).
- (2) The poset $(\mathcal{F}, \trianglelefteq)$ is λ -inductive, that is, every λ -small subchain of $(\mathcal{F}, \trianglelefteq)$ has an upper bound in \mathcal{F} .
- (3) For every $f \in \mathcal{F}$ and every $x \in M$, there exists $g \in \mathcal{F}$ such that $f \trianglelefteq g$ and $x \in \mathbf{d}(g)$.
- (4) For every $f \in \mathcal{F}$ and every $y \in N$, there exists $g \in \mathcal{F}$ such that $f \trianglelefteq g$ and $y \in \mathbf{r}(g)$.

We will sum this up by writing $\mathcal{F}: \mathbf{M} \triangleleft_\lambda \mathbf{N}$. Let $\mathbf{M} \triangleleft_\lambda \mathbf{N}$ hold if there exists a λ -BFS between \mathbf{M} and \mathbf{N} .

The condition, added by Karttunen in [18, Definition 3.2], that $(\mathcal{F}, \trianglelefteq)$ be a tree, is redundant, in the sense that any $\mathcal{F}: \mathbf{M} \triangleleft_\lambda \mathbf{N}$ gives rise to some $\mathcal{F}': \mathbf{M} \triangleleft_\lambda \mathbf{N}$ where \mathcal{F}' is a tree: indeed, let \mathcal{F}' consist of all \trianglelefteq -ascending sequences $(f_\xi \mid \xi \leq \alpha)$ from \mathcal{F} , with $\alpha < \lambda$ and $f_0 = e$ is a fixed element of \mathcal{F} , partially ordered under extension, and precompose the assignment $f \mapsto \bar{f}$ with the “projection” $\mathcal{F}' \rightarrow \mathcal{F}$, $(f_\xi \mid \xi \leq \alpha) \mapsto f_\alpha$.

The case where \trianglelefteq is function extension, $f = \bar{f}$, $\mathbf{d}(f) = \text{dom } f$, and $\mathbf{r}(f) = \text{rng } f$ whenever $f \in \mathcal{F}$, is Dickmann’s original $\simeq_\lambda^{p,e}$ from [6, Definition 4.2.3].

As many aspects of the present paper are category-theoretical, we should point the reader to Beke and Rosický’s categorical take on back-and-forth presented in [3].

Many instances of λ -BFSs arise from the functors considered in the author’s paper [37], as stated in the forthcoming Proposition 9.3.

Notation 9.2. For any set Ω , denote⁹ by $[\Omega]^{\text{inj}}$ the subcategory of **Set** whose objects are all the nonempty subsets of Ω and whose arrows are the one-to-one maps.

Proposition 9.3. *Let λ be a regular cardinal, let \mathbf{v} be a λ -ary vocabulary, let Ω be a set, let $X, Y \subseteq \Omega$, and let $\Gamma: [\Omega]^{\text{inj}} \rightarrow \mathbf{Str}(\mathbf{v})$ be a λ -continuous functor. If either $\text{card } X = \text{card } Y$ or $\lambda \leq \min\{\text{card } X, \text{card } Y\}$, then $\Gamma(X) \triangleleft_\lambda \Gamma(Y)$.*

Note. We are not assuming, in the statement of Proposition 9.3, that Γ turns embeddings to embeddings (that assumption would in fact trivialize the result). Nonetheless, by [37, Proposition 4.5], this is automatically the case for embeddings with domain of cardinality at least λ .

Proof. The conclusion is trivial if $\text{card } X = \text{card } Y$: let \mathcal{F} be a singleton, acting on $\Gamma(X)$ via $\Gamma(f)$ where f is any bijection from X onto Y .

Suppose now that $\lambda \leq \min\{\text{card } X, \text{card } Y\}$. We define \mathcal{F} as the set of bijections $f: U \rightarrow V$ where $U \in [X]^{<\lambda}$ and $V \in [Y]^{<\lambda}$, ordered by function extension (i.e., $f \trianglelefteq g$ if g extends f). Further, denoting by id_S^T the inclusion map from S into T ,

⁹In [37], $[\Omega]^{\text{inj}}$ is denoted by $\mathfrak{P}_{\text{inj}}(\Omega)$.

and setting $e_S^T \stackrel{\text{def}}{=} \Gamma(\text{id}_S^T)$, whenever $S \subseteq T \subseteq \Omega$, we set $\mathbf{d}(f) = \mathbf{M}_U \stackrel{\text{def}}{=} e_U^X[\Gamma(U)]$ and $\mathbf{r}(f) = \mathbf{N}_V \stackrel{\text{def}}{=} e_V^Y[\Gamma(V)]$. Trivially, $(\mathcal{F}, \trianglelefteq)$ is a λ -inductive poset.

Claim. *There exists a unique \mathbf{v} -isomorphism $\bar{f}: \mathbf{M}_U \rightarrow \mathbf{N}_V$ such that $\bar{f} \circ e_U^X = e_V^Y \circ \Gamma(f)$.*

Proof of Claim. Let $e_U^X(x_1) = e_U^X(x_2)$ where $x_1, x_2 \in \Gamma(U)$. Since Γ is a λ -continuous functor and X is the colimit of its λ -small subsets, we get

$$(\Gamma(X), e_S^X \mid S \in [X]^{<\lambda}) = \varinjlim (\Gamma(S), e_S^T \mid S \subseteq T \text{ in } [X]^{<\lambda}). \quad (9.1)$$

Hence, using Lemma 2.1, there exists $X' \in [X]^{<\lambda}$ containing U such that

$$e_U^{X'}(x_1) = e_U^{X'}(x_2). \quad (9.2)$$

Since $\text{card } X' < \lambda \leq \text{card } Y$, the bijection $f: U \rightarrow V$ extends to a one-to-one map $g: X' \rightarrow Y$. From $g \circ \text{id}_{X'}^X = \text{id}_Y^Y \circ f$ it follows that $\Gamma(g) \circ e_U^{X'} = e_V^Y \circ \Gamma(f)$. From (9.2) it thus follows that $e_V^Y \Gamma(f)(x_1) = e_V^Y \Gamma(f)(x_2)$.

This completes the proof of the existence of a unique map $\bar{f}: \mathbf{M}_U \rightarrow \mathbf{N}_V$ such that $\bar{f} \circ e_U^X = e_V^Y \circ \Gamma(f)$. A similar argument shows that \bar{f} is a \mathbf{v} -homomorphism.

By applying the above to the inverse map $f^{-1}: V \rightarrow U$, we obtain a unique \mathbf{v} -homomorphism $\bar{g}: \mathbf{N}_V \rightarrow \mathbf{M}_U$ such that $\bar{g} \circ e_V^Y = e_U^X \circ \Gamma(f^{-1})$. It follows that $\bar{g}\bar{f}e_U^X = \bar{g}e_V^Y \Gamma(f) = e_U^X \Gamma(f^{-1}) \Gamma(f) = e_U^X$, whence $\bar{g}\bar{f}$ is the identity on \mathbf{M}_U . Similarly, $\bar{f}\bar{g}$ is the identity on \mathbf{N}_V . Therefore, \bar{f} and \bar{g} are mutually inverse isomorphisms. \square Claim.

It is then straightforward to verify that $f \trianglelefteq g$ implies $\bar{f} \subseteq \bar{g}$, for all $f, g \in \mathcal{F}$. Let us verify Item (3) of Definition 9.1. Let $f: U \rightarrow V$ in \mathcal{F} and let $x \in \Gamma(X)$. We need to enlarge f to some $g \in \mathcal{F}$ such that $x \in \mathbf{d}(g)$. By (9.1) and Lemma 2.1, there exists a λ -small subset X' of X , containing U , such that $x \in M_{X'}$. Since $\text{card } X' < \lambda \leq \text{card } Y$, f extends to a bijection $g: X' \rightarrow Y'$ for some $Y' \in [Y]^{<\lambda}$. Now $f \trianglelefteq g$ and $x \in M_{X'} = \mathbf{d}(g)$, so g is as required. The verification of Item (4) of Definition 9.1 is similar. \square

9.2. The infinitely deep languages $\mathcal{M}_{\kappa\lambda}$. The infinitely deep language $\mathcal{M}_{\kappa\lambda}$ (here $\lambda \leq \kappa$ are either infinite cardinals or ∞) extends the infinitary language $\mathcal{L}_{\kappa\lambda}$ in the sense that infinite alternations of quantifiers are allowed; satisfaction is then defined in terms of existence of a winning strategy for the “semantic game”. For example, the satisfaction of a “formula” of the form $\forall x_0 \exists x_1 \forall x_2 \cdots \mathbf{E}(x_0, x_1, \dots)$ is expressed by the existence of a winning strategy for player \exists in a standard Gale-Stewart game. While the latter has “clock” ω , the games underlying the formulas from $\mathcal{M}_{\kappa\lambda}$ may be clocked by *trees*. Since a full formal description of the logics $\mathcal{M}_{\kappa\lambda}$ will not be needed here, we will omit it, and refer the reader to Hyttinen [13], Karttunen [18], Rantala [28], Tuuri [32] for more details. The main result of Tuuri [32], extending a result from Hyttinen [13], states as follows.

Proposition 9.4. *Let κ be a regular cardinal, set $\lambda \stackrel{\text{def}}{=} \kappa^{\kappa}$, let \mathbf{v} be a κ -ary vocabulary, and let \mathbf{E} and \mathbf{F} be sentences of $\mathcal{L}_{\kappa+\kappa}(\mathbf{v})$ such that the conjunction $\mathbf{E} \wedge \mathbf{F}$ has no \mathbf{v} -model. Then there exists a sentence \mathbf{G} of $\mathcal{M}_{\lambda+\lambda}$, with vocabulary the intersection of the vocabularies of \mathbf{E} and \mathbf{F} , such that every \mathbf{v} -structure satisfies both sentences $\mathbf{E} \Rightarrow \mathbf{G}$ and $\mathbf{F} \Rightarrow \sim \mathbf{G}$.*

Here, $\sim \mathbf{G}$ denotes the sentence obtained by interchanging \bigvee and \bigwedge , \exists and \forall , \mathbf{F} and $\neg \mathbf{F}$ in the expression of \mathbf{G} by a tree-clocked game; it implies the usual negation $\neg \mathbf{G}$. By a counterexample due to Malitz [21, Theorem 4.2], $\mathcal{M}_{\lambda+\lambda}$ cannot be replaced by $\mathcal{L}_{\infty\infty}$ in the statement of Proposition 9.4. Karttunen states in [18, Theorem 3.5] that whenever \mathbf{M} and \mathbf{N} are \mathbf{v} -structures, $\mathbf{M} \sqsubset_{\lambda} \mathbf{N}$ implies¹⁰ that \mathbf{M} and \mathbf{N} satisfy the same sentences in a logic, denoted there by $\mathcal{N}_{\infty\lambda}$, extending $\mathcal{M}_{\infty\lambda}$, with uniquely defined syntax but allowing for several (quite unwieldy) nonequivalent definitions of satisfaction. Since we will need that result only for the far better behaved $\mathcal{M}_{\infty\lambda}$, we record here the corresponding statement.

Proposition 9.5. *Let \mathbf{M} and \mathbf{N} be \mathbf{v} -structures. If $\mathbf{M} \sqsubset_{\lambda} \mathbf{N}$, then \mathbf{M} and \mathbf{N} are $\mathcal{M}_{\infty\lambda}$ -elementarily equivalent.*

10. APPLICATION TO INTRACTABILITY RESULTS

All intractability results stated in the author's paper [37] establish a property, called there *anti-elementarity*, for pairs of classes $\mathcal{C}_0, \mathcal{C}_1$ of \mathbf{v} -structures with $\mathcal{C}_0 \subseteq \mathcal{C}_1$. By definition, $(\mathcal{C}_0, \mathcal{C}_1)$ is anti-elementary if there are arbitrarily large infinite cardinals $\lambda < \kappa$, with λ regular and a λ -continuous functor $\Gamma: [\kappa]^{\text{inj}} \rightarrow \mathbf{Str}(\mathbf{v})$, such that $\Gamma(\lambda) \in \mathcal{C}_0$ and $\Gamma(\kappa) \notin \mathcal{C}_1$. Then every class \mathcal{C} with $\mathcal{C}_0 \subseteq \mathcal{C} \subseteq \mathcal{C}_1$ is anti-elementary, in the sense that $(\mathcal{C}, \mathcal{C})$ is anti-elementary. Since $\Gamma(\lambda)$ and $\Gamma(\kappa)$ are $\mathcal{L}_{\infty\lambda}$ -elementarily equivalent, it follows that \mathcal{C} is not closed under $\mathcal{L}_{\infty\lambda}$ -elementary equivalence. In fact, by Proposition 9.3, $\Gamma(\lambda) \sqsubset_{\lambda} \Gamma(\kappa)$, thus it follows from Proposition 9.5 that \mathcal{C} is not closed under $\mathcal{M}_{\infty\lambda}$ -elementary equivalence. In particular, by Proposition 9.4, \mathcal{C} cannot be simultaneously $\text{PC}(\mathcal{L}_{\infty\infty})$ and $\text{co-PC}(\mathcal{L}_{\infty\infty})$.

We shall illustrate that line of argument with a sample of extensions of various intractability results mostly arising from [37].

10.1. Posets of convex ℓ -subgroups in ℓ -groups. Denote by $\text{Cs}_c G$ the (distributive) lattice with zero consisting of all finitely generated convex ℓ -subgroups of an ℓ -group G . As we verified in [36], this lattice is *Cevian*, that is, it has a binary operation \setminus such that all inequalities $x \leq y \vee (x \setminus y)$, $(x \setminus y) \wedge (y \setminus x) = 0$, and $x \setminus z \leq (x \setminus y) \vee (y \setminus z)$ hold on that lattice. Denote by \mathbf{Cev} the full subcategory of $\mathbf{Str}(\vee, \wedge, 0)$ consisting of all Cevian lattices¹¹.

Our next result involves the (non-commutative) diagram \vec{A} , of Archimedean ℓ -groups and ℓ -embeddings, introduced in [36, § 4] (also described in [37, § 12]).

Theorem 10.1. *For any accessible category \mathcal{G} of ℓ -groups and ℓ -homomorphisms, containing all objects and arrows of \vec{A} and closed under products and λ -directed colimits of ℓ -groups, for some regular cardinal λ , there is no $\text{co-PC}(\mathcal{L}_{\infty\infty})$ class \mathcal{C} of Cevian lattices containing $\text{Cs}_c \mathcal{G}$. In particular, $\text{Cs}_c \mathcal{G}$ is not $\text{co-PC}(\mathcal{L}_{\infty\infty})$.*

Proof. By assumption, \mathcal{G} contains the category denoted by $\mathcal{A}(\theta, \vec{A})$ in [37, Notation 12.2], for some regular cardinal θ . By [37, Theorem 12.3], there are arbitrarily large regular cardinals $\lambda < \kappa$ with λ -continuous functors $\Gamma: [\kappa]^{\text{inj}} \rightarrow \mathbf{Str}(\vee, \wedge, 0)$ such that $\Gamma(\lambda) \in \text{Cs}_c \mathcal{G}$ and $\Gamma(\kappa) \notin \mathbf{Cev}$. Since λ is arbitrarily large, we may assume

¹⁰Strictly speaking, the results of [18] are stated there for *finitary* vocabularies. Nonetheless, it is not hard to verify that the part concerning Proposition 9.5 extends to λ -ary vocabularies. It is plausible that this would extend to $\mathcal{N}_{\infty\lambda}$ as well, but we did not verify this.

¹¹The class \mathbf{Cev} is obviously $\text{PC}(\mathcal{L}_{\omega\omega})$ -definable. However, this will not be needed in the paper.

that \mathcal{G} is closed under λ -directed colimits. Since the functor Cs_c is finitely continuous on all ℓ -groups, its restriction Φ to \mathcal{G} is λ -continuous, thus, by Theorem 7.5, the class $\text{Cs}_c \mathcal{G} = \Phi(\mathcal{G})$ is $\text{PC}(\mathcal{L}_{\infty\lambda})$. Since \mathcal{C} is $\text{co-PC}(\mathcal{L}_{\infty\infty})$ and by Proposition 9.4, there exists an $\mathcal{M}_{\infty\infty}(\vee, \wedge, 0)$ -sentence \mathbf{E} such that $\text{Cs}_c \mathcal{G} \subseteq \mathbf{Mod}_{(\vee, \wedge, 0)}(\mathbf{E}) \subseteq \mathcal{C}$. In particular, $\Gamma(\lambda) \models \mathbf{E}$ and $\Gamma(\kappa) \models \neg \mathbf{E}$.

On the other hand, it follows from Proposition 9.3 that $\Gamma(\lambda) \simeq_{\lambda} \Gamma(\kappa)$. By Proposition 9.5, $\Gamma(\lambda)$ and $\Gamma(\kappa)$ satisfy the same $\mathcal{M}_{\infty\lambda}(\vee, \wedge, 0)$ -sentences; a contradiction. \square

Theorem 10.1 applies, in particular, to \mathcal{G} defined as any variety of ℓ -groups containing all Archimedean ℓ -groups. The particular case, where \mathcal{G} is the category of all Abelian ℓ -groups, can thus be viewed as another negative solution to Mundici's *MV-spectrum Problem* [23, Problem 2] handled in [37]: it was proved there that the $\text{PC}(\mathcal{L}_{\infty\omega})$ class $\text{Cs}_c \mathcal{G}$ is not closed under any $\mathcal{L}_{\infty\lambda}$ -elementary equivalence; now we know that it is not $\text{co-PC}(\mathcal{L}_{\infty\infty})$ either, and that it is in fact not contained in any $\text{co-PC}(\mathcal{L}_{\infty\infty})$ class of Cevian lattices.

Our next result aims at extending Theorem 10.1 to the functor that to every ℓ -group G associates its $(\vee, 0)$ -semilattice $\text{Id}_c G$ of finitely generated ℓ -ideals; thus $\text{Id}_c G$ is isomorphic to the $(\vee, 0)$ -semilattice $\text{Con}_c G$ of all finitely generated congruences of G . This result will involve the forgetful functor Ψ introduced in Gillibert and Wehrung [10, § 5.1] (see also [37, § 13]), that to every “semilattice-metric space” (A, δ, \tilde{A}) of type 1, with \tilde{A} generated by the range of δ , associates its underlying $(\vee, 0)$ -semilattice \tilde{A} . Recalling the full definition of Ψ will not be needed here. However, the reader might find it helpful to record that $\text{im } \Psi$ contains all $(\vee, 0)$ -semilattices either of the form $\text{Id}_c R \stackrel{\text{def}}{=} \text{semilattice of all finitely generated two-sided ideals of a ring } R$, or $\text{Id}_c G$ for an ℓ -group G , or $\text{Sub}_c M \stackrel{\text{def}}{=} \text{semilattice of all finitely generated submodules of a module } M$ (cf. [37, Proposition 13.1]). Our statement will also involve the (non-commutative) diagrams $\vec{C}_{a,g}$, indexed by non-representable¹² ℓ -groups G and elements $a, g \in C$ such that $a > 0$ and $a \wedge (g + a - g) = 0$, introduced in [37, § 13].

Theorem 10.2. *For any accessible category \mathcal{G} of ℓ -groups and ℓ -homomorphisms, containing all objects and arrows of some $\vec{C}_{a,g}$ for some non-representable ℓ -group C , and closed under products and λ -directed colimits for some regular cardinal λ , there is no $\text{co-PC}(\mathcal{L}_{\infty\infty})$ class \mathcal{C} such that $\text{Id}_c \mathcal{G} \subseteq \mathcal{C} \subseteq \text{im } \Psi$. In particular, $\text{Id}_c \mathcal{G}$ is not $\text{co-PC}(\mathcal{L}_{\infty\infty})$.*

Note. The case of categories of representable ℓ -groups is covered by Theorem 10.1: indeed, whenever G is representable, $\text{Id}_c G$ is a lattice homomorphic image of $\text{Cs}_c G$; hence it is Cevian (cf. [36, Proposition 5.10]).

Proof. By assumption, \mathcal{G} contains some $\mathcal{A}(\theta, \vec{C}_{a,g})$ (cf. [37, Notation 12.2]) for some regular cardinal θ . By [37, Theorem 13.8], there are arbitrarily large regular cardinals $\lambda < \kappa$ with λ -continuous functors $\Gamma: [\kappa]^{\text{inj}} \rightarrow \mathbf{Str}(\vee, \wedge, 0)$ such that $\Gamma(\lambda) \in \text{Id}_c \mathcal{G}$ and $\Gamma(\kappa) \notin \text{im } \Psi$. Since λ is arbitrarily large, we may assume that \mathcal{G} is closed under λ -directed colimits. Since the functor Id_c is finitely continuous on all ℓ -groups, its restriction Φ to \mathcal{G} is λ -continuous, thus, by Theorem 7.5, the class $\text{Id}_c \mathcal{G} = \Phi(\mathcal{G})$ is $\text{PC}(\mathcal{L}_{\infty\lambda})$. Since \mathcal{C} is $\text{co-PC}(\mathcal{L}_{\infty\infty})$ and by Proposition 9.4, there exists an

¹²Recall that an ℓ -group is *representable* if it embeds into a product of totally ordered groups.

$\mathcal{M}_{\infty\infty}(V, 0)$ -sentence E such that $\text{Cs}_c \mathcal{G} \subseteq \mathbf{Mod}_{(V,0)}(E) \subseteq \mathcal{C}$. The desired conclusion then follows as in the last part of the argument of the proof of Theorem 10.2. \square

For an extension of the method of Subsection 10.1 to an *a priori* non-co-PC class, specific to the case of Abelian ℓ -groups, see Section 11.

10.2. Posets of ideals in rings. Our next result is an analogue of Theorem 10.2 for (two-sided) ideal lattices of rings. It involves the (non-commutative) diagrams $\vec{R}_{\mathbb{k}}$ of matrix algebras over fields \mathbb{k} , introduced in [37, § 13]. Its proof is, *mutatis mutandis*, identical to the one of Theorem 10.2, using [37, Theorem 13.15] in place of [37, Theorem 13.8], and we omit it.

Theorem 10.3. *For any accessible category \mathcal{R} of unital rings and unital ring homomorphisms, containing all objects and arrows of some $\vec{R}_{\mathbb{k}}$, and closed under products and λ -directed colimits for some regular cardinal λ , there is no co-PC($\mathcal{L}_{\infty\infty}$) class \mathcal{C} such that $\text{Id}_c \mathcal{R} \subseteq \mathcal{C} \subseteq \text{im } \Psi$. In particular, $\text{Id}_c \mathcal{R}$ is not co-PC($\mathcal{L}_{\infty\infty}$).*

Theorem 10.3 applies, in particular, to the category of all unital von Neumann regular rings, also to the one of all unital rings.

10.3. Nonstable \mathbf{K}_0 -theory of rings. We denote by $V(R)$ the (commutative) monoid of isomorphism classes $[X]$ of all finitely generated projective right R -modules X , for any unital ring R . A ring R is *V-semiprimitive* if for all idempotent matrices a and b of the same finite order over R , if aR and bR have no nontrivial isomorphic direct summands, then $ab = 0$ (this can be expressed as a left-right symmetric statement). Our next statement involves the class \mathcal{C} of all monoids that can be represented as images of some $V(R)$, for a V-semiprimitive ring R , by some “pre-V-homomorphism” as defined in [37, § 14] (this class of monoid homomorphisms includes all isomorphisms). Again, a full formal definition of \mathcal{C} will not be needed here. The proof of Theorem 10.4 is, *mutatis mutandis*, identical to the one of Theorem 10.2, using [37, Theorem 14.6] in place of [37, Theorem 13.8], and we omit it.

Theorem 10.4. *For any accessible category \mathcal{R} of V-semiprimitive unital rings and unital ring homomorphisms, containing all objects and arrows of some $\vec{R}_{\mathbb{k}}$, and closed under products and λ -directed colimits for some regular cardinal λ , there is no co-PC($\mathcal{L}_{\infty\infty}$) class \mathcal{C}' of monoids such that $V(\mathcal{R}) \subseteq \mathcal{C}' \subseteq \mathcal{C}$. In particular, $V(\mathcal{R})$ is not co-PC($\mathcal{L}_{\infty\infty}$).*

Theorem 10.4 applies, in particular, to the category \mathbf{vNRing} of all unital von Neumann regular rings with unital ring homomorphisms, and also to the category \mathbf{RR}_0 of all unital C^* -algebras of real rank zero and unit-preserving C^* -homomorphisms (take $\mathbb{k} \stackrel{\text{def}}{=} \mathbb{C}$ for the latter). Note that in the latter case, directed colimits are not preserved (the C^* -colimit is the topological closure of the algebraic colimit); nevertheless, λ -directed colimits are preserved whenever λ is uncountable.

10.4. Coordinatizability of sectionally complemented modular lattices. A lattice L is *coordinatizable* if it is isomorphic to the lattice $\mathbb{L}(R)$ of all principal right ideals of any (not necessarily unital) von Neumann regular ring R . This implies that L is sectionally complemented and modular. Conversely, if a sectionally complemented modular lattice L has a countable cofinal sequence, then the coordinatizability of L follows from an algebraic condition with geometric inspiration,

introduced in Jónsson [17], called *existence of a large (partial) 4-frame*. The category \mathcal{A} , introduced in [37, § 15], is a category of von Neumann regular rings, with additional matrix units structure ensuring that every member of $\mathbb{L}(\mathcal{A})$ has a large 4-frame. It is ω_1 -accessible, and the restriction Φ of the functor \mathbb{L} to \mathcal{A} is finitely continuous. Hence, using the $(\kappa, < \omega, \lambda) \rightarrow \rho$ notation from Erdős *et al.* [8], by the same token as the one used in the proofs of Theorem 10.1 above, and applying [37, Theorem 5.9], we obtain the following.

Theorem 10.5. *Assume that for every cardinal λ there exists a cardinal κ such that $(\kappa, < \omega, \lambda) \rightarrow \aleph_1$ (this holds if there is a proper class of Erdős cardinals). Then the class of all coordinatizable lattices with a large 4-frame is not $\text{co-PC}(\mathcal{L}_{\infty\infty})$.*

Let us now move to intractability results obtained from earlier sources than [37], not obtained *via* anti-elementarity.

Removing the large 4-frame condition but adding a unit element, we move to studying the class \mathcal{CL} of all coordinatizable, complemented, modular lattices. We stated in [33] that \mathcal{CL} is not the class of all models of any $\mathcal{L}_{\infty\infty}$ -sentence — in fact, the argument of the proof of [33, Theorem 9.4] shows that \mathcal{CL} is not closed under $\mathcal{L}_{\infty\lambda}$ -elementary equivalence, for any infinite cardinal λ (thus solving a problem from Jónsson [16]). Consider the class \mathcal{CL}_2 of all 2-distributive¹³ coordinatizable complemented modular lattices.

Theorem 10.6. *There is no $\text{co-PC}(\mathcal{L}_{\infty\infty})$ class \mathcal{C}' of lattices with $\mathcal{CL}_2 \subseteq \mathcal{C}' \subseteq \mathcal{CL}$. In particular, neither \mathcal{CL}_2 nor \mathcal{CL} is $\text{co-PC}(\mathcal{L}_{\infty\infty})$.*

Outline of proof. For any set X , denote by M_X the (2-distributive, complemented, modular) lattice $X \sqcup \{0, 1\}$ with the only nontrivial relations $0 < x < 1$ for $x \in X$. Let λ be a regular cardinal, set $\kappa \stackrel{\text{def}}{=} \lambda^+$. Due to the need for introducing the λ -BFS \mathcal{F}_λ below, we need to modify the definitions of the lattices \bar{L}_λ and L_λ introduced in [33, § 9], by changing “finite” to “ λ -small”. Our new lattices, for which we keep the same notation as in [33, § 9], are thus the following:

$$\begin{aligned} \bar{L}_\lambda &\stackrel{\text{def}}{=} \left\{ x \in {}^\kappa M_{\lambda+1} \mid (\exists X \in [\kappa]^{<\lambda})(x \upharpoonright_{\kappa \setminus X} \text{ is constant}) \right\}, \\ L_\lambda &\stackrel{\text{def}}{=} \{ x \in \bar{L}_\lambda \mid x_\infty \in M_\lambda \}. \end{aligned}$$

It is straightforward to verify that the lattices \bar{L}_λ and L_λ are both complemented, modular, and 2-distributive.

Claim. *The lattice \bar{L}_λ is coordinatizable whereas L_λ is not.*

Proof of Claim. The proof that \bar{L}_λ is coordinatizable follows the argument presented at the beginning of the proof of [33, Theorem 9.4]: if D denotes any division ring with λ elements, then \bar{L}_λ is coordinatized by the von Neumann regular ring

$$\left\{ x \in {}^\kappa M_2(D) \mid (\exists X \in [\kappa]^{<\lambda})(x \upharpoonright_{\kappa \setminus X} \text{ is constant}) \right\}.$$

Although the proof that L_λ is not coordinatizable is much harder, it works along the same lines as the one of [33, Theorem 9.3], with the following modifications. Change κ and κ^+ (from [33, Theorem 9.3]) to λ and κ , respectively; change “finite” to “ λ -small”. Moreover, redefine J_X , for $X \in [\kappa]^{<\lambda}$, as the ideal $\{x \in R \mid \varepsilon(xR) \upharpoonright_{\kappa \setminus X} = 0\}$, and denote by J the union of all J_X . We keep the ideal $I =$

¹³A lattice L is 2-distributive if $x \wedge (y_0 \vee y_1 \vee y_2) = \bigvee_{i < j} (x \wedge (y_i \vee y_j))$ for all $x, y_0, y_1, y_2 \in L$.

$\{\varepsilon(xR) \mid x \in J\} = \{x \in L_\lambda \mid x_\infty = 0\}$. The remainder of the argument is, *mutatis mutandis*, identical to the one of the proof of [33, Theorem 9.3]. \square Claim.

Since \mathbb{L} defines a finitely continuous functor on the finitely accessible category of all unital von Neumann regular rings, it follows from Theorem 7.5 that the class \mathcal{CL} is $\text{PC}(\mathcal{L}_{\infty\omega})$; thus so is also the class \mathcal{CL}_2 of all 2-distributive members of \mathcal{CL} .

We claim that $\bar{L}_\lambda \rightleftharpoons_\lambda L_\lambda$. To that end, denote by \mathcal{F}_λ the set of all κ -sequences $f = (f_\xi \mid \xi < \kappa)$, constant on the complement of a λ -small set, where each $f_\xi: \mathbf{d}(f_\xi) \rightarrow \mathbf{r}(f_\xi)$ is a bijection between λ -small subsets $\mathbf{d}(f_\xi)$ and $\mathbf{r}(f_\xi)$ of $\lambda + 1$, and such that $\mathbf{r}(f_\infty) \subseteq \lambda$. For $f, g \in \mathcal{F}_\lambda$, let $f \leq g$ hold if each g_ξ extends f_ξ . To each $f \in \mathcal{F}_\lambda$ we assign the map

$$\bar{f}: \bar{L}_\lambda \cap \prod_{\xi < \kappa} M_{\mathbf{d}(f_\xi)} \rightarrow L_\lambda \cap \prod_{\xi < \kappa} M_{\mathbf{r}(f_\xi)}, \quad (x_\xi \mid \xi < \kappa) \mapsto (\bar{f}_\xi(x_\xi) \mid \xi < \kappa)$$

where each $\bar{f}_\xi: M_{\mathbf{d}(f_\xi)} \rightarrow M_{\mathbf{r}(f_\xi)}$ is the unique 0, 1-preserving extension of f_ξ . This defines a λ -BFS $\mathcal{F}_\lambda: \bar{L}_\lambda \rightleftharpoons_\lambda L_\lambda$.

Since $\mathcal{CL}_2 \subseteq \mathcal{C}'$ and both \mathcal{CL}_2 and the complement of \mathcal{C}' are $\text{PC}(\mathcal{L}_{\infty\omega})$, there exists by Proposition 9.4 an $\mathcal{M}_{\infty\omega}(\vee, \wedge)$ -sentence \mathbf{E} with $\mathcal{CL}_2 \subseteq \mathbf{Mod}_{(\vee, \wedge)}(\mathbf{E}) \subseteq \mathcal{C}'$. Since $\bar{L}_\lambda \in \mathcal{CL}_2$ and $L_\lambda \notin \mathcal{C}'$, it follows that $\bar{L}_\lambda \models \mathbf{E}$ and $L_\lambda \models \neg \mathbf{E}$; a contradiction since $\bar{L}_\lambda \rightleftharpoons_\lambda L_\lambda$ and by Proposition 9.5. \square

10.5. Real spectra of commutative unital rings. A subset P in a commutative, unital ring A is a *cone* if it is both an additive and multiplicative submonoid of A such that $x^2 \in P$ whenever $x \in A$. The cone P is *prime* if $P \cup (-P) = A$ and $P \cap (-P)$ is a prime ideal of A . The *real spectrum* of A is the set $\text{Spec}_r A$ of all prime cones of A , endowed with the topology generated by the sets $\{P \in \text{Spec}_r A \mid a \notin P\}$ for $a \in A$. The topological space $\text{Spec}_r A$ is *spectral*, which is equivalent to saying that it is the Stone dual of a bounded distributive lattice (cf. Stone [29], Johnstone [15, § II.3], Grätzer [12, § II.5]). Mellor and Tressl proved in [22, Theorem 5.1] that the class \mathcal{S} of Stone duals of real spectra is not closed under strong λ -back-and-forth equivalence, for any infinite λ ; hence it is not the class of models of any class of $\mathcal{L}_{\infty\lambda}$ -sentences, for any infinite cardinal λ .

Theorem 10.7. *The class \mathcal{S} , of Stone duals of real spectra of commutative unital rings, is not $\text{co-PC}(\mathcal{L}_{\infty\omega})$.*

Proof. We first argue that \mathcal{S} is the image of a finitely continuous functor on a finitely accessible category. This can be done in more than one way. For example, the argument sketched at the end of the Introduction in [37], translated from spectra of Abelian ℓ -groups to *Brumfiel spectra* (known to be equivalent to real spectra), shows that \mathcal{S} is $\text{PC}(\mathcal{L}_{\omega_1\omega})$. A more direct way to do this follows from Johnstone [15, p. 218], where the Stone dual of $\text{Spec}_r A$, which we will denote by $\Phi(A)$, is expressed as the bounded distributive lattice defined by generators $D_r(a)$, with $a \in A$, subjected to specific relations (e.g., $D_r(a) \wedge D_r(b) \leq D_r(a + b) \leq D_r(a) \vee D_r(b)$, etc.). This entails that Φ is a finitely continuous functor from the (finitely accessible) category of all commutative unital rings to the (finitely accessible) category of all bounded distributive lattices. By Theorem 7.5, the class $\mathcal{S} = \text{im } \Phi$ is $\text{PC}(\mathcal{L}_{\infty\omega})$ -definable.

If \mathcal{S} were also $\text{co-PC}(\mathcal{L}_{\infty\omega})$, then, by Proposition 9.4, it would be the class of models of some $\mathcal{M}_{\infty\lambda}(\vee, 0)$ -sentence \mathbf{E} , for a regular cardinal λ . In particular, \mathcal{S} is closed under $\rightleftharpoons_\lambda$, in contradiction with Mellor and Tressl [22, Theorem 5.1]. \square

11. BEYOND co-PC: PLOŠČICA'S CONDITION AND ℓ -SPECTRA

Denote by \mathcal{A} the class of all Abelian ℓ -groups. By Theorem 10.1, $\text{Id}_c \mathcal{A}$ is not a co-PC($\mathcal{L}_{\infty\infty}$) class of lattices. On the other hand, $\text{Id}_c \mathcal{A}$ is closed under isomorphic copies, thus, using Lemma 3.2, it is the intersection of all co-PC($\mathcal{L}_{\infty\omega}$) classes containing it. We shall introduce a condition, originating in Ploščica [26], expressible as a conjunction, ranging over all infinite cardinals, of co-PC($\mathcal{L}_{\infty\infty}$)-sentences, satisfied by all members of $\text{Id}_c \mathcal{A}$. We shall also prove (cf. Theorem 11.6) that under the GCH, this condition, together with any co-PC($\mathcal{L}_{\infty\infty}$)-sentence satisfied by all members of $\text{Id}_c \mathcal{A}$, is still not sufficient to characterize $\text{Id}_c \mathcal{A}$.

Definition 11.1. A join-semilattice $(S, \vee, 0)$ has *countably based differences* (in abbreviation CBD; see Wehrung [35, § 10]) if every pair (x, y) has a *difference basis*, that is, a descending sequence $(z_n \mid n < \omega)$ of elements of S such that for every $z \in S$, $x \leq y \vee z$ iff $z_n \leq z$ for some $n < \omega$.

In particular, for every Abelian ℓ -group G , the lattice $\text{Id}_c G$ satisfies CBD.

Definition 11.2. For any cardinal number λ , define *Ploščica's Condition at λ* , denoted by Plo_λ , the statement of lattice theory stating that for every element a and every λ -sequence $(\mathfrak{m}_\xi \mid \xi < \lambda)$ of maximal ideals of $\downarrow a$, the quotient $\downarrow a / \bigcap_{\xi < \lambda} \mathfrak{m}_\xi$ has at most 2^λ elements (cf. Subsection 2.1 for the notation D/I).

We leave the verification of the following lemma as an exercise.

Lemma 11.3. CBD is equivalent to an $\mathcal{L}_{\omega_1\omega_1}$ -statement of lattice theory, and each Plo_λ is equivalent to a co-PC($\mathcal{L}_{\theta\theta}$)-statement of lattice theory where $\theta \stackrel{\text{def}}{=} (2^\lambda)^+$.

Lemma 11.4. The following statements hold, for any distributive lattice D with zero:

- (1) D satisfies Plo_n whenever $n < \omega$.
- (2) If D satisfies Plo_κ for every $\kappa \leq \text{card } D$, then it satisfies all Plo_κ .
- (3) Suppose that D has CBD. For every cardinal λ with uncountable cofinality, if $D \models \text{Plo}_\alpha$ whenever $\alpha < \lambda$, then $D \models \text{Plo}_\lambda$.
- (4) Suppose that D has CBD. For every infinite cardinal λ , $D \models \text{Plo}_\lambda$ implies $D \models \text{Plo}_{\lambda^+}$.

Proof. Ad (1). For every $a \in D$ and every finite sequence $(\mathfrak{m}_i \mid i < n)$ of maximal ideals of $\downarrow a$, it follows from the distributivity of D that the diagonal map embeds $\downarrow a / \bigcap_{i < n} \mathfrak{m}_i$ into $\prod_{i < n} (\downarrow a / \mathfrak{m}_i)$. Since each \mathfrak{m}_i is a maximal ideal of $\downarrow a$, $\downarrow a / \mathfrak{m}_i$ is the two-element lattice.

Ad (2). Every intersection of a collection of subsets of D (here maximal ideals of some $D \downarrow a$) is the intersection of a subcollection of size at most $\text{card } D$.

Ad (3). Setting $\mathfrak{b}_\alpha \stackrel{\text{def}}{=} \bigcap_{\xi < \alpha} \mathfrak{m}_\xi$ for $\alpha \leq \lambda$, $\text{card}(\downarrow a / \mathfrak{b}_\alpha) \leq 2^{\text{card } \alpha} \leq 2^\lambda$ whenever $\alpha < \lambda$, thus it suffices to prove that the diagonal map $\varepsilon: \downarrow a / \mathfrak{b}_\lambda \rightarrow \prod_{\alpha < \lambda} (\downarrow a / \mathfrak{b}_\alpha)$ is an order-embedding. Let $x, y \in \downarrow a$ and let $(z_n \mid n < \omega)$ be a difference basis of (x, y) . Then $\varepsilon(x / \mathfrak{b}_\lambda) \leq \varepsilon(y / \mathfrak{b}_\lambda)$ means that for every $\alpha < \lambda$ there exists $n < \omega$ such that $z_n \in \mathfrak{b}_\alpha$. Denote by n_α the least such n . Since the λ -sequence $(\mathfrak{b}_\alpha \mid \alpha < \lambda)$ and the ω -sequence $(z_n \mid n < \omega)$ are both descending, the λ -sequence $(n_\alpha \mid \alpha < \lambda)$ is ascending. Since cf $\lambda > \omega$, $\{n_\alpha \mid \alpha < \lambda\}$ is bounded above, say by an integer m . Then $z_m \in \mathfrak{b}_\lambda$, so $x \leq y \pmod{\mathfrak{b}_\lambda}$.

Ad (4) follows immediately from (3). □

Denote by $\text{Plo}_{\leq \kappa}$ (resp., $\text{Plo}_{< \infty}$) the conjunction of all Plo_α over $\alpha \leq \kappa$ (resp., over all α). A mild extension of the argument of the proof of Plošćica [26, Theorem 2.1] yields the following.

Proposition 11.5. *For every Abelian ℓ -group G , the lattice $\text{Id}_c G$ satisfies $\text{Plo}_{< \infty}$.*

Proof. By Lemma 11.4(1), we may assume that λ is infinite. The elements of $\text{Id}_c G$ are the $\langle u \rangle \stackrel{\text{def}}{=} \{x \in G \mid (\exists n < \omega)(|x| \leq nu)\}$ where $u \in G^+$, and then, for that element, $(\text{Id}_c G) \downarrow \langle u \rangle = \text{Id}_c \langle u \rangle$. Setting $H \stackrel{\text{def}}{=} \langle u \rangle$ (an ℓ -ideal of G), we thus need to verify that for every λ -sequence $\langle \mathfrak{m}_\xi \mid \xi < \lambda \rangle$ of maximal ideals of $\text{Id}_c H$, with intersection $\mathfrak{a} \stackrel{\text{def}}{=} \bigcap_{\xi < \lambda} \mathfrak{m}_\xi$, the quotient $(\text{Id}_c H)/\mathfrak{a}$ has at most 2^λ elements. Every $M_\xi \stackrel{\text{def}}{=} \{x \in H \mid \langle x \rangle \in \mathfrak{m}_\xi\}$ is a maximal ℓ -ideal of H , thus H/M_ξ can be embedded into the reals (cf. Anderson and Feil [2, Theorem 2.3]). Further, setting $A \stackrel{\text{def}}{=} \bigcap_{\xi < \lambda} M_\xi$, the diagonal map embeds H/A into $\prod_{\xi < \lambda} (H/M_\xi)$; it follows that $\text{card}(H/A) \leq (2^{\aleph_0})^\lambda = 2^\lambda$. Therefore $(\text{Id}_c H)/\mathfrak{a}$, which is isomorphic to $\text{Id}_c(H/A)$ (cf. Wehrung [35, Lemma 2.5]), has at most 2^λ elements. \square

By Theorem 10.1, for any co-PC($\mathcal{L}_{\infty\infty}$) sentence \mathbf{E} of lattice theory, there is no cardinal κ such that $\mathbf{E} \wedge \text{Plo}_{\leq \kappa}$ characterizes $\text{Id}_c \mathcal{A}$. We shall now verify that under the GCH, the same can be said about $\text{Plo}_{< \infty}$.

Theorem 11.6. *Suppose that the GCH holds. Then for every co-PC($\mathcal{L}_{\infty\infty}$) class \mathcal{C} of lattices containing the class $\text{Id}_c \mathcal{A}$, there exists a bounded distributive lattice in $\mathcal{C} \setminus \text{Id}_c \mathcal{A}$ that satisfies $\text{Plo}_{< \infty}$.*

Proof. Since \mathcal{C} is co-PC($\mathcal{L}_{\infty\infty}$), it is co-PC($\mathcal{L}_{\kappa^3 + \kappa^{2+}}$) for some regular cardinal κ (where κ^{n+} denotes the n -th cardinal successor of κ). Setting $\lambda \stackrel{\text{def}}{=} \kappa^{2+}$, it follows from [37, Theorem 12.3] that there exists a λ -continuous functor $\Delta: [\lambda^{2+}]^{\text{inj}} \rightarrow \mathbf{Str}(\vee, \wedge, 0)$ such that $\Delta(\lambda) \in \text{Id}_c \mathcal{A}$, $\Delta(\lambda^{2+})$ is not Cevian (thus it does not belong to $\text{Id}_c \mathcal{A}$), and (due to the GCH) $\text{card} \Delta(X) \leq \max\{\lambda, \text{card} X\}$ for every $X \subseteq \lambda^{2+}$. Since $\text{Id}_c \mathcal{A}$ and the complement of $\mathcal{C} \cap \mathbf{Mod}_{(\vee, \wedge, 0)}(\text{Plo}_{\leq \kappa})$ are both PC($\mathcal{L}_{\lambda+\lambda}$) (cf. Lemma 11.3), with $\text{Id}_c \mathcal{A} \subseteq \mathcal{C} \cap \mathbf{Mod}_{(\vee, \wedge, 0)}(\text{Plo}_{\leq \kappa})$, it follows from Tuuri's Interpolation Theorem (Proposition 9.4) that there exists an $\mathcal{M}_{\lambda+\lambda}(\vee, \wedge, 0)$ -sentence \mathbf{E} such that $\text{Id}_c \mathcal{A} \subseteq \mathbf{Mod}_{(\vee, \wedge, 0)}(\mathbf{E}) \subseteq \mathcal{C} \cap \mathbf{Mod}_{(\vee, \wedge, 0)}(\text{Plo}_{\leq \kappa})$. In particular, $\Delta(\lambda)$ is a bounded distributive lattice satisfying $\text{CBD} \wedge \mathbf{E} \wedge \text{Plo}_{\leq \kappa}$. Since $\Delta(\lambda) \hookrightarrow_\lambda \Delta(\lambda^{2+})$ (cf. Proposition 9.3) and by Proposition 9.5, it follows that $\Delta(\lambda^{2+})$ is also a bounded distributive lattice satisfying $\text{CBD} \wedge \mathbf{E} \wedge \text{Plo}_{\leq \kappa}$; whence $\Delta(\lambda^{2+}) \in \mathcal{C}$. By Lemma 11.4(4), $\Delta(\lambda^{2+})$ satisfies $\text{Plo}_{\leq \kappa^{n+}}$ for every $n < \omega$. Taking $n = 2$, we get $\Delta(\lambda^{2+}) \models \text{Plo}_{\leq \lambda^{2+}}$. By Lemma 11.4(2), it follows that $\Delta(\lambda^{2+})$ satisfies $\text{Plo}_{< \infty}$, and thus is as required. \square

In the proof above, $\Delta(\lambda^{2+}) = \Delta(\kappa^{4+})$ has cardinality κ^{4+} . In particular, applying Theorem 11.6 to the class \mathcal{C} of all distributive 0-lattices with CBD, we can take $\kappa = \aleph_0$, so we get the bounded distributive lattice $\Delta(\omega_4)$. Due to the GCH, this lattice has cardinality \aleph_4 , is completely normal, satisfies CBD together with $\text{Plo}_{< \infty}$, but does not belong to $\text{Id}_c \mathcal{A}$ (it is not Cevian). We do not know whether the conclusion of Theorem 11.6 holds in full generality (e.g., without the GCH). We find it plausible that this might follow from a closer look at the condensate construction from [37], within bounded distributive lattices.

12. A FINITE BOUNDED POSET FOR WHICH CONDENSATES DO NOT WORK

The poset of the title is represented in the left hand side of Figure 12.2 and denoted there by P_0 . The condensates are those of Gillibert and Wehrung [10], Wehrung [37]. This section brings a counterexample to [10, Problem 1].

Recall that a $(\vee, 0)$ -semilattice is an idempotent commutative monoid; in order to avoid any notational confusion between the semilattice operation and logical disjunction, we will in this section denote such structures additively. Letting $x \leq y$ stand for $x + y = y$, a $(\vee, 0)$ -semilattice S is *distributive* (cf. Grätzer [12]) if for all $a, b, c \in S$ such that $c \leq a + b$ there are $x \leq a$ and $y \leq b$ in S such that $c = x + y$.

Consider the category \mathbf{DSLat}^0 of all distributive $(\vee, 0)$ -semilattices with monoid homomorphisms. By the Ershov-Pudlák Lemma (cf. Ershov [9], Pudlák [27]; see Wehrung [34, Theorem 7-4.2] for a discussion), every distributive $(\vee, 0)$ -semilattice is the directed union of its finite distributive 0-subsemilattices. Hence, \mathbf{DSLat}^0 is a finitely accessible category; it has all products.

In this section we shall verify that \mathbf{DSLat}^0 does not have all pushouts or, for that matter, pullbacks (cf. Theorem 12.3); hence it is not locally presentable. To that end, consider the Boolean semilattice $\mathbf{2} \stackrel{\text{def}}{=} \{0, 1\}$ and define $(\vee, 0)$ -semilattice embeddings $\iota: \mathbf{2} \hookrightarrow \mathbf{2}\mathbf{2}$, $\alpha, \beta: \mathbf{2}\mathbf{2} \hookrightarrow \mathbf{3}\mathbf{2}$, and $\gamma_3, \gamma_4: \mathbf{3}\mathbf{2} \hookrightarrow \mathbf{4}\mathbf{2}$ by

$$\begin{aligned} \iota(x) &= (x, x); & \alpha(x, y) &\stackrel{\text{def}}{=} (x, x, y); & \beta(x, y) &\stackrel{\text{def}}{=} (y, x, x); \\ \gamma_3(x, y, z) &= (x, y, z, x + z); & \gamma_4(x, y, z) &= (x, y, z, x + y + z), \end{aligned}$$

for all $x, y, z \in \mathbf{2}$. It is straightforward to verify that the diagram \vec{D} represented in Figure 12.1 is commutative.





FIGURE 12.1. The commutative diagram \vec{D}

This diagram is indexed by the poset P_0 , with universe $\{0, 1, 2, 3, 4, 5\}$, represented on the left hand side of Figure 12.2. The intersection $L \stackrel{\text{def}}{=} \gamma_3[\mathbf{3}\mathbf{2}] \cap \gamma_4[\mathbf{3}\mathbf{2}] = \{0000, 1001, 0011, 1101, 0111, 1011, 1111\}$ is a 0, 1-subsemilattice of $\mathbf{4}\mathbf{2}$, represented on the right hand side of Figure 12.2.

Our next task will be representing \mathbf{DSLat}^0 as the image of an accessible functor (by Theorem 5.5 this is certainly possible). Moreover, the domain of that functor will be constructed as a (finitary) *variety of algebras*. This will be performed by a simple Skolemization argument. Consider the extension $\mathbf{v}^* \stackrel{\text{def}}{=} (+, 0, f)$ of the

FIGURE 12.2. The posets P_0 and $L \stackrel{\text{def}}{=} \gamma_3[{}^3\mathbf{2}] \cap \gamma_4[{}^3\mathbf{2}]$

vocabulary $\mathbf{v} \stackrel{\text{def}}{=} (+, 0)$, where the operation symbols $+$, 0 , and f have respective arities 2, 0, and 3. Denote by \mathcal{V} the variety of \mathbf{v}^* -algebras defined by the identities

$$x + x = x; \quad x + y = y + x; \quad x + (y + z) = (x + y) + z; \quad x + 0 = x; \quad (12.1)$$

$$f(x + y, x, x) = x; \quad (12.2)$$

$$f(x, y, z) + x = x; \quad (12.3)$$

$$f(x + y, z, z) = f(x, y, f(x + y, z, z)) + f(y, x, f(x + y, z, z)). \quad (12.4)$$

In particular, the identities (12.1) define the variety of all $(\vee, 0)$ -semilattices.

Proposition 12.1. *A $(\vee, 0)$ -semilattice S is distributive iff it expands to a member of \mathcal{V} .*

Proof. Let $(S, +, 0, f)$ be a member of \mathcal{V} and let $a, b, c \in S$ such that $c \leq a + b$. Since $(S, +, 0, f)$ satisfies (12.2), we get

$$c = f(c + (a + b), c, c) = f(a + b, c, c).$$

Since $(S, +, 0, f)$ satisfies (12.4), we thus get $c = f(a, b, c) + f(b, a, c)$. Since $(S, +, 0, f)$ satisfies (12.3), $f(a, b, c) \leq a$ and $f(b, a, c) \leq b$. Therefore, $(S, +, 0)$ is distributive.

Let, conversely, $(S, +, 0)$ be a distributive $(\vee, 0)$ -semilattice. For all $a, b, c \in S$ with $c \leq a + b$, there are $f_0(a, b, c) \leq a$ and $f_1(a, b, c) \leq b$ in S such that $c = f_0(a, b, c) + f_1(a, b, c)$. We may further assume that $a \leq b$ implies $f_0(b, a, a) = a$. The map $f: {}^3S \rightarrow S$ defined by the rule

$$f(a, b, c) \stackrel{\text{def}}{=} \begin{cases} f_0(a, b, c) + f_1(b, a, c), & \text{if } c \leq a + b, \\ a, & \text{otherwise} \end{cases}, \quad \text{for all } a, b, c \in S,$$

obviously satisfies the identities (12.2) and (12.3). Let $a, b, c \in S$ and set $c' \stackrel{\text{def}}{=} f(a + b, c, c)$. Then $c' \leq a + b$ by (12.3), thus

$$f(a, b, c') + f(b, a, c') = f_0(a, b, c') + f_1(a, b, c') + f_0(b, a, c') + f_1(b, a, c') = c',$$

so f satisfies (12.4). \square

By Proposition 12.1, the forgetful functor $\Phi: \mathcal{V} \rightarrow \mathbf{Str}(\mathbf{v})$, $(S, +, 0, f) \mapsto (S, +, 0)$ has range \mathbf{DSLat}^0 . This functor is faithful, finitely accessible, and it preserves all products.

We shall now argue that the surjectivity of Φ does not extend to diagrams, in a strong sense designed to provide some traction on Gillibert and Wehrung [10, Problem 1]. As in [10], say that a homomorphism $\varphi: S \rightarrow T$ of $(\vee, 0)$ -semilattices

is *weakly distributive* if for all $s \in S$ and $t_1, t_2 \in T$ such that $\varphi(s) \leq t_1 + t_2$ there are $s_1, s_2 \in S$ such that $s \leq s_1 + s_2$ and each $\varphi(s_i) \leq t_i$. Since the functor Φ preserves all directed colimits, and since any canonical projection $E \times F \rightarrow E$ is weakly distributive, Condition $\text{PROJ}(\Phi, \mathcal{S}^{\Rightarrow})$ of [10, Definition 3.4.1] is satisfied.

Let $P_1 \stackrel{\text{def}}{=} P_0 \cup \{\mu\}$ where μ is a new element greater than 1 and 2 but smaller than 3 and 4. Say that a P_0 -indexed commutative diagram is *mid-extendible* if it extends to a P_1 -indexed diagram in \mathbf{DSLat}^0 .

Proposition 12.2. *Every diagram in \mathbf{DSLat}^0 of the form $\Phi(\vec{E})$, where \vec{E} is a P_0 -indexed commutative diagram in \mathcal{V} , is mid-extendible.*

Proof. Write $\vec{E} = (E_p, \varepsilon_{p,q} \mid p \leq q \text{ in } P_0)$ and set

$$E_\mu \stackrel{\text{def}}{=} \{(x_3, x_4) \in E_3 \times E_4 \mid \varepsilon_{3,5}(x_3) = \varepsilon_{4,5}(x_4)\},$$

the pullback of $(\varepsilon_{j,5}: E_j \rightarrow E_5 \mid j \in \{3,4\})$. The missing maps are provided by $\varepsilon_{i,\mu} \stackrel{\text{def}}{=} (\varepsilon_{i,3}, \varepsilon_{i,4})$ ($i \in \{1,2\}$) and $\varepsilon_{\mu,j} \stackrel{\text{def}}{=} \text{projection } E_\mu \rightarrow E_j$ ($j \in \{3,4\}$). \square

Theorem 12.3. *There are no mid-extendible P_0 -indexed commutative diagram \vec{E} in \mathbf{DSLat}^0 and no natural transformation $\vec{\chi}: \vec{E} \rightarrow \vec{D}$ such that χ_0 is surjective and χ_1 and χ_2 are both weakly distributive. In particular, \vec{D} is not mid-extendible; thus it is not isomorphic to $\Phi(\vec{E})$ for any diagram \vec{E} in \mathcal{V} .*

Proof. Denote by $\vec{E}' \stackrel{\text{def}}{=} (E_p, \varepsilon_{p,q} \mid p \leq q \text{ in } P_0 \cup \{\mu\})$ the given extension of \vec{E} to a P_1 -indexed commutative diagram in \mathbf{DSLat}^0 . We represent it, together with the natural transformation $\vec{\chi}$, on the left hand side of Figure 12.3. Note that the ‘‘mid component’’ χ_μ is undefined at that point.




FIGURE 12.3. The natural transformation $\vec{\chi}$

Now we set $\chi_\mu \stackrel{\text{def}}{=} \chi_5 \varepsilon_{\mu,5}$. For each $j \in \{3,4\}$, $\chi_\mu = \chi_5 \varepsilon_{j,5} \varepsilon_{\mu,j} = \gamma_j \chi_j \varepsilon_{\mu,j}$ has range contained in $\gamma_j[\mathbf{3}\mathbf{2}]$; whence $\chi_\mu: E_\mu \rightarrow L$. Moreover, $\chi_\mu \varepsilon_{i,\mu} = \chi_5 \varepsilon_{i,5}$ whenever $i \in \{1,2\}$.

Since χ_0 is surjective, $1 = \chi_0(e)$ for some $e \in E_0$. Since

$$\chi_1 \varepsilon_{0,1}(e) = \iota \chi_0(e) = \iota(1) = (1, 1) = (1, 0) + (0, 1),$$

it follows from the weak distributivity of the map χ_1 and the distributivity of the semilattice E_1 that there are $a_1, b_1 \in E_1$ such that $\varepsilon_{0,1}(e) = a_1 + b_1$, $\chi_1(a_1) = (1, 0)$, and $\chi_1(b_1) = (0, 1)$. Likewise, there are $a_2, b_2 \in E_2$ such that $\varepsilon_{0,2}(e) = a_2 + b_2$, $\chi_2(a_2) = (1, 0)$, and $\chi_2(b_2) = (0, 1)$. We compute $\chi_5\varepsilon_{1,5}(a_1) = \delta_{1,5}\chi_1(a_1) = \delta_{1,5}(1, 0) = (1, 1, 0, 1)$. Similarly, $\chi_5\varepsilon_{1,5}(b_1) = (0, 0, 1, 1)$, $\chi_5\varepsilon_{2,5}(a_2) = (0, 1, 1, 1)$, and $\chi_5\varepsilon_{2,5}(b_2) = (1, 0, 0, 1)$. Since

$$\varepsilon_{1,\mu}(a_1) \leq \varepsilon_{1,\mu}(a_1 + b_1) = \varepsilon_{1,\mu}\varepsilon_{0,1}(e) = \varepsilon_{0,\mu}(e) = \varepsilon_{2,\mu}(a_2) + \varepsilon_{2,\mu}(b_2)$$

and E_μ is distributive, there are $x, y \in E_\mu$ such that $\varepsilon_{1,\mu}(a_1) = x + y$ whereas $x \leq \varepsilon_{2,\mu}(a_2)$ and $y \leq \varepsilon_{2,\mu}(b_2)$. By applying the homomorphism χ_μ to both sides of the equation $\varepsilon_{1,\mu}(a_1) = x + y$, we obtain that $(1, 1, 0, 1) = \chi_\mu(x) + \chi_\mu(y)$, with elements $\chi_\mu(x), \chi_\mu(y) \in L$ such that $\chi_\mu(x) \leq (0, 1, 1, 1)$ and $\chi_\mu(y) \leq (1, 0, 0, 1)$. Since $\chi_\mu(x)$ belongs to L , it follows (cf. Figure 12.2) that $\chi_\mu(x) = (0, 0, 0, 0)$, whence $(1, 1, 0, 1) \leq (1, 0, 0, 1)$, a contradiction. \square

Theorem 12.3 provides a negative solution for [10, Problem 1]: namely, although the functor Φ is finitely continuous with domain a variety, it is surjective on objects (cf. Proposition 12.1) but not on P_0 -indexed diagrams of $(\vee, 0)$ -embeddings, even up to weakly distributive natural transformations.

By using CLL (viz. [10, Theorem 3.4.2]), it can be verified that the counterexample underlying Theorem 12.3 has no analogue for diagrams indexed by finite lattices, or, more generally, finite *almost join-semilattices with zero* as defined in [10]. This means that for every finite almost join-semilattice P with zero and every P -indexed commutative diagram \vec{D} in \mathbf{DSLat}^0 , there are a P -indexed commutative diagram \vec{E} in \mathcal{V} and a natural transformation $\vec{\chi}: \Phi(\vec{E}) \rightarrow \vec{D}$ whose components are all *ideal-induced* in the sense of [10]; thus, *a fortiori*, both surjective and weakly distributive.

REFERENCES

- [1] Jiří Adámek and Jiří Rosický, *Locally Presentable and Accessible Categories*, London Mathematical Society Lecture Note Series, vol. 189, Cambridge University Press, Cambridge, 1994. MR 1294136 (95j:18001)
- [2] Marlow Anderson and Todd Feil, *Lattice-Ordered Groups*, Reidel Texts in the Mathematical Sciences, D. Reidel Publishing Co., Dordrecht, 1988, An introduction. MR 937703 (90b:06001)
- [3] Tibor Beke and Jiří Rosický, *Elementary equivalences and accessible functors*, Ann. Pure Appl. Logic **169** (2018), no. 7, 674–703. MR 3788743
- [4] Chen C. Chang, *Some remarks on the model theory of infinitary languages*, The Syntax and Semantics of Infinitary Languages. Edited by Jon Barwise, Springer, Berlin-New York, 1968, pp. 36–63. Lecture Notes in Mathematics, No. 72. MR 0234827
- [5] J. C. Cole and Maximo A. Dickmann, *Non-axiomatizability results in infinitary languages for higher-order structures*, Conference in Mathematical Logic—London '70 (Proc. Conf., Bedford Coll., London, 1970), Springer, Berlin, 1972, pp. 29–41. Lecture Notes in Math., Vol. 255. MR 0337562
- [6] Maximo A. Dickmann, *Larger infinitary languages*, Model-theoretic logics, Perspect. Math. Logic, Springer, New York, 1985, pp. 317–363. MR 819540
- [7] Heinz-Dieter Ebbinghaus, *Extended logics: the general framework*, Model-theoretic logics, Perspect. Math. Logic, Springer, New York, 1985, pp. 25–76. MR 819533
- [8] Paul Erdős, András Hajnal, Attila Máté, and Richard Rado, *Combinatorial Set Theory: Partition Relations for Cardinals*, Studies in Logic and the Foundations of Mathematics, vol. 106, North-Holland Publishing Co., Amsterdam, 1984. MR 795592

- [9] Ju. L. Eršov, *Teoriya Numeratsii*, “Nauka”, Moscow, 1977, *Matematicheskaya Logika i Osnovaniya Matematiki*. [Monographs in Mathematical Logic and Foundations of Mathematics]. MR 506676 (81j:03073)
- [10] Pierre Gillibert and Friedrich Wehrung, *From Objects to Diagrams for Ranges of Functors*, *Lecture Notes in Mathematics*, vol. 2029, Springer, Heidelberg, 2011. MR 2828735 (2012i:18001)
- [11] Kenneth R. Goodearl, *von Neumann regular rings and direct sum decomposition problems*, *Abelian groups and modules* (Padova, 1994), *Math. Appl.*, vol. 343, Kluwer Acad. Publ., Dordrecht, 1995, pp. 249–255. MR 1378203
- [12] George Grätzer, *Lattice Theory: Foundation*, Birkhäuser/Springer Basel AG, Basel, 2011. MR 2768581 (2012f:06001)
- [13] Tapani Hyttinen, *Model theory for infinite quantifier languages*, *Fund. Math.* **134** (1990), no. 2, 125–142. MR 1074640
- [14] Thomas Jech, *Set Theory*, *Springer Monographs in Mathematics*, Springer-Verlag, Berlin, 2003, The third millennium edition, revised and expanded. MR 1940513 (2004g:03071)
- [15] Peter T. Johnstone, *Stone Spaces*, *Cambridge Studies in Advanced Mathematics*, vol. 3, Cambridge University Press, Cambridge, 1982. MR 698074
- [16] Bjarni Jónsson, *Representations of complemented modular lattices*, *Trans. Amer. Math. Soc.* **97** (1960), 64–94. MR 0120175
- [17] ———, *Representations of relatively complemented modular lattices*, *Trans. Amer. Math. Soc.* **103** (1962), 272–302. MR 0140457
- [18] Maaret Karttunen, *Infinitary languages $N_{\infty\lambda}$ and generalized partial isomorphisms*, *Essays on mathematical and philosophical logic* (Proc. Fourth Scandinavian Logic Sympos. and First Soviet-Finnish Logic Conf., Jyväskylä, 1976), *Synthese Library*, vol. 122, Reidel, Dordrecht-Boston, Mass., 1979, pp. 153–168. MR 546435
- [19] Klaus Keimel, *Some trends in lattice-ordered groups and rings*, *Lattice theory and its applications* (Darmstadt, 1991), *Res. Exp. Math.*, vol. 23, Heldermann, Lemgo, 1995, pp. 131–161. MR 1366870
- [20] Michael Makkai and Robert Paré, *Accessible Categories: the Foundations of Categorical Model Theory*, *Contemporary Mathematics*, vol. 104, American Mathematical Society, Providence, RI, 1989. MR 1031717
- [21] Jerome Malitz, *Infinitary analogs of theorems from first order model theory*, *J. Symbolic Logic* **36** (1971), 216–228. MR 290943
- [22] Timothy Mellor and Marcus Tressl, *Non-axiomatizability of real spectra in $\mathcal{L}_{\infty\lambda}$* , *Ann. Fac. Sci. Toulouse Math. (6)* **21** (2012), no. 2, 343–358. MR 2978098
- [23] Daniele Mundici, *Advanced Łukasiewicz Calculus and MV-Algebras*, *Trends in Logic—Studia Logica Library*, vol. 35, Springer, Dordrecht, 2011. MR 2815182
- [24] Juha Oikkonen, *On PC- and RPC-classes in generalized model theory*, *Proceedings from 5th Scandinavian Logic Symposium* (Aalborg, 1979), Aalborg Univ. Press, Aalborg, 1979, pp. 257–270. MR 606603
- [25] ———, *Finite and infinite versions of the operations PC and RPC in infinitary languages*, *Workshop in Extended Model Theory* (Berlin, 1980), *Rep. MATH 1981*, vol. 3, Akad. Wiss. DDR, Berlin, 1981, pp. 114–135. MR 644812
- [26] Miroslav Ploščica, *Cevian properties in ideal lattices of Abelian ℓ -groups*, *Forum Math.* **33** (2021), no. 6, 1651–1658. MR 4333993
- [27] Pavel Pudlák, *On congruence lattices of lattices*, *Algebra Universalis* **20** (1985), no. 1, 96–114. MR 790908 (87d:06024)
- [28] Veikko Rantala, *Game-theoretical semantics and back-and-forth*, *Essays on mathematical and philosophical logic* (Proc. Fourth Scandinavian Logic Sympos. and First Soviet-Finnish Logic Conf., Jyväskylä, 1976), *Synthese Library*, vol. 122, Reidel, Dordrecht-Boston, Mass., 1979, pp. 119–151. MR 546434
- [29] Marshall H. Stone, *Topological representations of distributive lattices and Brouwerian logics*, *Čas. Mat. Fys.* **67** (1938), no. 1, 1–25.
- [30] Alfred Tarski, *Cardinal Algebras. With an Appendix: Cardinal Products of Isomorphism Types*, by Bjarni Jónsson and Alfred Tarski, Oxford University Press, New York, N. Y., 1949. MR 0029954 (10,686f)

- [31] Jiří Tůma and Friedrich Wehrung, *Congruence lifting of diagrams of finite Boolean semi-lattices requires large congruence varieties*, *Internat. J. Algebra Comput.* **16** (2006), no. 3, 541–550. MR 2241622
- [32] Heikki Matti Tuuri, *Relative separation theorems for $\mathcal{L}_{\kappa+\kappa}$* , *Notre Dame J. Formal Logic* **33** (1992), no. 3, 383–401. MR 1184069
- [33] Friedrich Wehrung, *Von Neumann coordinatization is not first-order*, *J. Math. Log.* **6** (2006), no. 1, 1–24. MR 2250951 (2007f:06011)
- [34] ———, *Schmidt and Pudlák’s Approaches to CLP*, *Lattice Theory: Special Topics and Applications*. Vol. 1, Birkhäuser/Springer, Cham, 2014, pp. 235–296. MR 3330600
- [35] ———, *Spectral spaces of countable Abelian lattice-ordered groups*, *Trans. Amer. Math. Soc.* **371** (2019), no. 3, 2133–2158. MR 3894048
- [36] ———, *Cevian operations on distributive lattices*, *J. Pure Appl. Algebra* **224** (2020), no. 4, 106202. MR 4021916
- [37] ———, *From noncommutative diagrams to anti-elementary classes*, *J. Math. Log.* **21** (2021), no. 2, Paper No. 2150011, 56. MR 4290500

NORMANDIE UNIVERSITÉ, UNICAEN, CNRS UMR 6139, LMNO, 14000 CAEN, FRANCE
E-mail address: friedrich.wehrung01@unicaen.fr
URL: <https://wehrungf.users.lmno.cnrs.fr>