



HAL
open science

Zeta Invariants of Morse Forms

Jesus Alvarez, Yuri Kordyukov, Eric Leichtnam

► **To cite this version:**

| Jesus Alvarez, Yuri Kordyukov, Eric Leichtnam. Zeta Invariants of Morse Forms. 2022. hal-03581090

HAL Id: hal-03581090

<https://hal.science/hal-03581090>

Preprint submitted on 19 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Zeta invariants of Morse forms

Jesús A. Álvarez López, Yuri A. Kordyukov and Eric Leichtnam

ABSTRACT

Given a closed real 1-form η on a closed Riemannian manifold (M, g) , let d_z , δ_z and Δ_z be the induced Witten's type perturbations of the de Rham derivative and coderivative and the Laplacian on differential forms on M , parametrized by $z \in \mathbb{C}$, and let $\zeta(s, z)$ be the zeta function of $s \in \mathbb{C}$ given by $\zeta(s, z) = \text{Tr}^s(\eta \wedge \delta_z \Delta_z^{-s})$ when $\Re s \gg 0$. For a class of Morse forms η , we prove that $\zeta(s, z)$ is smooth at $s = 1$ for $|\Re z| \gg 0$, and the zeta invariant $\zeta(1, z)$ converges to some $\mathbf{z} \in \mathbb{R}$ as $\Re z \rightarrow +\infty$, uniformly on $\Im z$. We describe \mathbf{z} in terms of the instantons of an auxiliary Smale gradient-like vector field X and the Mathai-Quillen current on TM defined by g . Any real cohomology class has a representative η satisfying the needed hypothesis. If n is even, we can prescribe any real value for \mathbf{z} by perturbing g , η and X ; if moreover M is oriented, we can also achieve the same limit as $\Re z \rightarrow -\infty$. This is used to define and describe certain tempered distributions induced by g and η . These distributions appear in another publication as the contributions from the compact leaves preserved by the flow in a trace formula for simple foliated flows on closed foliated manifolds, which gives a solution to a problem proposed by C. Deninger.

Contents

1	Introduction	2
2	Witten's perturbations	6
3	Small and large complexes	12
4	Zeta invariants of Morse forms	22
5	The small complex vs the Morse complex	32
6	Asymptotics of the large zeta invariant	42
7	Asymptotics of the small zeta-invariant	47
8	Prescription of the asymptotics of the zeta invariant	50
9	The switch of the order of integration	51
	Appendix A. Integrals along instantons	52
	References	55

2020 *Mathematics Subject Classification* 58A12, 58A14, 58J20, 57R58

Keywords: Witten's perturbation, Morse form, Morse complex, zeta function of operators, heat invariant, Ray-Singer metric.

The authors are partially supported by the grants MTM2017-89686-P and PID2020-114474GB-I00 (AEI/FEDER, UE) and ED431C 2019/10 (Xunta de Galicia, FEDER)

1. Introduction

1.1 Witten's perturbed operators

Let M be a closed n -manifold. For any smooth function h on M , Witten [Wit82] introduced a perturbed de Rham differential operator $d_\mu = d + \mu dh \wedge$, depending on a parameter $\mu \in \mathbb{R}$. Endowing M with a Riemannian metric g , we have a corresponding perturbed codifferential operator $\delta_\mu = \delta - \mu dh \lrcorner$, and a perturbed Laplacian $\Delta_\mu = d_\mu \delta_\mu + \delta_\mu d_\mu$. Since $d_\mu = e^{-\mu h} d e^{\mu h}$, it defines the same Betti numbers as d . However Δ_μ and the usual Laplacian Δ have different spectrum in general. In fact, if h is a Morse function and g is Euclidean with respect to Morse coordinates around the critical points, then the spectrum of Δ_μ develops a long gap as $\mu \rightarrow +\infty$, giving rise to the small and large spectrum. The eigenforms of the small/large eigenvalues generate the small/large subcomplex, $E_{\mu, \text{sm/la}}$. When h is a Morse function, Witten gave a beautiful analytic proof of the Morse inequalities by analyzing the small spectrum. This was refined by subsequent work of Helffer and Sjöstrand [HS85] and Bismut and Zhang [BZ92, BZ94], showing that, if moreover $X := -\text{grad } h$ is a Smale vector field, then the Morse complex $(\mathbf{C}^\bullet, \mathbf{d})$ of X can be considered as the limit of $(E_{\mu, \text{sm}}, d_\mu)$. More precisely, for certain perturbed Morse complex $(\mathbf{C}^\bullet, \mathbf{d}_\mu)$, isomorphic to $(\mathbf{C}^\bullet, \mathbf{d})$, there is a quasi-isomorphism $\Phi_\mu : (E_{z, \text{sm}}, d_\mu) \rightarrow (\mathbf{C}^\bullet, \mathbf{d}_\mu)$, defined by integration on the unstable cells of the zero points of X , which becomes an isomorphism for $\mu \gg 0$ and almost isometric as $\mu \rightarrow +\infty$ (after rescaling at every degree).

We can replace dh with any real closed 1-form η , obtaining a generalization of the Witten's perturbations, d_μ , δ_μ and Δ_μ . Now d_μ need not be gauge equivalent to d , obtaining new twisted Betti numbers β_μ^k . However the numbers β_μ^k have well defined ground values β_{No}^k , called the Novikov numbers, which depend upon the de Rham cohomology class $[\eta] \in H^1(M, \mathbb{R})$. Assume that:

- (a) η is a Morse form (it has Morse type zeros), and g is Euclidean with respect to Morse coordinates around the zero points of η .

(Some concepts used in this section are recalled in Sections 3.1 and 5.1.) Then Δ_μ also develops a long gap separating a small spectrum and a large spectrum, and the analysis of the small spectrum gives Morse inequalities for the Novikov numbers. Take any auxiliary vector field X such that:

- (b) X has Morse type zeros, and is gradient-like and Smale; and
- (c) η is Lyapunov for X , and η and g are in standard form with respect to X .

Then the small complex approaches a perturbed Morse complex of X . We refer to work by Novikov [Nov81, Nov82], Pajitnov [Paj87], Braverman and Farber [BF97], Burghelca and Haller [BH01, BH04, BH08], and Harvey and Minervini [HM06, Min15].

We can similarly define the perturbation $d_z = d + z \eta \wedge$ with parameter $z = \mu + i\nu \in \mathbb{C}$ ($\mu, \nu \in \mathbb{R}$ and $i = \sqrt{-1}$). Its adjoint is $\delta_z = \delta - \bar{z} \eta \lrcorner$, and we have a corresponding perturbed Laplacian $\Delta_z = d_z \delta_z + \delta_z d_z$. As a first step in our study, we prove extensions of the above results to this case, taking limits as $|\mu| \rightarrow \infty$, uniformly on ν . First, assuming (a), we get the long gap in the spectrum of Δ_z separating the small and large spectrum, which depends only on μ (Theorem 3.10). Second, assuming (a)–(c), we show that the quasi-isomorphism $\Phi_z : (E_{z, \text{sm}}, d_z) \rightarrow (\mathbf{C}^\bullet, \mathbf{d}_z)$ becomes an isomorphism for $|\mu| \gg 0$ and almost isometric as $|\mu| \rightarrow \infty$ (Theorem 5.3). To get that the convergence is uniform on ν , we show a version of a Sobolev inequality for a Sobolev norm defined by $\Delta_{i\nu}$, where the constant involved is independent of ν (Proposition 2.2). Then we adapt the arguments of Bismut and Zhang [BZ92, BZ94] (see also [Zha01]).

1.2 Zeta invariants of some Morse forms

Let Π_z^\perp and Π_z^1 be the orthogonal projections to the images of Δ_z and d_z , and let w be the degree involution. We consider the zeta function $\zeta(s, z) = \zeta(s, z, \eta) := \zeta(s, \Delta_z, \eta \wedge \delta_z w)$ [Gil95, Section 1.12.4]. As a function of s , this is the meromorphic extension to \mathbb{C} of the function $\text{Tr}^s(\eta \wedge \delta_z \Delta_z^{-s} \Pi_z^\perp) = \text{Tr}^s(\eta \wedge d_z^{-1} \Delta_z^{-s+1} \Pi_z^1)$, which is well defined and holomorphic for $\Re s \gg 0$. Then the zeta invariant $\zeta(1, z)$ would be a renormalized interpretation of the super-trace of $\eta \wedge d_z^{-1} \Pi_z^1$, which is not of trace class by the Weyl's law. However, according to the general theory of zeta functions of operators, $\zeta(s, z)$ might have a simple pole at $s = 1$. To study this zeta function, we decompose it as sum of the terms defined by the contributions from the small/large spectrum, $\zeta_{\text{sm/la}}(s, z) = \zeta_{\text{sm/la}}(s, z, \eta)$. As a function of s , $\zeta_{\text{sm}}(s, z)$ is always holomorphic on \mathbb{C} .

For a class of Morse forms, our first main theorem states that $\zeta(s, z)$ is smooth at $s = 1$ for $|\mu| \gg 0$, and describes the asymptotic behavior of $\zeta(1, z)$ as $\mu \rightarrow \pm\infty$, uniformly on ν . In fact, since

$$\zeta(s, z, \eta) = -\zeta(s, -z, -\eta), \quad \zeta_{\text{sm/la}}(s, z, \eta) = -\zeta_{\text{sm/la}}(s, -z, -\eta), \quad (1.1)$$

it is enough to consider the case where $\mu \gg 0$ and take the limit as $\mu \rightarrow +\infty$.

We use the current $\psi(M, \nabla^M)$ of degree $n - 1$ on TM constructed by Mathai and Quillen in [MQ86], depending on the Levi-Civita connection ∇^M . This current is smooth on the complement of the zero section, where it is given by the solid angle. It is also locally integrable, and its wave front set is contained in the conormal bundle in T^*TM of the zero section of TM . Since this set does not meet the conormal bundle of the map $X : M \rightarrow TM$ (assuming (b)), $X^*\psi(M, \nabla^M)$ is well defined as a current on M . Assuming (a)–(c), consider the real number

$$\mathbf{z}_{\text{la}} = \mathbf{z}_{\text{la}}(M, g, \eta) = \int_M \eta \wedge X^*\psi(M, \nabla^M),$$

which is known to be independent of X [BZ92, Proposition 6.1].

Now suppose also that:

- (d) for every zero point p of X , the maximum value of the integrals of η along the instantons of X with α -limit $\{p\}$ only depends on the Morse index k of p .

This maximum value is denoted by $-a_k$ for some $a_k > 0$. Let m_k^1 denote the dimension of $d_z(E_{z, \text{sm}}^{k-1})$ for $|\mu| \gg 0$, which is independent of z . Consider also the real number

$$\mathbf{z}_{\text{sm}} = \mathbf{z}_{\text{sm}}(M, g, \eta, X) = \sum_{k=1}^n (-1)^k (1 - e^{a_k}) m_k^1,$$

and let $\mathbf{z} = \mathbf{z}(M, g, \eta, X) = \mathbf{z}_{\text{sm}} + \mathbf{z}_{\text{la}}$.

THEOREM 1.1. *Let $M \equiv (M, g)$ be a closed Riemannian manifold of dimension n . Let η be a closed real 1-form on M satisfying (a).*

- (i) *For $\mu \gg 0$, $\zeta(s, z)$ is smooth for $s \neq (1 - k)/2$ ($k = 0, 1, \dots$), and*

$$\zeta(1, z) = \lim_{t \downarrow 0} \text{Tr}^s(\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1).$$

- (ii) *Let X be a vector field on M satisfying (b)–(c). Then*

$$\zeta_{\text{la}}(1, z) = \mathbf{z}_{\text{la}} + O(\mu^{-1/2})$$

as $\mu \rightarrow +\infty$, uniformly on ν .

(iii) *If moreover (d) holds, then*

$$\zeta_{\text{sm}}(1, z) = \mathbf{z}_{\text{sm}} + O(\mu^{-1})$$

as $\mu \rightarrow +\infty$, uniformly on ν .

The existence of the limit of Theorem 1.1 (i) is rather surprising because $\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1$ is weakly convergent to $\eta \wedge d_z^{-1} \Pi_z^1$. An expression similar to $\text{Tr}^s(\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1)$ was used by Mrowka, Ruberman and Saveliev to define a cyclic eta invariant [MRS16]. Theorem 1.1 (iii) shows that \mathbf{z}_{sm} and \mathbf{z} are also independent of X . Thus X will be omitted in their notation. In the notation of $\mathbf{z}_{\text{sm}/1a}$ and \mathbf{z} , we may also omit M or g if they are fixed.

By (1.1), if we take $\mu \rightarrow -\infty$ in Theorem 1.1, we have to replace $\mathbf{z}_{\text{sm}/1a}(\eta)$ with $-\mathbf{z}_{\text{sm}/1a}(-\eta)$. Descriptions of $-\mathbf{z}_{\text{sm}/1a}(-\eta)$ are given in (6.9) and (7.1).

Our second main theorem is about the prescription of $\mathbf{z} = \mathbf{z}(M, g, \eta)$ without changing the cohomology class of η .

THEOREM 1.2. *Let M be a smooth closed manifold of dimension n .*

- (i) *Let X be a vector field satisfying (b). For every $\xi \in H^1(M, \mathbb{R})$ and $\tau \gg 0$, there is some $\eta \in \xi$ and a Riemannian metric g satisfying (a), (c) and (d) with X such that $\mathbf{z}(M, g, \eta) = \tau$. If n is even, this property holds for all $\tau \in \mathbb{R}$.*
- (ii) *Assume M is oriented and n is even. Then, for every $\xi \in H^1(M, \mathbb{R})$ and $\tau \in \mathbb{R}$, there is some $\eta \in \xi$, a Riemannian metric g and a vector field X satisfying (a)–(d) such that $\pm \mathbf{z}(M, g, \pm \eta) = \tau$.*

1.3 A distribution associated to some Morse forms

A trace formula for simple foliated flows on closed foliated manifolds was conjectured by C. Deninger (see e.g. [Den08]). He was motivated by analogies with a formula in Arithmetics, and previous work of Guillemin and Sternberg [Gui77]. This trace formula is an expression for a Lefschetz distribution in terms of infinitesimal data of the flow at the fixed points and closed orbits. This Lefschetz distribution should be an analogue of the Lefschetz number for the action induced by the flow on some leafwise cohomology, whose value is a distribution on \mathbb{R} —the precise definition of these notions is part of the problem. In [ALK02, ALK08], the first two authors proved such a trace formula when the flow has no preserved leaves; see also the contributions [Lei08, Lei14] by the third author. The general case is considerably more involved. In [ALKL20], we propose a solution to this problem using a few additional ingredients. One of them is the b-trace introduced by Melrose [Mel93]. Since the b-trace is not really a trace, it produces an extra term, denoted by Z , in the same way as the eta invariant shows up in Index Theory on manifolds with boundary. In our trace formula, the term Z is a contribution from the compact leaves preserved by the flow, which depends on the choice of a form defining the foliation and a metric on the ambient manifold. But Z may not be well defined in general; it will be proved that appropriate choices of the form and the metric guarantee its existence.

Precisely, we would like to define

$$Z = Z(M, g, \eta) = \lim_{\mu \rightarrow +\infty} Z_\mu, \tag{1.2}$$

in the space of tempered distributions, where $Z_\mu = Z_\mu(M, g, \eta)$ ($\mu \gg 0$) should be a tempered distribution defined by

$$\langle Z_\mu, f \rangle = -\frac{1}{2\pi} \int_0^\infty \int_{-\infty}^\infty \text{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z}) \hat{f}(\nu) d\nu du, \tag{1.3}$$

for any Schwartz function f , where Tr^s denotes the supertrace and \hat{f} the Fourier transform of f .

Let δ_0 denote the Dirac distribution at 0 on \mathbb{R} . The problem about the definition of Z is solved in our third main theorem for the same class of Morse forms as before.

THEOREM 1.3. *Let $M \equiv (M, g)$ be a closed Riemannian manifold of dimension n . Let η be a closed 1-form on M satisfying (a), (c) and (d) with some vector field satisfying (b). Then (1.2) and (1.3) define the tempered distribution $Z = \mathbf{z}\delta_0$.*

According to Theorems 1.2 and 1.3, we can choose η and g in the trace formula for foliated flows so that $Z(M, g, \eta) = 0$ if n is even, or $Z(M, g, \pm\eta) = 0$ if moreover M is oriented, achieving the original expression of Deninger's conjecture.

It looks clear that extensions of Theorems 1.1 to 1.3 with coefficients in flat vector bundles could be similarly proved. We only consider complex coefficients for the sake of simplicity since this is enough for our application.

1.4 Some ideas of the proofs of Theorems 1.1 to 1.3

Theorem 1.1 (i) follows by using that the derived heat trace invariants up to order n of the elliptic complex d_z are independent of z , proved by Gilkey and the first author [ALG20] (Section 4.6).

Consider the meromorphic function

$$\theta(s, z) = -\zeta(s, \Delta_z, \mathbf{N}w), \quad (1.4)$$

where \mathbf{N} is the number operator, and write

$$\theta(s, z) = \theta_{\text{sm}}(s, z) + \theta_{\text{la}}(s, z),$$

where

$$\theta_{\text{sm/la}}(s, z) = -\zeta_{\text{sm/la}}(s, \Delta_z, \mathbf{N}w), \quad (1.5)$$

using the contributions from the small/large spectrum as above. Thus $e^{\theta'(0, z)/2}$ is the factor used to define the Ray-Singer metric on $\det H_z^\bullet(M)$ [BZ92], using a prime to denote ∂_s . We obtain (Corollary 4.16)

$$\zeta_{\text{la}}(1, z) = \partial_z \theta'_{\text{la}}(0, z). \quad (1.6)$$

This equality allows us to use the deep relation between the Ray-Singer metric and the Morse metric on $\det H_z^\bullet(M)$, proved by Bismut-Zhang [BZ92, BZ94]. In this way, using also that $\Phi_z : E_{z, \text{sm}} \rightarrow \mathbf{C}^\bullet$ is an isomorphism, we obtain that $\zeta_{\text{la}}(1, z)$ is asymptotic to \mathbf{z}_{la} as $\mu \rightarrow +\infty$ (Section 6.2). This proves Theorem 1.1 (ii).

When η is exact, we show this asymptotic expression of $\zeta_{\text{la}}(1, z)$ assuming only (a) (Section 4.7), without using (1.6) and the indicated strong result of Bismut-Zhang. Instead, we apply that the index density of the elliptic complex d_z is independent of z , proved by Gilkey and the first author [ALG21] and by the authors [ALKL20].

On the other hand, given any $\xi \in H^1(M, \mathbb{R})$ and a vector field X satisfying (b), we prove that there is some $\eta \in \xi$ and a metric g satisfying (a), (c) and (d) (Theorem 7.1). This can be considered as an extension of a theorem of Smale stating the existence of nice Morse functions [Sma61, Theorem B] (the case where $\xi = 0$). Its proof is relegated to Appendix A because of its different nature.

The properties (a)–(d) are used to give an asymptotic description of \mathbf{d}_z as $\mu \rightarrow +\infty$ (Section 7.2). From this asymptotic description and using that $\Phi_z : E_{z, \text{sm}} \rightarrow \mathbf{C}^\bullet$ is an isomorphism for $\mu \gg 0$, we get upper and lower bounds of the nonzero small spectrum of Δ_z (Theorem 7.5), which

are independent of ν . This is a partial extension of accurate descriptions of the nonzero small eigenvalues achieved in the case where η is exact and the parameter is real [LPNV13, Mic19]. With the same procedure and using the bounds of the nonzero small spectrum, it also follows that $\zeta_{\text{sm}}(1, z)$ is asymptotic to \mathbf{z}_{sm} as $\mu \rightarrow +\infty$ (Section 7.4), showing Theorem 1.1 (iii).

Next, by modifying η and X around its zero points of index 0 and n , without changing the cohomology class of η , we can achieve any real number as $\mathbf{z}(\eta)$, or as both $\pm\mathbf{z}(\pm\eta)$ if M is oriented and n even (Section 8). This shows Theorem 1.2.

If it is possible to switch the order of integration in (1.3),

$$\begin{aligned} \langle Z_\mu, f \rangle &= -\frac{1}{2\pi} \int_{-\infty}^{\infty} \int_0^{\infty} \text{Tr}^s (\eta \wedge \delta_z e^{-u\Delta_z}) \hat{f}(\nu) du d\nu \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \lim_{t \downarrow 0} \text{Tr}^s (\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1) \hat{f}(\nu) d\nu, \end{aligned} \quad (1.7)$$

then Theorem 1.3 is an easy consequence of Theorem 1.1. Thus it only remains to prove that both (1.3) and (1.7) define the same tempered distribution Z_μ . This follows from the Lebesgue's dominated convergence theorem and Fubini's theorem (Section 9). The verification of the hypothesis of the Fubini's theorem requires the above lower estimate of the nonzero spectrum.

2. Witten's perturbations

2.1 Preliminaries on the Witten's perturbations

2.1.1 Basic notation Let M be a closed Riemannian n -manifold. For any smooth Euclidean/Hermitean vector bundle E over M , let $C^m(M; E)$, $C^\infty(M; E)$, $L^2(M; E)$, $L^\infty(M; E)$ and $H^m(M; E)$ denote the spaces of distributional sections that are C^m , C^∞ , L^2 , L^∞ and of Sobolev order m , respectively; as usual, E is removed from this notation if it is the trivial line bundle. Consider the induced scalar product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$ on $L^2(M; E)$, and the induced norm $\|\cdot\|_{L^\infty}$ on $L^\infty(M; E)$. Fix also norms, $\|\cdot\|_m$ on every $H^m(M; E)$ and $\|\cdot\|_{C^m}$ on $C^m(M; E)$ with $\|\cdot\| = \|\cdot\|_0$ and $\|\cdot\|_{C^0} = \|\cdot\|_{L^\infty}$. If P is the orthogonal projection of $L^2(M; E)$ to some closed subspace V , then P^\perp denotes the orthogonal projection to V^\perp .

Let $T_{\mathbb{C}}M = TM \otimes \mathbb{C}$ and $T_{\mathbb{C}}^*M = T^*M \otimes \mathbb{C}$. The exterior bundle with coefficients in $\mathbb{K} = \mathbb{R}, \mathbb{C}$ is denoted by $\Lambda_{\mathbb{K}} = \Lambda_{\mathbb{K}}M$, and let $\Omega(M, \mathbb{K}) = C^\infty(M; \Lambda_{\mathbb{K}})$; in particular, $C^\infty(M, \mathbb{K}) = \Omega^0(M, \mathbb{K})$. The Levi-Civita connection is denoted by $\nabla = \nabla^M$. As usual, d and δ denote the de Rham derivative and coderivative, and let $D = d + \delta$ and $\Delta = D^2 = d\delta + \delta d$ (the Laplacian). Let $Z(M, \mathbb{K})$ and $B(M, \mathbb{K})$ denote the kernel and image of d in $\Omega(M, \mathbb{K})$. Thus $H^\bullet(M, \mathbb{K}) = Z(M, \mathbb{K})/B(M, \mathbb{K})$ is the de Rham cohomology with coefficients in \mathbb{K} . We typically consider complex coefficients, so we will omit \mathbb{K} from all of the above notation just when $\mathbb{K} = \mathbb{C}$. Take $\|\cdot\|_m$ and $\|\cdot\|_{C^m}$ given on $\Omega(M)$ by

$$\|\alpha\|_m = \sum_{k=0}^m \|D^k \alpha\|, \quad \|\alpha\|_{C^m} = \sum_{k=0}^m \|\nabla^k \alpha\|_{L^\infty}.$$

For any homogeneous linear operator between graded vector spaces, $T : V^\bullet \rightarrow W^\bullet$, the notation T_k means its precomposition with the canonical projection of V^\bullet to V^k . On any graded vector space V^\bullet , let w and \mathbf{N} be the degree involution and number operator; i.e., $w = (-1)^k$ and $\mathbf{N} = k$ on V^k .

For any $\eta \in \Omega^1(M, \mathbb{R})$ with $\eta^\sharp = X \in \mathfrak{X}(M) := C^\infty(M; TM)$, let \mathcal{L}_X and ι_X denote the Lie derivative and interior product with respect to X , and let $\eta \lrcorner = -(\eta \wedge)^* = -\iota_X$. Using the

identity $\text{Cl}(T^*M) \equiv \Lambda M$ defined by the symbol of filtered algebras, the left Clifford multiplication by η is $c(\eta) = \eta \wedge + \eta \lrcorner$, and the composition of w with the right Clifford multiplication by η is $\hat{c}(\eta) = \eta \wedge - \eta \lrcorner$; in particular, $c(\eta)^* = -c(\eta)$ and $\hat{c}(\eta)^* = \hat{c}(\eta)$. Recall that, for any $h \in C^\infty(M, \mathbb{R})$,

$$[D, h] = \hat{c}(dh). \quad (2.1)$$

In the whole paper, unless otherwise indicated, we will use the following notation without further comment. We use constants $C, c > 0$ without even mentioning their existence, and their precise values may change from line to line. We may add subindices or primes to these constants if needed. We also use a complex parameter $z = \mu + i\nu \in \mathbb{C}$ ($\mu, \nu \in \mathbb{R}$ and $i = \sqrt{-1}$). Recall that $\partial_z = (\partial_\mu - i\partial_\nu)/2$ and $\partial_{\bar{z}} = (\partial_\mu + i\partial_\nu)/2$.

2.1.2 Perturbations defined by a closed real 1-form For any $\omega \in Z^1(M)$, we have the Witten's type perturbations $d_\omega, \delta_\omega, D_\omega$ and Δ_ω of d, δ, D and Δ . Given $\eta \in Z^1(M, \mathbb{R})$ and $z \in \mathbb{C}$, we write $d_z = d_{z\eta}, \delta_z = \delta_\eta, D_z = D_{z\eta}$ and $\Delta_z = \Delta_{z\eta}$. These operators have the following expressions:

$$\left. \begin{aligned} d_z &= d + z\eta \wedge, & \delta_z &= d_z^* = \delta - \bar{z}\eta \lrcorner, \\ D_z &= d_z + \delta_z = D + \mu\hat{c}(\eta) + i\nu c(\eta) = D_{i\nu} + \mu\hat{c}(\eta), \\ \Delta_z &= D_z^2 = d_z\delta_z + \delta_z d_z = \Delta + \mu H_\eta + i\nu J_\eta + |z|^2|\eta|^2 = \Delta_{i\nu} + \mu H_\eta + \mu^2|\eta|^2, \end{aligned} \right\} \quad (2.2)$$

where, for $X = \eta^\sharp$,

$$H_\eta = D\hat{c}(\eta) + \hat{c}(\eta)D = \mathcal{L}_X^* + \mathcal{L}_X, \quad J_\eta = Dc(\eta) + c(\eta)D = \mathcal{L}_X^* - \mathcal{L}_X.$$

Note that H_η is of order zero and J_η of order one.

As families of operators, d_z and δ_z are holomorphic and anti-holomorphic functions of z , respectively. More precisely, it follows from (2.2) that

$$\left. \begin{aligned} \partial_z d_z &= \eta \wedge, & \partial_z \delta_z &= 0, & \partial_z \Delta_z &= \eta \wedge \delta_z + \delta_z \eta \wedge, \\ \partial_{\bar{z}} d_z &= 0, & \partial_{\bar{z}} \delta_z &= -\eta \lrcorner, & \partial_{\bar{z}} \Delta_z &= -\eta \lrcorner d_z - d_z \eta \lrcorner. \end{aligned} \right\} \quad (2.3)$$

The operator d_z defines an elliptic complex on $\Omega(M)$, whose cohomology is denoted by $H_z^\bullet(M)$. Since d_z has the same principal symbol as d , it is a generalized Dirac complex and Δ_z a generalized Laplacian [BGV04, Definition 2.2]. If $\theta = \eta + dh$ for some $h \in C^\infty(M, \mathbb{R})$, then the multiplication operator

$$e^{zh} : (\Omega(M), d_{z\theta}) \rightarrow (\Omega(M), d_{z\eta}) \quad (2.4)$$

is an isomorphism of differential complexes, and therefore it induces an isomorphism $H_{z\theta}^\bullet(M) \cong H_{z\eta}^\bullet(M)$. Thus the isomorphism class of $H_z^\bullet(M)$ only depends on $\xi := [\eta] \in H^1(M, \mathbb{R})$ and $z \in \mathbb{C}$. By ellipticity, D_z and Δ_z have a discrete spectrum, and there is a Hodge type decomposition

$$\Omega(M) = \ker \Delta_z \oplus \text{im } d_z \oplus \text{im } \delta_z, \quad (2.5)$$

as topological vector spaces. It induces a Hodge type isomorphism

$$H_z^\bullet(M) \cong \ker \Delta_z. \quad (2.6)$$

We also have

$$\ker \Delta_z = \ker D_z = \ker d_z \cap \ker \delta_z, \quad \text{im } \Delta_z = \text{im } D_z = \text{im } d_z \oplus \text{im } \delta_z.$$

The orthogonal projections of $\Omega(M)$ to $\ker \Delta_z, \text{im } d_z$ and $\text{im } \delta_z$ are denoted by $\Pi_z = \Pi_z^0, \Pi_z^1$ and Π_z^2 , respectively; thus $\Pi_z^\perp = \Pi_z^1 + \Pi_z^2$. The restrictions $d_z : \text{im } \delta_z \rightarrow \text{im } d_z, \delta_z : \text{im } d_z \rightarrow \text{im } \delta_z$ and

$D_z : \text{im } D_z \rightarrow \text{im } D_z$ are topological isomorphisms, and therefore the compositions $d_z^{-1}\Pi^1$, $\delta_z^{-1}\Pi^2$ and $D_z^{-1}\Pi^1$ are defined and continuous on $\Omega(M)$. Moreover the diagram

$$\begin{array}{ccc} \text{im } \delta_{z,k+1} & \xrightarrow{d_{z,k}} & \text{im } d_{z,k} \\ \Delta_{z,k} \downarrow & & \downarrow \Delta_{z,k+1} \\ \text{im } \delta_{z,k+1} & \xrightarrow{d_{z,k}} & \text{im } d_{z,k} \end{array} \quad (2.7)$$

is commutative—recall that $d_{z,k}$, $\delta_{z,k}$ and $\Delta_{z,k}$ are the compositions of the projection to $\Omega^k(M)$ with d_z , δ_z and Δ_z . The alternate sum of the dimensions $\beta_z^k = \beta_z^k(M, \xi) = \dim H_z^k(M)$ ($k = 0, \dots, n$) is the Euler characteristic [Far04, Proposition 1.40],

$$\sum_k (-1)^k \beta_z^k = \chi(M). \quad (2.8)$$

(This is also a consequence of the index theorem.) For every degree k , β_z^k is independent of z outside a discrete subset of \mathbb{C} , where β_z^k jumps (Mityagin and Novikov [Nov02, Theorem 1]). This ground value of β_z^k is called the k -th *Novikov Betti number*, and will be denoted by $\beta_{\text{No}}^k = \beta_{\text{No}}^k(M, \xi)$. Moreover it will be shown in Section 5.2.2 that

$$\beta_z^k = \beta_{\text{No}}^k \quad \text{for } |\mu| \gg 0. \quad (2.9)$$

(When z is real, this is proved in [Far95, Theorem 2.8], [BF97, Lemma 1.3], [BH04, Proposition 4].) Thus the discrete set of parameters $z \in \mathbb{C}$ with $\beta_z^k(M, \xi) > \beta_{\text{No}}^k(M, \xi)$ for some degree k is contained in a strip $|\mu| \leq C$.

By (2.2) and since η is real, for all $\alpha \in \Omega(M)$,

$$\overline{d_z \alpha} = d_{\bar{z}} \bar{\alpha}, \quad \overline{\delta_z \alpha} = \delta_{\bar{z}} \bar{\alpha}, \quad \overline{D_z \alpha} = D_{\bar{z}} \bar{\alpha}, \quad \overline{\Delta_z \alpha} = \Delta_{\bar{z}} \bar{\alpha}. \quad (2.10)$$

So conjugation induces \mathbb{C} -antilinear \mathbb{R} -isomorphisms

$$H_z^k(M) \cong H_{\bar{z}}^k(M), \quad \ker \Delta_{z,k} \cong \ker \Delta_{\bar{z},k},$$

yielding $\beta_z^k = \beta_{\bar{z}}^k$.

2.1.3 Case of an exact form When $\eta = dh$ for some $h \in C^\infty(M, \mathbb{R})$, we have the original Witten's perturbations, which satisfy

$$\left. \begin{array}{l} d_z = e^{-zh} d e^{zh} = e^{-ivh} d_\mu e^{ivh}, \quad \delta_z = e^{\bar{z}h} \delta e^{-\bar{z}h} = e^{-ivh} \delta_\mu e^{ivh}, \\ D_z = e^{-ivh} D_\mu e^{ivh}, \quad \Delta_z = e^{-ivh} \Delta_\mu e^{ivh}. \end{array} \right\} \quad (2.11)$$

Thus the multiplication operator

$$e^{zh} : (\Omega(M), d_z) \rightarrow (\Omega(M), d) \quad (2.12)$$

is an isomorphism of differential complexes, and therefore $H_z^\bullet(M) \cong H^\bullet(M)$; thus $\beta_z^k = \beta^k = \beta^k(M)$ (the k th Betti number) in this case. Moreover multiplication by e^{ivh} defines a unitary isomorphism $\ker \Delta_z \cong \ker \Delta_\mu$.

2.1.4 Interpretation of the closed form as a flat connection There is a unique flat connection $\nabla^{M \times \mathbb{C}}$ on the trivial complex line bundle $M \times \mathbb{C}$ so that $\nabla^{M \times \mathbb{C}} 1 = \eta$. The corresponding flat complex line bundle is denoted by $\mathcal{L} = \mathcal{L}_\eta$. Note that $\mathcal{L}_{z\eta} = \mathcal{L}^z$. Since every \mathcal{L}^z is canonically trivial as line bundle, it has a canonical Hermitian structure $g^{\mathcal{L}^z}$. Let $(\Omega(M, \mathcal{L}^z) = \Omega(M), d^{\mathcal{L}^z})$ be the de Rham complex with coefficients in \mathcal{L}^z .

Suppose $\eta = dh$ for a while. Then the horizontal leaves of \mathcal{L}^z are the graphs of the functions se^{-zh} ($s \in \mathbb{C}$). So $e^{zh} : \mathcal{L}^z \rightarrow \mathcal{L}^0$ is an isomorphism of flat bundles, which induces an isomorphism of differential complexes,

$$e^{zh} : (\Omega(M), d^{\mathcal{L}^z}) \rightarrow (\Omega(M), d) .$$

Comparing with the isomorphism (2.12), we get $d^{\mathcal{L}^z} = d_z$ on $\Omega(M)$. Furthermore, since $g^{\mathcal{L}^z}$ corresponds to $e^{2\mu h} g^{\mathcal{L}^0}$ via the isomorphism $e^{zh} : \mathcal{L}^z \rightarrow \mathcal{L}^0$, it follows that $\nabla^{\mathcal{L}^z} g^{\mathcal{L}^z}$ corresponds to

$$\nabla^{\mathcal{L}^0} (e^{2\mu h} g^{\mathcal{L}^0}) = 2\mu e^{2\mu h} dh \otimes g^{\mathcal{L}^0} ,$$

yielding

$$\nabla^{\mathcal{L}^z} g^{\mathcal{L}^z} = 2\mu dh \otimes g^{\mathcal{L}^z} . \quad (2.13)$$

A general η is locally exact, and therefore \mathcal{L}^z can be locally described as above. Thus $d_z = d^{\mathcal{L}^z}$ on $\Omega(M) = \Omega(M, \mathcal{L}^z)$, and (2.13) gives

$$\nabla^{\mathcal{L}^z} g^{\mathcal{L}^z} = 2\mu \eta \otimes g^{\mathcal{L}^z} . \quad (2.14)$$

2.1.5 Perturbed operators on oriented manifolds Using for instance the interpretation of d_z given in Section 2.1.4, the mapping $(\alpha, \beta) \mapsto \alpha \wedge \beta$ induces a bilinear map $H_z^k(M) \times H_{-z}^l(M) \rightarrow H^{k+l}(M)$, and the mapping $(\alpha, \beta) \mapsto \alpha \wedge \bar{\beta}$ induces a sesquilinear map $H_z^k(M) \times H_{-\bar{z}}^l(M) \rightarrow H^{k+l}(M)$.

Now assume M is oriented. Then the above maps and integration on M define a nondegenerate bilinear pairing $H_z^k(M) \times H_{-z}^{n-k}(M) \rightarrow \mathbb{C}$, and a nondegenerate sesquilinear pairing $H_z^k(M) \times H_{-\bar{z}}^{n-k}(M) \rightarrow \mathbb{C}$. Thus $\beta_z^k = \beta_{-z}^{n-k} = \beta_{-\bar{z}}^{n-k} = \beta_{\bar{z}}^k$.

Let \star denote the \mathbb{C} -linear extension to ΛM of the Hodge operator \star on $\Lambda_{\mathbb{R}} M$, which is unitary, and let $\bar{\star}$ denote its \mathbb{C} -anti-linear extension. These operators are determined by the conditions

$$\alpha \wedge \bar{\star} \beta = g(\alpha, \beta) \text{ vol} = \alpha \wedge \star \beta$$

for $\alpha, \beta \in \Omega(M)$, where $\text{vol} = \star 1$ is the volume form. The following equalities on $\Omega^k(M)$ follow from (2.2) and the usual equalities relating \star , d , δ , $\eta \wedge$ and $\eta \lrcorner$ (see e.g. [Roe98, Chapters 1 and 3], [Gil95, Section 1.5.2], [BGV04, Section 3.6]):

$$\left. \begin{aligned} d_z \star &= (-1)^k \star \delta_{-\bar{z}} , & \delta_z \star &= (-1)^{k+1} \star d_{-\bar{z}} , & \Delta_z \star &= \star \Delta_{-\bar{z}} , \\ d_z \bar{\star} &= (-1)^k \bar{\star} \delta_{-z} , & \delta_z \bar{\star} &= (-1)^{k+1} \bar{\star} d_{-z} , & \Delta_z \bar{\star} &= \bar{\star} \Delta_{-z} . \end{aligned} \right\} \quad (2.15)$$

Then we get a linear isomorphism $\star : \ker \Delta_z \rightarrow \ker \Delta_{-\bar{z}}$ and an anti-linear isomorphism $\bar{\star} : \ker \Delta_z \rightarrow \ker \Delta_{-z}$, inducing a linear isomorphism $H_z^k(M) \cong H_{-\bar{z}}^{n-k}(M)$ and an anti-linear isomorphism $H_z^k(M) \cong H_{-z}^{n-k}(M)$ by (2.6).

2.2 Perturbation of the Sobolev norms

For $m \in \mathbb{N}_0$ and $\omega \in Z^1(M)$, define the norm $\|\cdot\|_{m,\omega}$ on $H^m(M; \Lambda)$ by

$$\|\alpha\|_{m,\omega} = \sum_{k=0}^m \|D_\omega^k \alpha\| .$$

PROPOSITION 2.1. For all $\omega \in Z^1(M)$ and $\alpha \in H^m(M; \Lambda)$,

$$\|\alpha\|_{m,\omega} \leq C_m \sum_{k=0}^m \|\omega\|_{C^k}^{m-k} \|\alpha\|_k , \quad \|\alpha\|_m \leq C_m \sum_{k=0}^m \|\omega\|_{C^k}^{m-k} \|\alpha\|_{k,\omega} .$$

Proof. We proceed by induction on m . We have $\|\cdot\|_{0,\omega} = \|\cdot\|$. Now take $m > 0$ and assume these inequalities hold for $m - 1$. For $\eta \in Z^1(M, \mathbb{R})$ and $\alpha \in \Omega(M)$, we have

$$\|\hat{c}(\eta)\alpha\|_m, \|c(\eta)\alpha\|_m \leq C'_m \|\eta\|_{C^m} \|\alpha\|_m. \quad (2.16)$$

Applying these inequalities to the real and imaginary parts of ω , and using the induction hypothesis and (2.2), we get

$$\begin{aligned} \|\alpha\|_{m,\omega} &= \|\alpha\| + \|D\omega\alpha\|_{m-1,\omega} \leq \|\alpha\| + C_{m-1} \sum_{k=0}^{m-1} \|\omega\|_{C^k}^{m-1-k} \|D\omega\alpha\|_k \\ &\leq \|\alpha\| + C_{m-1} \sum_{k=0}^{m-1} \|\omega\|_{C^k}^{m-1-k} (\|D\alpha\|_k + C'_k \|\omega\|_{C^k} \|\alpha\|_k) \\ &\leq \|\alpha\| + C_{m-1} \sum_{k=0}^{m-1} \|\omega\|_{C^k}^{m-1-k} (\|\alpha\|_{k+1} + C'_k \|\omega\|_{C^k} \|\alpha\|_k) \\ &\leq C_m \sum_{l=0}^m \|\eta\|_{C^l}^{m-l} \|\alpha\|_l, \end{aligned}$$

and

$$\begin{aligned} \|\alpha\|_m &= \|\alpha\| + \|D\alpha\|_{m-1} \leq \|\alpha\| + \|D\omega\alpha\|_{m-1} + C'_{m-1} \|\omega\|_{C^{m-1}} \|\alpha\|_{m-1} \\ &\leq \|\alpha\| + C_{m-1} \sum_{k=0}^{m-1} (\|\omega\|_{C^k}^{m-1-k} \|D\omega\alpha\|_{k,\omega} + C'_{m-1} \|\omega\|_{C^k}^{m-k} \|\alpha\|_{k,\omega}) \\ &\leq \|\alpha\| + C_{m-1} \sum_{k=0}^{m-1} (\|\omega\|_{C^k}^{m-1-k} \|\alpha\|_{k+1,\omega} + C'_{m-1} \|\eta\|_{C^k}^{m-k} \|\alpha\|_{k,\omega}) \\ &\leq C_m \sum_{l=0}^m \|\omega\|_{C^l}^{m-l} \|\alpha\|_{l,\omega}. \quad \square \end{aligned}$$

Let $Z(M, \mathbb{Z}) \subset Z(M, \mathbb{R})$ denote the graded subspace of forms that represent cohomology classes in the image of the canonical homomorphism $H^\bullet(M, \mathbb{Z}) \rightarrow H^\bullet(M, \mathbb{R})$. Recall that we can consider $H^1(M, \mathbb{Z})$ as a lattice in $H^1(M, \mathbb{R})$ by the universal coefficient theorem for cohomology. Let θ be the multivalued angle function on \mathbb{S}^1 . Then $d\theta$ is the angular form on \mathbb{S}^1 with $\int_{\mathbb{S}^1} d\theta = 2\pi$. For $\eta \in Z^1(M, \mathbb{R})$, we have $\eta \in 2\pi Z^1(M, \mathbb{Z})$ if and only if there is some smooth map $h : M \rightarrow \mathbb{S}^1$ such that $\eta = h^*d\theta$ (see e.g. [Far04, Lemma 2.1]).

In Proposition 2.1, the dependence of the constants on ω cannot be avoided. For instance, for $M = \mathbb{S}^1$ with the standard metric $g = (d\theta)^2$, we have $\|1\|_m = \sqrt{2\pi}$, whereas $\|1\|_{m,i\eta} = \sqrt{2\pi} \sum_{k=0}^m |\nu|^k$ for $\eta = \nu d\theta$ ($\nu \in \mathbb{R}$). However, the following version of a Sobolev inequality for $\|\cdot\|_{m,i\eta}$ involves a constant independent of η .

PROPOSITION 2.2. *If $m > n/2$, for all $\eta \in Z^1(M, \mathbb{R})$ and $\alpha \in H^m(M; \Lambda)$,*

$$\|\alpha\|_{L^\infty} \leq C_m \|\alpha\|_{m,i\eta}.$$

Proof. By the Sobolev embedding theorem, we have

$$C_{m,i\eta} := \sup_{0 \neq \alpha \in \Omega(M)} \frac{\|\alpha\|_{L^\infty}}{\|\alpha\|_{m,i\eta}} > 0.$$

Take any $\eta \in Z^1(M, \mathbb{R})$ and $\omega \in 2\pi Z^1(M, \mathbb{Z})$, and let $\eta' = \eta + \omega$. Then $\omega = h^*d\theta$ for some smooth function $h : M \rightarrow \mathbb{S}^1$. Since the difference between the multiple values of θ at every point of \mathbb{S}^1 are in $2\pi\mathbb{Z}$, the functions $e^{\pm ih^*\theta}$ are well defined and smooth on M . Moreover, applying (2.11) locally, we get $D_{\eta'} = e^{-ih^*\theta} D_{\eta} e^{ih^*\theta}$. So, for $0 \neq \alpha \in \Omega(M)$,

$$\begin{aligned} \|\alpha\|_{L^\infty} &= \|e^{ih^*\theta} \alpha\|_{L^\infty} \leq C_{m,\eta} \|e^{ih^*\theta} \alpha\|_{m,\eta} \\ &= C_{m,\eta} \sum_{k=0}^m \|D_{\eta}^k e^{ih^*\theta} \alpha\| = C_{m,\eta} \sum_{k=0}^m \|e^{-ih^*\theta} D_{\eta}^k e^{ih^*\theta} \alpha\| \\ &= C_{m,\eta} \sum_{k=0}^m \|D_{\eta'}^k \alpha\| = C_{m,\eta} \|\alpha\|_{m,\eta'} . \end{aligned}$$

This shows that

$$\sup_{\eta' \in \eta + 2\pi Z^1(M, \mathbb{Z})} C_{m,\eta'} \leq C_{m,\eta} . \quad (2.17)$$

Since $2\pi H^1(M, \mathbb{Z})$ is a lattice in $H^1(M, \mathbb{R})$, there is a compact subset $K \subset H^1(M, \mathbb{R})$ such that

$$K + 2\pi H^1(M, \mathbb{Z}) = H^1(M, \mathbb{R}) . \quad (2.18)$$

Take a linear subspace $V \subset Z^1(M, \mathbb{R})$ such that the canonical projection $V \rightarrow H^1(M, \mathbb{R})$ is an isomorphism, and let $L \subset V$ be the compact subset that corresponds to K . By (2.18),

$$L + 2\pi Z^1(M, \mathbb{Z}) = Z^1(M, \mathbb{R}) . \quad (2.19)$$

Moreover L is bounded with respect to $\|\cdot\|_{C^m}$. Therefore, by Proposition 2.1, for all $\eta \in L$ and $\alpha \in \Omega(M)$,

$$\|\alpha\|_{L^\infty} \leq C_{m,0} \|\alpha\|_m \leq C_m \|\alpha\|_{m,\eta} ,$$

yielding

$$\sup_{\eta \in L} C_{m,\eta} \leq C_m . \quad (2.20)$$

The result follows from (2.17), (2.19) and (2.20). \square

Given $\eta \in Z^1(M, \mathbb{R})$, we write $\|\cdot\|_{m,z} = \|\cdot\|_{m,z\eta}$. Proposition 2.1 has the following direct consequence.

COROLLARY 2.3. *For all $\alpha \in H^m(M; \Lambda)$,*

$$\|\alpha\|_{m,z} \leq C_m \sum_{k=0}^m |z|^{m-k} \|\alpha\|_k , \quad \|\alpha\|_m \leq C_m \sum_{k=0}^m |z|^{m-k} \|\alpha\|_{k,z} .$$

PROPOSITION 2.4. *For all $\alpha \in H^1(M; \Lambda)$,*

$$\|\alpha\|_{1,z} \leq C(\|\alpha\|_{1,i\nu} + |\mu| \|\alpha\|) , \quad \|\alpha\|_{1,i\nu} \leq C(\|\alpha\|_{1,z} + |\mu| \|\alpha\|) .$$

Proof. By (2.2) and (2.16),

$$\begin{aligned} \|\alpha\|_{1,z} &= \|\alpha\| + \|D_z \alpha\| \leq \|\alpha\| + \|D_{i\nu} \alpha\| + C' |\mu| \|\alpha\| \leq C(\|\alpha\|_{1,i\nu} + |\mu| \|\alpha\|) , \\ \|\alpha\|_{1,i\nu} &= \|\alpha\| + \|D_{i\nu} \alpha\| \leq \|\alpha\| + \|D_z \alpha\| + C' |\mu| \|\alpha\| \leq C(\|\alpha\|_{1,z} + |\mu| \|\alpha\|) . \quad \square \end{aligned}$$

3. Small and large complexes

3.1 Preliminaries on Morse forms

Recall that a critical point p of any $h \in C^\infty(M, \mathbb{R})$ is called *nondegenerate* if $\text{Hess}_p h$ is a nondegenerate symmetric bilinear form on $T_p M$. The index of $\text{Hess}_p h$, denoted by $\text{ind}(p)$, is called the *index* of h at p . By the Morse lemma [Mil63, Lemma 2.2], this means that

$$h - h(p) = \frac{1}{2} \sum_{j=1}^n \epsilon_{p,j} (x_p^j)^2 \quad (3.1)$$

on the domain U_p of some coordinates $x_p = (x_p^1, \dots, x_p^n)$ (centered) at p , called *Morse coordinates*, where

$$\epsilon_{p,j} = \begin{cases} -1 & \text{if } j \leq \text{ind}(p) \\ 1 & \text{if } j > \text{ind}(p) . \end{cases} \quad (3.2)$$

Recall that h is called a *Morse function* when all of its critical points are nondegenerate. In this case, its critical points form a finite set denoted by $\text{Crit}(h)$. The Morse functions form an open and dense subset of $C^\infty(M, \mathbb{R})$ [Hir76, Theorem 6.1.2]. On every U_p , we can assume the metric is Euclidean with respect to Morse coordinates:

$$g = \sum_{j=1}^n (dx_p^j)^2 . \quad (3.3)$$

Now take any $\eta \in Z^1(M, \mathbb{R})$. A zero p of η is said to be *nondegenerate* if $\eta = dh_{\eta,p}$ around p for some local Morse function $h_{\eta,p}$, which is chosen so that $h_{\eta,p}(p) = 0$. The index $\text{ind}(p)$ of $h_{\eta,p}$ at p is also called the *index* of η at p . On the domain U_p of Morse coordinates $x_p = (x_p^1, \dots, x_p^n)$ for $h_{\eta,p}$ at p , $h_{\eta,p}$ is given by the right-hand sides of (3.1), and

$$\eta = \sum_{j=1}^n \epsilon_{p,j} x_p^j dx_p^j . \quad (3.4)$$

It is also said that $x_p = (x_p^1, \dots, x_p^n)$ are *Morse coordinates* for η at p . With the notation

$$x_p^- = (x_p^1, \dots, x_p^k) : U_p \rightarrow \mathbb{R}^k , \quad x_p^+ = (x_p^{k+1}, \dots, x_p^n) : U_p \rightarrow \mathbb{R}^{n-k} ,$$

for $k = \text{ind}(p)$, we can write

$$h_{\eta,p} = \frac{1}{2} (|x_p^+|^2 - |x_p^-|^2) . \quad (3.5)$$

If all zeros are nondegenerate, then η is called a *Morse form*. In this case, its zeros form a finite set, $\mathcal{X} = \text{Zero}(\eta)$; subsets of \mathcal{X} defined by conditions on the index are denoted by writing the conditions as subscripts; for instance, \mathcal{X}_k , \mathcal{X}_+ and $\mathcal{X}_{<k}$ are the subsets of zeros of index k , of positive index, and of index $< k$, respectively. For any $\xi \in H^1(M, \mathbb{R})$, the Morse representatives of ξ form a dense open subset of ξ , considered as a subspace of $\Omega^1(M, \mathbb{R})$ with the C^∞ topology (see e.g. [Paj06, Theorem 2.1.25]).

From now on, unless otherwise stated, we will use some $\eta \in Z^1(M, \mathbb{R})$ and a Riemannian metric g on M such that η is a Morse form, and g is Euclidean with respect to Morse coordinates around zero points of η ; i.e., η and g satisfy (a) (Section 1.1).

The Hopf index of $-\eta^\sharp$ at every $p \in \mathcal{X}_k$ is $(-1)^k$ (see Section 5.1.1). Thus, by the Hopf index

theorem for $-\eta^\sharp$,

$$\sum_{k=0}^n (-1)^k |\mathcal{X}_k| = \chi(M). \quad (3.6)$$

3.2 The small and large spectrum

Consider the perturbed operators (2.2) defined by η and g . Recall that \mathcal{X} denotes the zero set of η . We can suppose the closures of the domains of Morse coordinates, $\overline{U_p}$ ($p \in \mathcal{X}$), are disjoint from each other, and $x_p(U_p) = (-4r, 4r)^n$ for some $r > 0$ independent of p . Let $U = \bigcup_{p \in \mathcal{X}} U_p$.

Denoting also the coordinates of \mathbb{R}^n by (x_p^1, \dots, x_p^n) , consider the function $h_p \in C^\infty(\mathbb{R}^n)$ defined by the right-hand side of (3.1) (or (3.5)). Let $d'_{p,z}$, $\delta'_{p,z}$, $D'_{p,z}$ and $\Delta'_{p,z}$ ($z \in \mathbb{C}$) denote the corresponding Witten's operators on \mathbb{R}^n , whose restrictions to $(-4r, 4r)^n$ agree via x_p with the restrictions of d_μ , δ_μ , D_μ and Δ_μ to U_p .

PROPOSITION 3.1 See e.g. [Roe98, Chapters 9 and 14], [Zha01, Sections 4.5 and 4.7]. *The following holds for $\mu \in \mathbb{R}$:*

(i) *We have*

$$\Delta'_{p,\mu} = \sum_{j=1}^n \left(- \left(\frac{\partial}{\partial x_p^j} \right)^2 + \mu^2 (x_p^j)^2 + \mu \epsilon_{p,j} [dx_p^j \lrcorner, dx_p^j \wedge] \right). \quad (3.7)$$

Moreover, using multi-index notation,

$$[dx_p^j \lrcorner, dx_p^j \wedge] dx_p^J = \begin{cases} dx_p^J & \text{if } j \in J \\ -dx_p^J & \text{if } j \notin J. \end{cases}$$

(ii) $\Delta'_{p,\mu}$ is a non-negative selfadjoint operator in $L^2(\mathbb{R}^n; \Lambda)$ with a discrete spectrum, which consists of the eigenvalues

$$\mu \sum_{j=1}^n (1 + 2u_j + \epsilon_{p,j} v_j), \quad (3.8)$$

where $u_j \in \mathbb{N}_0$ and $v_j = \pm 1$. For the restriction of $\Delta'_{p,\mu}$ to k -forms, the spectrum has the additional requirement that exactly k of the numbers v_j are equal to 1. In particular, 0 is an eigenvalue of $\Delta'_{p,\mu}$ with multiplicity 1 (choosing $u_j = 0$ and $v_j = -\epsilon_{p,j}$ for all j), and the nonzero eigenvalues are of order μ as $\mu \rightarrow +\infty$. $D'_{p,\mu}$ is also a selfadjoint operator in $L^2(\mathbb{R}^n; \Lambda)$ with a discrete spectrum, which consists of the positive and negative square roots of (3.8).

(iii) The kernel of $D'_{p,\mu}$ and $\Delta'_{p,\mu}$ is generated by the normalized form

$$e'_{p,\mu} = \left(\frac{\mu}{\pi} \right)^{n/4} e^{-\mu|x_p|^2/2} dx_p^1 \wedge \dots \wedge dx_p^{\text{ind}(p)}.$$

For any $z \in \mathbb{C}$ with $\mu > 0$, let $\Delta'_{p,z} = e^{-i\nu h_p} \Delta'_{p,\mu} e^{i\nu h_p}$. Since the operator of multiplication by $e^{-i\nu h_p}$ is unitary, $\Delta'_{p,z}$ is also selfadjoint and non-negative in $L^2(\mathbb{R}^n; \Lambda)$, it has a discrete spectrum with the same eigenvalues and multiplicities as $\Delta'_{p,\mu}$, and its kernel is generated by the normalized form $e'_{p,z} := e^{-i\nu h_p} e'_{p,\mu}$. We will also use the notation

$$e'_{p,z} = x_p^* e'_{p,z} \in C^\infty(U_p; \Lambda^{\text{ind}(p)}).$$

The function $x_p^* h_p \in C^\infty(U_p)$ agrees with $h_{\eta,p}$, which is also denoted by h_p in this section.

Fix an even C^∞ function $\rho : \mathbb{R} \rightarrow [0, 1]$ such that $\rho = 1$ on $[-r, r]$ and $\text{supp } \rho \subset [-2r, 2r]$. For every $p \in \mathcal{X}$, let

$$\rho_p = \rho(x_p^1) \cdots \rho(x_p^n) \in C_c^\infty(U_p), \quad (3.9)$$

$$e_{p,\mu} = \frac{\rho_p}{a_\mu} e'_{p,\mu} \in C_c^\infty(U_p; \Lambda^{\text{ind}(p)}), \quad (3.10)$$

$$e_{p,z} = e^{-ivh_p} e_{p,\mu} = \frac{\rho_p}{a_\mu} e'_{p,z} \in C_c^\infty(U_p; \Lambda^{\text{ind}(p)}), \quad (3.11)$$

where

$$a_\mu = \left(\int_{-2r}^{2r} \rho(x)^2 e^{-\mu x^2} dx \right)^{n/2} = \left(\frac{\pi}{\mu} \right)^{n/4} + O(e^{-c\mu}), \quad (3.12)$$

as $\mu \rightarrow +\infty$. The extensions by zero of the forms $e_{p,z}$ to M are also denoted by $e_{p,z}$. They form an orthonormal basis of a graded subspace $E_z \subset \Omega(M)$ with $\dim E_z = |\mathcal{X}|$. Let P_z be the orthogonal projection of $L^2(M; \Lambda)$ to E_z ; thus P_z^\perp is the orthogonal projection to E_z^\perp .

Remark 3.2. For the sake of simplicity, most of our results are stated for $\mu \gg 0$ or as $\mu \rightarrow +\infty$, but they have obvious versions for $\mu \ll 0$ or as $\mu \rightarrow -\infty$, as follows by considering $-\eta$.

PROPOSITION 3.3. *If $\mu \gg 0$ and $\beta \in H^1(M; \Lambda)$ with $\text{supp } \beta \subset M \setminus U$, then*

$$\|D_z \beta\| \geq C\mu \|\beta\|.$$

Proof. This follows like [Zha01, Proposition 4.7], using that H_η is of order zero in (2.2). Actually, according to the statement of [Zha01, Proposition 4.7], this inequality would hold with $\sqrt{\mu}$ instead of μ , but its proof clearly shows that using μ is fine. \square

PROPOSITION 3.4. *The following properties hold:*

(i) $P_z D_z P_z = 0$.

(ii) *If $\mu \gg 0$, $\alpha \in E_z$ and $\beta \in E_z^\perp \cap H^1(M; \Lambda)$, then*

$$\|P_z^\perp D_z \alpha\| \leq e^{-c\mu} \|\alpha\|, \quad \|P_z D_z \beta\| \leq e^{-c\mu} \|\beta\|.$$

(iii) *If $\mu \gg 0$ and $\beta \in E_z^\perp \cap H^1(M; \Lambda)$, then*

$$\|P_z^\perp D_z \beta\| \geq C\sqrt{\mu} \|\beta\|.$$

Proof. This follows like [Zha01, Propositions 4.11, 4.12 and 5.6]. Property (i) is true because every $D_z e_{p,z}$ is supported in U_p and has homogeneous components of degree different from $\text{ind}(p)$; therefore it is orthogonal to $\ker \Delta_z$. The other properties are consequences of Propositions 3.1 and 3.3 and (3.9)–(3.12). According to [Zha01, Proposition 4.11], the inequalities of (ii) hold with $1/\mu$ instead of $e^{-c\mu}$, but its proof shows that indeed $e^{-c\mu}$ can be achieved. \square

PROPOSITION 3.5. *For all $m \in \mathbb{N}_0$, if $\mu \gg 0$, then*

$$\|D_z e_{p,z}\|_m, \|D_z e_{p,z}\|_{m,iv} \leq e^{-cm\mu}.$$

Proof. From Proposition 3.1 (iii), (2.1), (3.10) and (3.11), we get

$$D_z e_{p,z} = D_z \left(\frac{\rho_p}{a_\mu} e'_{p,z} \right) = e^{-ivh_p} \frac{1}{a_\mu} \left(\frac{\pi}{4} \right)^{n/4} \hat{c}(d\rho_p) e'_{p,\mu}. \quad (3.13)$$

Thus the stated estimate of $\|D_z e_{p,z}\|_m$ is true by (3.10) and (3.12), and since $d\rho_p = 0$ around p . (When $\nu = 0$, this is indicated in [Zha01, Eq. (6.17)].)

By (2.11), for all $k \in \mathbb{N}_0$ and $p \in \mathcal{X}$, the form $D_{iv}^k D_z e_{p,z}$ is the extension by zero of the form $e^{-ivh_p} D_\mu^k D_\mu e_{p,\mu}$ on U_p . Then the stated estimate of $\|D_z e_{p,z}\|_{m,iv}$ follows from the case $\nu = 0$. \square

COROLLARY 3.6. *If $\mu \gg 0$, then*

$$\|D_z e_{p,z}\|_{L^\infty} \leq e^{-c\mu}.$$

Proof. Apply Propositions 2.2 and 3.5. □

Consider the partition of $\text{spec } \Delta_z$ into its intersections with $[0, 1]$ and $(1, \infty)$, called the *small* and *large spectrum*; the term *small/large eigenvalues* may be also used. Let $E_{z,\text{sm}} \subset \Omega(M)$ denote the graded finite dimensional subspace generated by the eigenforms of the small eigenvalues, let $E_{z,\text{la}} = E_{z,\text{sm}}^\perp$ in $L^2(M; \Lambda)$, and let $P_{z,\text{sm/la}}$ be the orthogonal projections to $E_{z,\text{sm/la}}$. Note that $E_{z,\text{sm}}$ and $E_{z,\text{la}} \cap \Omega(M)$ are subcomplexes with d_z , called the *small* and *large complexes*, and the term *small/large projection* will be used for $P_{z,\text{sm/la}}$. Thus $(\Omega(M), d_z)$ splits into a topological direct sum of the subcomplexes $E_{z,\text{sm}}$ and $E_{z,\text{la}} \cap \Omega(M)$, and (2.5) gives

$$H^\bullet(E_{z,\text{sm}}, d_z) \cong H_z^\bullet(M), \quad H^\bullet(E_{z,\text{la}} \cap \Omega(M), d_z) = 0. \quad (3.14)$$

For any operator B defined on $\Omega(M)$ or $L^2(M; \Lambda)$, let $B_{z,\text{sm/la}} = BP_{z,\text{sm/la}}$.

PROPOSITION 3.7. *For all $m \in \mathbb{N}_0$, $\mu \gg 0$ and $\alpha \in E_z$,*

$$\|\alpha - P_{z,\text{sm}}\alpha\|_{m,i\nu} \leq e^{-c_m\mu}\|\alpha\|.$$

Proof. This follows like [Zha01, Lemma 5.8 and Theorem 6.7], using $\|\cdot\|_{m,i\nu}$ instead of $\|\cdot\|_m$. The following are the main steps of the proof.

Let $\mathbb{S}^1 = \{\omega \in \mathbb{C} \mid |\omega| = 1\}$. With the argument of the proof of [Zha01, Eq. (5.27)], using Proposition 3.4, we get that, for all $\alpha \in H^1(M; \Lambda)$, $w \in \mathbb{S}^1$ and $\mu \gg 0$,

$$\|(w - D_z)\alpha\| \geq C\|\alpha\|.$$

Thus $w - D_z : H^1(M; \Lambda) \rightarrow L^2(M; \Lambda)$ has a bounded inverse and, for all $\beta \in L^2(M; \Lambda)$, $w \in \mathbb{S}^1$ and $\mu \gg 0$,

$$\|(w - D_z)^{-1}\beta\| \leq C^{-1}\|\beta\|. \quad (3.15)$$

On the other hand, arguing like in the proof of [Zha01, Eq. (6.18)], it follows that, for all $\gamma \in H^m(M; \Lambda)$, $w \in \mathbb{S}^1$ and $\mu \gg 0$,

$$\|\gamma\|_{m,i\nu} \leq C_m(\|(w - D_z)\gamma\|_{m-1,i\nu} + \mu\|\gamma\|_{m-1,i\nu} + \|\gamma\|).$$

Continuing by induction on $m \in \mathbb{N}_0$, we obtain

$$\|\gamma\|_{m,i\nu} \leq C_m\left(\mu^m\|\gamma\| + \sum_{k=1}^m \mu^{k-1}\|(w - D_z)\gamma\|_{m-k,i\nu}\right).$$

In other words, for all $\beta \in H^{m-1}(M; \Lambda)$,

$$\|(w - D_z)^{-1}\beta\|_{m,i\nu} \leq C_m\left(\mu^m\|(w - D_z)^{-1}\beta\| + \sum_{k=1}^m \mu^{k-1}\|\beta\|_{m-k,i\nu}\right).$$

Applying (3.15) to this inequality, we get

$$\|(w - D_z)^{-1}\beta\|_{m,i\nu} \leq C_m\mu^m\|\beta\|_{m-1,i\nu}. \quad (3.16)$$

Then, by Proposition 3.5,

$$\|(w - D_z)^{-1}D_z e_{p,z}\|_{m,i\nu} = O(e^{-c_m\mu}) \quad (3.17)$$

as $\mu \rightarrow +\infty$, uniformly on $w \in \mathbb{S}^1$. But, endowing \mathbb{S}^1 with the counter-clockwise orientation, basic spectral theory gives (see e.g. [DS88a, Section VII.3])

$$\begin{aligned} P_{z,\text{sm}}e_{p,z} - e_{p,z} &= \frac{1}{2\pi i} \int_{\mathbb{S}^1} ((w - D_z)^{-1} - w^{-1})e_{p,z} dw \\ &= \frac{1}{2\pi i} \int_{\mathbb{S}^1} w^{-1}(w - D_z)^{-1}D_z e_{p,z} dw . \end{aligned} \quad (3.18)$$

The result follows using (3.17) in (3.18). \square

COROLLARY 3.8. For $\mu \gg 0$ and $\alpha \in E_z$,

$$\|\alpha - P_{z,\text{sm}}\alpha\|_{L^\infty} \leq e^{-c\mu} \|\alpha\| .$$

Proof. Apply Propositions 2.2 and 3.7.

Alternatively, the proof of Proposition 3.7 can be modified as follows to get this result (some step of this alternative argument will be used later). Iterating (3.16), we get

$$\|(w - D_z)^{-1}\beta\|_{m,i\nu} \leq C'_m \mu^{(m+1)m/2} \|\beta\| ,$$

for all $\beta \in L^2(M; \Lambda)$. Then, by Proposition 2.2,

$$\|(w - D_z)^{-1}\beta\|_{L^\infty} \leq C\mu^{(m+1)m/2} \|\beta\| . \quad (3.19)$$

Thus, by Proposition 3.5,

$$\|(w - D_z)^{-1}D_z e_{p,z}\|_{L^\infty} = O(e^{-c_m\mu})$$

as $\mu \rightarrow +\infty$. Finally, apply this expression in (3.18). \square

COROLLARY 3.9. If $\mu \gg 0$, then $P_{z,\text{sm}} : E_z \rightarrow E_{z,\text{sm}}$ is an isomorphism; in particular, $\dim E_{z,\text{sm}} = |\mathcal{X}|$ and $\dim E_{z,\text{sm}}^k = |\mathcal{X}_k|$.

Proof. The result follows from Proposition 3.7 for $m = 0$ like [Zha01, Proposition 5.5]. \square

When $\mu \gg 0$, (3.6) also follows from Corollary 3.9, (2.8) and (3.14).

THEOREM 3.10 Cf. [BH01, Theorem 3]. We have

$$\text{spec } \Delta_z \subset [0, e^{-c|\mu|}] \cup [C|\mu|, \infty) .$$

Proof. We can assume $\mu \geq 0$ according to Remark 3.2. By Propositions 2.4, 3.4 and 3.7, for all $\alpha \in E_z$,

$$\begin{aligned} \|D_z P_{z,\text{sm}}\alpha\| &\leq \|D_z\alpha\| + \|D_z(\alpha - P_{z,\text{sm}}\alpha)\| \leq \|D_z\alpha\| + \|\alpha - P_{z,\text{sm}}\alpha\|_{1,z} \\ &\leq \|P_z^\perp D_z\alpha\| + C(\mu\|\alpha - P_{z,\text{sm}}\alpha\| + \|\alpha - P_{z,\text{sm}}\alpha\|_{1,i\nu}) \\ &\leq (e^{-c\mu} + C(\mu e^{-c_0\mu} + e^{-c_1\mu}))\|\alpha\| . \end{aligned}$$

Hence, by Corollary 3.9, for all $\beta \in E_{z,\text{sm}}$,

$$0 \leq \langle \Delta_z \beta, \beta \rangle = \|D_z \beta\|^2 \leq e^{-c\mu} \|\beta\|^2 .$$

This shows that

$$\text{spec } \Delta_z \cap [0, 1] \subset [0, e^{-c\mu}] . \quad (3.20)$$

Now let $\phi \in E_{z,\text{la}} \cap H^1(M; \Lambda)$, and write $\alpha = P_z \phi \in E_z$ and $\beta = P_z^\perp \phi \in E_z^\perp \cap H^1(M; \Lambda)$. By Proposition 3.7,

$$\|\alpha\|^2 = \langle \alpha, \phi \rangle = \langle \alpha - P_{z,\text{sm}}\alpha, \phi \rangle \leq \|\alpha - P_{z,\text{sm}}\alpha\| \|\phi\| \leq e^{-c_0\mu} \|\alpha\| \|\phi\| ,$$

yielding

$$\|\alpha\| \leq e^{-c_0\mu} \|\phi\| .$$

So

$$\|\beta\| = \|\phi - \alpha\| \geq \|\phi\| - \|\alpha\| \geq (1 - e^{-c_0\mu}) \|\phi\| .$$

Then, by Proposition 3.4,

$$\begin{aligned} \|D_z\phi\| &\geq \|D_z\beta\| - \|D_z\alpha\| \geq \|P_z^\perp D_z\beta\| - e^{-c\mu} \|\alpha\| \\ &\geq C\sqrt{\mu} \|\beta\| - e^{-c\mu} \|\phi\| \geq (C\sqrt{\mu}(1 - e^{-c_0\mu}) - e^{-c\mu}) \|\phi\| . \end{aligned}$$

Therefore, for all $\phi \in E_{z,\text{la}} \cap H^1(M; \Lambda)$,

$$\langle \Delta_z\phi, \phi \rangle = \|D_z\phi\|^2 \geq C\mu \|\phi\|^2 .$$

This proves that

$$\text{spec } \Delta_z \cap (1, \infty) \subset [C\mu, \infty) . \quad (3.21)$$

The inclusions (3.20) and (3.21) give the result for $\mu \gg 0$. But, in those inclusions, we can take c and C so small that, if one of them is not true for some $\mu \geq 0$, then $C\mu \leq e^{-c\mu}$. \square

3.3 Ranks of some projections in the small complex

Recall that $(\Pi_z^\perp)_{\text{sm},k}$, $\Pi_{z,\text{sm},k}^1$ and $\Pi_{z,\text{sm},k}^2$ denote the orthogonal projections to the images of $\Delta_{z,\text{sm},k}$, $d_{z,\text{sm},k-1}$ and $\delta_{z,\text{sm},k+1}$, respectively. Let $m_{z,k}$, $m_{z,k}^1$ and $m_{z,k}^2$ be the corresponding ranks (or traces) of these projections. They satisfy

$$m_{z,k} = m_{z,k}^1 + m_{z,k}^2, \quad m_{z,0}^1 = m_{z,n}^2 = 0, \quad m_{z,k}^2 = m_{z,k+1}^1, \quad (3.22)$$

where the last equality is true because $d_z : \text{im } \delta_z \rightarrow \text{im } d_z$ is an isomorphism. For $\mu \gg 0$, we have $m_{z,k}, m_{z,k}^j \leq |\mathcal{X}_k|$ by Corollary 3.9 and (3.22).

LEMMA 3.11. *The numbers $m_{z,k}^j$ are determined by the numbers $m_{z,k}$.*

Proof. This follows from (3.22) with an easy induction argument on k . \square

LEMMA 3.12. *We have $m_{z,k} = |\mathcal{X}_k| - \beta_z^k$.*

Proof. This is a consequence of (2.5), (3.14) and Corollary 3.9. \square

COROLLARY 3.13. $\text{Tr}^s((\Pi_z^\perp)_{\text{sm}}) = 0$.

Proof. By (2.8), (3.6) and Lemma 3.12,

$$\text{Tr}^s((\Pi_z^\perp)_{\text{sm}}) = \sum_k (-1)^k |\mathcal{X}_k| - \sum_k (-1)^k \beta_z^k = \chi(M) - \chi(M) = 0 . \quad \square$$

LEMMA 3.14. *If M is oriented, then, for $k = 0, \dots, n$,*

$$m_{z,k} = m_{-\bar{z},n-k} = m_{-z,n-k}, \quad m_{z,k}^1 = m_{-\bar{z},n-k}^2 = m_{-z,n-k}^2 .$$

Proof. This is true because, by (2.15),

$$\begin{aligned} (\Pi_z^\perp)_{\text{sm},k} \star &= \star (\Pi_{-\bar{z}}^\perp)_{\text{sm},n-k}, & \Pi_{z,\text{sm},k}^1 \star &= \star \Pi_{-\bar{z},\text{sm},n-k}^2, \\ (\Pi_z^\perp)_{\text{sm},k} \bar{\star} &= \bar{\star} (\Pi_{-z}^\perp)_{\text{sm},n-k}, & \Pi_{z,\text{sm},k}^1 \bar{\star} &= \bar{\star} \Pi_{-z,\text{sm},n-k}^2 . \quad \square \end{aligned}$$

COROLLARY 3.15. *For $|\mu| \gg 0$, $m_{z,k}$ and $m_{z,k}^j$ only depend on $|\mathcal{X}_k|$ and the class $\xi = [\eta] \in H^1(M, \mathbb{R})$.*

Proof. Apply (2.9) and Lemmas 3.11 and 3.12. \square

By Corollary 3.15, we write $m_k = m_{z,k}$ and $m_k^j = m_{z,k}^j$ for $|\mu| \gg 0$.

COROLLARY 3.16. $\text{Tr}^s(\Pi_{z,\text{sm}}^j) = 0$ if M is oriented, n is even and $\mu \gg 0$.

Proof. By Corollaries 3.13 and 3.15 and Lemma 3.14, and since n is even,

$$\text{Tr}^s(\Pi_{z,\text{sm}}^1) = -\text{Tr}^s(\Pi_{z,\text{sm}}^2) = -\text{Tr}^s(\Pi_{z,\text{sm}}^1). \quad \square$$

3.4 Asymptotic properties of the small projection

Notation 3.17. Consider a function $f(x) > 0$ ($x > 0$). When referring to vectors in Banach spaces, the order notation $O(f(|\mu|))$ ($\mu \rightarrow \pm\infty$) will be used for a family of vectors $v = v(z)$ ($z \in \mathbb{C}$) with $\|v(z)\| = O(f(|\mu|))$. This notation applies e.g. to bounded operators between Banach spaces. We may also consider this notation when the Banach spaces depend on z .

PROPOSITION 3.18. For every $\tau \in \mathbb{R}$, on $L^2(M; \Lambda)$, as $\mu \rightarrow +\infty$,

$$P_{z,\text{sm}} = P_z + O(e^{-c\mu}) = P_{z,\text{sm}}P_{z+\tau,\text{sm}}P_{z,\text{sm}} + O(\mu^{-2}) = P_{z+\tau,\text{sm}} + O(\mu^{-1}).$$

Proof. By Corollary 3.9, for $\mu \gg 0$, the elements $P_{z,\text{sm}}e_{p,z}$ ($p \in \mathcal{X}$) form a base of $E_{z,\text{sm}}$. Applying the Gram-Schmidt process to this base, we get an orthonormal base $\tilde{e}_{p,z}$. By Proposition 3.7,

$$\tilde{e}_{p,z} = e_{p,z} + O(e^{-c\mu}). \quad (3.23)$$

This gives the first equality of the statement: for any $\alpha \in L^2(M; \Lambda)$,

$$P_z\alpha = \sum_{p \in \mathcal{X}} \langle \alpha, e_{p,z} \rangle e_{p,z} = \sum_{p \in \mathcal{X}} \langle \alpha, \tilde{e}_{p,z} \rangle \tilde{e}_{p,z} + O(e^{-c\mu})\|\alpha\| = P_{z,\text{sm}}\alpha + O(e^{-c\mu})\|\alpha\|.$$

Since the sets U_p ($p \in \mathcal{X}$) are disjoint one another, for $p \neq q$ in \mathcal{X} ,

$$\langle e_{p,z}, e_{q,z+\tau} \rangle = 0. \quad (3.24)$$

On the other hand, by (3.9)–(3.12), we can also assume

$$\begin{aligned} \langle e_{p,z}, e_{p,z+\tau} \rangle &= \langle e^{-i\nu h_p} e_{p,\mu}, e^{-i\nu h_p} e_{p,\mu+\tau} \rangle = \langle e_{p,\mu}, e_{p,\mu+\tau} \rangle \\ &= \frac{(\mu(\mu+\tau))^{n/4}}{\pi^{n/2}} \langle \rho_p e^{-\mu|x_p|^2/2}, \rho_p e^{-(\mu+\tau)|x_p|^2/2} \rangle + O(e^{-c\mu}) \\ &= \frac{(\mu(\mu+\tau))^{n/4}}{\pi^{n/2}} \int_{\mathbb{R}^n} e^{-(\mu+\tau/2)|x_p|^2} dx_p + O(e^{-c\mu}) \\ &= \frac{(\mu(\mu+\tau))^{n/4}}{(\mu+\tau/2)^{n/2}} + O(e^{-c\mu}) = 1 + O(\mu^{-2}), \end{aligned} \quad (3.25)$$

where $dx_p = dx_p^1 \dots dx_p^n = \text{vol}(x_p)$. Combining (3.23) for z and $z+\tau$ with (3.24) and (3.25), we obtain

$$\begin{aligned} P_{z+\tau,\text{sm}}\tilde{e}_{p,z} &= \sum_{q \in \mathcal{X}} \langle \tilde{e}_{p,z}, \tilde{e}_{q,z+\tau} \rangle \tilde{e}_{q,z+\tau} = \sum_{q \in \mathcal{X}} \langle e_{p,z}, e_{q,z+\tau} \rangle e_{q,z+\tau} + O(e^{-c\mu}) \\ &= e_{p,z+\tau} + O(\mu^{-2}) = \tilde{e}_{p,z+\tau} + O(\mu^{-2}). \end{aligned} \quad (3.26)$$

Repeating (3.26) interchanging the roles of z and $z+\tau$, we get

$$P_{z,\text{sm}}P_{z+\tau,\text{sm}}\tilde{e}_{p,z} = P_{z,\text{sm}}\tilde{e}_{p,z+\tau} + O(\mu^{-2}) = \tilde{e}_{p,z} + O(\mu^{-2}).$$

This gives the second equality of the statement: for any $\alpha \in L^2(M; \Lambda)$,

$$\begin{aligned} P_{z,\text{sm}}\alpha &= \sum_{p \in \mathcal{X}} \langle \alpha, \tilde{e}_{p,z} \rangle \tilde{e}_{p,z} = P_{z,\text{sm}} P_{z+\tau,\text{sm}} \sum_{p \in \mathcal{X}} \langle \alpha, \tilde{e}_{p,z} \rangle \tilde{e}_{p,z} + O(\mu^{-2}) \|\alpha\| \\ &= P_{z,\text{sm}}\alpha + O(\mu^{-2}) \|\alpha\|. \end{aligned}$$

By (3.26),

$$\begin{aligned} \|\tilde{e}_{p,z} - \tilde{e}_{p,z+\tau}\|^2 &= \|\tilde{e}_{p,z}\|^2 - 2\Re\langle \tilde{e}_{p,z}, \tilde{e}_{p,z+\tau} \rangle + \|\tilde{e}_{p,z+\tau}\|^2 = 2 - 2\Re\langle P_{z+\tau,\text{sm}}\tilde{e}_{p,z}, \tilde{e}_{p,z+\tau} \rangle \\ &= 2 - 2\Re\langle \tilde{e}_{p,z+\tau}, \tilde{e}_{p,z+\tau} \rangle + O(\mu^{-2}) = O(\mu^{-2}), \end{aligned}$$

which means

$$\tilde{e}_{p,z} = \tilde{e}_{p,z+\tau} + O(\mu^{-1}). \quad (3.27)$$

The last stated equality follows from (3.26) and (3.27): for any $\alpha \in L^2(M; \Lambda)$,

$$P_{z,\text{sm}}\alpha = \sum_{p \in \mathcal{X}} \langle \alpha, \tilde{e}_{p,z} \rangle \tilde{e}_{p,z} = \sum_{p \in \mathcal{X}} \langle \alpha, \tilde{e}_{p,z+\tau} \rangle \tilde{e}_{p,z+\tau} + O(\mu^{-1})\alpha = P_{z+\tau,\text{sm}}\alpha + O(\mu^{-1})\alpha. \quad \square$$

COROLLARY 3.19. *For every $\tau \in \mathbb{R}$, on $L^2(M; \Lambda)$,*

$$d_{z+\tau,\text{sm}} - d_{z+\tau} P_{z,\text{sm}} = O(\mu^{-1}) \quad (\mu \rightarrow +\infty).$$

Proof. Since $d_{z+\tau} = d_z + \tau \eta \wedge$, it follows from Theorem 3.10 that $d_{z+\tau}$ is bounded on $E_{z,\text{sm}} + E_{z+\tau,\text{sm}}$, uniformly on $\mu \gg 0$. Hence, by Proposition 3.18,

$$d_{z+\tau,\text{sm}} - d_{z+\tau} P_{z,\text{sm}} = d_{z+\tau} (P_{z+\tau,\text{sm}} - P_{z,\text{sm}}) = O(\mu^{-1}). \quad \square$$

PROPOSITION 3.20. *On $L^2(M; \Lambda)$,*

$$P_{z,\text{sm}} \eta \wedge, \eta \wedge P_{z,\text{sm}} = O(\mu^{-1/2}) \quad (\mu \rightarrow +\infty).$$

Proof. Let $\alpha \in L^2(M; \Lambda^k)$ and $p \in \mathcal{X}_{k+1}$. Using multi-index notation, write $\alpha = \sum_{|J|=k} \alpha_J dx_p^J$ on U_p with $\alpha_J \in L^2(U_p)$. Let $J_j = \{1, \dots, k+1\} \setminus \{j\}$ ($j = 1, \dots, k+1$). By Proposition 3.1 (iii), (3.10) and (3.11),

$$\begin{aligned} \langle \eta \wedge \alpha, e_{p,z} \rangle &= -\frac{1}{a_\mu} \sum_{j=1}^{k+1} \langle \alpha_{J_j} x_p^j dx_p^{J_j} \wedge dx_p^j, e^{-i\nu h_p} \rho_p e^{-\mu|x_p|^2/2} dx_p^1 \wedge \dots \wedge dx_p^{k+1} \rangle \\ &= (-1)^j \frac{1}{a_\mu} \sum_{j=1}^{k+1} \langle \alpha_{J_j} x_p^j, e^{-i\nu h_p} \rho_p e^{-\mu|x_p|^2/2} \rangle. \end{aligned}$$

Hence

$$|\langle \eta \wedge \alpha, e_{p,z} \rangle| \leq \frac{1}{a_\mu} \sum_{j=1}^{k+1} |\langle \alpha_{J_j}, e^{-i\nu h_p} \rho_p x_p^j e^{-\mu|x_p|^2/2} \rangle| \leq \|\alpha\| \frac{1}{a_\mu} \sum_{j=1}^{k+1} \|\rho_p x_p^j e^{-\mu|x_p|^2/2}\|, \quad (3.28)$$

and

$$\begin{aligned}
 \|\rho_p x_p^j e^{-\mu|x_p|^2/2}\| &= \left(\int_{U_p} (\rho_p(x_p) x_p^j)^2 e^{-\mu|x_p|^2} \text{vol}(x_p) \right)^{1/2} \\
 &= \left(\int_{-2r}^{2r} \rho(x)^2 x^2 e^{-\mu x^2} dx \right)^{\frac{1}{2}} \left(\int_{-2r}^{2r} \rho(y)^2 e^{-\mu y^2} dy \right)^{\frac{n-1}{2}} \\
 &= \left(\int_{-2r}^{2r} x^2 e^{-\mu x^2} dx \right)^{\frac{1}{2}} \left(\int_{-2r}^{2r} e^{-\mu y^2} dy \right)^{\frac{n-1}{2}} + O(e^{-c\mu}) \\
 &= \frac{1}{(2\mu)^{1/2}} \left(\frac{\pi}{\mu} \right)^{n/4} + O(e^{-c\mu}) .
 \end{aligned} \tag{3.29}$$

By (3.12), (3.28) and (3.29),

$$|\langle \eta \wedge \alpha, e_{p,z} \rangle| = \|\alpha\| O(\mu^{-1/2}) .$$

Since

$$P_z(\eta \wedge \alpha) = \sum_{p \in \mathcal{X}_{k+1}} \langle \eta \wedge \alpha, e_{p,z} \rangle e_{p,z} ,$$

it follows that

$$\|P_z(\eta \wedge \alpha)\| = \|\alpha\| O(\mu^{-1/2}) .$$

This shows that $P_z \eta \wedge \alpha = O(\mu^{-1/2})$, yielding $P_{z,\text{sm}} \eta \wedge \alpha = O(\mu^{-1/2})$ by Proposition 3.18.

On the other hand, for $q \in \mathcal{X}_k$,

$$\eta \wedge e_{q,z} = \frac{\rho_q}{a_\mu} e^{-\mu|x_q|^2/2} \sum_{j=k+1}^n x_q^j dx_q^j \wedge dx_q^1 \wedge \cdots \wedge dx_q^k .$$

So

$$\|\eta \wedge e_{q,z}\| = \frac{1}{a_\mu} \left(\sum_{j=k+1}^n \int_{U_q} (\rho_q(x_q) x_q^j)^2 e^{-\mu|x_q|^2} \text{vol}(x_q) \right)^{1/2} ,$$

which becomes $O(\mu^{-1/2})$ like in the previous part of the proof. Since

$$\eta \wedge P_z \alpha = \sum_{q \in \mathcal{X}_k} \langle \alpha, e_{q,z} \rangle \eta \wedge e_{q,z} ,$$

we get $\eta \wedge P_z \alpha = O(\mu^{-1/2})$, yielding $\eta \wedge P_{z,\text{sm}} \alpha = O(\mu^{-1/2})$ as before. \square

3.5 Derivatives of the small projection

Remark 3.21. For reasons of brevity, most of the results about derivatives are stated for ∂_z , which may be simply denoted with a dot. But there are obvious versions of those results for $\partial_{\bar{z}}$ with analogous proofs.

PROPOSITION 3.22. *We have*

$$\text{rank } \partial_z P_{z,\text{sm}} \leq 2|\mathcal{X}| \quad (\mu \gg 0) , \quad \partial_z P_{z,\text{sm}} = O(\mu^{-1/2}) \quad (\mu \rightarrow +\infty) .$$

Proof. By (2.3) and Theorem 3.10, for $\mu \gg 0$ and every $\omega \in \mathbb{S}^1$, a standard computation gives

$$\partial_z((w - D_z)^{-1}) = (w - D_z)^{-1} \eta \wedge (w - D_z)^{-1} . \tag{3.30}$$

Then, by (3.15), $\partial_z((w - D_z)^{-1})$ defines an operator on $L^2(M; \Lambda)$, bounded uniformly on $w \in \mathbb{S}^1$ and $z \in \mathbb{C}$. By (3.15) and Proposition 3.20, we also get

$$P_{z, \text{la}/\text{sm}} \partial_z((w - D_z)^{-1}) P_{z, \text{sm}/\text{la}} = (w - D_z)^{-1} P_{z, \text{la}/\text{sm}} \eta \wedge P_{z, \text{sm}/\text{la}} (w - D_z)^{-1} = O(\mu^{-1/2}),$$

uniformly on $w \in \mathbb{S}^1$.

On the other hand, applying again basic spectral theory, we obtain

$$P_{z, \text{sm}} = \frac{1}{2\pi i} \int_{\mathbb{S}^1} (w - D_z)^{-1} dw$$

for $\mu \gg 0$, yielding

$$\dot{P}_{z, \text{sm}} = \frac{1}{2\pi i} \int_{\mathbb{S}^1} \partial_z((w - D_z)^{-1}) dw, \quad (3.31)$$

which defines an operator on $L^2(M; \Lambda)$, bounded uniformly on z .

Using that $P_{z, \text{sm}}$ is an orthogonal projection, the argument of the proof of [BGV04, Proposition 9.37] shows that

$$\dot{P}_{z, \text{sm}} = P_{z, \text{la}} \dot{P}_{z, \text{sm}} P_{z, \text{sm}} + P_{z, \text{sm}} \dot{P}_{z, \text{sm}} P_{z, \text{la}}. \quad (3.32)$$

So $\text{rank } \dot{P}_{z, \text{sm}} \leq 2 \text{rank } P_{z, \text{sm}} \leq 2|\mathcal{X}|$ by Corollary 3.9, and

$$\begin{aligned} \dot{P}_{z, \text{sm}} &= \frac{1}{2\pi i} \int_{\mathbb{S}^1} P_{z, \text{la}} \partial_z((w - D_z)^{-1}) P_{z, \text{sm}} dw \\ &\quad + \frac{1}{2\pi i} \int_{\mathbb{S}^1} P_{z, \text{sm}} \partial_z((w - D_z)^{-1}) P_{z, \text{la}} dw = O(\mu^{-1/2}). \quad \square \end{aligned}$$

LEMMA 3.23. For all $p \in \mathcal{X}$,

$$\partial_z e_{p, z} = \left(\frac{n}{8\mu} - \frac{|x_p^+|^2}{2} + O(e^{-c\mu}) \right) e_{p, z} \quad (\mu \rightarrow +\infty).$$

Proof. Using integration by parts, and since ρ is an even function and ρ' vanishes on $[-r, r]$, we obtain

$$\begin{aligned} \int_{-2r}^{2r} \rho(x)^2 x^2 e^{-\mu x^2} dx &= \frac{1}{2\mu} \int_{-2r}^{2r} (2\rho(x)\rho'(x)x + \rho(x)^2) e^{-\mu x^2} dx \\ &= \frac{1}{2\mu} \left(\frac{\pi}{\mu} \right)^{\frac{1}{2}} + O(e^{-c\mu}). \end{aligned} \quad (3.33)$$

So

$$\begin{aligned} \partial_\mu a_\mu &= \partial_\mu \left(\left(\int_{-2r}^{2r} \rho(x)^2 e^{-\mu x^2} dx \right)^{\frac{n}{2}} \right) \\ &= -\frac{n}{2} \left(\int_{-2r}^{2r} \rho(x)^2 e^{-\mu x^2} dx \right)^{\frac{n}{2}-1} \int_{-2r}^{2r} \rho(x)^2 x^2 e^{-\mu x^2} dx \\ &= -\frac{n}{2} \left(\frac{\pi}{\mu} \right)^{\frac{n}{4}-\frac{1}{2}} \frac{1}{2\mu} \left(\frac{\pi}{\mu} \right)^{\frac{1}{2}} + O(e^{-c\mu}) = -\frac{n}{4\mu} \left(\frac{\pi}{\mu} \right)^{\frac{n}{4}} + O(e^{-c\mu}). \end{aligned}$$

Hence, by (3.12),

$$\partial_\mu \left(\frac{1}{a_\mu} \right) = -\frac{\partial_\mu a_\mu}{a_\mu^2} = \frac{n}{4\mu} \left(\frac{\pi}{\mu} \right)^{\frac{n}{4}} \left(\frac{\mu}{\pi} \right)^{\frac{n}{2}} + O(e^{-c\mu}) = \frac{n}{4\mu} \left(\frac{\mu}{\pi} \right)^{\frac{n}{4}} + O(e^{-c\mu}). \quad (3.34)$$

It also follows from Proposition 3.1 (iii), (3.10), (3.12) and (3.34) that

$$\begin{aligned}\partial_\mu e_{p,\mu} &= \partial_\mu \left(\frac{\rho_p}{a_\mu} e^{-\mu|x_p|^2/2} dx_p^1 \wedge \cdots \wedge dx_p^{\text{ind}(p)} \right) = \left(\partial_\mu \left(\frac{1}{a_\mu} \right) a_\mu - \frac{|x_p|^2}{2} \right) e_{p,\mu} \\ &= \left(\frac{n}{4\mu} - \frac{|x_p|^2}{2} + O(e^{-c\mu}) \right) e_{p,\mu}.\end{aligned}\quad (3.35)$$

So, by (3.11),

$$\partial_\mu e_{p,z} = \left(\frac{n}{4\mu} - \frac{|x_p|^2}{2} + O(e^{-c\mu}) \right) e_{p,z}, \quad \partial_\nu e_{p,z} = -i h_p e_{p,z}.\quad (3.36)$$

Then the result follows using (3.5). \square

PROPOSITION 3.24. *For all $p \in \mathcal{X}$,*

$$\|\partial_z(D_z e_{p,z})\|_{L^\infty} = O(e^{-c\mu}) \quad (\mu \rightarrow +\infty).$$

Proof. From (3.13), we get

$$\begin{aligned}\partial_z(D_z e_{p,z}) &= \frac{1}{2} \left(e^{-i\nu h_p} \partial_\mu \left(\frac{1}{a_\mu} \left(\frac{\pi}{\mu} \right)^{n/4} \right) \hat{c}(d\rho_p) e_{p,\mu} + e^{-i\nu h_p} \frac{1}{a_\mu} \left(\frac{\pi}{\mu} \right)^{n/4} \hat{c}(d\rho_p) \partial_\mu e_{p,\mu} \right. \\ &\quad \left. - h_p e^{-i\nu h_p} \frac{1}{a_\mu} \left(\frac{\pi}{\mu} \right)^{n/4} \hat{c}(d\rho_p) e_{p,\mu} \right).\end{aligned}\quad (3.37)$$

By (3.12) and (3.34),

$$\begin{aligned}\partial_\mu \left(\frac{1}{a_\mu} \left(\frac{\pi}{\mu} \right)^{n/4} \right) &= \partial_\mu \left(\frac{1}{a_\mu} \right) \left(\frac{\pi}{\mu} \right)^{n/4} - \frac{n\pi}{4a_\mu \mu^2} \left(\frac{\pi}{\mu} \right)^{n/4-1} \\ &= \frac{n}{4\mu} \left(\frac{\mu}{\pi} \right)^{n/4} \left(\frac{\pi}{\mu} \right)^{n/4} - \frac{n\pi}{4\mu^2} \left(\frac{\pi}{\mu} \right)^{n/4-1} \left(\frac{\mu}{\pi} \right)^{n/4} + O(e^{-c\mu}) = O(e^{-c\mu}).\end{aligned}\quad (3.38)$$

The result follows applying Proposition 3.1 (iii), (3.10), (3.12), (3.35) and (3.38) to (3.37), and using that $d\rho_p = 0$ around p . \square

PROPOSITION 3.25. *For every $p \in \mathcal{X}$,*

$$\|\partial_z(P_{z,\text{sm}} e_{p,z} - e_{p,z})\|_{L^\infty} = O(e^{-c\mu}) \quad (\mu \rightarrow +\infty).$$

Proof. By (3.18),

$$\begin{aligned}\partial_z(P_{z,\text{sm}} e_{p,z} - e_{p,z}) &= \frac{1}{2\pi i} \int_{\mathbb{S}^1} w^{-1} \partial_z((w - D_z)^{-1}) D_z e_{p,z} dw \\ &\quad + \frac{1}{2\pi i} \int_{\mathbb{S}^1} w^{-1} (w - D_z)^{-1} \partial_z(D_z e_{p,z}) dw.\end{aligned}$$

Now apply (3.19), (3.30), Corollary 3.6 and Proposition 3.24. \square

4. Zeta invariants of Morse forms

4.1 Preliminaries on asymptotic expansions of heat kernels

Let A be a positive semi-definite symmetric elliptic differential operator of order a , and B a differential operator of order b ; both of them are defined in $C^\infty(M; E)$ for some Hermitian vector bundle E over M . Then $B e^{-tA}$ is a smoothing operator with Schwartz kernel $K_t(x, y)$ in

$C^\infty(M^2; E \boxtimes E^*)$ (omitting the Riemannian density $\text{vol}(y)$ of the second factor). On the diagonal, there is an asymptotic expansion (as $t \downarrow 0$) [Gil95, Lemma 1.9.1], [BGV04, Theorem 2.30],

$$K_t(x, x) \sim \sum_{l=0}^{\infty} e_l(x) t^{(l-n-b)/a}, \quad (4.1)$$

where $e_l \in C^\infty(M; E \otimes E^*)$ is locally computable in terms of the jets of the local coefficients of A and B . Hence the function

$$h(t) = \text{Tr}(Be^{-tA}) = \int_M \text{tr} K_t(x, x) \text{vol}(x)$$

has an asymptotic expansion

$$h(t) \sim \sum_{l=0}^{\infty} a_l t^{(l-n-b)/a}, \quad (4.2)$$

where

$$a_l = \int_M \text{tr} e_l(x) \text{vol}(x). \quad (4.3)$$

Given any $\lambda \geq 0$, let $P_{A,\lambda}$ be the spectral projection of A corresponding to $[0, \lambda]$; thus $P_{A,\lambda}^\perp$ is the spectral projection corresponding to (λ, ∞) . By ellipticity, $P_{A,\lambda}$ is of finite rank, and $Be^{-tA}P_{A,\lambda}$ is a smoothing operator defined for all $t \in \mathbb{R}$. Take any orthonormal frame $\phi_1, \dots, \phi_\kappa$ of $\text{im} P_{A,\lambda}$, consisting of eigensections with corresponding eigenvalues $0 \leq \lambda_1 \leq \dots \leq \lambda_\kappa \leq \lambda$. Then the Schwartz kernel $H_t(x, y)$ of $Be^{-tA}P_{A,\lambda}$ ($t \geq 0$) is given by

$$H_t(x, y) = \sum_{j=1}^{\kappa} e^{-t\lambda_j} (B\phi_j)(x) \otimes \phi_j(y),$$

using the identity $E \equiv E^*$ given by the Hermitian structure. Thus $H_t(x, y)$ is defined for all $t \in \mathbb{R}$ and smooth. So

$$\text{Tr}(Be^{-tA}P_{A,\lambda}) = \int_M \text{tr} H_t(x, x) \text{vol}(x).$$

In particular, for $t = 0$, we have

$$H_0(x, x) = \sum_{j=1}^{\kappa} (B\phi_j)(x) \otimes \phi_j(x), \quad (4.4)$$

$$\text{Tr}(BP_{A,\lambda}) = \int_M \text{tr} H_0(x, x) \text{vol}(x). \quad (4.5)$$

The Schwartz kernel of $Be^{-tA}P_{A,\lambda}^\perp$ is $\tilde{K}_t(x, y) = K_t(x, y) - H_t(x, y)$ ($t > 0$), which has an asymptotic expansion

$$\tilde{K}_t(x, x) \sim \sum_{l=0}^{\infty} \tilde{e}_l(x) t^{(l-n-b)/a}, \quad (4.6)$$

where the first $n + b$ sections \tilde{e}_l are given by

$$\tilde{e}_l(x) = \begin{cases} e_l(x) & \text{if } l < n + b \\ e_l(x) - H_0(x, x) & \text{if } l = n + b. \end{cases}$$

Then the function

$$\tilde{h}_\lambda(t) = \text{Tr}(Be^{-tA}P_{A,\lambda}^\perp) = \text{Tr}(Be^{-tA}) - \text{Tr}(Be^{-tA}P_{A,\lambda}) \quad (4.7)$$

has an asymptotic expansion

$$\tilde{h}_\lambda(t) = \int_M \tilde{K}_t(x, x) \operatorname{vol}(x) \sim \sum_{l=0}^{\infty} \tilde{a}_{\lambda,l} t^{(l-n-b)/a}, \quad (4.8)$$

where the first $n + b$ coefficients \tilde{a}_l are given by

$$\tilde{a}_{\lambda,l} = \begin{cases} a_l & \text{if } l < n + b \\ a_l - \operatorname{Tr}(BP_{A,\lambda}) & \text{if } l = n + b. \end{cases} \quad (4.9)$$

Consider also smooth families of such operators, $\{A_\epsilon\}$ and $\{B_\epsilon\}$, for ϵ in some parameter space. Then $\operatorname{Tr}(B_\epsilon e^{-tA_\epsilon})$ is smooth in (t, ϵ) , and we add ϵ to the above notation, writing for instance $K_t(x, y, \epsilon)$, $e_l(x, \epsilon)$, $h(t, \epsilon)$, $a_l(\epsilon)$, $\tilde{K}_t(x, y, \epsilon)$, $\tilde{e}_{\lambda,l}(x, \epsilon)$, $\tilde{h}_\lambda(t, \epsilon)$ and $\tilde{a}_{\lambda,l}(\epsilon)$ in (4.1), (4.2), (4.6) and (4.8). The operator $B_\epsilon P_{A_\epsilon, \lambda}$ may not be smooth in ϵ when some non-constant spectral branch of $\{A_\epsilon\}$ reaches the value λ . If the values of the non-constant spectral branches of $\{A_\epsilon\}$ stay uniformly away from λ , then $\tilde{h}_\lambda(t, \epsilon)$ is smooth in (t, ϵ) .

4.2 Preliminaries on zeta functions of operators

PROPOSITION 4.1 See e.g. [Gil95, Theorems 1.12.2 and 1.12.5], [BGV04, Propositions 9.35–9.37]. *The following holds:*

- (i) For every $\lambda \geq 0$, there is a meromorphic function $\zeta(s, A, B, \lambda)$ on \mathbb{C} such that, for $\Re s \gg 0$,

$$\zeta(s, A, B, \lambda) = \operatorname{Tr}(BA^{-s}P_{A,\lambda}^\perp) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \tilde{h}_\lambda(t) dt. \quad (4.10)$$

- (ii) The meromorphic function $\Gamma(s)\zeta(s, A, B, \lambda)$ has simple poles at the points $s = (n + b - l)/a$, for $l \in \mathbb{N}_0$ with $\tilde{a}_l \neq 0$. The corresponding residues are \tilde{a}_l , and $\zeta(s, A, B, \lambda)$ is smooth away from these exceptional values of s .
- (iii) For $\mu > \lambda \geq 0$, let $\lambda_1 \leq \dots \leq \lambda_k$ denote the eigenvalues of A in $(\lambda, \mu]$, taking multiplicities into account, and let ψ_1, \dots, ψ_k be corresponding orthonormal eigensections. Then, for all s ,

$$\zeta(s, A, B, \mu) - \zeta(s, A, B, \lambda) = \sum_{j=1}^k \lambda_k^{-s} \langle B\psi_j, \psi_j \rangle.$$

- (iv) For smooth families $\{A_\epsilon\}$ and $\{B_\epsilon\}$ of such operators, if the non-constant branches of eigenvalues of $\{A_\epsilon\}$ stay uniformly away from the value λ , then $\zeta(s, A_\epsilon, B_\epsilon, \lambda)$ is smooth in (s, ϵ) away from the exceptional values of s given in (ii).
- (v) Consider the conditions of (iv) for ϵ in some open neighborhood of 0 in \mathbb{R} . If A_0 and B_0 commute, then

$$\partial_\epsilon \zeta(s, A_\epsilon, B_\epsilon, \lambda) \Big|_{\epsilon=0} = \zeta(s, A_0, \dot{B}_0, \lambda) - s\zeta(s+1, A_0, \dot{A}_0 B_0, \lambda),$$

where the dot denotes ∂_ϵ .

The last expression of (4.10) is the Mellin transform of the function $\tilde{h}_\lambda(t)$ divided by $\Gamma(s)$. This function $\zeta(s, A, B, \lambda)$ is called the *zeta function* of (A, B, λ) . If $B = 1$ or $\lambda = 0$, they may be omitted from the notation.

We will also use $\zeta(s, A, B, \lambda)$ when B is not a differential operator, with the same definition. Then the asymptotic expansion (4.8) and the properties stated in Proposition 4.1 need to be

checked. With this generality, we can write

$$\zeta(s, A, B, \lambda) = \zeta(s, A, BP_{A,\lambda}^\perp) = \zeta(s, A, P_{A,\lambda}^\perp B) ,$$

and we have

$$\zeta(s, A, B) = \zeta(s, A, BP_{A,\lambda}) + \zeta(s, A, B, \lambda) .$$

The function $\zeta(s, A, BP_{A,\lambda})$ is always defined and holomorphic on \mathbb{C} because $P_{A,\lambda}$ is of finite rank.

In particular, when A is Δ_z or $\Delta_{z,k}$ ($k = 0, \dots, n$) and B is another operator in $L^2(M; \Lambda)$, we will use $\zeta(s, A, B, \lambda)$ if it is defined. We get

$$\zeta(s, \Delta_z, B) = \zeta_{\text{sm}}(s, \Delta_z, B) + \zeta_{\text{la}}(s, \Delta_z, B) ,$$

where, with the notation of Section 3.2,

$$\zeta_{\text{sm/la}}(s, \Delta_z, B) = \zeta(s, \Delta_z, B_{z,\text{sm/la}}) .$$

These are the contributions from the small/large spectrum to $\zeta(s, \Delta_z, B)$, which are called the *small/large zeta functions* of (Δ_z, B) .

4.3 Zeta invariants of Morse forms

By Proposition 4.1 (i), as a function of s , $\zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is meromorphic on \mathbb{C} . Moreover, for $\Re s \gg 0$,

$$\begin{aligned} \zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w}) &= \text{Tr}^s (\eta \wedge D_z \Delta_z^{-s} \Pi_z^\perp) = \text{Tr}^s (\eta \wedge \delta_z \Delta_z^{-s} \Pi_z^\perp) \\ &= \text{Tr}^s (\eta \wedge D_z^{-1} \Delta_z^{-s+1} \Pi_z^\perp) = \text{Tr}^s (\eta \wedge d_z^{-1} \Delta_z^{-s+1} \Pi_z^\perp) , \end{aligned}$$

using that $\eta \wedge d_z$ and $\eta \wedge \delta_z^{-1}$ change the degree of homogeneous forms. So, when $\zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is regular at $s = 1$, the value $\zeta(1, \Delta_z, \eta \wedge D_z \mathbf{w})$ is a renormalized version of the super-trace of $\eta \wedge d_z^{-1} \Pi_z^\perp$, which is called the *zeta invariant* of (M, g, η) for the scope of this paper. Unfortunately, according to Proposition 4.1 (ii) and since $\Gamma(s)$ is regular at $s = 1$, $\zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ might have a simple pole at $s = 1$. It will be shown that we can choose η in the given class $\xi \in H^1(M, \mathbb{R})$ such that $\zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is regular at $s = 1$ for $\mu \gg 0$. To achieve this task, we consider its decomposition into small/large zeta functions (Section 4.2),

$$\zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w}) = \zeta_{\text{sm}}(s, \Delta_z, \eta \wedge D_z \mathbf{w}) + \zeta_{\text{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w}) .$$

The values $\zeta_{\text{sm/la}}(1, \Delta_z, \eta \wedge D_z \mathbf{w})$ will be called the *small/large zeta invariant* of (M, g, η) , if they are defined— $\zeta_{\text{sm}}(1, \Delta_z, \eta \wedge D_z \mathbf{w})$ is always defined.

4.4 Heat invariants of perturbed operators

For $k = 0, \dots, n$, the Schwartz kernels of $e^{-t\Delta_{z,k}}$ and $e^{-t\Delta_{z,k}} P_{z,\text{la},k}$ are denoted by $K_{z,k,t}(x, y)$ and $\tilde{K}_{z,k,t}(x, y)$, respectively. Their restrictions to the diagonal have asymptotic expansions (as $t \downarrow 0$),

$$\begin{aligned} K_{z,k,t}(x, x) &\sim \sum_{l=0}^{\infty} e_{k,l}(x, z) t^{(l-n)/2} , \\ \tilde{K}_{z,k,t}(x, x) &\sim \sum_{l=0}^{\infty} \tilde{e}_{k,l}(x, z) t^{(l-n)/2} , \end{aligned} \tag{4.11}$$

where $e_{k,l}(x, z)$ is locally given by smooth expressions involving z and the jets of the local coefficients of g and η . Take $\mu \gg 0$ so that $P_{z,la,k}$ depends smoothly on z (Theorem 3.10) and $\dim E_{z,sm}^k = |\mathcal{X}_k|$ (Corollary 3.9). Thus $\tilde{e}_{k,l}(x, z)$ also depends smoothly on z (Section 4.1). Then

$$\tilde{e}_{k,l}(x, z) = \begin{cases} e_{k,l}(x, z) & \text{if } l < n \\ e_{k,n}(x, z) - H_{z,k,0}(x, x) & \text{if } l = n, \end{cases} \quad (4.12)$$

where $H_{z,k,t}(x, y)$ is the Schwartz kernel of $e^{-t\Delta_{z,k}}P_{z,sm,k}$, which is defined for all $t \in \mathbb{R}$ and smooth. According to Section 4.2, the corresponding functions

$$h_k(t, z) = \text{Tr} \left(e^{-t\Delta_{z,k}} \right), \quad \tilde{h}_k(t, z) = \text{Tr} \left(e^{-t\Delta_{z,k}} P_{z,la,k} \right),$$

have asymptotic expansions

$$h_k(t, z) \sim \sum_{l=0}^{\infty} a_{k,l}(z) t^{(l-n)/2}, \quad \tilde{h}_k(t, z) \sim \sum_{l=0}^{\infty} \tilde{a}_{k,l}(z) t^{(l-n)/2}, \quad (4.13)$$

where, by (4.3)–(4.5) and (4.9),

$$a_{k,l}(z) = \int_M \text{tr}^s e_{k,l}(x, z) \text{vol}(x), \quad (4.14)$$

$$\tilde{a}_{k,l}(z) = \int_M \text{tr}^s \tilde{e}_{k,l}(x, z) \text{vol}(x) = \begin{cases} a_{k,l}(z) & \text{if } l < n \\ a_{k,l}(z) - |\mathcal{X}_k| & \text{if } l = n. \end{cases} \quad (4.15)$$

Consider the operators $e^{-t\Delta_z} \mathbf{w}$ and $e^{-t\Delta_z} P_{z,la} \mathbf{w}$, whose respective Schwartz kernels are

$$K_{z,t}(x, y) = \sum_{k=0}^n (-1)^k K_{z,k,t}(x, y), \quad \tilde{K}_{z,t}(x, y) = \sum_{k=0}^n (-1)^k \tilde{K}_{z,k,t}(x, y).$$

We have induced asymptotic expansions

$$K_{z,t}(x, x) \sim \sum_{l=0}^{\infty} e_l(x, z) t^{(l-n)/2}, \quad \tilde{K}_{z,t}(x, x) \sim \sum_{l=0}^{\infty} \tilde{e}_l(x, z) t^{(l-n)/2},$$

where

$$e_l(x, z) = \sum_{k=0}^n (-1)^k e_{k,l}(x, z), \quad \tilde{e}_l(x, z) = \sum_{k=0}^n (-1)^k \tilde{e}_{k,l}(x, z).$$

The corresponding functions,

$$h(t, z) = \text{Tr}^s \left(e^{-t\Delta_z} \right) = \sum_{k=0}^n (-1)^k h_k(t, z),$$

$$\tilde{h}(t, z) = \text{Tr}^s \left(e^{-t\Delta_z} P_{z,la} \right) = \sum_{k=0}^n (-1)^k \tilde{h}_k(t, z),$$

have asymptotic expansions

$$h(t, z) \sim \sum_{l=0}^{\infty} a_l(z) t^{(l-n)/2}, \quad \tilde{h}(t, z) \sim \sum_{l=0}^{\infty} \tilde{a}_l(z) t^{(l-n)/2},$$

where

$$a_l(z) = \sum_{k=0}^n (-1)^k a_{k,l}(z), \quad \tilde{a}_l(z) = \sum_{k=0}^n (-1)^k \tilde{a}_{k,l}(z).$$

THEOREM 4.2 [ALG21, Theorem 1.5], [ALKL20]. *We have:*

- (i) $e_l(x, z) = 0$ for $l < n$; and,
- (ii) if n is even, then $e_n(x, z) = e(M, \nabla^M)(x)$.

Remark 4.3. Actually, [ALG21, Theorem 1.5] gives Theorem 4.2 when z is real. But, since the functions $e_l(x, z)$ have local expressions, we can assume η is exact. Then the result can be extended to non-real z using (2.11).

4.5 Derived heat invariants of perturbed operators

For $k = 0, \dots, n$ and $j = 1, 2$, let

$$h_k^j(t, z) = \text{Tr} \left(e^{-t\Delta_{z,k}} \Pi_{z,k}^j \right), \quad \tilde{h}_k^j(t, z) = \text{Tr} \left(e^{-t\Delta_{z,k}} \Pi_{z,1a,k}^j \right).$$

LEMMA 4.4. *We have*

$$h_{k+1}^1(t, z) = h_k^2(t, z) = \sum_{p=0}^k (-1)^{k-p} h_p(t, z) = \sum_{q=k+1}^n (-1)^{q-k-1} h_q(t, z).$$

Proof. This follows by induction on k , using that

$$h_0^1(t, z) = h_n^2(t, z) = 0, \quad h_k(t, z) = h_k^1(t, z) + h_k^2(t, z), \quad h_k^2(t, z) = h_{k+1}^1(t, z).$$

The last equality holds because (2.7) is commutative. □

Let

$$\begin{aligned} h^j(t, z) &= \sum_{k=0}^n (-1)^k h_k^j(t, z) = \text{Tr}^s \left(e^{-t\Delta_z} \Pi_z^j \right), \\ \tilde{h}^j(t, z) &= \sum_{k=0}^n (-1)^k \tilde{h}_k^j(t, z) = \text{Tr}^s \left(e^{-t\Delta_z} \Pi_{z,1a}^j \right). \end{aligned}$$

Thus

$$h(t, z) = h^1(t, z) + h^2(t, z), \quad \tilde{h}(t, z) = \tilde{h}^1(t, z) + \tilde{h}^2(t, z). \quad (4.16)$$

COROLLARY 4.5. *We have* $h(t, z) = 0$.

Proof. This is a direct consequence of Lemma 4.4 and (4.16). □

COROLLARY 4.6. *We have*

$$-(-1)^j h^j(t, z) = \sum_{k=0}^n (-1)^k k h_k(t, z) = \text{Tr}^s \left(\mathbf{N} e^{-t\Delta_z} \Pi_z^\perp \right).$$

Proof. By Lemma 4.4 and Corollary 4.5,

$$\begin{aligned} h^1(t, z) &= \sum_{k=0}^n (-1)^k \sum_{q=k}^n (-1)^{q-k} h_q(t, z) = \sum_{q=0}^n (-1)^q (q+1) h_q(t, z) \\ &= h(t, z) + \sum_{q=0}^n (-1)^q q h_q(t, z) = \sum_{q=0}^n (-1)^q q h_q(t, z). \quad \square \end{aligned}$$

Applying (4.13) and Lemma 4.4, we get

$$h_k^j(t, z) \sim \sum_{l=0}^{\infty} a_{k,l}^j(z) t^{(l-n)/2}, \quad h^j(t, z) \sim \sum_{l=0}^{\infty} a_l^j(z) t^{(l-n)/2}, \quad (4.17)$$

where

$$\begin{aligned} a_{k+1,l}^1(z) &= a_{k,l}^2(z) = \sum_{p=0}^k (-1)^{k-p} a_{p,l}(t, z) = \sum_{q=k+1}^n (-1)^{q-k-1} a_{q,l}(t, z), \\ &- (-1)^j a_l^j(z) = \sum_{k=0}^n (-1)^k k a_{k,l}(z). \end{aligned}$$

This $a_l^1(z)$ is sometimes called the *derived heat invariant* of Δ_z [GS77], [RS71], [Gil95, page 181], [ALG20].

For $\mu \gg 0$, Lemma 4.4, Corollary 4.5 and (4.17) have obvious versions for $\tilde{h}_k^j(t, z)$ and $\tilde{h}^j(t, z)$, with the similar proofs. The coefficients of the corresponding asymptotic expansions are denoted by $\tilde{a}_{k,l}^j(z)$ and $\tilde{a}_l^j(z)$.

THEOREM 4.7 [ALG20, Theorem 1.3 (1b)]. *For all $l \leq n$, $a_l^1(z)$ and $\tilde{a}_l^1(z)$ are independent of z .*

Remark 4.8. Again, [ALG20, Theorem 1.3 (1b)] gives Theorem 4.7 in case of $a_l^1(z)$ for real z , but this can be extended for non-real z like in Remark 4.3. Then the case of $\tilde{a}_l^1(z)$ follows from (4.15). In Theorem 4.7, the case of $a_l^1(z)$ is true for any $\eta \in Z^1(M, \mathbb{R})$.

Remark 4.9. Theorem 4.2 and the case of $a_l^1(z)$ in Theorem 4.7 hold for any $\eta \in Z^1(M, \mathbb{R})$ (there is no need of other conditions). This is also true for all properties of Sections 4.4 and 4.5 concerning $K_{z,k,t}(x, y)$, $e_{k,l}(x, z)$, $h_k(t, z)$, $h(t, z)$, $h^j(t, z)$, $h_k^j(t, z)$, $a_{k,l}(z)$, $a_l(z)$, $a_{k,l}^j(z)$, $\zeta(s, \Delta_z, \Pi_z^j \mathbf{w})$ and $\theta(s, z)$.

4.6 Regularity

By (4.8), we have an asymptotic expansion of the form

$$\mathrm{Tr}^s (\eta \wedge D_z e^{-t\Delta_z} P_{z, \mathrm{la}}) \sim \sum_{l=0}^{\infty} \tilde{b}_l(z) t^{(l-n-1)/2}. \quad (4.18)$$

LEMMA 4.10. *The function $\zeta_{\mathrm{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ has a simple pole at every $s = (n+1-l)/2$ with $\tilde{b}_l(z) \neq 0$, for $1 \leq l \leq n$ and for even/odd $l \geq n+2$ if n is even/odd, whose residue is $\tilde{b}_l(z)$, and it is smooth away from these values of s . Moreover the value of $\zeta_{\mathrm{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ at every regular point $s = (n+1-l)/2$, for odd $l \geq n+1$, is $(l-n-1)! \tilde{b}_l(z)$.*

Proof. By Proposition 4.1 (ii), the product $\Gamma(s) \zeta_{\mathrm{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ has a simple pole at every $s = (n+1-l)/2$ with $\tilde{b}_l(z) \neq 0$ ($l \in \mathbb{N}_0$), whose residue is $\tilde{b}_l(z)$, and $\zeta_{\mathrm{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is smooth away from these exceptional values of s . Then the result follows because $\Gamma(s)$ has a simple pole at every point $s = -k$ ($k \in \mathbb{N}_0$), whose residue is $(-1)^k/k!$, and it is smooth on $\mathbb{C} \setminus (-\mathbb{N}_0)$. \square

By Corollary 4.6 and using the notation (1.4) and (1.5), we get

$$\begin{aligned} (-1)^j \zeta(s, \Delta_z, \Pi_z^j \mathbf{w}) &= \theta(s, z), \\ (-1)^j \zeta_{\mathrm{sm}/\mathrm{la}}(s, \Delta_z, \Pi_z^j \mathbf{w}) &= \theta_{\mathrm{sm}/\mathrm{la}}(s, z). \end{aligned} \quad (4.19)$$

Thus the following result follows like Lemma 4.10.

LEMMA 4.11. *The function $\zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w})$ has a simple pole at every $s = (n-l)/2$ with $\tilde{a}_l^1(z) \neq 0$, for $1 \leq l \leq n-1$ and for odd/even $l \geq n+1$ if n is even/odd, whose residue is $\tilde{a}_l^1(z)$, and it is smooth away from these values of s . Moreover the value of $\zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w})$ at every regular point $s = (n-l)/2$, for even $l \geq n$, is $(l-n)! \tilde{a}_l^1(z)$.*

According to Lemmas 4.10 and 4.11, $\zeta_{\text{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ and $\zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w})$ are smooth at $s = 0$ [See67], but they might have a simple pole at $s = 1$.

PROPOSITION 4.12. *We have*

$$\partial_z \zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w}) = -s \zeta_{\text{la}}(s+1, \Delta_z, \eta \wedge D_z \mathbf{w}) .$$

Proof. Recall that a dot may be used to denote ∂_z . Like in (3.32),

$$\dot{\Pi}_{z,\text{la}}^1 = (\Pi_{z,\text{la}}^1)^\perp \dot{\Pi}_{z,\text{la}}^1 \Pi_{z,\text{la}}^1 + \Pi_{z,\text{la}}^1 \dot{\Pi}_{z,\text{la}}^1 (\Pi_{z,\text{la}}^1)^\perp . \quad (4.20)$$

Therefore, for $\Re s \gg 0$,

$$\zeta_{\text{la}}(s, \Delta_z, \dot{\Pi}_z^1 \mathbf{w}) = \text{Tr}^s (\dot{\Pi}_{z,\text{la}}^1 \Delta_z^{-s} \Pi_{z,\text{la}}^1) = 0 ,$$

yielding $\zeta_{\text{la}}(s, \Delta_z, \dot{\Pi}_z^1 \mathbf{w}) = 0$ for all s because this is a meromorphic function of s . Hence, since Δ_z and $\Pi_{z,\text{la}}^1 \mathbf{w}$ commute, Proposition 4.1 (i),(v) give

$$\partial_z \zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w}) = -s \zeta_{\text{la}}(s+1, \Delta_z, \dot{\Delta}_z \Pi_z^1 \mathbf{w}) = -s \text{Tr}^s (\dot{\Delta}_z \Delta_z^{-s-1} \Pi_{z,\text{la}}^1) .$$

Next, by (2.3),

$$\dot{\Delta}_z \Pi_{z,\text{la}}^1 = (\eta \wedge \delta_z + \delta_z \eta \wedge) \Pi_{z,\text{la}}^1 = \eta \wedge \delta_z \Pi_{z,\text{la}}^1 + \delta_z \eta \wedge \Pi_{z,\text{la}}^1 . \quad (4.21)$$

But, since $\Pi_z^1 \delta_z = 0$,

$$\text{Tr}^s (\delta_z \eta \wedge \Delta_z^{-s-1} \Pi_{z,\text{la}}^1) = -\text{Tr}^s (\eta \wedge \Delta_z^{-s-1} \Pi_{z,\text{la}}^1 \delta_z) = 0 . \quad (4.22)$$

Combining (4.6)–(4.22) and Proposition 4.1 (i), we get

$$\begin{aligned} \partial_z \zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w}) &= -s \text{Tr}^s (\eta \wedge \delta_z \Delta_z^{-s-1} \Pi_{z,\text{la}}^1) = -s \text{Tr}^s (\eta \wedge D_z \Delta_z^{-s-1} \Pi_{z,\text{la}}^1) \\ &= -s \zeta_{\text{la}}(s+1, \Delta_z, \eta \wedge D_z \mathbf{w}) . \quad \square \end{aligned}$$

THEOREM 4.13. *For $\mu \gg 0$, $\zeta_{\text{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is smooth on $\mathbb{C} \setminus ((1 - \mathbb{N}_0)/2)$.*

Proof. By Lemma 4.10, it is enough to prove that $\zeta_{\text{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is regular at $s = s_l = (n+1-l)/2$ for $l \leq n-1$, which means $\tilde{b}_l(z) = 0$.

First, take $l \leq n-2$. By Lemmas 4.10 and 4.11, Theorem 4.7, and Proposition 4.12, for $\mu \gg 0$,

$$0 = \partial_z \tilde{a}_{l+1}^1(z) = -(s_l - 1) \tilde{b}_l(z) .$$

Thus $\tilde{b}_l(z) = 0$ because $s_l - 1 = (n-1-l)/2 > 0$.

Next, take $l = n-1$. (The previous argument does not apply in this case because $s_{n-1} - 1 = 0$.) By Theorem 4.7 and Lemma 4.11, for $\mu \gg 0$,

$$\partial_z \zeta_{\text{la}}(0, \Delta_z, \Pi_z^1 \mathbf{w}) = \partial_z \tilde{a}_n^1(z) = 0 .$$

So there is a holomorphic function $\phi_z(s)$, defined around $s = 0$ and depending smoothly on z , such that

$$\partial_z \zeta_{\text{la}}(s, \Delta_z, \Pi_z^1 \mathbf{w}) = s \phi_z(s) .$$

Hence, by Proposition 4.12,

$$\phi_z(s) = -\zeta_{\text{la}}(s+1, \Delta_z, \eta \wedge D_z \mathbf{w}) ,$$

obtaining that $\zeta_{\text{la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w})$ is regular at $s = s_{n-1} = 1$. \square

From now on, we use the simpler notation

$$\zeta(s, z) = \zeta(s, \Delta_z, \eta \wedge D_z \mathbf{w}) , \quad \zeta_{\text{sm/la}}(s, z) = \zeta_{\text{sm/la}}(s, \Delta_z, \eta \wedge D_z \mathbf{w}) .$$

We may also add η to this notation, writing $\zeta(s, z, \eta)$ and $\zeta_{\text{sm/la}}(s, z, \eta)$.

COROLLARY 4.14. *If $\Re s > 1/2$ and $\mu \gg 0$, then*

$$\zeta_{\text{la}}(s, z) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \text{Tr}^s (\eta \wedge D_z e^{-t\Delta_z} P_{z,\text{la}}) dt ,$$

where the integral is absolutely convergent.

Proof. By Lemma 4.10, Theorem 4.13 and (4.18), for $\mu \gg 0$,

$$\text{Tr}^s (\eta \wedge D_z e^{-t\Delta_z} P_{z,\text{la}}) = O(t^{-1/2}) \quad (t \rightarrow 0) . \quad (4.23)$$

Moreover there is some $c > 0$ such that

$$\text{Tr}^s (\eta \wedge D_z e^{-t\Delta_z} P_{z,\text{la}}) = O(e^{-ct}) \quad (t \rightarrow \infty) . \quad (4.24)$$

So the stated integral is absolutely convergent for $\Re s > 1/2$, defining a holomorphic function of s on this half-plane. Then the stated equality is true because it holds for $\Re s \gg 0$. \square

COROLLARY 4.15. *For $\mu \gg 0$,*

$$\begin{aligned} \zeta_{\text{sm}}(1, z) &= \text{Tr}^s (\eta \wedge D_z^{-1} (\Pi_z^\perp)_{\text{sm}}) , \\ \zeta_{\text{la}}(1, z) &= \lim_{t \downarrow 0} \text{Tr}^s (\eta \wedge D_z^{-1} e^{-t\Delta_z} P_{z,\text{la}}) , \\ \zeta(1, z) &= \lim_{t \downarrow 0} \text{Tr}^s (\eta \wedge D_z^{-1} e^{-t\Delta_z} \Pi_z^\perp) . \end{aligned}$$

Proof. By Corollary 4.14, (4.23) and (4.24), and since

$$\text{Tr}^s (\eta \wedge D_z^{-1} e^{-t\Delta_z} P_{z,\text{la}}) = O(e^{-ct}) \quad (t \rightarrow \infty) ,$$

we get

$$\begin{aligned} \zeta_{\text{la}}(1, z) &= \int_0^\infty \text{Tr}^s (\eta \wedge D_z e^{-u\Delta_z} P_{z,\text{la}}) du = \lim_{t \downarrow 0} \int_t^\infty \text{Tr}^s (\eta \wedge D_z e^{-u\Delta_z} P_{z,\text{la}}) du \\ &= \lim_{t \downarrow 0} \text{Tr}^s (\eta \wedge D_z^{-1} e^{-t\Delta_z} P_{z,\text{la}}) . \end{aligned}$$

The expressions of $\zeta(1, z)$ and $\zeta_{\text{sm}}(1, z)$ follow from the expression of $\zeta_{\text{la}}(1, z)$ and Proposition 4.1 (iii). \square

Theorem 4.13 and Corollary 4.15 give Theorem 1.1 (i).

COROLLARY 4.16. *If $\mu \gg 0$, then (1.6) is true.*

Proof. Apply Proposition 4.12 and Theorem 4.13. \square

4.7 The case of exact forms

Let us consider the special case where $\eta = dh$ for a Morse function h . These results will be extended later, but the arguments are much simpler in this case, with weaker conditions and less ingredients.

LEMMA 4.17. *For $\mu \gg 0$,*

$$\begin{aligned} \mathrm{Tr}^s (\eta \wedge d_z^{-1} \Pi_{z,\mathrm{sm}}^1) &= -\mathrm{Tr}^s (h (\Pi_z^\perp)_{\mathrm{sm}}) , \\ \mathrm{Tr}^s (\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_{z,\mathrm{la}}^1) &= -\mathrm{Tr}^s (h e^{-t\Delta_z} P_{z,\mathrm{la}}) , \\ \mathrm{Tr}^s (\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1) &= -\mathrm{Tr}^s (h e^{-t\Delta_z} \Pi_z^\perp) . \end{aligned}$$

Proof. We prove the first equality, the other ones being analogous. Since $\eta \wedge = [d, h]$,

$$\begin{aligned} \mathrm{Tr}^s (\eta \wedge d_z^{-1} \Pi_{z,\mathrm{sm}}^1) &= \mathrm{Tr}^s ([d_z, h] d_z^{-1} \Pi_{z,\mathrm{sm}}^1) = \mathrm{Tr}^s (d_z h d_z^{-1} \Pi_{z,\mathrm{sm}}^1) - \mathrm{Tr}^s (h d_z d_z^{-1} \Pi_{z,\mathrm{sm}}^1) \\ &= -\mathrm{Tr}^s (h d_z^{-1} \Pi_{z,\mathrm{sm}}^1 d_z) - \mathrm{Tr}^s (h \Pi_{z,\mathrm{sm}}^1) \\ &= -\mathrm{Tr}^s (h d_z^{-1} d_z \Pi_{z,\mathrm{sm}}^2) - \mathrm{Tr}^s (h \Pi_{z,\mathrm{sm}}^1) \\ &= -\mathrm{Tr}^s (h \Pi_{z,\mathrm{sm}}^2) - \mathrm{Tr}^s (h \Pi_{z,\mathrm{sm}}^1) = -\mathrm{Tr}^s (h (\Pi_z^\perp)_{\mathrm{sm}}) . \quad \square \end{aligned}$$

REMARK 4.18. The last equality of Lemma 4.17 does not require any condition on η or g (h can be any smooth real-valued function).

COROLLARY 4.19. *For $\mu \gg 0$,*

$$\begin{aligned} \zeta_{\mathrm{sm}}(1, z) &= -\mathrm{Tr}^s (h (\Pi_z^\perp)_{\mathrm{sm}}) , \\ \zeta_{\mathrm{la}}(1, z) &= -\lim_{t \downarrow 0} \mathrm{Tr}^s (h e^{-t\Delta_z} P_{z,\mathrm{la}}) , \\ \zeta(1, z) &= -\lim_{t \downarrow 0} \mathrm{Tr}^s (h e^{-t\Delta_z} \Pi_z^\perp) . \end{aligned}$$

Proof. Apply Corollary 4.15 and Lemma 4.17. □

COROLLARY 4.20. *The value $\zeta_{\mathrm{sm}}(1, z)$ is uniformly bounded on z for $\mu \gg 0$.*

Proof. The operator $h (\Pi_z^\perp)_{\mathrm{sm}}$ is uniformly bounded and, for $\mu \gg 0$, has uniformly bounded rank. So $\mathrm{Tr}^s (h (\Pi_z^\perp)_{\mathrm{sm}})$ is uniformly bounded on z for $\mu \gg 0$, and therefore the result follows from Corollary 4.19. □

COROLLARY 4.21. *If $\mu \gg 0$, then $\zeta(1, z), \zeta_{\mathrm{sm}/\mathrm{la}}(1, z) \in \mathbb{R}$.*

Proof. We consider the case of $\zeta(1, z)$, the other cases being similar. By Corollary 4.19, it is enough to prove that $\mathrm{Tr}^s (h e^{-t\Delta_z} \Pi_z^\perp) \in \mathbb{R}$. This is true because, taking adjoints,

$$\mathrm{Tr}^s (h e^{-t\Delta_z} \Pi_z^\perp) = \overline{\mathrm{Tr}^s (\Pi_z^\perp e^{-t\Delta_z} h)} = \overline{\mathrm{Tr}^s (h \Pi_z^\perp e^{-t\Delta_z})} = \overline{\mathrm{Tr}^s (h e^{-t\Delta_z} \Pi_z^\perp)} . \quad \square$$

COROLLARY 4.22. *If M is oriented, n is even and $|\mu| \gg 0$, then*

$$\begin{aligned} \zeta(1, z) &= \zeta(1, -\bar{z}) = \zeta(1, -z) = \zeta(1, \bar{z}) , \\ \zeta_{\mathrm{sm}/\mathrm{la}}(1, z) &= \zeta_{\mathrm{sm}/\mathrm{la}}(1, -\bar{z}) = \zeta_{\mathrm{sm}/\mathrm{la}}(1, -z) = \zeta_{\mathrm{sm}/\mathrm{la}}(1, \bar{z}) . \end{aligned}$$

Proof. We prove the case of $\zeta_{\mathrm{sm}}(1, z)$, the other cases being similar. By (2.15) and since n is even,

$$\begin{aligned} \mathrm{Tr}^s (h (\Pi_z^\perp)_{\mathrm{sm}}) &= \mathrm{Tr}^s (\star \star^{-1} h (\Pi_z^\perp)_{\mathrm{sm}}) = \mathrm{Tr}^s (\star^{-1} h (\Pi_z^\perp)_{\mathrm{sm}} \star) \\ &= \mathrm{Tr}^s (\star^{-1} \star h (\Pi_{-\bar{z}}^\perp)_{\mathrm{sm}}) = \mathrm{Tr}^s (h (\Pi_{-\bar{z}}^\perp)_{\mathrm{sm}}) . \end{aligned}$$

Thus the first equality of the statement follows by Corollary 4.19. The second equality follows with a similar argument, using $\bar{\star}$ instead of \star . The third equality is a consequence of the other ones. \square

THEOREM 4.23. *The following limit holds uniformly on ν :*

$$\lim_{\mu \rightarrow +\infty} \zeta_{\text{la}}(1, z) = - \int_M h e(M, \nabla^M) \text{vol} + \sum_{p \in \mathcal{X}}^n (-1)^{\text{ind}(p)} h(p) .$$

Proof. By (4.11), (4.12), Theorem 4.2 and Corollary 4.19, for $\mu \gg 0$,

$$\begin{aligned} \zeta_{\text{la}}(1, z) &= - \lim_{t \downarrow 0} \text{Tr}^s (h e^{-t\Delta_z} P_{z, \text{la}}) = - \int_M h(x) \text{tr}^s \tilde{e}_n(x, z) \text{vol}(x) \\ &= - \int_M h(x) \text{tr}^s e_n(x, z) \text{vol}(x) + \text{Tr}^s(hP_{z, \text{sm}}) \\ &= - \int_M h e(M, \nabla^M) \text{vol} + \text{Tr}^s(hP_{z, \text{sm}}) . \end{aligned}$$

According to Corollary 3.9, the elements $P_{z, \text{sm}} e_{p, z}$ ($p \in \mathcal{X}$) form a base of $E_{z, \text{sm}}^k$ when $\mu \gg 0$. Applying the Gram-Schmidt process to this base, we get an orthonormal frame $\tilde{e}_{p, z}$ ($p \in \mathcal{X}$) of $E_{z, \text{sm}}$. By Proposition 3.7 for $m = 0$ and (3.9)–(3.12), we get

$$\lim_{\mu \rightarrow +\infty} \langle h \tilde{e}_{p, z}, \tilde{e}_{q, z} \rangle = \lim_{\mu \rightarrow +\infty} \langle h e_{p, z}, e_{q, z} \rangle = h(p) \delta_{pq} .$$

Hence

$$\lim_{\mu \rightarrow +\infty} \text{Tr}^s(hP_{z, \text{sm}}) = \sum_{k=0}^n (-1)^k \sum_{p \in \mathcal{X}_k} h(p) . \quad \square$$

5. The small complex vs the Morse complex

5.1 Preliminaries on Smale vector fields

5.1.1 Vector fields with Morse type zeros Let X be a real smooth vector field on M with flow $\phi = \{\phi^t\}$. Let $\mathcal{X} = \text{Zero}(X)$ denote the set of zeros of X (or rest points ϕ). It is said that a zero p of X is of *Morse type* with (*Morse*) *index* of $\text{ind}(p)$ if, using the notation (3.2),

$$X = - \sum_{j=1}^n \epsilon_{p, j} x_p^j \frac{\partial}{\partial x_p^j} \quad (5.1)$$

on the domain U_p of some coordinates $x_p = (x_p^1, \dots, x_p^n)$ at p , also called *Morse coordinates*. This condition means that $X = - \text{grad}_g h_{X, p}$ on U_p , where $h_{X, p}$ and g are given on U_p by the right-hand side of (3.1) and (3.3). The coordinates x_p used in (5.1) are not unique; that expression is invariant by taking positive multiples of the coordinates (contrary to the expressions (3.1), (3.3) and (3.4)). But $\text{ind}(p)$ is independent of x_p . If (3.3), (3.4) and (5.1) hold with the same coordinates, then η and g are said to be in *standard form* with respect to X around p . In this case, $C\eta$ and Cg ($C > 0$) are also in standard form with respect to X around p ; indeed, $C\eta$, X and Cg satisfy (3.3), (3.4) and (5.1) with the coordinates $\sqrt{C}x_p$. When g is defined on M , if η and g are in standard form with respect to X around every $p \in \mathcal{X}$, then η and g are said to be in *standard form* with respect to X . This concept is also applied to any Morse function h on M referring to dh and g . The reference to g may be omitted in this terminology.

The vector field X is fixed in most of the paper and plays an auxiliary role. Unless otherwise indicated, we assume from now on that X has Morse type zeros. Then \mathcal{X} is finite, and the sets \mathcal{X}_k , \mathcal{X}_+ and $\mathcal{X}_{<k}$ are defined like in Section 3.1.

5.1.2 Stable and unstable manifolds For $k = 0, \dots, n$ and $p \in \mathcal{X}_k$, the *stable/unstable manifolds* of p are smooth injective immersions, $\iota_p^\pm : W_p^\pm \rightarrow M$, where the images $\iota_p^\pm(W_p^\pm)$ consist of the points satisfying $\phi^t(x) \rightarrow p$ as $t \rightarrow \pm\infty$, and the manifolds W_p^+ and W_p^- are diffeomorphic to \mathbb{R}^{n-k} and \mathbb{R}^k , respectively [Sma63, Theorem 9.1]. In particular, $p \in \iota_p^\pm(W_p^\pm)$, and the maps ι_p^+ and ι_p^- meet transversely at p . Let $p^\pm = (\iota_p^\pm)^{-1}(p)$. Assume every U_p is connected, and let U_p^\pm be the connected component of $(\iota_p^\pm)^{-1}(U_p)$ that contains p^\pm . The restriction $\iota_p^\pm : U_p^\pm \rightarrow (x_p^\pm)^{-1}(0)$ is a diffeomorphism, and therefore $(U_p^\pm, x_p^\pm \iota_p^\pm)$ is a coordinate system of W_p^\pm centered at p^\pm .

5.1.3 Gradient-like vector fields Given a Morse function h on M in standard form with respect to X , we have $X = -\text{grad}_g h$ on M for some Riemannian metric g if and only if $Xh < 0$ on $M \setminus \mathcal{X}$ [BFK10, Lemma 2.1], [Lau12, Section 6.1.3]; in this case, X is said to be *gradient-like* (with respect to h). If X is gradient-like, then the maps ι_p^\pm are embeddings [Sma60b, Lemma 3.8], [BFK10, Lemma 2.2], and their images cover M [Sma61, Lemma 1.1], [BFK10, Corollary 2.5]. Thus, in this case, we will write $W_p^\pm = \iota_p^\pm(W_p^\pm)$ and $p^\pm = p$, and ι_p^\pm becomes the inclusion map.

Unless otherwise indicated, we also assume in the rest of the paper that X is gradient-like.

5.1.4 Smale vector fields X is said to be *Smale* if $W_p^+ \pitchfork W_q^-$ for all $p, q \in \mathcal{X}$. In this case, $\mathcal{M}(p, q) := W_p^+ \cap W_q^-$ is a ϕ -saturated smooth submanifold of dimension $\text{ind}(p) - \text{ind}(q)$. In particular, $\mathcal{M}(p, p) = \{p\}$, and define $\mathcal{T}(p, p) = \emptyset$. If $p \neq q$, then the induced \mathbb{R} -action on $\mathcal{M}(p, q)$ is free and proper, and therefore the orbit space $\mathcal{T}(p, q) := \mathcal{M}(p, q)/\mathbb{R}$ is a smooth manifold of dimension $\text{ind}(p) - \text{ind}(q) - 1$. The elements of $\mathcal{T}(p, q)$ are the (unparameterized) trajectories with α -limit $\{p\}$ and ω -limit $\{q\}$, which are oriented by X . If $\text{ind}(p) \leq \text{ind}(q)$, then $\mathcal{T}(p, q) = \emptyset$. If $\text{ind}(p) - \text{ind}(q) = 1$, then $\mathcal{T}(p, q)$ consists of isolated points, each of them representing a trajectory in M . Let $\mathcal{T} = \bigcup_{p, q \in \mathcal{X}} \mathcal{T}(p, q)$, and

$$\mathcal{T}_p^1 = \bigcup_{q \in \mathcal{X}_{\text{ind}(p)-1}} \mathcal{T}(p, q), \quad \mathcal{T}_k^1 = \bigcup_{p \in \mathcal{X}_k} \mathcal{T}_p^1, \quad \mathcal{T}^1 = \bigcup_{k=0}^n \mathcal{T}_k^1.$$

The elements of \mathcal{T}^1 are called *instantons*.¹

Unless otherwise indicated, besides the above conditions, we assume from now on that X is Smale; i.e., we assume (b) (Section 1.1). Thus the α - and ω -limits of the orbits of X are zero points [Sma61, Theorem B and Lemma 1.1].

5.1.5 Lyapunov forms Any $\eta \in Z^1(M, \mathbb{R})$ is said to be *Lyapunov* for X if $\eta(X) < 0$ on $M \setminus \mathcal{X}$ [BH08, Definition 2.3]. By (b), every class in $H^1(M, \mathbb{R})$ has a representative η which is Lyapunov for X and $\eta^\sharp = -X$ for some Riemannian metric g on M , with η and g in standard form with respect to X [BH04, Proposition 16 (i)], [BH08, Observations 2.5 and 2.6], [HM06, Lemma 3.7], [Lau12, Section 6.1.3]. Moreover X can be C^∞ -approximated by gradient-like Smale vector fields that agree with X around \mathcal{X} [BH08, Proposition 2.4] (this follows from [Sma61, Theorem A]). A well known consequence is that, for any Morse function h , there is a C^∞ -dense set of Riemannian metrics g on M such that $-\text{grad}_g h$ is Smale; this density is also true in the subspace of metrics

¹Sometimes, the elements of \mathcal{T} are called *instantons*, and the elements of \mathcal{T}^1 *proper instantons* [Bot88].

that are Euclidean with respect to Morse coordinates on given neighborhoods of the critical points.

5.1.6 Completion of the unstable manifolds

PROPOSITION 5.1 [BZ92, Appendix by F. Laudenbach, Proposition 2], [Lat94, Chapter 2], [Bur97, Theorem 2.1], [BH01, Theorem 1], [BFK10, Theorem 4.4], [Lau12, Sections A.2 and A.8], [Min15, Corollary 2.3.2]. *The following holds for every $p \in \mathcal{X}_k$ ($k = 0, \dots, n$):*

- (i) $\overline{W_p^-}$ is a C^1 submanifold with conic singularities (in the sense of [BZ92, Appendix by F. Laudenbach, Section a]) and [Lau12, Appendix A.1]) and a Whitney stratified subspace. Its strata are the submanifolds W_q^- for $q \in \mathcal{X}_{<k}$ with $\mathcal{T}(p, q) \neq \emptyset$. As a consequence, W_p^- has finite volume, and

$$\overline{W_q^-} \cap \overline{W_p^-} \subset \bigcup_{x \in \widehat{\mathcal{X}}_{<k}} W_x^-$$

if $q \neq p$ in \mathcal{X}_k ; in particular, $p \notin \overline{W_q^-}$.

- (ii) There is a compact k -manifold with corners² \widehat{W}_p^- whose l -corner³ ($0 \leq l \leq k$) is

$$\partial_l \widehat{W}_p^- = \bigsqcup_{(q_0, \dots, q_l) \in \{p\} \times \mathcal{X}^l} \left(\prod_{j=1}^l \mathcal{T}(q_{j-1}, q_j) \right) \times W_{q_l}^-.$$

In particular, the interior of \widehat{W}_p^- is $\partial_0 \widehat{W}_p^- = W_p^-$, and the set $\mathcal{T}(p, q)$ is finite if $q \in \mathcal{X}_{k-1}$.

- (iii) There is a smooth map $\hat{\iota}_p^- : \widehat{W}_p^- \rightarrow M$ whose restriction to every component of $\partial_l \widehat{W}_p^-$ in (ii) is given by the factor projection to $W_{q_l}^-$; in particular, $\hat{\iota}_p^- = \iota_p^-$ on W_p^- . Moreover $\hat{\iota}_p^- : \widehat{W}_p^- \rightarrow \overline{W_p^-}$ is a stratified map.

By Proposition 5.1 (i), we can choose the open sets U_p ($p \in \mathcal{X}_k$, $k = 0, \dots, n$) so small that $U_p \cap \overline{W_q^-} = \emptyset$ if $q \neq p$ in \mathcal{X}_k .

For every $q \in \mathcal{X}_{k-1}$ and $\gamma \in \mathcal{T}(p, q)$, the closure $\bar{\gamma}$ in M is a compact oriented submanifold with boundary of dimension one, and $\partial \bar{\gamma} = \{p, q\}$. We may also consider $\bar{\gamma}$ as the closure of γ in \widehat{W}_p^- .

5.2 Preliminaries on the Morse complex

5.2.1 *The Morse complex* Fix an orientation \mathcal{O}_p^- of every unstable manifold W_p^- ($p \in \mathcal{X}_k$, $k = 0, \dots, n$), which can be also considered as an orientation of \widehat{W}_p^- . Then $W_p^- \equiv (W_p^-, \mathcal{O}_p^-)$ defines a current of dimension k on M , also denoted by W_p^- ; namely, for $\alpha \in \Omega^k(M)$,

$$\langle W_p^-, \alpha \rangle = \int_{W_p^-} \alpha = \int_{\widehat{W}_p^-} (\hat{\iota}_p^-)^* \alpha.$$

Let $\partial_1 \mathcal{O}_p^-$ be the orientation of $\partial_1 \widehat{W}_p^-$ induced by \mathcal{O}_p^- like in the Stokes' theorem. The restriction of $\partial_1 \mathcal{O}_p^-$ to every component $\mathcal{T}(p, q) \times W_q^-$ ($q \in \mathcal{X}_{k'}$) of $\partial_1 \widehat{W}_p^-$ is of the form $\mathcal{O}_{p,q} \otimes \mathcal{O}_q^-$ for a unique orientation $\mathcal{O}_{p,q}$ of $\mathcal{T}(p, q)$. If $k' = k - 1$, then $\mathcal{O}_{p,q}$ can be represented by a unique function $\epsilon_{p,q} : \mathcal{T}(p, q) \rightarrow \{\pm 1\}$; combining these functions, we get a map $\epsilon : \mathcal{T}^1 \rightarrow \{\pm 1\}$. By

²In the sense of [Mel96, Section 1.1.8].

³The union of the interiors of the boundary faces of codimension l .

the descriptions of $\partial_1 \widehat{W}_p^-$ and $\hat{\iota}_p^- : \partial_1 \widehat{W}_p^- \rightarrow M$, and by the Stokes' theorem for manifolds with corners, we have [BZ92, Appendix by F. Laudenbach], [HM06, Remark 1.9], [BFK10, Theorem 3.6 and Proposition 5.3], [Lau12, Section 6.5.3]

$$\partial W_p^- = \sum_{q \in \mathcal{X}_{k-1}, \gamma \in \mathcal{T}(p,q)} \epsilon(\gamma) W_q^- . \quad (5.2)$$

Thus the currents W_p^- ($p \in \mathcal{X}$) generate over \mathbb{C} a finite dimensional subcomplex $(C_\bullet(X, W^-), \partial)$ of the complex $(\Omega(M)', \partial)$ of currents on M , called the *Morse complex*. The simpler notation $\mathbf{C}_\bullet = \mathbf{C}_\bullet(X) = C_\bullet(X, W^-)$ may be also used. Moreover $\mathbf{C}_\bullet \hookrightarrow \Omega(M)'$ induces an isomorphism⁴ $H_\bullet(\mathbf{C}_\bullet, \partial) \cong H_\bullet(M, \mathbb{C})$ [Tho49, Sma60a, Mil65] (see also [Flo89, Sch93, Sch99], [HS85, Theorem 0.1], [BZ92, Appendix by F. Laudenbach, Proposition 7], [Lau12, Section 6.6.5]).

The *dual Morse complex* is the dual $(\mathbf{C}^\bullet(X, W^-), \mathbf{d})$ of $(\mathbf{C}_\bullet, \partial)$; namely, $C^k(X, W^-) = (\mathbf{C}_k)^\bullet \equiv \mathbb{C}^{\mathcal{X}_k}$ ($k = 0, \dots, n$). We will usually denote $\mathbf{C}^\bullet = \mathbf{C}^\bullet(X) = C^\bullet(X, W^-)$. Moreover boldface notation will be used for elements of \mathbf{C}^\bullet and operators on \mathbf{C}^\bullet . Let \mathbf{e}_p ($p \in \mathcal{X}$) denote the elements of the canonical base of \mathbf{C}^\bullet , determined by $\mathbf{e}_p(q) = \delta_{pq}$, using the Kronecker delta. Then, for $q \in \mathcal{X}_{k-1}$,

$$\mathbf{d}\mathbf{e}_q = \sum_{p \in \mathcal{X}_k, \gamma \in \mathcal{T}(p,q)} \epsilon(\gamma) \mathbf{e}_p . \quad (5.3)$$

5.2.2 The perturbed Morse complex Take any $\eta \in Z^1(M, \mathbb{R})$ defining a class $\xi \in H^1(M, \mathbb{R})$ (there is no need of any condition on η or g in Sections 5.2.2 to 5.2.4). For reasons of brevity, write $\eta(\gamma) = \int_\gamma \eta$ for every $\gamma \in \mathcal{T}^1$. According to [BH01, BH04, BH08], $(\mathbf{C}^\bullet, \mathbf{d})$ has an analog of the Witten's perturbation, $(\mathbf{C}^\bullet, \mathbf{d}_z = \mathbf{d}_{z\eta})$ ($z \in \mathbb{C}$), where, for $q \in \mathcal{X}_{k-1}$ ($k = 1, \dots, n$),

$$\mathbf{d}_z \mathbf{e}_q = \sum_{p \in \mathcal{X}_k, \gamma \in \mathcal{T}(p,q)} \epsilon(\gamma) e^{z\eta(\gamma)} \mathbf{e}_p . \quad (5.4)$$

If $\eta = dh$ for some $h \in C^\infty(M, \mathbb{R})$, then $\mathbf{d}_z = e^{-zh} \mathbf{d} e^{zh}$ on \mathbf{C}^\bullet because $\eta(\gamma) = h(q) - h(p)$ for $p \in \mathcal{X}_k$, $q \in \mathcal{X}_{k-1}$ and $\gamma \in \mathcal{T}(p, q)$; here, $e^{\pm zh}$ also denotes the operator of multiplication by the restriction of this function to \mathcal{X} . It will be said that $(\mathbf{C}^\bullet, \mathbf{d}_z)$ ($z \in \mathbb{C}$) is the *perturbed dual Morse complex* defined by X and η . A perturbation $(\mathbf{C}_\bullet, \partial^z)$ is similarly defined, adding the factor $e^{z\eta(\gamma)}$ to the terms of the right-hand side of (5.2).

Since W_p^- ($p \in \mathcal{X}_k$, $k = 0, \dots, n$) is diffeomorphic to \mathbb{R}^k , there is a unique $h_{\eta,p}^- \in C^\infty(W_p^-, \mathbb{R})$ such that $h_{\eta,p}^-(\hat{p}^-) = 0$ and $dh_{\eta,p}^- = (\iota_p^-)^* \eta$. Indeed $h_{\eta,p}^-$ has a smooth extension $\hat{h}_{\eta,p}^-$ to \widehat{W}_p^- because \widehat{W}_p^- is contractile. By Proposition 5.1 (ii), for all $q \in \mathcal{X}_{k-1}$ and $\gamma \in \mathcal{T}(p, q)$, we have $\hat{h}_{\eta,p}^-(\gamma, \hat{q}^-) = \eta(\gamma)$, yielding

$$\hat{h}_{\eta,q}^- \equiv \hat{h}_{\eta,p}^- - \eta(\gamma) \quad \text{on} \quad \widehat{W}_q^- \equiv \{\gamma\} \times \widehat{W}_q^- \subset \partial_1 \widehat{W}_p^- .$$

According to [BH01, Proposition 4], [BH04, Proposition 10], [BH08, Propositions 2.15 and 2.16 and Section 6.2], a surjective homomorphism of complexes,

$$\Phi_z : (\Omega(M), d_z) \rightarrow (\mathbf{C}^\bullet, \mathbf{d}_z) ,$$

⁴Actually, $H_\bullet(M, \mathbb{Z})$ is isomorphic to the homology of the complex of free Abelian groups generated by the currents W_p^- .

is defined by

$$\Phi_z(\omega)(p) = \int_{W_p^-} e^{zh_{\eta,p}^-} \omega = \int_{\widehat{W}_p^-} e^{z\hat{h}_{\eta,p}^-} (\hat{i}_p^-)^* \omega .$$

Moreover Φ_z is a quasi-isomorphism for all $z \in \mathbb{C}$ [BZ92, Proposition 1 in the Appendix by F. Laudenbach] (see also [BZ92, Theorem 2.9], [BZ94, Theorem 1.6], [BH08, Proposition 2.17 and Section 6.2]). Then, by (3.14), the same is true for

$$\Phi_z : (E_{z,\text{sm}}, d_z) \rightarrow (\mathbf{C}^\bullet, \mathbf{d}_z) .$$

Since direct adaptation of [BH04, Appendix A] shows that, for $k = 0, \dots, n$, $\dim H^k(\mathbf{C}^\bullet, \mathbf{d}_z)$ is independent of $z \in \mathbb{C}$ with $|\mu| \gg 0$, we get (2.9).

5.2.3 Morse complex with coefficients in a flat vector bundle With more generality [BZ92, Section 1c)], for a flat vector bundle F , we may consider $(C^\bullet(X, W^-, F), \mathbf{d}^F)$, where $C^k(X, W^-, F) = \bigoplus_{p \in \mathcal{X}_k} F_p$, and $\mathbf{d}^F \mathbf{e}$ ($\mathbf{e} \in F_q$, $q \in \mathcal{X}_{k-1}$) is defined like in the right-hand side of (5.3), replacing \mathbf{e}_p with the parallel transport of \mathbf{e} along $\bar{\gamma}^{-1}$. This is the dual of the complex $(C_\bullet(X, W^-, F^*), \partial^{F^*})$, where $C_k(X, W^-, F^*) = \bigoplus_{p \in \mathcal{X}_k} F_p^*$, and $\partial^F f$ ($f \in F_p^*$, $p \in \mathcal{X}_k$) is defined like in the right-hand side of (5.2), replacing W_q^- with the parallel transport of f along $\bar{\gamma}$. A homomorphism

$$\Phi^F = \Phi^{X,F} : (\Omega(M, F), d) \rightarrow (C^\bullet(X, W^-, F), \mathbf{d}^F)$$

can be defined like Φ_z , using the isomorphism

$$\Omega^\bullet(\widehat{W}_p^-, (\hat{i}_p^-)^* F) \cong \Omega^\bullet(\widehat{W}_p^-) \otimes F_p$$

given by the parallel transport of $(\hat{i}_p^-)^* F$. With this generality, Φ^F is also induces a quasi-isomorphism [BZ92, Theorem 2.9]. If $F = \mathcal{L}^z$ (Section 2.1.4), then

$$(C^\bullet(X, W^-, \mathcal{L}^z), \mathbf{d}^{\mathcal{L}^z}) \equiv (\mathbf{C}^\bullet, \mathbf{d}_z) , \quad \Phi^{\mathcal{L}^z} \equiv \Phi_z .$$

5.2.4 Hodge theory of the Morse complex Consider the Hermitian scalar product on \mathbf{C}^\bullet so that the canonical base \mathbf{e}_p ($p \in \mathcal{X}$) is orthonormal. All operators induced by \mathbf{d}_z and this Hermitian structure are called *perturbed Morse operators*. For instance, besides \mathbf{d}_z , we have the perturbed Morse operators

$$\delta_z = \mathbf{d}_z^* , \quad \mathbf{D}_z = \mathbf{d}_z + \delta_z , \quad \mathbf{\Delta}_z = \mathbf{D}_z^2 = \mathbf{d}_z \delta_z + \delta_z \mathbf{d}_z .$$

In particular, it will be said that $\mathbf{\Delta}_z$ is the *perturbed Morse Laplacian*, and its eigenvalues will be called *perturbed Morse eigenvalues*. If $z = 0$, we omit the subscript “ z ” and the word “perturbed”. From (5.4), we easily get

$$\delta_z \mathbf{e}_p = \sum_{q \in \mathcal{X}_{k-1}, \gamma \in \mathcal{T}(p,q)} e^{\bar{z}\eta(\gamma)} \epsilon(\gamma) \mathbf{e}_q , \quad (5.5)$$

for $p \in \mathcal{X}_k$. We also have

$$\begin{aligned} \mathbf{C}^\bullet &= \ker \mathbf{\Delta}_z \oplus \text{im } \mathbf{d}_z \oplus \text{im } \delta_z , \\ \ker \mathbf{\Delta}_z &= \ker \mathbf{D}_z = \ker \mathbf{d}_z \cap \ker \delta_z , \quad \text{im } \mathbf{\Delta}_z = \text{im } \mathbf{D}_z = \text{im } \mathbf{d}_z \oplus \text{im } \delta_z . \end{aligned}$$

The orthogonal projections of \mathbf{C}^\bullet to $\ker \mathbf{\Delta}_z$, $\text{im } \mathbf{d}_z$ and $\text{im } \delta_z$ are denoted by $\mathbf{\Pi}_z = \mathbf{\Pi}_z^0$, $\mathbf{\Pi}_z^1$ and $\mathbf{\Pi}_z^2$, respectively. The compositions $\mathbf{d}_z^{-1} \mathbf{\Pi}^1$, $\delta_z^{-1} \mathbf{\Pi}^2$ and $\mathbf{D}_z^{-1} \mathbf{\Pi}^1$ are defined like in Section 2.1.2, and there is an obvious version of the commutative diagram (2.7).

5.3 The small complex vs the Morse complex

Our main objects of interest are the form $\eta \in Z^1(M; \mathbb{R})$ and the Riemannian metric g ; X plays an auxiliary role. Unless otherwise indicated, assume from now on that η is Lyapunov for X , and η and g are in standard form with respect to X ; i.e., we assume (c) besides (a) and (b). Since every $\xi \in H^1(M, \mathbb{R})$ is Lyapunov for X by (b), we can choose some $\eta \in \xi$ and g satisfying (a) and (c).

For every $p \in \mathcal{X}$, consider the functions $h_{\eta,p}$, $h_{X,p}$, $h_{\eta,p}^-$ and $\hat{h}_{\eta,p}^-$ defined in Sections 3.1, 5.1.1 and 5.2.2. Since η and g are in standard form with respect to X , we have $h_{\eta,p} = h_{X,p}$ on U_p , and

$$h_{\eta,p}^- = h_{\eta,p} = -\frac{1}{2}|x_p^-|^2 \quad (5.6)$$

on U_p^- . From now on, the simpler notation $h_p = h_{\eta,p} = h_{X,p}$, $h_p^- = h_{\eta,p}^-$ and $\hat{h}_p^- = \hat{h}_{\eta,p}^-$ will be used. Since η is Lyapunov for X ,

$$h_p^- < 0 \quad \text{on} \quad W_p^- \setminus \{p\}. \quad (5.7)$$

Consider the notation of Section 5.2.2. Let $J_z : \mathbf{C}^\bullet \rightarrow E_z$ be the \mathbb{C} -linear isometry given by $J_z(\mathbf{e}_p) = e_{p,z}$, and let $\Psi_z = P_{z,\text{sm}} J_z : \mathbf{C}^\bullet \rightarrow E_{z,\text{sm}}$, which is an isomorphism for $\mu \gg 0$ (Corollary 3.9). By Proposition 3.7,

$$\|\Psi_z \mathbf{e}\| = (1 + O(e^{-c\mu})) \|\mathbf{e}\| \quad (\mu \rightarrow +\infty)$$

for all $\mathbf{e} \in \mathbf{C}^\bullet$. Using polarization (see e.g. [Kat95, Section I.6.2]) and conjugation, this means that, as $\mu \rightarrow +\infty$,

$$\Psi_z^* \Psi_z = 1 + O(e^{-c\mu}), \quad \Psi_z \Psi_z^* = 1 + O(e^{-c\mu}). \quad (5.8)$$

Notation 5.2. Consider functions $u(z)$ and $v(z)$ ($z \in \mathbb{C}$) with values in Banach spaces. The notation $u(z) \asymp_0 v(z)$ ($\mu \rightarrow \pm\infty$) means

$$u(z) = v(z) + O(e^{-c|\mu|}) \quad (\mu \rightarrow \pm\infty).$$

We may also consider this asymptotic relation when the Banach spaces also depend on z .

THEOREM 5.3 Cf. [BZ94, Theorem 6.11], [Zha01, Theorem 6.9], [BH01, Theorem 4]. *For every $\tau \in \mathbb{R}$, as $\mu \rightarrow +\infty$,*

$$\Phi_{z+\tau} \Psi_z \asymp_0 \left(\frac{\pi}{\mu + \tau/2} \right)^{N/2} \left(\frac{\mu}{\pi} \right)^{n/4}.$$

Proof. We adapt the proof of [Zha01, Theorem 6.9] to the case of complex parameter. For every $p \in \mathcal{X}_k$,

$$\Phi_{z+\tau} \Psi_z \mathbf{e}_p = \sum_{q \in \mathcal{X}_k} \mathbf{e}_q \int_{\widehat{W}_q^-} e^{(z+\tau)\hat{h}_q^-} (\hat{t}_q^-)^* P_{z,\text{sm}} e_{p,z}. \quad (5.9)$$

Then the result follows by checking the asymptotics of these integrals.

In the case $q = p$, by (5.7) and Corollary 3.8,

$$\int_{\widehat{W}_p^-} e^{(z+\tau)\hat{h}_p^-} (\hat{t}_p^-)^* (P_{z,\text{sm}} - 1) e_{p,z} \asymp_0 0.$$

But, by Proposition 3.1 (iii), (3.9)–(3.12) and (5.6),

$$\begin{aligned}
 \int_{\widehat{W}_p^-} e^{(z+\tau)\hat{h}_p^-} (\hat{t}_p^-)^* e_{p,z} &= \int_{\widehat{W}_p^-} e^{(z+\tau)\hat{h}_p^-} (\hat{t}_p^-)^* (e^{-i\nu h_p} e_{p,\mu}) = \int_{\widehat{W}_p^-} e^{(\mu+\tau)\hat{h}_p^-} (\hat{t}_p^-)^* e_{p,\mu} \\
 &= \frac{1}{a_\mu} \left(\int_{-2r}^{2r} \rho(x) e^{-(2\mu+\tau)x^2/2} dx \right)^k \\
 &= \left(\frac{\pi}{\mu + \tau/2} \right)^{k/2} \left(\frac{\mu}{\pi} \right)^{n/4} (1 + O(e^{-c\mu})) .
 \end{aligned} \tag{5.10}$$

(When $\tau = 0$, the last equality is the same as [Zha01, Eq. (6.30)].)

For $q \neq p$ in \mathcal{X}_k , since $e_{p,z} = 0$ on $\overline{W_q^-}$ because $U_p \cap \overline{W_q^-} = \emptyset$ (Section 5.1.6), like in the previous case, we get

$$\int_{\widehat{W}_q^-} e^{(z+\tau)\hat{h}_q^-} (\hat{t}_q^-)^* P_{z,\text{sm}} e_{p,z} \asymp_0 0 . \quad \square$$

COROLLARY 5.4. For every $\tau \in \mathbb{R}$, if $\mu \gg 0$, then $\Phi_{z+\tau} : E_{z,\text{sm}} \rightarrow \mathbf{C}^\bullet$ is an isomorphism.

Proof. Apply Theorem 5.3 and Corollary 3.9. □

Remark 5.5. The argument of the proof of Theorem 5.3 shows that

$$\Phi_z J_z = \left(\frac{\pi}{\mu} \right)^{N/2-n/4} + O(e^{-c\mu}) \quad (\mu \rightarrow +\infty) .$$

So $\Phi_z : E_z \rightarrow \mathbf{C}^\bullet$ is an isomorphism for $\mu \gg 0$ (see also [BH08, Lemma 5.2]).

Let

$$\tilde{\Psi}_z = \left(\frac{\mu}{\pi} \right)^{N/2-n/4} \Psi_z : \mathbf{C}^\bullet \rightarrow E_{z,\text{sm}} .$$

COROLLARY 5.6. Consider $\tilde{\Psi}_z^* : E_{z,\text{sm}} \rightarrow \mathbf{C}^\bullet$. As $\mu \rightarrow +\infty$,

$$\tilde{\Psi}_z^* \tilde{\Psi}_z = \left(\frac{\mu}{\pi} \right)^{N-n/2} + O(e^{-c\mu}) , \quad \tilde{\Psi}_z \tilde{\Psi}_z^* = \left(\frac{\mu}{\pi} \right)^{N-n/2} + O(e^{-c\mu}) .$$

Proof. This is a direct consequence of (5.8). □

COROLLARY 5.7. For any $\tau \in \mathbb{R}$, consider $\Phi_{z+\tau} : E_{z,\text{sm}} \rightarrow \mathbf{C}^\bullet$. As $\mu \rightarrow +\infty$,

$$\Phi_{z+\tau} \tilde{\Psi}_z \asymp_0 \left(\frac{\mu}{\mu + \tau/2} \right)^{N/2} , \quad \tilde{\Psi}_z \Phi_{z+\tau} \asymp_0 \left(\frac{\mu}{\mu + \tau/2} \right)^{N/2} .$$

Proof. The first relation is a restatement of Theorem 5.3. The second relation follows by conjugating the first one by $\tilde{\Psi}_z$ and using Corollary 5.6. □

COROLLARY 5.8. As $\mu \rightarrow +\infty$, $\tilde{\Psi}_z^{-1} \asymp_0 \Phi_z$ on $E_{z,\text{sm}}$.

Proof. By Corollaries 5.6 and 5.7, on $E_{z,\text{sm}}$,

$$\tilde{\Psi}_z^{-1} \asymp_0 \tilde{\Psi}_z^{-1} \tilde{\Psi}_z \Phi_z = \Phi_z . \quad \square$$

In the rest of this section, we consider $\Phi_z : E_{z,\text{sm}} \rightarrow \mathbf{C}^\bullet$ unless otherwise indicated.

COROLLARY 5.9. As $\mu \rightarrow +\infty$,

$$\Phi_z^* \Phi_z \asymp_0 \left(\frac{\pi}{\mu} \right)^{N-n/2} , \quad \Phi_z \Phi_z^* \asymp_0 \left(\frac{\pi}{\mu} \right)^{N-n/2} .$$

Proof. We show the first relation, the other one being similar. By Corollaries 5.6 and 5.8, on $E_{z,\text{sm}}$,

$$\Phi_z^* \Phi_z \asymp_0 (\tilde{\Psi}_z^{-1})^* \tilde{\Psi}_z^{-1} = (\tilde{\Psi}_z^*)^{-1} \tilde{\Psi}_z^{-1} = (\tilde{\Psi}_z \tilde{\Psi}_z^*)^{-1} \asymp_0 \left(\frac{\pi}{\mu}\right)^{N-n/2}. \quad \square$$

COROLLARY 5.10. As $\mu \rightarrow +\infty$,

$$\tilde{\Psi}_z \asymp_0 \left(\frac{\mu}{\pi}\right)^{N-n/2} \Phi_z^*.$$

Proof. By Corollaries 5.7 and 5.9,

$$\tilde{\Psi}_z \asymp_0 \left(\frac{\mu}{\pi}\right)^{N-n/2} \tilde{\Psi}_z \Phi_z \Phi_z^* \asymp_0 \left(\frac{\mu}{\pi}\right)^{N-n/2} \Phi_z^*. \quad \square$$

COROLLARY 5.11. For every $\tau \in \mathbb{R}$, as $\mu \rightarrow +\infty$,

$$\Phi_{z+\tau} P_{z+\tau,\text{sm}} \tilde{\Psi}_z \asymp_0 \left(\frac{\mu}{\mu + \tau/2}\right)^{N/2} + O(\mu^{-1}).$$

Proof. By Corollaries 5.6, 5.7 and 5.9 and Proposition 3.18,

$$\Phi_{z+\tau} P_{z+\tau,\text{sm}} \tilde{\Psi}_z = \Phi_{z+\tau} (P_{z+\tau,\text{sm}} - P_{z,\text{sm}}) \tilde{\Psi}_z + \Phi_{z+\tau} \tilde{\Psi}_z \asymp_0 O(\mu^{-1}) + \left(\frac{\mu}{\mu + \tau/2}\right)^{N/2}. \quad \square$$

COROLLARY 5.12. As $\mu \rightarrow +\infty$,

$$d_{z,\text{sm}} \asymp_0 \tilde{\Psi}_z \mathbf{d}_z \Phi_z, \quad \delta_{z,\text{sm}} \asymp_0 \tilde{\Psi}_z \delta_z \Phi_z.$$

Proof. By Theorem 3.10 and Corollary 5.7,

$$d_{z,\text{sm}} \asymp_0 \tilde{\Psi}_z \Phi_z d_{z,\text{sm}} = \tilde{\Psi}_z \mathbf{d}_z \Phi_z.$$

Now, taking adjoints and using Corollaries 5.6, 5.9 and 5.10, we obtain

$$\delta_{z,\text{sm}} = \Phi_{z,\text{sm}}^* \delta_z \tilde{\Psi}_z^* \asymp_0 \tilde{\Psi}_z \delta_z \Phi_z. \quad \square$$

The orthogonal projections of \mathbf{C}^\bullet to the subspaces $\Phi_z(\ker \Delta_{z,\text{sm}})$, $\Phi_z(\text{im } d_{z,\text{sm}})$ and $\Phi_z(\text{im } \delta_{z,\text{sm}})$ are denoted by $\tilde{\Pi}_z = \tilde{\Pi}_z^0, \tilde{\Pi}_z^1$ and $\tilde{\Pi}_z^2$, respectively. The inclusion $\Phi_z(\text{im } d_{z,\text{sm}}) \subset \text{im } \mathbf{d}_z$

$$\tilde{\Pi}_z^1 = \tilde{\Pi}_z^1 \Pi_z^1. \quad (5.11)$$

COROLLARY 5.13. For $j = 0, 1, 2$, as $\mu \rightarrow +\infty$,

$$\Phi_z \Pi_{z,\text{sm}}^j \asymp_0 \tilde{\Pi}_z^j \Phi_{z,\text{sm}}, \quad \Pi_{z,\text{sm}}^j \asymp_0 \tilde{\Psi}_z \tilde{\Pi}_z^j \Phi_{z,\text{sm}}, \quad \Pi_{z,\text{sm}}^j \tilde{\Psi}_z \asymp_0 \tilde{\Psi}_z \tilde{\Pi}_{z,\text{sm}}^j.$$

Proof. We only prove the case of $\tilde{\Pi}_z^2$, the other cases being similar. Let $\alpha_{z,1}, \dots, \alpha_{z,p_z}$ be an orthonormal frame of $\delta_z(E_{z,\text{sm}}^{k+1})$. So $\Phi_z \alpha_{z,1}, \dots, \Phi_z \alpha_{z,p_z}$ is a base of $\Phi_z \delta_z(E_{z,\text{sm}}^{k+1})$ for $\mu \gg 0$ by Corollary 5.4. Applying the Gram-Schmidt process to this base, we get an orthonormal base $\mathbf{f}_{z,1}, \dots, \mathbf{f}_{z,p_z}$ of $\Phi_z \delta_z(E_{z,\text{sm}}^{k+1})$. By Corollary 5.9,

$$\langle \Phi_z \alpha_{z,a}, \Phi_z \alpha_{z,b} \rangle \asymp_0 \left(\frac{\pi}{\mu}\right)^{k-n/2} \delta_{ab},$$

for $1 \leq a, b \leq p_z$. So

$$\mathbf{f}_{z,a} \asymp_0 \left(\frac{\mu}{\pi}\right)^{k/2-n/4} \Phi_z \alpha_{z,a}.$$

Hence, by Corollary 5.9, for any $\beta \in E_{z,\text{sm}}^k$,

$$\begin{aligned} \tilde{\Pi}_z^2 \Phi_z \beta &= \sum_{a=1}^{p_z} \langle \Phi_z \beta, \mathbf{f}_{z,a} \rangle \mathbf{f}_{z,a} \asymp_0 \left(\frac{\mu}{\pi} \right)^{k-n/2} \sum_{a=1}^{p_z} \langle \Phi_z \beta, \Phi_z \alpha_{z,a} \rangle \Phi_z \alpha_{z,a} \\ &\asymp_0 \sum_{a=1}^m \langle \beta, \alpha_{z,a} \rangle \Phi_z \alpha_{z,a} = \Phi_z \Pi_{z,\text{sm}}^2 \beta. \end{aligned}$$

This shows the first relation of the statement because $\dim E_{z,\text{sm}}^k < \infty$. Then the other stated relations follow using Corollaries 5.6, 5.7 and 5.9. \square

According to Corollary 5.4, in the following corollaries, we take $\mu \gg 0$ so that $\Phi_z : E_{z,\text{sm}} \rightarrow \mathbf{C}^\bullet$ is an isomorphism.

COROLLARY 5.14. As $\mu \rightarrow +\infty$,

$$(\Phi_z^{-1})^* \Phi_z^{-1} \asymp_0 \left(\frac{\mu}{\pi} \right)^{N-n/2}, \quad \Phi_z^{-1} (\Phi_z^{-1})^* \asymp_0 \left(\frac{\mu}{\pi} \right)^{N-n/2}.$$

Proof. By Corollary 5.9, for $\mathbf{e} \in \mathbf{C}^k$ with $\|\mathbf{e}\| = 1$,

$$\|\Phi_z^{-1} \mathbf{e}\| \asymp_0 \left(\frac{\mu}{\pi} \right)^{k/2-n/4} \|\Phi_z \Phi_z^{-1} \mathbf{e}\| = \left(\frac{\mu}{\pi} \right)^{k/2-n/4},$$

yielding the first stated relation. The second one has a similar proof. \square

COROLLARY 5.15. As $\mu \rightarrow +\infty$,

$$\Phi_z^* \asymp_0 \left(\frac{\pi}{\mu} \right)^{N-n/2} \Phi_z^{-1}, \quad \tilde{\Psi}_z \asymp_0 \Phi_z^{-1}.$$

Proof. By Corollaries 5.9 and 5.14,

$$\Phi_z^* = \Phi_z^* \Phi_z \Phi_z^{-1} \asymp_0 \left(\frac{\pi}{\mu} \right)^{N-n/2} \Phi_z^{-1}, \quad \tilde{\Psi}_z = \tilde{\Psi}_z \Phi_z \Phi_z^{-1} \asymp_0 \Phi_z^{-1}. \quad \square$$

COROLLARY 5.16. We have $\tilde{\Pi}_z^1 = \Pi_z^1$ for $\mu \gg 0$, and $\tilde{\Pi}_z^2 \asymp_0 \Pi_z^2$ as $\mu \rightarrow +\infty$.

Proof. Since $\Phi_z(\text{im } d_{z,\text{sm}}) = \text{im } \mathbf{d}_z$ for $\mu \gg 0$, we get $\tilde{\Pi}_z^1 = \Pi_z^1$.

To prove $\tilde{\Pi}_z^2 \asymp_0 \Pi_z^2$ as $\mu \rightarrow +\infty$, consider the notation of the proof of Corollary 5.13. We have $\alpha_{z,a} = \delta_z \beta_{z,a}$ ($a = 1, \dots, p_z$) for some base $\beta_{z,1}, \dots, \beta_{z,p_z}$ of $\text{im } d_{z,\text{sm},k}$. Hence, by Corollaries 5.7, 5.9 and 5.12,

$$\Phi_z \alpha_{z,a} = \Phi_z \delta_z \beta_{z,a} \asymp_0 \Phi_z \tilde{\Psi}_z \delta_z \Phi_z \beta_{z,a} \asymp_0 \delta_z \Phi_z \beta_{z,a}, \quad (5.12)$$

and $\delta_z \Phi_z \beta_{z,1}, \dots, \delta_z \Phi_z \beta_{z,p_z}$ is a base of $\text{im } \delta_{z,k+1}$. Applying the Gram-Schmidt process to this base, we get an orthonormal base $\mathbf{g}_{z,1}, \dots, \mathbf{g}_{z,p_z}$ of $\text{im } \delta_{z,k+1}$ satisfying $\mathbf{g}_{z,a} \asymp_0 \mathbf{f}_{z,a}$ by (5.12). Then, for any $\mathbf{e} \in \mathbf{C}^k$ with $\|\mathbf{e}\| = 1$,

$$\tilde{\Pi}_z^2 \mathbf{e} = \sum_{a=1}^{p_z} \langle \mathbf{e}, \mathbf{g}_{z,a} \rangle \mathbf{g}_{z,a} \asymp_0 \sum_{a=1}^{p_z} \langle \mathbf{e}, \mathbf{f}_{z,a} \rangle \mathbf{f}_{z,a} = \Pi_z^2 \mathbf{e}. \quad \square$$

COROLLARY 5.17. We have

$$d_{z,\text{sm}} = \Phi_z^{-1} \mathbf{d}_z \Phi_{z,\text{sm}}, \quad d_{z,\text{sm}}^{-1} \Pi_{z,\text{sm}}^1 = \Pi_{z,\text{sm}}^2 \Phi_z^{-1} \mathbf{d}_z^{-1} \Phi_z \Pi_{z,\text{sm}}^1.$$

Proof. The first equality follows like the first relation of Corollary 5.12, using Φ_z^{-1} instead of $\tilde{\Psi}_z$. To prove the second one, take any $\alpha \in \text{im } d_{z,\text{sm}}$. Since

$$d_z \Pi_{z,\text{sm}}^2 \Phi_z^{-1} \mathbf{d}_z^{-1} \Phi_z \alpha = d_z \Phi_z^{-1} \mathbf{d}_z^{-1} \Phi_z \alpha = \Phi_z^{-1} \mathbf{d}_z \mathbf{d}_z^{-1} \Phi_z \alpha = \alpha$$

with $\Pi_{z,\text{sm}}^2 \Phi_z^{-1} \mathbf{d}_z^{-1} \Phi_z \alpha \in \text{im } \delta_{z,\text{sm}}$, we obtain

$$\Pi_{z,\text{sm}}^2 \Phi_z^{-1} \mathbf{d}_z^{-1} \Phi_z \alpha = d_{z,\text{sm}}^{-1} \alpha . \quad \square$$

5.4 Derivatives of some homomorphisms

THEOREM 5.18. As $\mu \rightarrow +\infty$,

$$\partial_z(\Phi_z \Psi_z), \partial_{\bar{z}}(\Phi_z \Psi_z) \asymp_0 \left(\frac{n}{8\mu} - \frac{N}{4\mu} \right) \left(\frac{\pi}{\mu} \right)^{N/2-n/4} .$$

Proof. By (5.9),

$$\partial_z(\Phi_z \Psi_z e_p) = \sum_{q \in \mathcal{X}_k} e_q \left(\int_{\widehat{W}_q^-} \hat{h}_q^- e^{z\hat{h}_q^-} (\hat{l}_q^-)^* P_{z,\text{sm}} e_{p,z} + \int_{\widehat{W}_q^-} e^{z\hat{h}_q^-} (\hat{l}_q^-)^* \partial_z(P_{z,\text{sm}} e_{p,z}) \right) , \quad (5.13)$$

for every $p \in \mathcal{X}_k$ ($k = 0, \dots, n$). We estimate each of these integrals.

Like in the proof of Theorem 5.3, we get, for any $q \neq p$ in \mathcal{X}_k ,

$$\int_{\widehat{W}_p^-} \hat{h}_p^- e^{z\hat{h}_p^-} (\hat{l}_p^-)^* (P_{z,\text{sm}} - 1) e_{p,z} \asymp_0 0 , \quad (5.14)$$

$$\int_{\widehat{W}_q^-} \hat{h}_q^- e^{z\hat{h}_q^-} (\hat{l}_q^-)^* P_{z,\text{sm}} e_{p,z} \asymp_0 0 . \quad (5.15)$$

Moreover, by Proposition 3.1 (iii), (3.9)–(3.12) and (3.33),

$$\begin{aligned} \int_{\widehat{W}_p^-} \hat{h}_p^- e^{z\hat{h}_p^-} (\hat{l}_p^-)^* e_{p,z} &= -\frac{k}{2a_\mu} \left(\int_{-2r}^{2r} \rho(x) e^{-\mu x^2/2} dx \right)^{k-1} \int_{-2r}^{2r} \rho(x) x^2 e^{-\mu x^2/2} dx \\ &= -\frac{k}{4\mu} \left(\frac{\pi}{\mu} \right)^{\frac{k}{2} - \frac{n}{4}} + O(e^{-c\mu}) . \end{aligned} \quad (5.16)$$

On the other hand, by (5.7) and Proposition 3.25 ,

$$\int_{\widehat{W}_q^-} e^{z\hat{h}_q^-} (\hat{l}_q^-)^* \partial_z(P_{z,\text{sm}} e_{p,z} - e_{p,z}) \asymp_0 0 ,$$

for all $q \in \mathcal{X}_k$. In the case $q = p$, by (5.10) and Lemma 3.23,

$$\begin{aligned} \int_{\widehat{W}_p^-} e^{z\hat{h}_p^-} (\hat{l}_p^-)^* \partial_z e_{p,z} &= \left(\frac{n}{8\mu} + O(e^{-c\mu}) \right) \int_{\widehat{W}_p^-} e^{z\hat{h}_p^-} (\hat{l}_p^-)^* e_{p,z} \\ &= \left(\frac{n}{8\mu} + O(e^{-c\mu}) \right) \left(\left(\frac{\pi}{\mu} \right)^{\frac{k}{2} - \frac{n}{4}} + O(e^{-c\mu}) \right) \\ &= \frac{n}{8\mu} \left(\frac{\pi}{\mu} \right)^{\frac{k}{2} - \frac{n}{4}} + O(e^{-c\mu}) . \end{aligned} \quad (5.17)$$

In the case $q \neq p$, using Lemma 3.23 and arguing again like in the proof of Theorem 5.3, we get

$$\int_{\widehat{W}_q^-} e^{z\hat{h}_q^-} (\hat{l}_q^-)^* \partial_z e_{p,z} \asymp_0 0 \quad (\mu \rightarrow +\infty) . \quad (5.18)$$

Now the result for ∂_z follows from (5.13)–(5.16), (5.17) and (5.18).

If we consider $\partial_{\bar{z}}$, the proof has to be modified as follows. In the expression analogous to (5.13), the first term of the right-hand side must be removed. In the analogue of Lemma 3.23, we get $|x_p^-|$ instead of $|x_p^+|^2$ by (3.5) and (3.36). So $\partial_{\bar{z}}(\Psi_z^* \Psi_z)$ has the same final expression as $\partial_z(\Psi_z^* \Psi_z)$ by (5.16). \square

THEOREM 5.19. As $\mu \rightarrow +\infty$,

$$\partial_z((\Psi_z^* \Psi_z)^{\pm 1}), \partial_{\bar{z}}((\Psi_z^* \Psi_z)^{\pm 1}) = O(\mu^{-1/2}) .$$

Proof. We only show the case of ∂_z . Consider $P_{z,\text{sm}} : E_z \rightarrow E_{z,\text{sm}}$, whose adjoint is $P_z : E_{z,\text{sm}} \rightarrow E_z$. Then, since $J_z : \mathbf{C}^\bullet \rightarrow E_z$ is an isometry,

$$\Psi_z^* \Psi_z = (P_{z,\text{sm}} J_z)^* P_{z,\text{sm}} J_z = J_z^{-1} P_z P_{z,\text{sm}} J_z .$$

It follows that, for every $p \in \mathcal{X}_k$ ($k = 0, \dots, n$),

$$\Psi_z^* \Psi_z \mathbf{e}_p = \sum_{q \in \mathcal{X}_k} \langle P_{z,\text{sm}} e_{p,z}, e_{q,z} \rangle \mathbf{e}_q .$$

Therefore

$$\partial_z(\Psi_z^* \Psi_z) \mathbf{e}_p = \sum_{q \in \mathcal{X}_k} (\langle \partial_z(P_{z,\text{sm}}) e_{p,z}, e_{q,z} \rangle + \langle P_{z,\text{sm}} \partial_z(e_{p,z}), e_{q,z} \rangle + \langle P_{z,\text{sm}} e_{p,z}, \partial_z(e_{q,z}) \rangle) \mathbf{e}_q .$$

Then, by Propositions 3.18 and 3.22, Lemma 3.23 and its analogue for $\partial_{\bar{z}}$,

$$\begin{aligned} \partial_z(\Psi_z^* \Psi_z) \mathbf{e}_p &= O(\mu^{-1/2}) + \left(\frac{n}{8\mu} - \frac{1}{2} \langle |x_p^+|^2 e_{p,z}, e_{p,z} \rangle \right) \mathbf{e}_p + O(e^{-c\mu}) \\ &= \left(\frac{n}{8\mu} - \frac{1}{2} \langle |x_p^+|^2 e_{p,z}, e_{p,z} \rangle \right) \mathbf{e}_p + O(\mu^{-1/2}) . \end{aligned}$$

But, by (3.12) and (3.33),

$$\begin{aligned} \langle |x_p^+|^2 e_{p,z}, e_{p,z} \rangle &= \left(\int_{-2r}^{2r} \rho(x)^2 e^{-\mu x^2} dx \right)^{n-1} (n-k) \int_{-2r}^{2r} y^2 \rho(y)^2 e^{-\mu y^2} dy \\ &= \frac{n-k}{2\mu} \left(\frac{\pi}{\mu} \right)^{\frac{n}{2}} + O(e^{-c\mu}) . \end{aligned}$$

Hence

$$\partial_z(\Psi_z^* \Psi_z) \mathbf{e}_p = \left(\frac{n}{8\mu} - \frac{n-k}{4\mu} \left(\frac{\pi}{\mu} \right)^{\frac{n}{2}} \right) \mathbf{e}_p + O(\mu^{-1/2}) = O(\mu^{-1/2}) ,$$

yielding the stated expression for $\partial_z(\Psi_z^* \Psi_z)$.

Now, arguing like in the proof of (3.30) and using (5.8), we get

$$\begin{aligned} \partial_z((\Psi_z^* \Psi_z)^{-1}) &= -(\Psi_z^* \Psi_z)^{-1} \partial_z(\Psi_z^* \Psi_z) (\Psi_z^* \Psi_z)^{-1} \\ &= -(1 + O(e^{-c\mu})) O(\mu^{-1/2}) (1 + O(e^{-c\mu})) = O(\mu^{-1/2}) . \quad \square \end{aligned}$$

6. Asymptotics of the large zeta invariant

6.1 Preliminaries on Quillen metrics

6.1.1 *Case of a finite dimensional complex* All vector spaces considered here are over \mathbb{C} . For a line λ , its dual λ^* is also denoted by λ^{-1} . For a vector space V of finite dimension, let $\det V = \bigwedge^{\dim V} V$. For a graded vector space V^\bullet of finite dimension, let $\det V^\bullet = \bigotimes_k (\det V^k)^{(-1)^k}$.

Now consider a finite dimensional cochain complex (V^\bullet, ∂) , whose cohomology is denoted by $H^\bullet(V)$. Then there is a canonical isomorphism [KM76], [BGS88, Section 1 a)]

$$\det V^\bullet \cong \det H^\bullet(V) . \quad (6.1)$$

Given a Hermitian metric on V^\bullet so that the homogeneous components V^k are orthogonal one another, the corresponding norm $\| \cdot \|_{V^\bullet}$ on V^\bullet induces a metric $\| \cdot \|_{\det V^\bullet}$ on $\det V^\bullet$, which corresponds to a metric $\| \cdot \|_{\det H^\bullet(V)}$ on $\det H^\bullet(V)$ via (6.1).

On the other hand, consider the induced Laplacian, $\square = (\partial + \partial^*)^2 = \partial\partial^* + \partial^*\partial$, whose kernel is a graded vector subspace \mathcal{H}^\bullet . Then finite dimensional Hodge theory gives an isomorphism $H^\bullet(V) \cong \mathcal{H}^\bullet$, which induces an isomorphism

$$\det H^\bullet(V) \cong \det \mathcal{H}^\bullet . \quad (6.2)$$

The restriction of $\| \cdot \|_{V^\bullet}$ to \mathcal{H}^\bullet induces a metric $\| \cdot \|_{\det \mathcal{H}^\bullet}$ on $\det \mathcal{H}^\bullet$, which corresponds to another metric $| \cdot |_{\det H^\bullet(V)}$ on $\det H^\bullet(V)$ via (6.2).

Let \square' denote the restriction $\square : \text{im } \square \rightarrow \text{im } \square$. For $s \in \mathbb{C}$, let

$$\theta(s) = \theta(s, \square) = - \text{Tr}^s(\mathbf{N}(\square')^{-s}) . \quad (6.3)$$

(Do not confuse the superscript “s” of the supertrace with the complex variable s .) This defines a holomorphic function on \mathbb{C} . Then the above metrics on $\det H^\bullet(V)$ satisfy [BGS88, Proposition 1.5], [BZ92, Theorem 1.1], [BZ94, Theorem 1.4]

$$\| \cdot \|_{\det H^\bullet(V)} = | \cdot |_{\det H^\bullet(V)} e^{\theta'(0)/2} . \quad (6.4)$$

If $H^\bullet(V) = 0$, then $\det H^\bullet(V) \equiv \mathbb{C}$ is canonically generated by 1, and we have $\| 1 \|_{\det H^\bullet(V)} = e^{\theta'(0)/2}$. Using the orthogonal projection $\Pi^1 : V \rightarrow \text{im } \partial$, we can write (6.3) as

$$\theta(s) = - \text{Tr}^s((\square')^{-s} \Pi^1) . \quad (6.5)$$

Let $(\tilde{V}^\bullet, \tilde{\partial})$ be another finite dimensional cochain complex, endowed with a Hermitian metric so that the homogeneous components are orthogonal to each other, and let $\phi : (V, \partial) \rightarrow (\tilde{V}^\bullet, \tilde{\partial})$ be an isomorphism of cochain complexes, which may not be unitary. Then (see the proof of [BZ94, Theorem 6.17])

$$\log \left(\frac{\| \cdot \|_{\det H^\bullet(\tilde{V})}}{\| \cdot \|_{\det H^\bullet(V)}} \right)^2 = \text{Tr}^s(\log(\phi^* \phi)) . \quad (6.6)$$

6.1.2 Case of an elliptic complex Some of the concepts of Section 6.1.1 extend to the case where $V^\bullet = C^\infty(M; E^\bullet)$, for some graded Hermitian vector bundle E^\bullet over M , and ∂ is an elliptic differential complex of order one. Then $\det H^\bullet(V)$ is defined because $\dim H^\bullet(V) < \infty$. Moreover Hodge theory for the Laplacian \square gives the isomorphism (6.2). Thus at least the norm $| \cdot |_{\det H^\bullet(V)}$ is defined in this setting. Now the expression (6.3) only defines $\theta(s) = \theta(s, \square)$ when $\Re s > n/2$, but it has a meromorphic extension to \mathbb{C} , denoted in the same way; indeed, (6.3) becomes

$$\theta(s) = \theta(s, \square) = -\zeta(s, \square, \mathbf{Nw}) ,$$

for $\Re s > n/2$, and therefore this equality also holds for the meromorphic extensions. Furthermore $\theta(s)$ is smooth at $s = 0$ [See67], and $\theta'(0)$ can be considered as a renormalized version of the super-trace of the operator $\mathbf{N} \log(\square')$, which is not of trace class. Thus the right-hand side of (6.4) is defined in this way and plays the role of an analytic version of the metric $\| \cdot \|_{\det H^\bullet(V)}$, which is not directly defined. This kind of metrics were introduced by D. Quillen [Qui85] for the case

of the Dolbeault complex. The expression (6.5) also holds in this case for $\Re s \gg 0$; in fact, it becomes

$$\theta(s) = -\zeta(s, \square, \Pi^1 \mathbf{w}) ,$$

where this zeta function can be shown to define a meromorphic function on \mathbb{C} , even though Π^1 is not a differential operator, and this equality holds as meromorphic functions.

6.1.3 Reidemeister, Milnor and Ray-Singer metrics Let F be a flat vector bundle over M , defined by a representation ρ of $\pi_1 M$, and let ∇^F denote its covariant derivative. Consider a smooth triangulation K of M and the corresponding cochain complex $C^\bullet(K, F)$ with coefficients in F , whose cohomology is isomorphic to $H^\bullet(M, F)$ via the quasi-isomorphism

$$\Omega(M; F) \rightarrow C^\bullet(K, F) = C_\bullet(K, F^*)^*$$

defined by integration of differential forms on smooth simplices. Given a Hermitian structure g^F on F , we get an induced metric on $C^\bullet(K, F)$, and the concepts of Section 6.1.1 can be applied. In this case, the left-hand side of (6.4) is called the *Reidemeister metric*, denoted by $\| \cdot \|_{\det H^\bullet(M, F)}^R$.

If $\nabla^F g^F = 0$ (ρ is unitary) and $H^\bullet(M, F) = 0$, then the Reidemeister torsion $\tau_M(\rho)$ is defined using K , and it is a topological invariant of M [Fra35, Rei35, dR50]. Moreover $\tau_M(\rho) = \|1\|_{\det H^\bullet(M, F)}^R$ is the exponential factor of the right-hand side of (6.4) [RS71, Proposition 1.7]. If we only assume $\nabla^F g^F = 0$, then $\| \cdot \|_{\det H^\bullet(M, F)}^R$ is still a topological invariant of M .

Next, given a vector field X on M satisfying (b), consider the complex $(C^\bullet(-X, W^-, F), \mathbf{d}^F)$, whose cohomology is also isomorphic to $H^\bullet(M, F)$ via the quasi-isomorphism

$$\Phi^{-X, F} : \Omega(M, F) \rightarrow C^\bullet(-X, W^-, F) = C_\bullet(-X, W^-, F^*)^* .$$

This complex has a metric induced by g^F , like in Section 5.2.2, and the concepts of Section 6.1.1 can be also applied. In this case, the left-hand side of (6.4) is called the *Milnor metric*, denoted by $\| \cdot \|_{\det H^\bullet(M, F)}^{M, X}$, and the metric factor of the right-hand side of (6.4) is denoted by $\| \cdot \|_{\det H^\bullet(M, F)}^{M, X}$. If $\nabla^F g^F = 0$, then $\| \cdot \|_{\det H^\bullet(M, F)}^{M, X} = \| \cdot \|_{\det H^\bullet(M, F)}^R$ [Mil66, Theorem 9.3].

Finally, the concepts of Section 6.1.2 can be applied to $(\Omega(M, F), d^F)$, whose cohomology is again $H^\bullet(M, F)$. In this case, the right-hand side of (6.4) is called the *Ray-Singer metric*, denoted by $\| \cdot \|_{\det H^\bullet(M, F)}^{\text{RS}}$, and the metric factor of the right-hand side of (6.4) is denoted by $\| \cdot \|_{\det H^\bullet(M, F)}^{\text{RS}}$. If $H^\bullet(M, F) = 0$, then the exponential factor of the right-hand side of (6.4) is called the *analytic torsion* or *Ray-Singer torsion*, denoted by $T_M(\rho)$. These concepts were introduced by Ray and Singer [RS71], who conjectured that $T_M(\rho) = \tau_M(\rho)$ if $\nabla^F g^F = 0$ and $H^\bullet(M, F) = 0$, and $\| \cdot \|_{\det H^\bullet(M, F)}^{\text{RS}} = \| \cdot \|_{\det H^\bullet(M, F)}^R$ assuming only that $\nabla^F g^F = 0$. Independent proofs of this conjecture were given by Cheeger [Che79] and Müller [M78]. Actually, this equality still holds if the induced Hermitian structure $g^{\det F}$ on $\det F$ is flat [M93].

In the case where $g^{\det F}$ is not assumed to be flat, Bismut and Zhang [BZ92, BZ94] extended the above results by introducing an additional term. The first ingredient of this extra term is the 1-form

$$\theta(F, g^F) = \text{tr} \left((g^F)^{-1} \nabla^F g^F \right) , \tag{6.7}$$

which vanishes if and only if $g^{\det F}$ is flat. Moreover $\theta(F, g^F)$ is closed and its cohomology class of $\theta(F, g^F)$ is independent of the choice of g^F [BZ92, Proposition 4.6]; this class measures the obstruction to the existence of a flat Hermitian structure on $\det F$.

Let $e(M, \nabla^M)$ be the representative of the Euler class of M given by the Chern-Weil theory

using g^M , and let $\psi(M, \nabla^M)$ be the current of degree $n - 1$ on TM constructed in [MQ86] (see also [BGS90, Section 3], [BZ92, Section 3], [BH06, Section 2], [BH08, Section 4]). Identify the image of the zero section of TM with M , and identify the conormal bundle of M in TM with T^*M . Let δ_M be the current on TM defined by integration on M , and let $\pi : TM \rightarrow M$ be the vector bundle projection. Since M may not be oriented, the form $e(M, \nabla^M)$ is valued in the orientation line bundle $o(M)$ of M , and the currents are the elements of $\Omega(M, o(M))'$.

PROPOSITION 6.1 Bismut-Zhang [BZ92, Theorem 3.7]. *The following holds:*

- (i) *For any smooth function $\lambda : TM \rightarrow \mathbb{R}^\pm$, under the mapping $v \mapsto \lambda v$, $\psi(M, \nabla^M)$ is changed into $(\pm 1)^n \psi(M, \nabla^M)$.*
- (ii) *The current $\psi(M, \nabla^M)$ is locally integrable, and its wave front set is contained in T^*M (thus $\psi(M, \nabla^M)$ is smooth on $TM \setminus M$).*
- (iii) *The restriction of $-\psi(M, \nabla^M)$ to the fibers of $TM \setminus M$ coincides with the solid angle defined by g^M .*
- (iv) *We have*

$$d\psi(M, \nabla^M) = \pi^*e(M, \nabla^M) - \delta_M .$$

Remark 6.2. In Proposition 6.1, observe that (i) and (iv) are compatible because $e(M, \nabla^M) = 0$ if n is odd. By (ii)–(iv), the restriction of $\psi(M, \nabla^M)$ to $TM \setminus M$ is induced by a smooth differential form on the sphere bundle which transgresses $e(M, \nabla^M)$ (such a differential form was already defined and used in [Che44]).

THEOREM 6.3 Bismut-Zhang [BZ92, Theorem 0.2], [BZ94, Theorem 0.2]. *We have*

$$\log \left(\frac{\| \det_{M, X}^{\text{RS}} H^\bullet(M, F) \|}{\| \det_{M, X} H^\bullet(M, F) \|} \right)^2 = - \int_M \theta(F, g^F) \wedge X^* \psi(M, \nabla^M) .$$

Remark 6.4. By (b), $X = -\text{grad}_{g'} h$ for some Morse function h and some Riemannian metric g' on M , which may not be the given metric g^M . If we fix h , the right-hand side of the equality in Theorem 6.3 is independent of the choice of X satisfying $X = -\text{grad}_{g'} h$ for some g' [BZ92, Proposition 6.1].

Theorem 6.3 will be applied to the case of the flat complex line bundle \mathcal{L}^z with a Hermitian structure $g^{\mathcal{L}^z}$ (Section 2.1.2). By (2.14) and (6.7),

$$\theta(\mathcal{L}^z, g^{\mathcal{L}^z}) = 2\mu\eta . \tag{6.8}$$

6.2 Asymptotics of the large zeta invariant

We prove Theorem 1.1 (ii) here. With the notation of Section 6.1.2, consider the meromorphic function $\theta(s, z) = \theta(s, \Delta_z)$, also defined in (1.4), as well as its components $\theta_{\text{sm/la}}(s, z)$ defined in (1.5). Consider also the current $\psi(M, \nabla^M)$ of degree $n - 1$ on TM (Section 6.1.3). By Proposition 6.1 (i),

$$- \mathbf{z}_{\text{la}}(-\eta) = (-1)^n \mathbf{z}_{\text{la}}(\eta) . \tag{6.9}$$

Notation 6.5. Let \asymp_1 be defined like \asymp_0 in Notation 5.2, using $O(|\mu|^{-1/2})$ instead of $O(e^{-c|\mu|})$.

Take some Morse function h on M such that $Xh < 0$ on $M \setminus \mathcal{X}$, and h is in standard form with respect to X . Then $X = -\text{grad}_{g'} h$ for some Riemannian metric g' (Section 5.1.3), which

may not be the given metric g . Consider the flat complex line bundle $\mathcal{L}_{dh-z\eta}$ with the Hermitian structure $g^{F_{dh-z\eta}}$ (Section 2.1.2). Note that $\mathbf{d}_{-dh}^{\mathcal{L}_{dh-z\eta}} \equiv \mathbf{d}_{-z\eta}$ on $C^\bullet(-X, W^-, \mathcal{L}_{dh-z\eta}) \equiv \mathbf{C}^\bullet(-X)$. So, by (6.8), Theorem 6.3 and Remark 6.4,

$$\log \frac{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS}}}{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{M},X}} = - \int_M (dh - \mu\eta) \wedge X^* \psi(M, \nabla^M), \quad (6.10)$$

where $H_{-z}^\bullet(M) = H_{-z\eta}^\bullet(M)$. With the notation of Section 6.1.3, let

$$\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS,sm}} = \| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS}} e^{\theta'_{\text{sm}}(0, -z)/2}.$$

By (6.4),

$$\log \frac{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS}}}{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{M},X}} = \log \frac{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS,sm}}}{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{M},X}} + \frac{\theta'_{\text{la}}(0, -z)}{2}. \quad (6.11)$$

By (6.6) and Corollary 5.4, for $\mu \gg 0$,

$$\begin{aligned} \log \left(\frac{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS,sm}}}{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{M},X}} \right)^2 &= -\text{Tr}^s(\log(\Phi_{-z}^* \Phi_{-z})) = -\text{Tr}^s(\log(\Psi_{-z}^{-1} \Phi_{-z}^* \Phi_{-z} \Psi_{-z})) \\ &= -\text{Tr}^s(\log((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z})). \end{aligned} \quad (6.12)$$

From (5.8) and Theorems 5.3, 5.18 and 5.19, we obtain

$$((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z})^{-1} = \left(\frac{\pi}{\mu} \right)^{\frac{n}{2} - N} + O(e^{-c\mu})$$

and

$$\begin{aligned} \partial_z &((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z}) \\ &= \partial_z((\Psi_{-z}^* \Psi_{-z})^{-1}) (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z} + (\Psi_{-z}^* \Psi_{-z})^{-1} (\partial_z(\Phi_{-z} \Psi_{-z}))^* \Phi_{-z} \Psi_{-z} \\ &\quad + (\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \partial_z(\Phi_{-z} \Psi_{-z}) \\ &= \left(O(\mu^{-1/2}) + (1 + O(e^{-c\mu})) \left(\frac{n}{4\mu} - \frac{N}{2\mu} \right) \right) \left(\frac{\pi}{\mu} \right)^{N - \frac{n}{2}} + O(e^{-c\mu}) \\ &= \left(O(\mu^{-1/2}) + \left(\frac{n}{4\mu} - \frac{N}{2\mu} \right) \right) \left(\frac{\pi}{\mu} \right)^{N - \frac{n}{2}} + O(e^{-c\mu}). \end{aligned}$$

So

$$\begin{aligned} \partial_z \text{Tr}^s(\log((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z})) \\ &= \text{Tr}^s((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z})^{-1} \partial_z((\Psi_{-z}^* \Psi_{-z})^{-1} (\Phi_{-z} \Psi_{-z})^* \Phi_{-z} \Psi_{-z}) \\ &= O(\mu^{-1/2}) + \text{Tr}^s\left(\frac{n}{4\mu} - \frac{N}{2\mu}\right) + O(e^{-c\mu}) = O(\mu^{-1/2}). \end{aligned}$$

Then, by (6.12),

$$\partial_z \log \frac{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{RS,sm}}}{\| \! \|_{\det H_{-z}^\bullet(M)}^{\text{M},X}} = O(\mu^{-1/2}). \quad (6.13)$$

By taking the derivative with respect to z of both sides of (6.10), and using (6.11), (6.13) and Corollary 4.16, we get $\zeta_{\text{la}}(1, -z) \asymp_1 \mathbf{z}_{\text{la}}$. Then Theorem 1.1 (ii) follows because \mathbf{z}_{la} is independent

of z .

Remark 6.6. In the case where $\eta = dh$, Theorem 1.1 (ii) agrees with Theorem 4.23. In fact, by Proposition 6.1 (iv), Theorem 1.1 (ii) and the Stokes formula,

$$\begin{aligned} \zeta_{\text{la}}(1, z) &\asymp_1 - \int_M h X^* d\psi(M, \nabla^M) = - \int_M h X^*(\pi^* e(M, \nabla^M) - \delta_M) \\ &= - \int_M h e(M, \nabla^M) + \sum_{p \in \mathcal{X}} (-1)^{\text{ind}(p)} h(p). \end{aligned}$$

7. Asymptotics of the small zeta-invariant

7.1 Condition on the integrals along instantons

Let

$$\begin{aligned} \mathcal{M}_p &= \mathcal{M}_p(\eta, X) = - \max\{\eta(\gamma) \mid \gamma \in \mathcal{T}_p^1\} \quad (p \in \mathcal{X}_+), \\ \mathcal{M}_k &= \mathcal{M}_k(\eta, X) = \min_{p \in \mathcal{X}_k} \mathcal{M}_p \quad (k = 1, \dots, n). \end{aligned}$$

Thus (d) means that $\mathcal{M}_p = \mathcal{M}_k$ for all $k = 1, \dots, n$ and $p \in \mathcal{X}_k$. The following result will be proved in Appendix A.

THEOREM 7.1. *For every $\xi \in H^1(M, \mathbb{R})$ and numbers $a_n \geq \dots \geq a_1 \gg 0$ or $a_1 \geq \dots \geq a_n \gg 0$, there is some $\eta \in \xi$, satisfying (a) and (c) with the given X and some metric g , such that $\mathcal{M}_p(\eta, X) = a_k$ for all $k = 1, \dots, n$ and $p \in \mathcal{X}_k$.*

Remark 7.2. If $\xi \neq 0$, for $p \in \mathcal{X}_k$, $q \in \mathcal{X}_{k-1}$ and $\gamma, \delta \in \mathcal{T}(p, q) \subset \mathcal{T}_p^1$, the period $\langle \xi, \bar{\gamma} \bar{\delta}^{-1} \rangle = \eta(\gamma) - \eta(\delta)$ may not be zero. Hence it may not be possible to get $\eta(\gamma) = -a_k$ for all $\gamma \in \mathcal{T}_p^1$, contrary to the case where $\xi = 0$.

From now on, we assume η satisfies (d), besides (a) and (c). By Theorem 7.1, this is possible for any prescription of the class $\xi = [\eta] \in H^1(M, \mathbb{R})$. Let $a_k = \mathcal{M}_k(\eta, X)$ ($k = 1, \dots, n$). Then $-\eta$ also satisfies (a), (c) and (d) with $-X$ and g , and $\mathcal{M}_k(-\eta, -X) = a_{n-k+1}$. So, by Corollary 3.15,

$$- \mathbf{z}_{\text{sm}}(-\eta) = - \sum_{k=1}^n (-1)^k (1 - e^{a_{n-k+1}}) m_k^1. \quad (7.1)$$

LEMMA 7.3. *Suppose M is oriented and n is even. Then*

$$\mathbf{z}_{\text{sm}} = \sum_{k=1}^n (-1)^k e^{a_k} m_k^1.$$

If moreover all numbers a_k are equal one another, then $\mathbf{z}_{\text{sm}} = 0$.

Proof. Use Lemma 3.14 and Corollary 3.16. □

7.2 Asymptotics of the perturbed Morse operators

Consider the notation of Section 5.2.2. By (5.4),

$$\mathbf{d}_{z, k-1} = e^{-a_k z} (\mathbf{d}'_{k-1} + \mathbf{d}''_{z, k-1}), \quad (7.2)$$

for $k = 1, \dots, n$, where

$$\mathbf{d}'_{k-1} \mathbf{e}_q = \sum_{p \in \mathcal{X}_k, \gamma \in \mathcal{T}(p,q), \eta(\gamma) = -a_k} \epsilon(\gamma) \mathbf{e}_p, \quad (7.3)$$

$$\mathbf{d}''_{z,k-1} \mathbf{e}_q = \sum_{p \in \mathcal{X}_k, \gamma \in \mathcal{T}(p,q), \eta(\gamma) < -a_k} e^{z(a_k + \eta(\gamma))} \epsilon(\gamma) \mathbf{e}_p, \quad (7.4)$$

for $q \in \mathcal{X}_{k-1}$. Observe that

$$e^{a_k z} \mathbf{d}_{z,k-1} = \mathbf{d}'_{k-1} + O(e^{-c\mu}) \quad (\mu \rightarrow +\infty). \quad (7.5)$$

So

$$\mathbf{d}'_k \mathbf{d}'_{k-1} = \lim_{\mu \rightarrow +\infty} e^{(a_{k+1} + a_k)z} \mathbf{d}_{z,k} \mathbf{d}_{z,k-1} = 0.$$

Hence the operator $\mathbf{d}' = \sum_k \mathbf{d}'_k$ on \mathbf{C}^\bullet satisfies $(\mathbf{d}')^2 = 0$. Taking adjoints in (7.2)–(7.4), or using (5.5), we also get

$$\boldsymbol{\delta}_{z,k} = e^{-a_k \bar{z}} (\boldsymbol{\delta}'_k + \boldsymbol{\delta}''_{z,k}), \quad (7.6)$$

for $k = 1, \dots, n$, where

$$\boldsymbol{\delta}'_k \mathbf{e}_p = \sum_{q \in \mathcal{X}_{k-1}, \gamma \in \mathcal{T}(p,q), \eta(\gamma) = -a_k} \epsilon(\gamma) \mathbf{e}_q, \quad (7.7)$$

$$\boldsymbol{\delta}''_{z,k} \mathbf{e}_p = \sum_{q \in \mathcal{X}_{k-1}, \gamma \in \mathcal{T}(p,q), \eta(\gamma) = -a_k} e^{\bar{z}(a_k + \eta(\gamma))} \epsilon(\gamma) \mathbf{e}_q, \quad (7.8)$$

for $p \in \mathcal{X}_k$. Moreover (7.5) yields

$$e^{a_k \bar{z}} \boldsymbol{\delta}_{z,k} = \boldsymbol{\delta}'_k + O(e^{-c\mu}) \quad (\mu \rightarrow +\infty). \quad (7.9)$$

Let $\boldsymbol{\delta}' = \sum_k \boldsymbol{\delta}'_k = (\mathbf{d}')^*$, and let

$$\mathbf{D}' = \mathbf{d}' + \boldsymbol{\delta}', \quad \boldsymbol{\Delta}' = (\mathbf{D}')^2 = \mathbf{d}' \boldsymbol{\delta}' + \boldsymbol{\delta}' \mathbf{d}'.$$

We have

$$\begin{aligned} \mathbf{C}^\bullet &= \ker \boldsymbol{\Delta}' \oplus \operatorname{im} \mathbf{d}' \oplus \operatorname{im} \boldsymbol{\delta}', \\ \operatorname{im} \boldsymbol{\Delta}' &= \operatorname{im} \mathbf{D}' = \operatorname{im} \mathbf{d}' \oplus \operatorname{im} \boldsymbol{\delta}', \quad \ker \boldsymbol{\Delta}' = \ker \mathbf{D}' = \ker \mathbf{d}' \cap \ker \boldsymbol{\delta}'. \end{aligned}$$

The orthogonal projections of \mathbf{C}^\bullet to $\ker \boldsymbol{\Delta}'$, $\operatorname{im} \mathbf{d}'$ and $\operatorname{im} \boldsymbol{\delta}'$ are denoted by $\boldsymbol{\Pi}' = \boldsymbol{\Pi}'^0$, $\boldsymbol{\Pi}'^1$ and $\boldsymbol{\Pi}'^2$, respectively. Like in Sections 2.1.2 and 5.2.4, the composition $(\mathbf{d}')^{-1} \boldsymbol{\Pi}'^1$ is defined on \mathbf{C}^\bullet . From (7.5) and (7.9), we easily get that, as $\mu \rightarrow +\infty$,

$$\boldsymbol{\Pi}'^j_{z,k} = \boldsymbol{\Pi}'^j_k + O(e^{-c\mu}) \quad (j = 0, 1, 2), \quad (7.10)$$

$$e^{-a_k z} (\mathbf{d}_{z,k-1})^{-1} \boldsymbol{\Pi}'^1_{z,k} = (\mathbf{d}'_{k-1})^{-1} \boldsymbol{\Pi}'^1_k + O(e^{-c\mu}). \quad (7.11)$$

By (7.5) and (7.9), on $\operatorname{im} \boldsymbol{\delta}_{z,k} + \operatorname{im} \mathbf{d}_{z,k-1}$,

$$\boldsymbol{\Delta}_z = e^{-2a_k \mu} \boldsymbol{\Delta}' + O(e^{-(2a_k + c)\mu}) \quad (\mu \rightarrow +\infty). \quad (7.12)$$

PROPOSITION 7.4. *For $k = 0, \dots, n$ and $\mu \gg 0$, the spectrum of $\boldsymbol{\Delta}_z$ on $\operatorname{im} \boldsymbol{\delta}_{z,k} + \operatorname{im} \mathbf{d}_{z,k-1}$ is contained in an interval of the form*

$$[C e^{-2a_k \mu}, C' e^{-2a_k \mu}] \quad (C' \geq C).$$

Proof. The positive eigenvalues of $\boldsymbol{\Delta}'$ are contained in an interval $[C_0, C'_0]$ ($C'_0 \geq C_0 > 0$).

By (7.12), for $\mu \gg 0$ and $\mathbf{e} \in \text{im } \delta_{z,k} + \text{im } \mathbf{d}_{z,k-1}$,

$$\begin{aligned} \langle \Delta_z \mathbf{e}, \mathbf{e} \rangle &\geq e^{2a_k \mu} \langle \Delta' \mathbf{e}, \mathbf{e} \rangle - C_1 e^{-(2a_k+c)\mu} \|\mathbf{e}\|^2 \geq (C_0 e^{-2a_k \mu} - C_1 e^{-(2a_k+c)\mu}) \|\mathbf{e}\|^2, \\ \langle \Delta_z \mathbf{e}, \mathbf{e} \rangle &\leq e^{2a_k \mu} \langle \Delta' \mathbf{e}, \mathbf{e} \rangle + C_1 e^{-(2a_k+c)\mu} \|\mathbf{e}\|^2 \leq (C'_0 e^{-2a_k \mu} + C_1 e^{-(2a_k+c)\mu}) \|\mathbf{e}\|^2. \end{aligned}$$

Then result follows taking $0 < C < C_0$ and $C' > C'_0$. \square

7.3 Estimates of the nonzero small spectrum

THEOREM 7.5. *If $\mu \gg 0$, the spectrum of $\Delta_{z,\text{sm}}$ on $\text{im } \delta_{z,\text{sm},k} + \text{im } d_{z,\text{sm},k-1}$ is contained in an interval of the form*

$$[C\mu e^{-2a_k \mu}, C'\mu e^{-2a_k \mu}] \quad (C' \geq C).$$

Proof. By the commutativity of (2.7), for every eigenvalue λ of $\Delta_{z,\text{sm}}$ on $\text{im } \delta_{z,\text{sm},k} + \text{im } d_{z,\text{sm},k-1}$, there are normalized λ -eigenforms, $e \in \text{im } \delta_{z,\text{sm},k}$ and $e' \in \text{im } d_{z,\text{sm},k-1}$, so that $d_z e = \lambda^{1/2} e'$ and $\delta_z e' = \lambda^{1/2} e$. Hence the maximum and minimum of the spectrum of $\Delta_{z,\text{sm}}$ on $\text{im } \delta_{z,\text{sm},k} + \text{im } d_{z,\text{sm},k-1}$ is $\|d_{z,\text{sm},k-1}\|^2$ and $\|d_{z,\text{sm},k-1}^{-1} \Pi_{z,\text{sm},k}^1\|^{-2}$, respectively. Similarly, the maximum and minimum of the spectrum of Δ_z on $\text{im } \delta_{z,k} + \text{im } \mathbf{d}_{z,k-1}$ is $\|\mathbf{d}_{z,k-1}\|^2$ and $\|\mathbf{d}_{z,k-1}^{-1} \mathbf{\Pi}_{z,k}^1\|^{-2}$, respectively. Then the result follows from Corollaries 5.9, 5.14 and 5.17 and Proposition 7.4:

$$\begin{aligned} \|d_{z,\text{sm},k-1}\|^2 &\leq \|\Phi_{z,k}^{-1}\|^2 \|\mathbf{d}_{z,k-1}\|^{-2} \|\Phi_{z,\text{sm},k-1}\|^{-2} \\ &\leq \left(\left(\frac{\mu}{\pi} \right)^{k-n/2} + O(e^{-c\mu}) \right) C'_0 e^{-2a_k \mu} \left(\left(\frac{\pi}{\mu} \right)^{k-1-n/2} + O(e^{-c\mu}) \right) \\ &\leq C' \mu e^{-2a_k \mu}, \\ \|d_{z,\text{sm},k-1}^{-1} \Pi_{z,\text{sm},k-1}^1\|^{-2} &\geq \|\Phi_{z,k-1}^{-1}\|^{-2} \|\mathbf{d}_{z,k-1}^{-1} \mathbf{\Pi}_{z,k}^1\|^{-2} \|\Phi_{z,\text{sm},k}\|^{-2} \\ &\geq \left(\left(\frac{\pi}{\mu} \right)^{k-1-n/2} + O(e^{-c\mu}) \right) C_0 e^{-2a_k \mu} \left(\left(\frac{\mu}{\pi} \right)^{k-n/2} + O(e^{-c\mu}) \right) \\ &\geq C \mu e^{-2a_k \mu}. \quad \square \end{aligned}$$

7.4 Asymptotics of the small zeta invariant

Theorem 1.1 (iii) is proved here.

Notation 7.6. Let \asymp_2 be defined like \asymp_0 in Notation 5.2, using $O(|\mu|^{-1})$ instead of $O(e^{-c|\mu|})$.

THEOREM 7.7. *As $\mu \rightarrow +\infty$,*

$$\eta \wedge d_z^{-1} \Pi_{z,\text{sm},k}^1 \asymp_2 \pm (1 - e^{a_k}) \Pi_{z,\text{sm},k}^1.$$

Proof. Consider the notation of Sections 5.3 and 7.2. By Corollary 5.13 and (5.11),

$$\Pi_{z,\text{sm}}^1 \asymp_0 \tilde{\Psi}_z \tilde{\Pi}_z^1 \Phi_{z,\text{sm}} = \tilde{\Psi}_z \tilde{\Pi}_z^1 \Pi_z^1 \Phi_{z,\text{sm}} = \tilde{\Psi}_z \Pi_z^1 \Phi_{z,\text{sm}} - \tilde{\Psi}_z (\tilde{\Pi}_z^1)^\perp \Pi_z^1 \Phi_{z,\text{sm}}.$$

But $(\tilde{\Pi}_z^1)^\perp \Pi_z^1 = 0$ if $\mu \gg 0$ by Corollary 5.16. Hence

$$\Pi_{z,\text{sm}}^1 \asymp_0 \tilde{\Psi}_z \Pi_z^1 \Phi_{z,\text{sm}}. \quad (7.13)$$

For brevity, let $R_z = P_{z-1,\text{sm}} P_{z,\text{sm}}$ on $L^2(M; \Lambda)$, and $S_z = \Phi_z \tilde{\Psi}_{z-1}$ and $T_z = \Phi_{z-1} P_{z-1,\text{sm}} \tilde{\Psi}_z$ on \mathbf{C}^\bullet . By (7.5)–(7.11), (7.13), Propositions 3.18 and 7.4, Corollaries 3.19, 5.6, 5.7, 5.9, 5.11,

5.13, 5.15 and 5.17, and Theorem 7.5,

$$\begin{aligned}
 e^{a_k} \Pi_{z,\text{sm},k}^1 &\asymp_0 e^{a_k} \tilde{\Psi}_z \Pi_{z,k}^1 \Phi_{z,\text{sm}} \asymp_2 e^{a_k} \tilde{\Psi}_z \Pi_k'^1 \Phi_{z,\text{sm}} = e^{a_k} \tilde{\Psi}_z \mathbf{d}'_{k-1} (\mathbf{d}'_{k-1})^{-1} \Pi_k'^1 \Phi_{z,\text{sm}} \\
 &\asymp_2 e^{a_k} R_z \tilde{\Psi}_z S_z \mathbf{d}'_{k-1} T_z (\mathbf{d}'_{k-1})^{-1} \Pi_k'^1 \Phi_{z,\text{sm}} \\
 &\asymp_2 e^{a_k} R_z \tilde{\Psi}_{z-1} \mathbf{d}'_{k-1} T_z (\mathbf{d}'_{k-1})^{-1} \Pi_k'^1 \Phi_{z,\text{sm}} \\
 &\asymp_2 e^{a_k} \tilde{\Psi}_{z-1} \mathbf{d}'_{k-1} T_z (\mathbf{d}'_{k-1})^{-1} \Pi_k'^1 \Phi_{z,\text{sm}} \\
 &\asymp_2 e^{a_k} \tilde{\Psi}_{z-1} e^{a_k(z-1)} \mathbf{d}_{z-1,k-1} T_z e^{-a_k z} \mathbf{d}_{z,k-1}^{-1} \Pi_{z,k}^1 \Phi_{z,\text{sm}} \\
 &= \tilde{\Psi}_{z-1} \mathbf{d}_{z-1,k-1} T_z \mathbf{d}_{z,k-1}^{-1} \Pi_{z,k}^1 \Phi_{z,\text{sm}} \\
 &\asymp_0 \tilde{\Psi}_{z-1} \mathbf{d}_{z-1,k-1} T_z \tilde{\Pi}_{z,k-1}^2 \mathbf{d}_{z,k-1}^{-1} \tilde{\Pi}_{z,k}^1 \Phi_{z,\text{sm}} \\
 &\asymp_0 \tilde{\Psi}_{z-1} \mathbf{d}_{z-1,k-1} \Phi_{z-1} P_{z-1,\text{sm}} \Pi_{z,k-1}^2 \tilde{\Psi}_z \mathbf{d}_{z,k-1}^{-1} \Phi_z \Pi_{z,\text{sm}}^1 \\
 &\asymp_0 \tilde{\Psi}_{z-1} \mathbf{d}_{z-1,k-1} \Phi_{z-1} P_{z-1,\text{sm}} \Pi_{z,k-1}^2 \Phi_z^{-1} \mathbf{d}_{z,k-1}^{-1} \Phi_z \Pi_{z,\text{sm}}^1 \\
 &= \tilde{\Psi}_{z-1} \Phi_{z-1} d_{z-1,\text{sm},k-1} d_{z,\text{sm},k-1}^{-1} \Pi_{z,\text{sm},k}^1 \asymp_0 d_{z-1} d_z^{-1} \Pi_{z,\text{sm},k}^1 .
 \end{aligned}$$

Therefore

$$\eta \wedge d_z^{-1} \Pi_{z,\text{sm},k}^1 = (d_z - d_{z-1}) d_z^{-1} \Pi_{z,\text{sm},k}^1 \asymp_2 (1 - e^{a_k}) \Pi_{z,\text{sm},k}^1 . \quad \square$$

Theorem 1.1 (iii) follows from Corollaries 3.9 and 4.15 and Theorem 7.7.

Remark 7.8. Theorem 1.1 (iii) agrees with Corollaries 4.20 to 4.22 by (7.1) and Lemma 7.3.

8. Prescription of the asymptotics of the zeta invariant

We prove Theorem 1.2 here. By Theorem 7.1, given $a \gg 0$, there is some $\eta_0 \in \xi$ and some metric g satisfying (a) and (d) with the given X , and so that $\mathcal{M}_k(\eta_0, X) = a$ for all $k = 1, \dots, n$. Using the notation of Section 3.1, we are going to modify η_0 only in every U_p for $p \in \mathcal{X}_0 \cup \mathcal{X}_n$.

Fix any $\epsilon > 0$ such that, for every $p \in \mathcal{X}_0 \cup \mathcal{X}_n$, the open ball $B(p, 3\epsilon)$ is contained in U_p . Let

$$V = \bigcup_{p \in \mathcal{X}_0 \cup \mathcal{X}_n} B(p, \epsilon) , \quad V' = \bigcup_{p \in \mathcal{X}_0 \cup \mathcal{X}_n} B(p, 2\epsilon) .$$

Take a smooth function $\sigma : [0, 3\epsilon] \rightarrow [0, 1]$ so that

$$\sigma' \leq 0 , \quad \sigma([0, \epsilon]) = 1 , \quad \sigma([2\epsilon, 3\epsilon]) = 0 .$$

Let $f_j \in C^\infty(M, \mathbb{R})$ ($j = 0, n$) be the extension by zero of the combination of the functions $\sigma(|x_p|) \in C_c^\infty(B(p, 3\epsilon), \mathbb{R})$ ($p \in \mathcal{X}_j$). We have

$$df_j \subset V_j' \setminus V_j , \quad f_j(V_j) = 1 , \quad f_j(M \setminus V_j') = 0 , \quad Xf_0 \geq 0 , \quad Xf_n \leq 0 .$$

For any $c_0, c_n \geq 0$, let $\eta = \eta(c_0, c_n) = \eta_0 - c_0 df_0 + c_n df_n$. This closed 1-form satisfies (a) and (d) with X and g , and we have

$$\mathcal{M}_1(\eta, X) = a + c_0 , \quad \mathcal{M}_n(\eta, X) = a + c_n , \quad \mathcal{M}_k(\eta, X) = a \quad (1 < k < n) .$$

Hence, by Corollary 3.15,

$$\mathbf{z}_{\text{sm}}(\eta) - \mathbf{z}_{\text{sm}}(\eta_0) = e^a (e^{c_0} - 1) m_1^1 + (-1)^n e^a (1 - e^{c_n}) m_n^1 . \quad (8.1)$$

By (a), $e(M, \nabla^M) = 0$ on every U_p ($p \in \mathcal{X}$). So, using the Stokes formula,

$$\begin{aligned}
 \mathbf{z}_{\text{la}}(\eta) - \mathbf{z}_{\text{la}}(\eta_0) &= \int_M (c_0 df_0 - c_n df_n) \wedge X^* \psi(M, \nabla^M) \\
 &= \int_M (c_n f_n - c_0 f_0) X^* d\psi(M, \nabla^M) \\
 &= \int_M (c_n f_n - c_0 f_0) e(M, \nabla^M) - \sum_{p \in \mathcal{X}} (-1)^{\text{ind}(p)} (c_n f_n - c_0 f_0)(p) \\
 &= c_0 |\mathcal{X}_0| - (-1)^n c_n |\mathcal{X}_n|, \tag{8.2}
 \end{aligned}$$

Combining (8.1) and (8.2), we obtain

$$\mathbf{z}(\eta) - \mathbf{z}(\eta_0) = e^a (e^{c_0} - 1) m_1^1 + (-1)^n e^a (1 - e^{c_n}) m_n^1 + c_0 |\mathcal{X}_0| - (-1)^n c_n |\mathcal{X}_n|. \tag{8.3}$$

So, if n is even (respectively, odd), given any $\tau \in \mathbb{R}$ (respectively, $\tau \gg 0$), we get $\mathbf{z}(\eta(c_0, c_n)) = \tau$ for some $c_0, c_n \geq 0$ since $|\mathcal{X}_0|, |\mathcal{X}_n| > 0$ by (b).

Now assume M is oriented and n is even. Then $\mathbf{z}(\eta_0) = -\mathbf{z}(-\eta_0)$ by (6.9) and Lemma 7.3. Using local changes of X and applying [Sma61, Lemmas 1.1 and 1.2], we can increase $|\mathcal{X}_0|$ or $|\mathcal{X}_n|$ as much as desired. So we can assume $|\mathcal{X}_0| - |\mathcal{X}_n| \in (\tau - \mathbf{z}(\eta_0)) \cdot \mathbb{R}^+$ and $m_0, m_n > 0$ by Lemma 3.12, yielding $m_1^1, m_n^1 > 0$ by (3.22). Moreover $m_1^1 = m_n^1$ by (3.22) and Lemma 3.14. Thus, taking $c_0 = c_n =: c$, the expression (8.3) becomes

$$\mathbf{z}(\eta) - \mathbf{z}(\eta_0) = c(|\mathcal{X}_0| - |\mathcal{X}_n|).$$

Hence $\mathbf{z}(\eta) = \tau$ for some $c \geq 0$.

9. The switch of the order of integration

The proof of Theorem 1.3 is given in this section. Let \mathcal{S} be the Schwartz space on \mathbb{R} . Recall that the space of tempered distributions is the continuous dual space \mathcal{S}' , with the topology of uniform convergence on bounded sets. Suppose first that (1.7) is used as definition of Z_μ . By Theorem 1.1, the expression (1.7) defines a tempered distribution Z_μ for $\mu \gg 0$. Moreover, using also the formula of the inverse Fourier transform, we get, for $f \in \mathcal{S}$,

$$\langle Z_\mu, f \rangle = \frac{1}{2\pi} \int_{-\infty}^{\infty} \zeta(1, z) \hat{f}(\nu) d\nu \rightarrow \frac{\mathbf{z}}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\nu) d\nu = \mathbf{z}f(0),$$

as $\mu \rightarrow +\infty$, uniformly on ν . For every $C > 0$, this convergence is also uniform on $f \in \mathcal{S}$ with $|\hat{f}(\nu)|, |\nu^2 \hat{f}(\nu)| \leq C$. So $Z_\mu \rightarrow \mathbf{z}\delta_0$ in \mathcal{S}' as $\mu \rightarrow +\infty$. To get Theorem 1.3, it only remains to prove the following.

THEOREM 9.1. *Both (1.3) and (1.7) define the same tempered distribution Z_μ for $\mu \gg 0$.*

PROPOSITION 9.2. *For $\mu \gg 0, t > 0$ and $f \in \mathcal{S}$,*

$$\int_{-\infty}^{\infty} \int_t^{\infty} |\text{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z})| |\hat{f}(\nu)| du d\nu < \infty.$$

Proof. By [DS88b, Corollary XI.9.8 and Lemma XI.9.9 (d)],

$$\begin{aligned}
 |\text{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z})| &\leq |\eta \wedge \delta_z e^{-u\Delta_z}|_1 \leq \|\eta \wedge\| |\delta_z e^{-u\Delta_z}|_1 \\
 &= \|\eta\|_{L^\infty} \text{Tr}((d_z \delta_z)^{1/2} e^{-u\Delta_z}) \leq \|\eta\|_{L^\infty} \text{Tr}(\Delta_z^{1/2} e^{-u\Delta_z}),
 \end{aligned}$$

where $|\cdot|_1$ denotes the trace norm. Hence

$$\int_t^\infty |\mathrm{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z})| du \leq \|\eta\|_{L^\infty} \int_t^\infty \mathrm{Tr}(\Delta_z^{1/2} e^{-u\Delta_z}) du = \|\eta\|_{L^\infty} \mathrm{Tr}(\Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp).$$

The operator $(I + D^2)^{-N}$ is of trace class for any $N > n$. Therefore

$$\mathrm{Tr}(\Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp) \leq |(I + D^2)^{-N}|_1 \|(I + D^2)^N \Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp\|.$$

By Corollary 2.3 and Theorem 7.5, for $\mu \gg 0$ and $\alpha \in L^2(M; \Lambda)$,

$$\begin{aligned} & \|(I + D^2)^N \Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp \alpha\| \\ & \leq C_0 \|\Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp \alpha\|_{2N} \leq C_1 |z|^{2N} \|\Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp \alpha\|_{2N, z} \\ & = C_2 |z|^{2N} \sum_{k=0}^{2N} \|D_z^k \Delta_z^{-1/2} e^{-t\Delta_z} \Pi_z^\perp \alpha\| \leq C_3 |z|^{2N} \sum_{k=0}^{2N} \frac{1}{t^{k/2}} \|\Delta_z^{-1/2} \Pi_z^\perp \alpha\| \\ & \leq C |z|^{2N} (1 + t^{-N}) e^{c\mu} \|\alpha\|. \end{aligned}$$

Thus, since $f \in \mathcal{S}$,

$$\begin{aligned} & \int_{-\infty}^\infty \int_t^\infty |\mathrm{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z})| |\hat{f}(\nu)| du d\nu \\ & \leq C \|\eta\|_{L^\infty} |(I + D^2)^{-N}|_1 (1 + t^{-N}) e^{c\mu} \int_{-\infty}^\infty |z|^{2N} |\hat{f}(\nu)| d\nu < \infty. \quad \square \end{aligned}$$

Proof of Theorem 9.1. Using Theorem 1.1 and Proposition 9.2 to apply the Lebesgue's dominated convergence theorem and Fubini's theorem, we compute

$$\begin{aligned} -\frac{1}{2\pi} \int_{-\infty}^\infty \lim_{t \downarrow 0} \mathrm{Tr}^s(\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1) \hat{f}(\nu) d\nu &= -\frac{1}{2\pi} \lim_{t \downarrow 0} \int_{-\infty}^\infty \mathrm{Tr}^s(\eta \wedge d_z^{-1} e^{-t\Delta_z} \Pi_z^1) \hat{f}(\nu) d\nu \\ &= \frac{1}{2\pi} \lim_{t \downarrow 0} \int_{-\infty}^\infty \int_t^\infty \mathrm{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z}) \hat{f}(\nu) du d\nu \\ &= \frac{1}{2\pi} \lim_{t \downarrow 0} \int_t^\infty \int_{-\infty}^\infty \mathrm{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z}) \hat{f}(\nu) d\nu du \\ &= \frac{1}{2\pi} \int_0^\infty \int_{-\infty}^\infty \mathrm{Tr}^s(\eta \wedge \delta_z e^{-u\Delta_z}) \hat{f}(\nu) d\nu du. \quad \square \end{aligned}$$

Appendix A. Integrals along instantons

Theorem 7.1 is proved here. We show the case where $a_n \geq \dots \geq a_1 \gg 0$. Then the case where $a_1 \geq \dots \geq a_n \gg 0$ follows by using $-X$ and $-\xi$.

By [Sma61, Theorem B], there is some Morse function h on M such that $h(\mathcal{X}_k) = \{k\}$ ($k = 0, \dots, n$), $Xh < 0$ on $M \setminus \mathcal{X}$, and h is in standard form with respect to X ; in particular, $\mathrm{Crit}_k(h) = \mathcal{X}_k$. Now we proceed like in the proof of [BH04, Proposition 16 (i)]. Since \mathcal{X} is finite, there is some $\eta' \in \xi$ such that $\eta' = 0$ on some open neighborhood U_p of every $p \in \mathcal{X}$. Let $U_k = \bigcup_{p \in \mathcal{X}_k} U_p$ and $U = \bigcup_k U_k$. We can assume $h(U_k) \subset (k - 1/4, k + 1/4)$ for all $k = 0, \dots, n$. If $C \gg 0$, then the representative $\eta'' := \eta' + C dh$ of ξ satisfies $\eta''(X) < 0$ on $M \setminus \mathcal{X}$.

For $k = 0, \dots, n$, let $I_k^\pm \subset \mathbb{R}$ be the closed interval with boundary points $k \pm 1/4$ and $k \pm 1/2$. Since there are no critical values of h in I_k^\pm , every $T_k^\pm := h^{-1}(I_k^\pm)$ is compact submanifold with boundary of dimension n , every $\Sigma_k^\pm := h^{-1}(k \pm 1/2)$ is a closed submanifold of codimension 1,




FIGURE 1. A representation of the sets T_k^\pm , Σ_k^\pm , T_{k-1}^+ and T_{k+1}^- , taking $\mathcal{X}_k = \{p\}$.

and there are identities $T_k^\pm \equiv \Sigma_k^\pm \times I_k^\pm$ given by $x \equiv (\pi_k^\pm(x), h(x))$ ($x \in T_k^\pm$), where $\pi_k^\pm(x)$ is the unique point of Σ_k^\pm that meets the ϕ -orbit of x . Of course, $\Sigma_k^- = \Sigma_{k-1}^+$ ($k = 1, \dots, n$) and $T_0^- = \Sigma_0^- = T_n^+ = \Sigma_n^+ = \emptyset$. (See Figure 1.)

We have $\Sigma_k^\pm \cap \iota_p^\pm(W_p^\pm)$ for $p \in \mathcal{X}_k$. Let $K_p^\pm = \Sigma_k^\pm \cap \iota_p^\pm(W_p^\pm)$ and $K_k^\pm = \bigcup_{p \in \mathcal{X}_k} K_p^\pm$, which are closed submanifolds of Σ_k^\pm ; K_k^- is of codimension k in Σ_k^- , and K_k^+ of codimension $n - k$ in Σ_k^+ . Since the α - and ω -limits of the orbits of X are zero points, the orbit of ϕ through every point $x \in \Sigma_k^+ \setminus K_k^+$ meets $\Sigma_k^- \setminus K_k^-$ at a unique point $\psi_k(x) := \phi^{\tau_k(x)}(x)$ ($\tau_k(x) > 0$). This defines a diffeomorphism $\psi_k : \Sigma_k^+ \setminus K_k^+ \rightarrow \Sigma_k^- \setminus K_k^-$ and a smooth function $\tau_k : \Sigma_k^+ \setminus K_k^+ \rightarrow \mathbb{R}^+$. Moreover the sets K_p^\pm ($p \in \mathcal{X}_k$) have corresponding open neighborhoods V_p^\pm in Σ_k^\pm , with disjoint closures, such that $\psi_k(V_p^+ \setminus K_p^+) = V_p^- \setminus K_p^-$. Take smooth functions λ_p^\pm ($p \in \mathcal{X}_k$) on Σ_k^\pm so that $0 \leq \lambda_p^\pm \leq 1$, $\text{supp } \lambda_p^\pm \subset V_p^\pm$, $\lambda_p^\pm = 1$ on K_p^\pm , and $\lambda_p^+ = \psi_k^* \lambda_p^-$ on $\Sigma_k^+ \setminus K_k^+$. Moreover let

$$\begin{aligned} \tilde{T}_k &= h^{-1}([k - 1/2, k + 1/2]), \quad \tilde{K}_p = \tilde{T}_k \cap (\iota_p^+(W_p^+) \cup \iota_p^-(W_p^-)), \\ \tilde{V}_p &= \{ \phi^t(x) \mid x \in V_p^+ \setminus K_p^+, 0 \leq t \leq \tau_k(x) \} \cup \tilde{K}_p, \\ \tilde{K}_k &= \bigcup_{p \in \mathcal{X}_k} \tilde{K}_p, \quad \tilde{V}_k = \bigcup_{p \in \mathcal{X}_k} \tilde{V}_p, \quad M_k = h^{-1}((-\infty, k + 1/2]). \end{aligned}$$

Thus $M_k = \tilde{T}_0 \cup \dots \cup \tilde{T}_k$. Note that \tilde{T}_k and M_k are compact submanifolds with boundary of dimension n , and every \tilde{V}_p (respectively, \tilde{K}_p) is open (respectively, closed) in \tilde{T}_k . We also get smooth functions $\tilde{\lambda}_p$ ($p \in \mathcal{X}_k$) on \tilde{T}_k determined by the condition $\tilde{\lambda}_p(\phi^t(x)) = \lambda_p^+(x)$ for all $x \in \Sigma_k^+ \setminus K_k^+$ and $0 \leq t \leq \tau_k(x)$. They satisfy $0 \leq \tilde{\lambda}_p \leq 1$, $\text{supp } \tilde{\lambda}_p \subset \tilde{V}_p$, and $\tilde{\lambda}_p = 1$ on \tilde{K}_p .

Let

$$A_p = \max\{|\eta'(\gamma)| \mid \gamma \in \mathcal{T}_p^1\} \quad (p \in \mathcal{X}_+),$$

$$A_k = \max_{p \in \mathcal{X}_k} A_p \quad (k = 1, \dots, n), \quad A = \max\{A_1, \dots, A_n\}.$$

We can suppose $C > A$ and $a_1 > C + A > 0$. For $p \in \mathcal{X}_k$, $q \in \mathcal{X}_{k-1}$ and $\gamma \in \mathcal{T}(p, q)$, we have

$$dh(\gamma) = h(q) - h(p) = -1.$$

Therefore

$$0 > \eta''(\gamma) = \eta'(\gamma) + C dh(\gamma) \geq -A - C > -a_1 \quad (\gamma \in \mathcal{T}^1). \quad (\text{A.1})$$

Claim 1. For $k = 0, \dots, n$, there is a smooth function f_k on M such that

$$df_k(X) \leq 0, \quad (\text{A.2})$$

$$\text{supp } df_k \subset \overset{\circ}{M}_k, \quad (\text{A.3})$$

$$\max\{(\eta'' + df_k)(\gamma) \mid \gamma \in \mathcal{T}_p^1\} = -a_l \quad (p \in \mathcal{X}_l, 1 \leq l \leq k), \quad (\text{A.4})$$

$$(\eta'' + df_k)(\delta) > -a_k \quad (\delta \in \mathcal{T}_{k+1}^1). \quad (\text{A.5})$$

The statement follows directly from Claim 1 taking $\eta = \eta'' + df_n$. So we only have to prove this assertion.

We proceed by induction on k . For $k = 0$, we choose $f_0 = 0$. Then (A.4) is vacuous, (A.2) and (A.3) are trivial, and (A.5) is given by (A.1).

Now take any $k \geq 1$ and assume f_{k-1} is defined and satisfies (A.2)–(A.5). Let

$$b_p = -\max\{(\eta'' + df_{k-1})(\gamma) \mid \gamma \in \mathcal{T}_p^1\} \quad (p \in \mathcal{X}_k), \quad (\text{A.6})$$

$$b_k = \min\{b_p \mid p \in \mathcal{X}_k\}.$$

For every $p \in \mathcal{X}_k$, we have $b_p < a_{k-1} \leq a_k$ because f_{k-1} satisfies (A.5). So there is a smooth function h_p^- on I_k^- such that $(h_p^-)' \geq 0$, $h_p^- = 0$ around $k - 1/2$, and $h_p^- = a_k - b_p$ around $k - 1/4$. Let \tilde{h}_p^- be the function on $V_p^- \times I_k^- \subset \Sigma_k^- \times I_k^- \equiv T_k^-$ given by $\tilde{h}_p^-(x, s) = h_p^-(s)$. We have $\tilde{h}_p^- = 0$ around $V_p^- \times \{k - 1/2\}$ and $\tilde{h}_p^- = a_k - b_p$ around $V_p^- \times \{k - 1/4\}$. Thus \tilde{h}_p^- has a smooth extension to \tilde{V}_p , also denoted by \tilde{h}_p^- , which is equal to $a_k - b_p$ on $\tilde{V}_p \setminus T_k^-$. The function $\tilde{\lambda}_p \tilde{h}_p^-$ on \tilde{V}_p can be extended by zero to get a smooth function on \tilde{T}_k , also denoted by $\tilde{\lambda}_p \tilde{h}_p^-$. Let $\tilde{h}_k^- = \sum_{p \in \mathcal{X}_k} \tilde{\lambda}_p \tilde{h}_p^-$ on \tilde{T}_k .

On the other hand, let ρ_k be a smooth function on I_k^+ such that $\rho_k' \geq 0$, $\rho_k = 0$ around $k + 1/4$, and $\rho_k = 1$ around $k + 1/2$. Let $\tilde{\rho}_k$ be the smooth function on $T_k^+ \equiv \Sigma_k^+ \times I_k^+$ given by $\tilde{\rho}_k(x, s) = \rho_k(s)$, and let

$$\tilde{h}_k^+ = \tilde{h}_k^-(1 - \tilde{\rho}_k) + (a_k - b_k)\tilde{\rho}_k$$

on T_k^+ . This smooth function is equal to \tilde{h}_k^- around $\Sigma_k^+ \times \{k + 1/4\}$, and is equal to $a_k - b_k$ around $\Sigma_k^+ \times \{k + 1/2\} \equiv \Sigma_k^+$. So the functions, \tilde{h}_k^- on $\tilde{T}_k \setminus T_k^+$ and \tilde{h}_k^+ on T_k^+ , can be combined to produce a smooth function \tilde{h}_k on \tilde{T}_k . Since $\tilde{h}_k = 0$ around Σ_k^- and $\tilde{h}_k = a_k - b_k$ around Σ_k^+ , there is a smooth extension of \tilde{h}_k to M , also denoted by \tilde{h}_k , which is constant on $M \setminus \tilde{T}_k$.

Let $f_k = f_{k-1} + \tilde{h}_k$ on M . This smooth function satisfies (A.2) because f_{k-1} satisfies (A.2), and X induces the opposite of the standard orientation on every fiber $\{x\} \times I_k^\pm \equiv I_k^\pm$ of T_k^\pm ($x \in \Sigma_k^\pm$). It also satisfies (A.3) and (A.4) for $p \in \mathcal{X}_l$ with $1 \leq l < k$ because f_{k-1} satisfies these properties and $d\tilde{h}_k$ is supported in the interior of \tilde{T}_k .

Next, take any $p \in \mathcal{X}_k$, $q \in \mathcal{X}_{k-1}$ and $\gamma \in \mathcal{T}(p, q) \subset \mathcal{T}_p^1$. We have $\gamma \cap T_k^- \equiv \{x\} \times I_k^-$ for some $x \in K_p^- \cap K_q^+ \subset \Sigma_k^- = \Sigma_{k-1}^+$, and the orientation of $\gamma \cap T_k^-$ agrees with the opposite of the standard orientation of $\{x\} \times I_k^- \equiv I_k^-$. Then

$$\begin{aligned} (\eta'' + df_k)(\gamma) &= (\eta'' + df_{k-1} + d\tilde{h}_k)(\gamma) \leq -b_p + \lambda_p^-(x) d\tilde{h}_p^-(\gamma) \\ &= -b_p - \int_{I_k^-} dh_p^- = -b_p - (a_k - b_p) = -a_k. \end{aligned}$$

Here, the equality holds when the maximum of (A.6) is achieved at γ . Hence f_k also satisfies (A.4) for $p \in \mathcal{X}_k$.

Finally, take any $p \in \mathcal{X}_k$, $u \in \mathcal{X}_{k+1}$ and $\delta \in \mathcal{T}(u, p) \subset \mathcal{T}_u^1 \subset \mathcal{T}_{k+1}^1$. Thus $\delta \cap T_k^+ \equiv \{y\} \times I_k^+$ for some $y \in K_p^+ \cap K_u^- \subset \Sigma_k^+ = \Sigma_{k+1}^-$, and the orientation of $\delta \cap T_k^+$ agrees with the opposite of the standard orientation of $\{y\} \times I_k^+ \equiv I_k^+$. Then

$$\begin{aligned} (\eta'' + df_k)(\delta) &= (\eta'' + df_{k-1} + d\tilde{h}_k)(\delta) = \eta''(\delta) + d\tilde{h}_k^+(\delta) \\ &= \eta''(\delta) + (\tilde{h}_k^-(y) - (a_k - b_k)) \int_{I_k^-} d\rho_k = \eta''(\delta) + \tilde{\lambda}_p(y) \tilde{h}_p^-(y) + b_k - a_k \\ &= \eta''(\delta) + a_k - b_p + b_k - a_k = \eta''(\delta) + b_k - b_p \geq \eta''(\delta) > -a_k, \end{aligned}$$

where the second equality is true because f_{k-1} satisfies (A.3), and the last inequality holds by (A.1). So f_k satisfies (A.5).

REFERENCES

- ALG20 J.A. Álvarez López and P. Gilkey, *Derived heat trace asymptotics for the de Rham and Dolbeault complexes*, Pure Appl. Funct. Anal., to appear, arXiv:2010.01146, 2020.
- ALG21 ———, *The local index density of the perturbed de Rham complex*, Czechoslovak Math. J. **71** (2021), no. 3, 901–932.
- ALK02 J.A. Álvarez López and Y.A. Kordyukov, *Distributional Betti numbers of transitive foliations of codimension one*, Foliations: geometry and dynamics. Proceedings of the Euroworkshop, Warsaw, Poland, May 29–June 9, 2000 (Singapore), World Sci. Publ., 2002, pp. 159–183. MR 1882768
- ALK08 ———, *Lefschetz distribution of Lie foliations, C^* -algebras and elliptic theory II* (Basel), Trends Math., Birkhäuser, 2008, http://dx.doi.org/10.1007/978-3-7643-8604-7_1, pp. 1–40. MR 2408134
- ALKL20 J.A. Álvarez López, Y.A. Kordyukov, and E. Leichtnam, *A trace formula for foliated flows*, in preparation, 2020.
- BF97 M. Braverman and M. Farber, *Novikov type inequalities for differential forms with non-isolated zeros*, Math. Proc. Cambridge Philos. Soc. **122** (1997), 357–375. MR 1458239
- BFK10 D. Burghilea, L. Friedlander, and T. Kappeler, *On the space of trajectories of a generic gradient like vector field*, An. Univ. Vest Timiș. Ser. Mat.-Inform. (2010), no. 1-2, 45–126. MR 2849328
- BGS88 J.-M. Bismut, H. Gillet, and C. Soulé, *Analytic torsion and holomorphic determinant bundles. I. Bott-Chern forms and analytic torsion*, Comm. Math. Phys. **115** (1988), no. 1, 49–78. MR 929146
- BGS90 ———, *Complex immersions and Arakelov geometry*, The Grothendieck Festschrift, Vol. I, Progr. Math., vol. 86, Birkhäuser Boston, Boston, MA, 1990. MR 1086887
- BGV04 N. Berline, E. Getzler, and M. Vergne, *Heat kernels and Dirac operators*, Grundlehren Text Editions, Springer-Verlag, Berlin, 2004, Corrected reprint of the 1992 original. MR 2273508

- BH01 D. Burghelca and S. Haller, *On the topology and analysis of a closed one form. I. (Novikov's theory revisited)*, Essays on geometry and related topics, Vol. 1, 2, Monogr. Enseign. Math., vol. 38, Enseignement Math., Geneva, 2001, pp. 133–175. MR 1929325
- BH04 ———, *Laplace transform, dynamics, and spectral geometry*, arXiv:math/0405037, 2004.
- BH06 ———, *Euler structures, the variety of representations and the Milnor-Turaev torsion*, Geom. Topol. **10** (2006), 1185–1238. MR 2255496
- BH08 ———, *Dynamics, Laplace transform and spectral geometry*, J. Topol. **1** (2008), no. 1, 115–151. MR 2365654
- Bot88 R. Bott, *Morse theory indomitable*, Publ. Math. Inst. Hautes Études Sci. **68** (1988), 99–114. MR 1001450
- Bur97 D. Burghelca, *Lectures on Witten-Helffer-Sjöstrand theory*, Proceedings of the Third International Workshop on Differential Geometry and its Applications and the First German-Romanian Seminar on Geometry (Sibiu, 1997), vol. 5, 1997, pp. 85–99. MR 1723597
- BZ92 J.-M. Bismut and W. Zhang, *An extension of a theorem by Cheeger and Müller*, Astérisque **205** (1992), 235 pp., with an appendix by F. Laudenbach. MR 1185803
- BZ94 ———, *Milnor and Ray-Singer metrics on the equivariant determinant of a flat vector bundle*, Geom. Funct. Anal. **4** (1994), 136–212. MR 1262703
- Che44 S. Chern, *A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds*, Ann. Math. (2) **45** (1944), 741–752. MR 1458239
- Che79 J. Cheeger, *Analytic torsion and the heat equation*, Ann. of Math. (2) **109** (1979), no. 2, 259–322. MR 528965
- Den08 C. Deninger, *Analogies between analysis on foliated spaces and arithmetic geometry*, Groups and analysis, London Math. Soc. Lecture Note Ser., vol. 354, Cambridge Univ. Press, Cambridge, 2008, <https://doi.org/10.1017/CB09780511721410.010>, pp. 174–190. MR 2528467
- dr50 G. de Rham, *Complexes à automorphismes et homéomorphie différentiable*, Ann. Inst. Fourier (Grenoble) **2** (1950), 51–67. MR 43468
- DS88a N. Dunford and J. T. Schwartz, *Linear operators. Part I: General theory*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988. MR 1009162
- DS88b ———, *Linear operators. Part II: Spectral theory. Selfadjoint operators in Hilbert space*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988. MR 1009163
- Far95 M. Farber, *Singularities of the analytic torsion*, J. Differential Geom. **41** (1995), no. 3, 528–572. MR 1338482
- Far04 ———, *Topology of closed one-forms*, Mathematical Surveys and Monographs, vol. 108, Amer. Math. Soc., Providence, RI, 2004. MR 2034601
- Flo89 A. Floer, *Witten's complex and infinite dimensional Morse theory*, J. Differential Geom. **30** (1989), no. 1, 207–221. MR 1001276
- Fra35 W. Franz, *Über die Torsion einer Überdeckung*, J. Reine Angew. Math. **173** (1935), 245–254. MR 1581473
- Gil95 P.B. Gilkey, *Invariance theory, the heat equation, and the Atiyah-Singer index theorem*, second ed., Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1995. MR 1396308
- GS77 P. Günther and R. Schimming, *Curvature and spectrum of compact Riemannian manifolds*, J. Differential Geom. **12** (1977), no. 4, 599–618. MR 512929
- Gui77 V. Guillemin, *Lectures on spectral theory of elliptic operators*, Duke Math. J. **44** (1977), no. 3, 485–517, <http://projecteuclid.org/euclid.dmj/1077312384>. MR 0448452
- Hir76 M.W. Hirsch, *Differential topology*, Graduate Texts in Mathematics, vol. 33, Springer-Verlag, New York, Heidelberg, Berlin, 1976.
- HM06 F.R. Harvey and G. Minervini, *Morse Novikov theory and cohomology with forward supports*, Math. Ann. **335** (2006), no. 4, 787–818. MR 2232017

ZETA INVARIANTS OF MORSE FORMS

- HS85 B. Helffer and J. Sjöstrand, *Puits multiples en mécanique semi-classique. IV. étude du complexe de Witten*, Comm. Partial Differential Equations **10** (1985), 245–340. MR 780068
- Kat95 T. Kato, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition. MR 1335452
- KM76 F.F. Knudsen and D. Mumford, *The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”*, Math. Scand. **39** (1976), no. 1, 19–55. MR 437541
- Lat94 F. Latour, *Existence de 1-formes fermées non singulières dans une classe de cohomologie de de Rham*, Inst. Hautes Études Sci. Publ. Math. **80** (1994), 135–194. MR 1320607
- Lau12 F. Laudenbach, *Transversalité, courants et théorie de Morse*, Les Éditions de l’École Polytechnique, Palaiseau, 2012, Un cours de topologie différentielle. Exercices proposés by François Labourie. MR 3088239
- Lei08 E. Leichtnam, *On the analogy between arithmetic geometry and foliated spaces*, Rend. Mat. Appl. (7) **28** (2008), no. 2, 163–188. MR 2463936
- Lei14 ———, *On the analogy between L-functions and Atiyah-Bott-Lefschetz trace formulas for foliated spaces*, Rend. Mat. Appl. (7) **35** (2014), no. 1-2, 1–34. MR 3241361
- LPNV13 D. Le Peutrec, F. Nier, and C. Viterbo, *Precise Arrhenius law for p-forms: the Witten Laplacian and Morse-Barannikov complex*, Ann. Henri Poincaré **14** (2013), no. 3, 567–610. MR 3035640
- M78 W Müller, *Analytic torsion and R-torsion of Riemannian manifolds*, Adv. in Math. **28** (1978), no. 3, 233–305. MR 498252
- M93 W. Müller, *Analytic torsion and R-torsion for unimodular representations*, J. Amer. Math. Soc. **6** (1993), no. 3, 721–753. MR 1189689
- Mel93 R.B. Melrose, *The Atiyah-Patodi-Singer index theorem*, Research Notes in Mathematics, vol. 4, A.K. Peters, Ltd., Wellesley, MA, 1993. MR 1348401
- Mel96 ———, *Differential analysis on manifolds with corners*, <http://www-math.mit.edu/~rbm/book.html>, 1996.
- Mic19 L. Michel, *About small eigenvalues of the Witten Laplacian*, Pure Appl. Anal. **1** (2019), no. 2, 149–206. MR 3949372
- Mil63 J. Milnor, *Morse theory*, Annals of Mathematics Studies, vol. 51, Princeton University Press, Princeton, New Jersey, 1963, Based on lecture notes by M. Spivak and R. Wells. MR 0163331
- Mil65 J.W. Milnor, *Lectures on the h-cobordism theorem. Notes by L. Siebenmann and J. Sondow*, Princeton University Press, Princeton, N.J., 1965. MR 0190942
- Mil66 J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. **72** (1966), 358–426. MR 196736
- Min15 G. Minervini, *A current approach to Morse and Novikov theories*, Rend. Mat. Appl. (7) **36** (2015), no. 3-4, 95–195. MR 3533253
- MQ86 V. Mathai and D. Quillen, *Superconnections, Thom classes, and equivariant differential forms*, Topology **25** (1986), no. 1, 85–110. MR 836726
- MRS16 T. Mrowka, D. Ruberman, and N. Saveliev, *An index theorem for end-periodic operators*, Compos. Math. **152** (2016), no. 2, 399–444. MR 3462557
- Nov81 S.P. Novikov, *Multivalued functions and functionals. An analog of the Morse theory*, Soviet. Math., Dokl. **24** (1981), 222–226. MR 630459
- Nov82 ———, *The Hamiltonian formalism and a multivalued analogue of Morse theory*, Russian Math. Surveys **37** (1982), 1–56. MR 676612
- Nov02 ———, *On the exotic De-Rham cohomology. Perturbation theory as a spectral sequence*, arXiv:math-ph/0201019, 2002.
- Paj87 A.V. Pajitnov, *An analytic proof of the real part of Novikov’s inequalities*, Soviet Math., Dokl. **35** (1987), 456–457. MR 891557
- Paj06 ———, *Circle-valued Morse theory*, de Gruyter Studies in Mathematics, vol. 32, Walter de Gruyter & Co., Berlin, 2006. MR 2319639

- Qui85 D. Quillen, *Determinants of Cauchy-Riemann operators on Riemann surfaces*, *Funct. Anal. Appl.* **19** (1985), no. 1, 37–41. MR 783704
- Rei35 K. Reidemeister, *Homotopieringe und Linsenräume*, *Abh. Math. Sem. Univ. Hamburg* **11** (1935), no. 1, 102–109. MR 3069647
- Roe98 J. Roe, *Elliptic operators, topology and asymptotic methods*, second ed., *Pitman Research Notes in Mathematics*, vol. 395, Longman, Harlow, 1998. MR 1670907
- RS71 D.B. Ray and I.M. Singer, *R-torsion and the Laplacian on Riemannian manifolds*, *Advances in Math.* **7** (1971), 145–210. MR 295381
- Sch93 M. Schwarz, *Morse homology*, *Progress in Mathematics*, vol. 111, Birkhäuser Verlag, Basel, 1993. MR 1239174
- Sch99 ———, *Equivalences for Morse homology*, *Geometry and Topology in Dynamics* (Winston-Salem, NC, 1998/San Antonio, TX, 1999) (Providence, RI), *Contemporary Mathematics*, vol. 246, American Mathematical Society, 1999, pp. 197–216. MR 1732382
- See67 R.T. Seeley, *Complex powers of an elliptic operator*, *Singular Integrals* (Proc. Sympos. Pure Math., Chicago, Ill., 1966) (Providence, R.I.), vol. 10, Amer. Math. Soc., 1967, pp. 288–307. MR 0237943
- Sma60a S. Smale, *The generalized Poincaré conjecture in higher dimensions*, *Bull. Amer. Math. Soc.* **66** (1960), 373–375. MR 124912
- Sma60b ———, *Morse inequalities for a dynamical system*, *Bull. Amer. Math. Soc.* **66** (1960), 43–49. MR 117745
- Sma61 ———, *On gradient dynamical systems*, *Ann. of Math. (2)* **74** (1961), 199–206. MR 0133139
- Sma63 ———, *Stable manifolds for differential equations and diffeomorphisms*, *Ann. Scuola Norm. Sup. Pisa (3)* **17** (1963), no. 1–2, 97–116. MR 0165537
- Tho49 R. Thom, *Sur une partition en cellules associée à une fonction sur une variété*, *C.R. Acad. Sci. Paris* **228** (1949), 973–975. MR 29160
- Wit82 E. Witten, *Supersymmetry and Morse theory*, *J. Differ. Geom.* **17** (1982), 661–692. MR 683171
- Zha01 W. Zhang, *Lectures on Chern-Weil theory and Witten deformations*, *Nankai Tracts in Mathematics*, vol. 4, World Scientific Publishing Co., Inc., River Edge, NJ, 2001. MR 1864735

Jesús A. Álvarez López jesus.alvarez@usc.es

Department of Mathematics, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain

Yuri A. Kordyukov yurikor@matem.anrb.ru

Institute of Mathematics, Ufa Federal Research Center, Russian Academy of Sciences, 112 Chernyshevsky street, 450008 Ufa, Russia

Eric Leichtnam eric.leichtnam@imj-prg.fr

Institut de Mathématiques de Jussieu-PRG, CNRS, Batiment Sophie Germain (bureau 740), Case 7012, 75205 Paris Cedex 13, France