



## Non symmetry of the function $\frac{\zeta(s)}{s}$

W Oukil

### ► To cite this version:

| W Oukil. Non symmetry of the function  $\frac{\zeta(s)}{s}$ . 2022. hal-03581978v10

HAL Id: hal-03581978

<https://hal.science/hal-03581978v10>

Preprint submitted on 28 Apr 2022 (v10), last revised 26 Nov 2023 (v25)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Non symmetry of the function $\frac{\zeta(s)}{s}$

W. Oukil

Laboratory of Dynamical Systems, Faculty of Mathematics.  
University of Science and Technology Houari Boumediene,  
BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

April 28, 2022

## Abstract

We prove that  $\frac{\zeta(s)}{s} \neq \frac{\zeta(1-s)}{1-s}$  for every  $\Re(s) \in (0, \frac{1}{2})$  and  $\Im(s) \in \mathbb{R}^*$ , where  $\zeta$  is the Riemann Zeta function. In the end of the paper, we give a discussion about the Riemann hypothesis.

**Keywords:** Riemann functional equation, Zeta function.

## 1 Main result

Let be  $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$  be a bounded function. We study the function  $\psi[\eta] : (0, 1) \times \mathbb{R}^* \rightarrow \mathbb{C}$  defined as

$$\forall x \in (0, 1), \forall \tau \in \mathbb{R}^* : \quad \psi[\eta](x, \tau) := \int_0^{+\infty} \frac{\eta(t)}{t^{1+x+i\tau}} dt, \quad (1)$$

We focus about the function  $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$  satisfying the following hypotheses

$$H_1 : \quad 0 \leq \eta(t) \leq 1 \text{ for all } t \geq 0,$$

$$H_2 : \quad \text{the function } \eta \text{ is a } C^1 \text{ piecewise function.}$$

We prove in this paper the following Main result

**Main Result.** *Consider the function  $\psi$  defined by Equation (1). Then, for every function  $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$  that satisfies the hypotheses  $H_1$  and  $H_2$  we have*

$$\forall \tau \in \mathbb{R}^*, \forall x \in (0, \frac{1}{2}) : \quad \|\psi[\eta](x, \tau) - \psi[\eta](1-x, \tau)\| \neq 0.$$

**Notation 1.** Let  $\eta : \mathbb{R}_+ \mapsto \mathbb{R}$  be a function that satisfies  $H_1$  and  $H_2$ . For every  $r \in (0, \frac{1}{2})$  define the functions  $t \mapsto h_r(\eta, t)$  as

$$h_r(\eta, t) := 2 \exp(-\frac{1}{2}t)[\eta(\exp(t)) + 1] \sinh(rt), \quad \forall t \geq 0.$$

In order to simplify the notation we denote  $h_r(t) := h_r(\eta, t)$  for all  $t \geq 0$ .

*Proof of the Main result.* Let be  $r \in (0, \frac{1}{2})$  and  $\tau \in \mathbb{R}^*$ . Use the change of variable,  $t \rightarrow \exp(u)$ , in Equation (1), we obtain

$$\begin{aligned} \|\psi[\eta](\frac{1}{2} - r, \tau) - \psi[\eta](\frac{1}{2} + r, \tau)\| &= \left\| \int_0^{+\infty} h_r(s) \exp(i\tau s) ds \right\|, \\ \forall r \in (0, \frac{1}{2}), \quad \forall \tau \in \mathbb{R}^*. \end{aligned}$$

Then

$$\|\psi[\eta](\frac{1}{2} - r, \tau) - \psi[\eta](\frac{1}{2} + r, \tau)\| = 0,$$

implies

$$\left\| \int_0^{+\infty} \sin(\tau s) h_r(s) ds \right\| = 0.$$

Use the hypothesis  $H_2$  and the integration part formula, we get

$$\lim_{n \rightarrow +\infty} \int_0^{\frac{2n\pi}{\tau}} \cos(\tau u) \int_0^u h_r(v) dv du = 0.$$

Since  $r \in (0, \frac{1}{2})$ , there exists  $\alpha_r > 0$  such that

$$\lim_{u \rightarrow +\infty} \int_0^u h_r(v) dv = \alpha_r.$$

Then

$$\lim_{n \rightarrow +\infty} \int_0^{\frac{2n\pi}{\tau}} \cos(\tau u) [\alpha_r - \int_0^u h_r(v) dv] du = 0.$$

In order to symplify the notation, denote

$$\tilde{h}_r(t) := \alpha_r - \int_0^t h_r(v) dv, \quad \forall t \geq 0.$$

Use again the integration part formula,

$$\lim_{n \rightarrow +\infty} \left[ \int_0^{\frac{2n\pi}{\tau}} \tilde{h}_r(v) dv + \tau \int_0^{\frac{2n\pi}{\tau}} \sin(\tau u) \int_0^u \tilde{h}_r(v) dv du \right] = 0. \quad (2)$$

By hypothesis  $H_1$  the function  $v \mapsto h_r(v)$  is positive, which implies that

$$u \mapsto \int_0^u \tilde{h}_r(v)dv,$$

is positive, increasing, concave on  $\mathbb{R}_+$  and converge. Then there exists a sequence  $(t_n)_{n \in \mathbb{N}} \subset \mathbb{R}_+$  such that

- $t_n \in [\frac{(2n-1)\pi}{\tau}, \frac{2n\pi}{\tau}]$ ,  $\forall n \geq 0$ ,
- $\lim_{n \rightarrow +\infty} \left| t_n - [\frac{2n\pi}{\tau} - \frac{\pi}{2\tau}] \right| \neq 0$ ,
- $\int_0^{t_n} \sin(\tau u) \int_0^u \tilde{h}_r(v)dv du = 0$ .

Use the Cauchy mean value Theorem on Equation (2), for all  $n \geq 0$  there exists  $\eta_{\tau,r,n} \in (t_n, \frac{2n\pi}{\tau})$  such that

$$\lim_{n \rightarrow +\infty} \left[ \int_0^{\frac{2n\pi}{\tau}} \tilde{h}_r(v)dv + \int_0^{\eta_{\tau,r,n}} \tilde{h}_r(v)dv \int_{t_n}^{\frac{2n\pi}{\tau}} \tau \sin(\tau u)du \right] = 0$$

which can be written as

$$\lim_{n \rightarrow +\infty} \left[ 1 + \int_{t_n}^{\frac{2n\pi}{\tau}} \tau \sin(\tau u)du \right] = 0.$$

Contradiction with  $\lim_{n \rightarrow +\infty} \left| t_n - [\frac{2n\pi}{\tau} - \frac{\pi}{2\tau}] \right| \neq 0$ .  $\square$

## 2 Discussion about the Riemann hypothesis

Consider the representation of the Riemann Zeta function  $\zeta$  defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -s \int_0^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*, \quad (3)$$

and where  $\{t\}$  is the fractional part of the real  $t$ . Define the *non-trivial zeros* of the function  $\zeta$  in the following sense

**Definition 2.** Let be  $s \in \mathbb{C}$ . We say that  $s$  is a *non-trivial zero of the function  $\zeta$*  if

$$\zeta(s) = 0 \quad \text{and} \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*.$$

Denote  $\eta_*(t) := \{t\}$ . For every  $\tau \in \mathbb{R}^*$  and every  $x \in (0, 1)$  we have

$$\psi[\eta_*](x, \tau) = \frac{\zeta(s)}{s} \quad \text{and} \quad \psi[\eta_*](1-x, \tau) = -\frac{\zeta(1-s)}{1-s}, \quad s := x + i\tau.$$

Since

$$0 \leq \eta_*(t) \leq 1, \quad \forall t \geq 0,$$

Then  $\eta_*$  satisfies the hypotheses  $H_1$ . The function  $\eta_*$  is  $C^1$  on  $\mathbb{R}_+/\mathbb{N}$ . Then  $\eta_*$  satisfies the hypotheses  $H_2$ . By the Riemann functional equation [|1], page 13 Theorem 2.1], the non-trivial zeros of the function  $\zeta$  are symmetric about the line  $\Re(z) = \frac{1}{2}$ . By the Main result, the non-trivial zeros  $s \in \mathbb{C}$  symmetric about the line  $\Re(z) = \frac{1}{2}$  satisfies  $\Re(s) = \frac{1}{2}$ .

## References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).