



Non symmetry of the function $\frac{\zeta(s)}{s}$

W Oukil

► To cite this version:

| W Oukil. Non symmetry of the function $\frac{\zeta(s)}{s}$. 2022. hal-03581978v13

HAL Id: hal-03581978

<https://hal.science/hal-03581978v13>

Preprint submitted on 10 Jul 2022 (v13), last revised 26 Nov 2023 (v25)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non symmetry of the function $\Im\left(\frac{\zeta(s)}{s}\right)$

W. Oukil

Laboratory of Dynamical Systems, Faculty of Mathematics.
 University of Science and Technology Houari Boumediene,
 BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

July 10, 2022

Abstract

We prove that $\Im\left(\frac{\zeta(s)}{s}\right) + \Im\left(\frac{\zeta(1-s)}{1-s}\right) \neq 0$ for every $\Re(s) \in (0, \frac{1}{2})$ and $\Im(s) \in \mathbb{R}^*$, where ζ is the Riemann Zeta function. In the end of the paper, we give a discussion about the Riemann hypothesis.

Keywords: Riemann functional equation, Zeta function.

1 Main result

Consider the representation of the Riemann Zeta function ζ defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -s \int_0^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*, \quad (1)$$

and where $\{t\}$ is the fractional part of the real t . We prove in this paper the following Main result

Main Result. *Consider the function ζ defined by Equation (1). Then*

$$\Im\left(\frac{\zeta(\frac{1}{2} - r + i\tau)}{\frac{1}{2} - r + i\tau}\right) - \Im\left(\frac{\zeta(\frac{1}{2} + r + i\tau)}{\frac{1}{2} + r + i\tau}\right) \neq 0, \quad \forall r \in (0, \frac{1}{2}), \quad \forall \tau \in \mathbb{R}^*.$$

2 Main Proposition

Notation 1. Let be $m \in \mathbb{N}$ and $q \in \mathbb{N}^*$. We denote Λ_q^m as the set of q-combinations of the set $\{0, 1, \dots, m\}$;

$$\Lambda_q^m := \left\{ \{l_1, \dots, l_q\} \subset \{0, 1, \dots, m\} : \quad l_j \neq l_{j'}, \quad \forall 1 \leq j \neq j' \leq q \right\}.$$

In order to simplify the notation, we denote

$$\alpha_\tau := \frac{1}{2} + i\tau, \quad \forall \tau \in \mathbb{R}^*.$$

Notation 2. For every $r \in (0, \frac{1}{2})$ and $\tau \in \mathbb{R}^*$ denote

$$\phi_\tau(r) := \frac{\zeta(\alpha_\tau - r)}{\alpha_\tau - r} - \frac{\zeta(\alpha_\tau + r)}{\alpha_\tau + r}.$$

We have the following Main proposition

Proposition 3. *For every $\tau \in \mathbb{R}^*$ and $n \in \mathbb{N}$ we have*

$$\begin{aligned} \frac{1}{(2n+1)!} \frac{d^{2n+1}}{dr^{2n+1}} \phi_\tau(0) &= -\frac{3}{2} \alpha_\tau^{-2n-1} - (-1)^n \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(n+1)}} \\ &\quad - \sum_{k=n+2}^{+\infty} (-1)^k \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}} \right] \sum_{A \in \Lambda_{2(k-n)}^{2k+1}} \Pi_{j \in A}(\alpha_\tau + j) \end{aligned}$$

3 Main Lemma

Lemma 4. *We have*

$$\int_1^t (\{v\} + 1) dv = t \frac{3}{2} - \frac{1}{12} + \frac{1}{2} \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2} \exp(ij2\pi t), \quad \forall t \geq 0,$$

and

$$\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(m+1)}} \leq \frac{1}{2} \frac{1}{m} \frac{1}{(2m)!}, \quad \forall m \geq 2.$$

Proof. The function $v \mapsto \{v\} + 1$ is 1-periodic, then there exists a continuous 1-periodic function $p : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\int_0^t (\{v\} + 1) dv = t \int_0^1 (\{v\} + 1) dv + p(t), \quad \forall t \in \mathbb{R}.$$

Since

$$\int_0^1 (\{v\} + 1) dv = \frac{3}{2},$$

we get

$$\int_0^t (\{v\} + 1) dv = t \frac{3}{2} + p(t), \quad \forall t \in \mathbb{R}.$$

The function p is a piecewise C^∞ , continuous on \mathbb{R} and 1-periodic. By Dirichlet Theorem, the Fourier series

$$n \mapsto \sum_{k=-n}^n a_k \exp(ik2\pi t),$$

converge uniformly on \mathbb{R}_+ to the function $t \mapsto p(t)$, where $(a_k)_k \subset \mathbb{C}$ are the Fourier coefficients of the function p .

$$\int_0^t (\{v\} + 1) dv = t \frac{3}{2} + \sum_{j \in \mathbb{Z}} a_j \exp(ij2\pi t), \forall t \geq 0.$$

By definition of the Fourier coefficients we have

$$\begin{aligned} a_j &= \int_0^1 \exp(-ij2\pi u) \left(\int_0^u (\{v\} + 1) dv - u \frac{3}{2} \right) du \\ &= \frac{1}{2} \int_0^1 \exp(-ij2\pi u) u(u-1) du = \frac{1}{2} \frac{1}{(2j\pi)^2}, \quad \forall j \in \mathbb{Z}^*, \end{aligned}$$

and

$$a_0 = \frac{1}{2} \int_0^1 u(u-1) du = -\frac{1}{12}$$

Now prove the second item of the present Lemma. Let $m \geq 2$ and consider the 1-periodic function $g_m : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$g_m(t) := \frac{1}{m!} \frac{1}{2} (t^2 - t)^m, \quad \forall t \in [0, 1].$$

The function g_m is a piecewise C^∞ , continuous on \mathbb{R} and 1-periodic. By Dirichlet Theorem,

$$g_m(t) = \sum_{j \in \mathbb{Z}} a_{m,j} \exp(ij2\pi t), \quad \forall t \geq 0, \tag{2}$$

where the Fourier coefficients satisfies

$$a_{m,j} = \frac{1}{m!} \frac{1}{2} \int_0^1 \exp(-ij2\pi u) (u^2 - u)^m du.$$

Use a successive integration part formula, to obtain

$$a_{m,j} = \frac{1}{(ij2\pi)^{m+1}}, \quad \forall m \geq 0,$$

and replace its in Equation (2) , we get

$$g_m(t) = \sum_{j \in \mathbb{Z}} \frac{1}{(ij2\pi)^{m+1}} \exp(ij2\pi t), \quad \forall t \geq 0,$$

Then

$$\int_0^t g_m(u) du = \sum_{j \in \mathbb{Z}} \frac{1}{(ij2\pi)^{m+2}} \left(\exp(ij2\pi t) - 1 \right), \quad \forall t \geq 0,$$

implies

$$\int_0^1 \int_0^t g_m(u) du dt = \sum_{j \in \mathbb{Z}} \frac{1}{(ij2\pi)^{m+2}}, \quad \forall t \geq 0,$$

By definition of g_m we get

$$\begin{aligned} \sum_{j \in \mathbb{Z}} \frac{1}{(j2\pi)^{2(m+1)}} &= \left| \int_0^1 \int_0^t g_{2m}(u) du dt \right| \\ &\leq \frac{1}{2} \frac{1}{(2m)!} \int_0^1 (1-t)^m du dt \leq \frac{1}{2} \frac{1}{m} \frac{1}{(2m)!}, \quad \forall m \geq 2. \end{aligned}$$

□

Lemma 5. Let be $n \in \mathbb{N}$ and $\tau \in \mathbb{R}^*$. For every $k \geq 0$ define the function $V_{k,\tau}(n, u) : \mathbb{R} \rightarrow \mathbb{C}$ as

$$V_{0,\tau}(n, u) := u^n \exp(-\alpha_\tau u), \quad V_{k+1,\tau}(n, u) = \exp(-u) \frac{d}{du} V_{k,\tau}(n, u), \quad \forall k \geq 0.$$

Then

$$V_{k,\tau}(n, 0) = 0, \quad \forall k \in \llbracket 0, n-1 \rrbracket, \quad V_{n,\tau}(n, 0) = n!,$$

and

$$V_{k,\tau}(n, 0) = n! (-1)^{k-n} \sum_{A \in \Lambda_{k-n}^{k-1}} \Pi_{j \in A} (\alpha_\tau + j), \quad \forall k \geq n+1.$$

where we recall that Λ_{k-n}^{k-1} is defined in the Notation 1. In addition,

$$\int_0^{+\infty} \left| \frac{d}{du} V_{k,\tau}(n, u) \right| du \leq \frac{n}{k^{n+1}} \max\{|\alpha_\tau|, 1\} \Pi_{m=1}^{k-1} |\alpha_\tau + m|, \quad \forall k \geq 2.$$

Proof. Let be $n \in \mathbb{N}$ and $\tau \in \mathbb{R}^*$. By recurrence, for every $k \geq 0$ there exists a polynomial $P_k(n, u)$ such that

$$V_{k,\tau}(n, u) = \exp(-(\alpha_\tau + k)u)P_k(n, u).$$

We have

$$V_{k+1,\tau}(n, u) = \exp(-u) \frac{d}{du} \left[\exp(-(\alpha_\tau + k)u)P_k(n, u) \right]. \quad (3)$$

Then

$$P_{k+1}(n, u) = \frac{d}{du} P_k(n, u) - (\alpha_\tau + k)P_k(n, u). \quad (4)$$

Write

$$P_k(n, u) = \sum_{j=0}^n c_{k,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u), \quad (5)$$

and replace it in Equation (4),

$$\begin{aligned} \sum_{j=0}^n c_{k+1,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u) &= \sum_{j=0}^n c_{k,j}(n, \tau) \frac{d^{j+1}}{du^{j+1}} P_0(n, u) \\ &\quad - (\alpha_\tau + k) \sum_{j=0}^n c_{k,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u). \end{aligned}$$

Since $P_0(n, u) = u^n$ then $\frac{d^{n+1}}{du^{n+1}} P_0(n, u) = 0$, we get

$$\begin{aligned} \sum_{j=0}^n c_{k+1,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u) &= \sum_{j=0}^{n-1} c_{k,j}(n, \tau) \frac{d^{j+1}}{du^{j+1}} P_0(n, u) \\ &\quad - (\alpha_\tau + k) \sum_{j=0}^n c_{k,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u), \end{aligned}$$

which can be written as

$$\begin{aligned} \sum_{j=0}^n c_{k+1,j} \frac{d^j}{du^j} P_0(n, u) &= \sum_{j=1}^n c_{k,j-1}(n, \tau) \frac{d^j}{du^j} P_0(n, u) \\ &\quad - (\alpha_\tau + k) \sum_{j=0}^n c_{k,j}(n, \tau) \frac{d^j}{du^j} P_0(n, u). \end{aligned}$$

Implies

$$\begin{pmatrix} c_{k+1,0}(n, \tau) \\ c_{k+1,1}(n, \tau) \\ \vdots \\ c_{k+1,n}(n, \tau) \end{pmatrix} = \left(C_n - (\alpha_\tau + k)Id_n \right) \begin{pmatrix} c_{k,0}(n, \tau) \\ c_{k,1}(n, \tau) \\ \vdots \\ c_{k,n}(n, \tau) \end{pmatrix}$$

where Id_n is the $n \times n$ identity matrix and where C_n is the $n \times n$ matrix defined as

$$C_n := \begin{pmatrix} 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & 0 \end{pmatrix}.$$

Then

$$\begin{pmatrix} c_{k+1,0}(n, \tau) \\ c_{k+1,1}(n, \tau) \\ \vdots \\ c_{k+1,n}(n, \tau) \end{pmatrix} = \left[\prod_{m=0}^k \left(C_n - (\alpha_\tau + m)Id_n \right) \right] \begin{pmatrix} c_{0,0}(n, \tau) \\ c_{0,1}(n, \tau) \\ \vdots \\ c_{0,n}(n, \tau) \end{pmatrix}$$

By definition of $P_k(n, u)$ (Resp. $V_{k+1,\tau}(n, u)$) in Equation (5) (Resp. Equation (3)) we have $P_0(n, u) = u^n$, we get

$$\begin{pmatrix} c_{k+1,0}(n, \tau) \\ c_{k+1,1}(n, \tau) \\ \vdots \\ c_{k+1,n}(n, \tau) \end{pmatrix} = \left[\prod_{m=0}^k \left(C_n - (\alpha_\tau + m)Id_n \right) \right] \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

By Equation (3), and (5), for all $u \geq 0$ we find

$$\begin{aligned} & \left| \frac{d}{du} V_{k+1,\tau}(n, u) \right| \\ & \leq n! \max\{|\alpha_\tau|, 1\} \left[\prod_{m=1}^k |\alpha_\tau + m| \right] \exp(-(\frac{1}{2} + k + 1)u) u^n, \end{aligned}$$

Since

$$\int_0^{+\infty} \exp(-(\frac{1}{2} + k + 1)u) u^n du = \frac{1}{(\frac{1}{2} + k + 1)^{n+1}} \frac{n!}{n!}$$

Then

$$\int_0^{+\infty} \left| \frac{d}{du} V_{k+1,\tau}(n, u) \right| du \leq n \frac{\max\{|\alpha_\tau|, 1\} \left[\prod_{m=1}^k |\alpha_\tau + m| \right]}{(\frac{1}{2} + k + 1)^{n+1}}.$$

Use the fact C_n is a nilpotent matrix of degree n , we get $c_{n,n}(n, \tau) = 1$ and

$$V_{k,\tau}(n, 0) = P_k(n, 0) = n! c_{k,n}(n, \tau) = n! \sum_{A \in \Lambda_{k-n}^{k-1}} (-1)^{k-n} \Pi_{j \in A} (\alpha_\tau + j).$$

□

4 Proof of the Main Proposition

Lemma 6. *Let be $\tau \in \mathbb{R}^*$. Consider the function ϕ_τ defined in the Notation 2. Then $\frac{d^{2n}}{dr^{2n}}\phi_\tau(0) = 0$ for all $n \geq 0$ and*

$$\frac{d^{2n+1}}{dr^{2n+1}}\phi_\tau(0) = 2 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) (\{\exp(u)\} + 1) du, \quad \forall n \geq 0.$$

Proof. Let be $x \in (0, 1)$ and $\tau \in \mathbb{R}^*$. From Equation (1), we have

$$\begin{aligned} \frac{\zeta(x + i\tau)}{x + i\tau} &= - \int_0^{+\infty} \frac{\{t\}}{t^{1+x+i\tau}} dt \\ &= - \int_0^{+\infty} \exp(-i\tau \ln(t)) \frac{\{t\}}{t^{1+x}} dt \\ &= - \int_0^1 \exp(-i\tau \ln(t)) \frac{1}{t^x} dt - \int_1^{+\infty} \exp(-i\tau \ln(t)) \frac{\{t\}}{t^{1+x}} dt \end{aligned}$$

Use the change of variable, $t \rightarrow \exp(-u)$ for the integral

$$\int_0^1 \exp(-i\tau \ln(t)) \frac{1}{t^x} dt,$$

and $t \rightarrow \exp(u)$ for the integral

$$\int_1^{+\infty} \exp(-i\tau \ln(t)) \frac{\{t\}}{t^{1+x}} dt,$$

we obtain

$$\begin{aligned} \frac{\zeta(x + i\tau)}{x + i\tau} &= \int_0^{+\infty} \exp(-i\tau u) \exp(-(1-x)u) du \\ &\quad - \int_0^{+\infty} \exp(-i\tau u) \exp(-xu) \{\exp(u)\} du. \end{aligned}$$

Implies

$$\begin{aligned}\frac{\zeta(\alpha_\tau - r)}{\alpha_\tau - r} &= \int_0^{+\infty} \exp(-i\tau u) \exp(-(\frac{1}{2} + r)u) du \\ &\quad - \int_0^{+\infty} \exp(-i\tau u) \exp(-(\frac{1}{2} - r)u) \{\exp(u)\} du,\end{aligned}$$

and

$$\begin{aligned}\frac{\zeta(\alpha_\tau + r)}{\alpha_\tau + r} &= \int_0^{+\infty} \exp(-i\tau u) \exp(-(\frac{1}{2} - r)u) du \\ &\quad - \int_0^{+\infty} \exp(i\tau u) \exp(-(\frac{1}{2} + r)u) \{\exp(u)\} du.\end{aligned}$$

We get

$$\begin{aligned}\phi_\tau(r) &= - \int_0^{+\infty} \exp(-i\tau u) [\exp(-(\frac{1}{2} + r)u) - \exp(-(\frac{1}{2} - r)u)] du \\ &\quad + \int_0^{+\infty} \exp(-i\tau u) [\exp((\frac{1}{2} + r)u) - \exp((\frac{1}{2} - r)u)] \{\exp(u)\} du.\end{aligned}$$

Implies

$$\phi_\tau(r) = 2 \int_0^{+\infty} \exp(-\alpha_\tau u) \sinh(ru) (\{\exp(u)\} + 1) du.$$

Since $r \in (0, \frac{1}{2})$ then

$$\frac{d^{2n}}{d^{2n}} \phi_\tau(r) = 2 \int_0^{+\infty} \exp(-\alpha_\tau u) u^{2n} \sinh(ru) (\{\exp(u)\} + 1) du.$$

implies $\frac{d^{2n}}{d^{2n}} \phi_\tau(0) = 0$ and

$$\frac{d^{2n+1}}{d^{2n+1}} \phi_\tau(r) = 2 \int_0^{+\infty} \exp(-\alpha_\tau u) u^{2n+1} \cosh(ru) (\{\exp(u)\} + 1) du,$$

implies

$$\frac{d^{2n+1}}{d^{2n+1}} \phi_\tau(0) = 2 \int_0^{+\infty} \exp(-\alpha_\tau u) u^{2n+1} (\{\exp(u)\} + 1) du.$$

□

Proof of the Proposition 3. Let be $n \geq 0$ and $\tau \in \mathbb{R}^*$. By Lemma 6 , we have

$$\frac{d^{2n+1}}{dr^{2n+1}}\phi_\tau(0) = 2 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) [\{\exp(u)\} + 1] du,$$

Then

$$\begin{aligned} \frac{d^{2n+1}}{dr^{2n+1}}\phi_\tau(0) &= 3 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) du \\ &\quad + 2 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) \left[\{\exp(u)\} - \frac{1}{2} \right] du, \end{aligned}$$

we get

$$\begin{aligned} \frac{d^{2n+1}}{dr^{2n+1}}\phi_\tau(0) &= 3 \frac{(2n+1)!}{\alpha_\tau^{2n+1}} \\ &\quad + 2 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) \left[\{\exp(u)\} - \frac{1}{2} \right] du \end{aligned}$$

In order to simplify the notation, denote for every $m \in \mathbb{N}$,

$$J_{n,\tau} := 2 \int_0^{+\infty} u^{2n+1} \exp(-\alpha_\tau u) \left[\{\exp(u)\} - \frac{1}{2} \right] du$$

Then

$$\frac{d^{2n+1}}{dr^{2n+1}}\phi_\tau(0) = -3(2n+1)!\alpha_\tau^{-2n-1} + 2J_{n,\tau}. \quad (6)$$

Use the integration part formula,

$$\begin{aligned} J_{n,\tau} &= 2 \int_0^{+\infty} u^{2n+1} \exp(-(\alpha_\tau + 1)u) \exp(u) \left[\{\exp(u)\} - \frac{1}{2} \right] du \\ &= -2 \int_0^{+\infty} \frac{d}{du} \left[u^{2n+1} \exp(-(\alpha_\tau + 1)u) \right] \left[\int_0^{\exp(u)} (\{v\} - \frac{1}{2}) dv \right] du. \end{aligned}$$

By Lemma 4,

$$J_{n,\tau} = - \int_0^{+\infty} \frac{d}{du} \left[u^{2n+1} \exp(-(\alpha_\tau + 1)u) \right] \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2} \exp(ij2\pi \exp(u)) du$$

Use the fact

$$\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2} \exp(ij2\pi t) = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \cos(2j\pi t), \quad \forall t \in \mathbb{R},$$

Use a successive integration part formula, by induction we deduce that

$$J_{n,\tau} = - \sum_{k=1}^m (-1)^k \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}} \right] V_{2k,\tau}(2n+1,0) + R_{\tau,n}(m), \quad (7)$$

where

$$\begin{aligned} V_{0,\tau}(2n+1, u) &:= u^{2n+1} \exp(-(\alpha_\tau + 1)u), \\ V_{k+1,\tau}(2n+1, u) &= \exp(-u) \frac{d}{du} V_{k,\tau}(2n+1, u), \quad \forall k \geq 0, \end{aligned} \quad (8)$$

and where

$$R_{\tau,n}(m) := \int_0^{+\infty} \left[\frac{d}{du} V_{2(m+1),\tau}(2n+1, u) \right] p_m(u) du,$$

where

$$p_m(u) := \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(ij2\pi)^{2(m+2)}} \exp(ij2\pi \exp(u)) \right].$$

By Lemma 4,

$$\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(m+2)}} \leq \frac{1}{2} \frac{1}{m} \frac{1}{2m!}, \quad \forall m \geq 2.$$

and by Lemma 5,

$$\begin{aligned} |R_{\tau,n}(m)| &\leq (2n+1) \frac{1}{2} \frac{1}{m} \frac{1}{(2m)!} \frac{\max\{|\alpha_\tau|, 1\} \left[\prod_{j=1}^{2m+1} |\alpha_\tau + j| \right]}{(2(m+1))^{2(n+1)}} \\ &= (2n+1) \frac{1}{2} \max\{|\alpha_\tau|, 1\} \frac{1}{m} \frac{|\alpha_\tau + 2m+1|}{(2(m+1))^{2(n+1)}} \left[\prod_{j=1}^{2m} \left| \frac{\alpha_\tau}{j} + 1 \right| \right] \end{aligned}$$

We have

$$\forall j > \theta_\tau := [\frac{1}{2} + \tau^2 - 1] + 1 : \quad \left| \frac{\alpha_\tau}{j} + 1 \right| < 1 + \frac{1}{j}$$

where $[\frac{1}{2} + \tau^2 - 1]$ is the enter part of $\frac{1}{2} + \tau^2 - 1$. Then

$$\begin{aligned} \prod_{j=\theta_\tau}^{2(m+1)} \left| \frac{\alpha_\tau}{j} + 1 \right| &\leq \prod_{j=\theta_\tau}^{2m} \frac{1+j}{j} = \frac{1}{\theta_\tau} 2(m+2) \left[\prod_{j=\theta_\tau+1}^{2m} \frac{j}{j+1} \right] \\ &\leq \frac{1}{\theta_\tau} 2(m+2) \end{aligned}$$

Then

$$\left| R_{\tau,n}(m) \right| \leq C_{n,\tau} \frac{1}{m} \frac{2(m+2) |\alpha_\tau + 2m + 1|}{(2(m+1))^{2(n+1)}},$$

where

$$C_{n,\tau} := (2n+1) \frac{1}{2\theta_\tau} \left[\prod_{j=1}^{\theta_\tau} \left| \frac{\alpha_\tau}{j} + 1 \right| \right] \max\{|\alpha_\tau|, 1\}.$$

Since $n \geq 0$ we deduce that $\lim_{m \rightarrow +\infty} |R_{\tau,n}(m)| = 0$. Equation (7), can be written as

$$J_{n,\tau} = - \sum_{k=1}^{+\infty} (-1)^k \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}} \right] V_{2(k+1),\tau}(2n+1, 0)$$

In order to simplify the notation denote

$$\rho_{2(k+1)} = \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}},$$

Then

$$J_{n,\tau} = - \sum_{k=0}^{+\infty} (-1)^k \rho_{2(k+1)} V_{2(k+1),\tau}(2n+1, 0)$$

By Equation (8) and Lemma 5

$$\begin{aligned} J_{n,\tau} &= -(2n+1)!(-1)^n \rho_{2(n+1)} \\ &\quad - (2n+1)! \sum_{k=n+2}^{+\infty} (-1)^k \rho_{2(k+1)} \sum_{A \in \Lambda_{2(k-n)}^{2k+1}} (-1)^{2(k-n)} \Pi_{j \in A}(\alpha_\tau + j). \end{aligned}$$

By Equation (6)

$$\begin{aligned} \frac{1}{(2n+1)!} \frac{d^{2n+1}}{dr^{2n+1}} \phi_\tau(0) &= -\frac{3}{2} \alpha_\tau^{-2n-1} - (-1)^n \rho_{2(n+1)} \\ &\quad - \sum_{k=n+2}^{+\infty} (-1)^k \rho_{2(k+1)} \sum_{A \in \Lambda_{2(k-n)}^{2k+1}} (-1)^{2(k-n)} \Pi_{j \in A}(\alpha_\tau + j) \end{aligned}$$

□

5 Proof of the Main result

Proof of the Main result. Let be $\tau \in \mathbb{R}^*$ fixed. By the Proposition 3, we have

$$\begin{aligned} \frac{1}{(2n+1)!} \frac{d^{2n+1}}{dr^{2n+1}} \phi_\tau(0) &= -\frac{3}{2} \alpha_\tau^{-2n-1} - (-1)^n \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(n+1)}} \right] \\ &\quad - \sum_{k=n+2}^{+\infty} (-1)^k \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}} \right] \sum_{A \in \Lambda_{2(k-n)}^{2k+1}} \Pi_{j \in A} (\alpha_\tau + j). \end{aligned}$$

By Lemma 6, we havet

$$\phi_\tau(r) = \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} \frac{d^{2n+1}}{dr^{2n+1}} \phi_\tau(0) r^{2n+1},$$

Then

$$\begin{aligned} \phi_\tau(r) &= -\frac{3}{2} \sum_{n=0}^{\infty} r^{2n+1} \alpha_\tau^{-2n-1} - \sum_{n=0}^{\infty} (-1)^n r^{2n+1} \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(n+1)}} \right] \\ &\quad - \sum_{n=0}^{\infty} r^{2n+1} \sum_{k=n+2}^{+\infty} (-1)^k \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(k+1)}} \right] \sum_{A \in \Lambda_{2(k-n)}^{2k+1}} \Pi_{j \in A} (\alpha_\tau + j), \end{aligned}$$

implies

$$\begin{aligned} \phi_\tau(r) &= -\frac{3}{2} \sum_{n=0}^{\infty} r^{2n+1} \alpha_\tau^{-2n-1} - \sum_{n=0}^{\infty} (-1)^n r^{2n+1} \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(n+1)}} \right] \\ &\quad - \sum_{n=0}^{\infty} (-1)^n \left[\sum_{k=0}^n r^{2k+1} \right] \left[\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^{2(n+1)}} \right] \sum_{A \in \Lambda_2^{2n+3}} \Pi_{j \in A} (\alpha_\tau + j), \end{aligned}$$

since $\alpha_\tau = \frac{1}{2} + i\tau$ and since

$$\sum_{A \in \Lambda_2^{2n+3}} \Pi_{j \in A} (\alpha_\tau + j) = \sum_{0 \leq j \neq j' \leq 2n+3} \left(\frac{1}{2} + j + i\tau \right) \left(\frac{1}{2} + j' + i\tau \right),$$

we get

$$\Im(\phi_\tau(r)) = \tau r \left[\frac{3}{2} \frac{\frac{1}{4} + r^2 + \tau^2}{(\frac{1}{4} - r^2)^2 + 2(\frac{1}{4} - r^2)\tau^2 + \tau^4} - \psi(r) \right],$$

where

$$\psi(r) := \frac{1}{1-r} \sum_{j \in \mathbb{Z}^*} \sum_{n=0}^{\infty} \frac{(-1)^n (1-r^{2n})}{(j2\pi)^{2(n+1)}} \sum_{0 \leq m \neq m' \leq 2n+3} (1+m+m').$$

We have

$$\begin{aligned} \sum_{0 \leq m \neq m' \leq 2n+3} (1+m+m') &= 2(n+2) + \sum_{m=1}^{2(n+1)} (2n+3-m)m \\ &= 2n^2 + 3n - 1 \end{aligned}$$

Since the sequence

$$n \mapsto \frac{(1-r^{2n})}{(j2\pi)^{2(n+1)}} (2n^2 + 3n - 1), \quad n \geq 1,$$

is positive and decreasing for every $r \in (0, \frac{1}{2})$, then

$$\psi(r) = \frac{1}{1-r} \sum_{j \in \mathbb{Z}^*} \sum_{n=0}^{\infty} \frac{(-1)^n (1-r^{2n})}{(j2\pi)^{2(n+1)}} (2n^2 + 3n - 1) < 0, \quad \forall r \in (0, \frac{1}{2}).$$

We deduce that $\tau \Im(\phi_\tau(r)) > 0$ for every $r \in (0, \frac{1}{2})$ and $\tau \in \mathbb{R}^*$. \square

6 Discussion about the Riemann hypothesis

Consider the representation of the Riemann Zeta function ζ defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -s \int_0^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*, \quad (9)$$

and where $\{t\}$ is the fractional part of the real t . Define the τ_* -non trivial zeros of the function ζ in the following sense

Definition 7. Let be $s \in \mathbb{C}$. We say that s is a *non trivial zero of the function ζ* if

$$\zeta(s) = 0 \quad \text{and} \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*.$$

By the Riemann functional equation [[1], page 13 Theorem 2.1], the non trivial zeros of the function ζ are symmetric about the line $\Re(z) = \frac{1}{2}$. By the Main result, the non trivial zeros $s \in \mathbb{C}$ symmetric about the line $\Re(z) = \frac{1}{2}$ satisfies $\Re(s) = \frac{1}{2}$.

References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).