



HAL
open science

Differential equations properties and ψ function

W Oukil

► **To cite this version:**

| W Oukil. Differential equations properties and ψ function. 2022. hal-03581978v5

HAL Id: hal-03581978

<https://hal.science/hal-03581978v5>

Preprint submitted on 5 Apr 2022 (v5), last revised 26 Nov 2023 (v25)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Differential equations properties and ψ function

W. Oukil

Laboratory of Dynamical Systems, Faculty of Mathematics.
University of Science and Technology Houari Boumediene,
BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

April 5, 2022

Abstract

Using the differential equations properties, we study the function ψ which is a function defined by an integral. The particular case of this function coincide with the function $s \mapsto \zeta(s)/s$, where ζ is the Riemann Zeta function. In the end of the paper, we find that $\frac{\zeta(s)}{s} \neq -\frac{\zeta(1-s)}{1-s}$ for every $\Re(s) \in (0, \frac{1}{2})$ and $\Im(s) \in \mathbb{R}^*$. Thanks to the Riemann functional equation, we deduce that any *non-trivial zero* of ζ is of real part equal to $\frac{1}{2}$.

Keywords: Linear differential equation, Zeta function, Riemann hypothesis.

1 Main result

Let be $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a bounded function. We study the function $\psi[\eta] : (0, 1) \times \mathbb{R}^* \rightarrow \mathbb{C}$ defined as

$$\forall x \in (0, 1), \forall \tau \in \mathbb{R}^* : \quad \psi[\eta](x, \tau) := \int_0^{+\infty} \frac{\eta(t)}{t^{1+x+i\tau}} dt, \quad (1)$$

We focus about the function $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$ satisfying the following hypotheses

H_1 : $\exists \omega > 0$: $0 \leq \eta(t) < \omega$ for all $t \geq 0$,

H_2 : there exists an at most countable set $A \subset \mathbb{R}_+$ such that η is C^1 on \mathbb{R}_+/A .

We prove in this paper the following Main result

Main Result. Consider the function ψ defined by Equation (1). Then, for every function $\eta : \mathbb{R}_+ \rightarrow \mathbb{R}$ that satisfies the hypotheses H_1 and H_2 we have

$$\forall \tau \in \mathbb{R}^*, \forall x \in (0, \frac{1}{2}) : \quad \|\psi[\eta](x, \tau) - \psi[\eta](1-x, \tau)\| \neq 0.$$

2 Main proposition

In this Section we study the function ψ , defined in Equation (1), by transforming the problem to some non-homogeneous linear differential equation.

Proposition 1. Let $\eta : \mathbb{R}_+ \mapsto \mathbb{R}$ be a function that satisfies H_1 and H_2 . For every $r \in (0, \frac{1}{2})$ and $\tau > 0$ define the $h_{\tau,r} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ as

$$h_{\tau,r}(t) := \tau \exp(-\frac{1}{2}t)[\eta(\exp(t)) + 1] \sinh(rt), \quad \forall t \geq 0.$$

Then for every $r \in (0, \frac{1}{2})$ and every $\tau > 0$, the unique continuous solution $t \mapsto \delta_{\tau,r}(t)$ of the following linear differential equation

$$\frac{d}{dt}\delta_{\tau,r}(t) = \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \delta_{\tau,r}(t) + \begin{pmatrix} 0 \\ h_{\tau,r}(t) \end{pmatrix}, \quad t \in \mathbb{R}/\ln(A), \quad \delta_{\tau,r}(0) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (2)$$

satisfies

$$\lim_{t \rightarrow +\infty} \|\delta_{\tau,r}(t)\| \neq 0.$$

Lemma 2. For every $t \geq 0$ we have

$$\exp\left(\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} t\right) = \begin{pmatrix} \cos(\tau t) & \frac{1}{\tau} \sin(\tau t) \\ -\tau \sin(\tau t) & \cos(\tau t) \end{pmatrix}.$$

Proof. We have

$$\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix}^2 = -\tau^2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then

$$\begin{aligned} \exp\left(\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} t\right) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \sum_{k=0}^{+\infty} (-1)^k \frac{(\tau t)^{2k}}{(2k)!} \\ &\quad + \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \sum_{k=0}^{+\infty} (-1)^k (\tau)^{2k} \frac{t^{2k+1}}{(2k+1)!}. \end{aligned}$$

Implies

$$\begin{aligned} \exp\left(\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} t\right) &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \cos(\tau t) + \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \frac{1}{\tau} \sin(\tau t) \\ &= \begin{pmatrix} \cos(\tau t) & \frac{1}{\tau} \sin(\tau t) \\ -\tau \sin(\tau t) & \cos(\tau t) \end{pmatrix}. \end{aligned}$$

□

Proof of the Proposition 1. By contradiction, suppose that there exist $r \in (0, \frac{1}{2})$ and $\tau > 0$ such that

$$\lim_{t \rightarrow +\infty} \|\delta_{\tau,r}(t)\| = 0. \quad (3)$$

By the hypotheses H_1 , integrate Equation (2) using the fact $\delta_{\tau,r}$ is the continuous solution and the fact $\delta_{\tau,r}(0) = 0$, we get

$$\delta_{\tau,r}(t) = \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \int_0^t \delta_{\tau,r}(u) du + \begin{pmatrix} 0 \\ \int_0^t h_{\tau,r}(u) du \end{pmatrix}, \quad \forall t \geq 0. \quad (4)$$

Since

$$\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix}^2 = -\tau^2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

then

$$\int_0^t \delta_{\tau,r}(u) du = \frac{-1}{\tau^2} \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \left[\delta_{\tau,r}(t) - \begin{pmatrix} 0 \\ \int_0^t h_{\tau,r}(u) du \end{pmatrix} \right], \quad \forall t \geq 0. \quad (5)$$

By hypothesis H_1 the function η is bounded and positive. Since $r \in (0, \frac{1}{2})$ then there exists $\alpha_r > 0$ such that

$$\lim_{t \rightarrow +\infty} \int_0^t h_{\tau,r}(u) du = \tau \alpha_r.$$

By Equation (5) and the hypothesis (3), we get

$$\lim_{t \rightarrow +\infty} \int_0^t \delta_{\tau,r}(u) du = \frac{1}{\tau^2} \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ \tau \alpha_r \end{pmatrix} = \begin{pmatrix} \frac{1}{\tau} \alpha_r \\ 0 \end{pmatrix} \quad (6)$$

Equation (4), can be written as

$$\delta_{\tau,r}(t) = \begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} \left[\int_0^t \delta_{\tau,r}(u) du - \begin{pmatrix} \frac{1}{\tau} \alpha_r \\ 0 \end{pmatrix} \right] + \begin{pmatrix} 0 \\ -\int_t^{+\infty} h_{\tau,r}(u) du \end{pmatrix}, \quad \forall t \geq 0.$$

Integrate,

$$\int_0^t \delta_{\tau,r}(u) du = \begin{pmatrix} 1 - \cos(\tau t) & \frac{1}{\tau} - \sin(\tau t) \\ \tau \sin(\tau t) & 1 - \cos(\tau t) \end{pmatrix} \begin{pmatrix} \frac{1}{\tau} \alpha_r \\ 0 \end{pmatrix} \quad (7)$$

$$+ \begin{pmatrix} \cos(\tau t) & \frac{1}{\tau} \sin(\tau t) \\ -\tau \sin(\tau t) & \cos(\tau t) \end{pmatrix} \begin{pmatrix} \int_0^t \frac{1}{\tau} \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du \\ -\int_0^t \cos(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du \end{pmatrix}. \quad (8)$$

By definition of the function $h_{\tau,r}$ and by the hypothesis H_1 , the function $u \mapsto \int_u^{+\infty} h_{\tau,r}(v) dv$ is positive and increasing. Using the symmetry of the function $\sin(\tau u)$ on $[0, \frac{2\pi}{\tau}]$ about the point $\frac{\pi}{2\tau}$ we get,

$$\int_0^t \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du > 0, \quad \forall t > 0.$$

In particular,

$$\int_0^t \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du > \int_0^t \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du, \quad \forall t > \frac{2\pi}{\tau}. \quad (9)$$

Then the function

$$t \mapsto \frac{\int_0^t \cos(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du}{\int_0^t \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du}$$

is continuous and uniformly bounded on $(\frac{2\pi}{\tau}, +\infty)$. In addition, the limit

$$\lim_{t \rightarrow +\infty} \frac{\int_0^t \cos(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du}{\int_0^t \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du}$$

exists. Then there exists sequence $t_n \rightarrow +\infty$ such that

$$\frac{\cos(\tau t_n)}{\sin(\tau t_n)} := \frac{\int_0^{t_n} \cos(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du}{\int_0^{t_n} \sin(\tau u) \int_u^{+\infty} h_{\tau,r}(v) dv du},$$

and such that

$$\lim_{n \rightarrow +\infty} [t_n - (\frac{n\pi}{\tau} + \frac{\pi}{2\tau})] \neq 0, \quad (10)$$

Implies that, there exists a sub-sequence $(t_{n_k})_k \subset (t_n)_n$ such that $\lim_{k \rightarrow +\infty} \cos(\tau t_{n_k}) := \alpha \neq 0$. By Equation (7), we get

$$\lim_{k \rightarrow +\infty} \int_0^{t_{n_k}} \delta_{2,\tau,r}(u) du = \lim_{k \rightarrow +\infty} \frac{1}{\tau} \alpha_r \left(1 - \cos(\tau t_{n_k}) \right) \neq \frac{1}{\tau} \alpha_r,$$

where $(\delta_{1,\tau,r}, \delta_{2,\tau,r}) := \delta_{\tau,r}$. Contradiction with Equation (6). \square

3 Proof of the Main results

Proof of the Main result. Integrate Equation (2) using the fact $\delta_{\tau,r}$ is the continuous solution and the fact $\delta_{\tau,r}(0) = 0$, we get

$$\delta_{\tau,r}(t) = \int_0^t \exp \left(\begin{pmatrix} 0 & 1 \\ -\tau^2 & 0 \end{pmatrix} (t-u) \right) \begin{pmatrix} 0 \\ h_{\tau,r}(u) \end{pmatrix} du.$$

By Lemma 2, we obtain,

$$\delta_{\tau,r}(t) = \begin{pmatrix} \cos(\tau t) & \frac{1}{\tau} \sin(\tau t) \\ -\tau \sin(\tau t) & \cos(\tau t) \end{pmatrix} \begin{pmatrix} -\int_0^t \frac{1}{\tau} \sin(\tau u) h_{\tau,r}(u) du \\ \int_0^t \cos(\tau u) h_{\tau,r}(u) du \end{pmatrix}.$$

Use the change of variable, $t \rightarrow \exp(u)$, in Equation (1), we obtain

$$\begin{pmatrix} \frac{1}{\tau} \Im[\psi(\frac{1}{2} - r, \tau) - \psi(\frac{1}{2} + r, \tau)] \\ \Re[\psi(\frac{1}{2} - r, \tau) - \psi(\frac{1}{2} + r, \tau)] \end{pmatrix} = \lim_{t \rightarrow +\infty} \begin{pmatrix} \cos(\tau t) & \frac{1}{\tau} \sin(\tau t) \\ -\tau \sin(\tau t) & \cos(\tau t) \end{pmatrix}^{-1} \delta_{\tau,r}(t).$$

By the Main Proposition 1 we have $\|\psi(\frac{1}{2} - r, \tau) - \psi(\frac{1}{2} + r, \tau)\| \neq 0$ for every $r \in (0, \frac{1}{2})$ and $\tau \in \mathbb{R}^*$. \square

4 Discussion about the Riemann hypothesis

Consider the representation of the Riemann Zeta function ζ defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -s \int_0^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*, \quad (11)$$

and where $\{t\}$ is the fractional part of the real t . Define the *non-trivial zeros* of the function ζ in the following sense

Definition 3. Let be $s \in \mathbb{C}$. We say that s is a *non-trivial zero* of the function ζ if

$$\zeta(s) = 0 \quad \text{and} \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*.$$

Denote $\eta_*(t) := \{t\}$. For every $\tau \in \mathbb{R}^*$ and every $x \in (0, 1)$ we have

$$\psi[\eta_*](x, \tau) = \frac{\zeta(s)}{s} \quad \text{and} \quad \psi[\eta_*](1-x, \tau) = -\frac{\zeta(1-s)}{1-s}, \quad s := x + i\tau.$$

Since

$$0 \leq \eta_*(t) \leq 1, \quad \forall t \geq 0,$$

Then η_* satisfies the hypotheses H_1 . The function η_* is C^1 on \mathbb{R}_+/\mathbb{N} . Then η_* satisfies the hypotheses H_2 . By the Riemann functional equation [[1], page 13 Theorem 2.1], the non-trivial zeros of the function ζ are symmetric about the line $\Re(z) = \frac{1}{2}$. By the Main result, the non-trivial zeros $s \in \mathbb{C}$ symmetric about the line $\Re(z) = \frac{1}{2}$ satisfies $\Re(s) = \frac{1}{2}$.

References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).