



HAL
open science

A new kinetics database of organic oxidation reactions for use in structure-activity relationship studies

M R McGillen, J J Orlando, A Mellouki, B Picquet-Varrault, T J Wallington,
W P L Carter

► To cite this version:

M R McGillen, J J Orlando, A Mellouki, B Picquet-Varrault, T J Wallington, et al.. A new kinetics database of organic oxidation reactions for use in structure-activity relationship studies. 11th International Conference on Chemical Kinetics, Jun 2019, Orléans, France. hal-03582140

HAL Id: hal-03582140

<https://hal.science/hal-03582140>

Submitted on 2 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new kinetics database of organic oxidation reactions for use in structure-activity relationship studies

M.R. McGillen^{1,2}, J.J. Orlando³, A. Mellouki¹, B. Picquet-Varrault⁴, T.J. Wallington⁵, W.P.L. Carter⁶


¹CNRS-INSIS, Institut de Combustion, Aérothermique, Réactivité et Environnement 1C, Avenue de la recherche scientifique, 45071 Orléans cedex 2, France. ²Le Studium Loire Valley Institute for Advanced Studies, Orléans, France. ³Atmospheric Chemistry Observations and Modeling Laboratory, National Center for Atmospheric Research, Boulder, CO, 80307, USA. ⁴Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR 7583 CNRS, Universités Paris-Est Créteil et Paris Diderot, Institut Pierre-Simon Laplace, Créteil Cedex, France. ⁵Research & Advanced Engineering, Ford Motor Company, Dearborn, MI, 48121-2053, USA. ⁶College of Engineering, Center for Environmental Research and Technology (CE-CERT), University of California, Riverside, CA, 92521, USA.

Introduction:

Several compendia of kinetic data exist, including thoroughly evaluated datasets provided by the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation,¹⁻⁸ the JPL Panel for Data Evaluation,⁹ and the Calvert et al. reviews.¹⁰⁻¹⁴ The NIST Chemical Kinetics database¹⁵ is another extensive (unevaluated) database. Despite the many useful aspects of these resources, each possess several drawbacks:


1. The frequency with which kinetic data are published is much faster than that with which these data are compiled and reviewed.
2. The number of reaction rates that have been evaluated is much smaller than the number that have been determined. It is therefore inevitable that such evaluations do not currently capture the full chemical diversity available in the experimental literature.
3. The data within these reviews are currently not downloadable in a digital, searchable format.
4. Non-downloadable databases that cannot be accessed offline are subject to downtime (e.g. NIST was recently out of commission for 38 days as a consequence of the 2018–19 US federal government shutdown)

Methods:



Decision tree:
The process by which our panel reviews kinetic data.



Temperature parameterization:
Where extensive temperature dependent data are reported, new parameterizations are provided, which usually take the form of Kooij's equation, such is the case for dimethyl ether.



Temperature dependence:
Although a large amount of temperature-dependent data is available, the majority of species are only represented by room temperature rate coefficients. This is especially true of the NO₃ dataset, where temperature dependence has been reported for only 12% of compounds.

Overlap with atmospheric species:
An Euler diagram¹⁶ showing the number of compounds formed from oxidation of α -pinene and *n*-octane according to the automatic mechanism generator of GECKO-A.¹⁷ The blue circle represents the total number of compounds for which a rate coefficient has been measured. It is obvious that there are relatively few compounds that have measurements, and that the overlap is small.



Free energy relationships:
Free energy relationships of reaction rates between oxidants. Strong relationships are observed in some functionalities and oxidants (e.g. alkanes/ethers/halocarbons in the OH–Cl comparison). Sometimes very different relationships are observed depending on the functionality (e.g. alkanes vs other compounds in the NO₃–Cl comparison).



Relative coverage of databases:
Euler diagrams,¹⁶ where the size of each curve is proportional to the number of species studied with respect to each oxidant, the number of compounds available in each review, and the overlap between these reviews

Database statistics:
Number of measurements, temperature dependence and functionality are three criteria that vary a lot between oxidants. *hydrocarbons with no higher-order bonds are considered as 0, for halocarbons with several X substitutions, these are grouped as a single substitution.

Oxidant	# species	# Arrhenius	# Kooij	# functional groups*						
				0	1	2	3	4	5	6
OH	1354	464	70	73	467	572	190	38	13	1
O ₃	310	60	0	0	91	171	39	7	2	0
NO ₃	389	47	0	18	154	161	46	9	1	0
Cl	708	156	3	33	271	333	56	9	5	1



Conclusions:

1. This new database provides an updated and near-exhaustive collection of rate coefficients for the reactions of OH, Cl, NO₃ and O₃ with organic compounds.
2. This new resource will soon be available online and will be updated as new data become available.
3. Further improvements and extensions to this database are envisaged, including data for site-of-attack and reactions of other oxidants, and further experimental metadata where available.

References:

- (1) Ammann, M. et al. *Atmos. Chem. Phys.* **2013**, *13* (16), 8045–8228. <https://doi.org/10.5194/acp-13-8045-2013>.
- (2) Atkinson, R. et al. *Atmos. Chem. Phys.* **2004**, *4* (6), 1461–1738. <https://doi.org/10.5194/acp-4-1461-2004>.
- (3) Atkinson, R. et al. *Atmos. Chem. Phys.* **2006**, *6* (11), 3625–4055. <https://doi.org/10.5194/acp-6-3625-2006>.
- (4) Atkinson, R. et al. *Atmos. Chem. Phys.* **2007**, *7* (4), 981–1191. <https://doi.org/10.5194/acp-7-981-2007>.
- (5) Atkinson, R. et al. *Atmos. Chem. Phys.* **2008**, *8* (15), 4141–4496. <https://doi.org/10.5194/acp-8-4141-2008>.
- (6) Crowley, J. N. et al. *Atmos. Chem. Phys.* **2010**, *10* (18), 9059–9223. <https://doi.org/10.5194/acp-10-9059-2010>.
- (7) Crowley, J. N. et al. *Atmos. Chem. Phys.* **2013**, *13* (15), 7359–7359. <https://doi.org/10.5194/acp-13-7359-2013>.
- (8) IUPAC. Task Group on Atmospheric Chemical Kinetic Data Evaluation <http://iupac.pole-ether.fr/>.
- (9) Burkholder, J. B. et al. Evaluation No. 18. JPL Publication 15-10, Jet Propulsion Laboratory, Pasadena 2015.
- (10) Calvert, J. G. et al. *The Mechanisms of Atmospheric Oxidation of the Alkenes*; Oxford University Press: New York, 2000.
- (11) Calvert, J. G. et al. *The Mechanisms of Atmospheric Oxidation of Aromatic Hydrocarbons*; Oxford University Press: New York, 2002.
- (12) Calvert, J. G. et al. *Mechanisms of the Atmospheric Oxidations of the Alkanes*; Oxford University Press, 2008.
- (13) Calvert, J. G. et al. *The Mechanisms of the Atmospheric Oxidations of the Oxygenates*; Oxford University Press, 2011.
- (14) Calvert, J. G. et al. *The Mechanisms of Reactions Influencing Atmospheric Ozone*; Oxford University Press: New York, 2015.
- (15) Manion, J. A. et al. NIST Chemical Kinetics Database, Standard Reference Database 17, Version 7.0 (Web Version), Release 1.6.8 Data Version 2015.09, A compilation of kinetics data on gas-phase reactions <http://kinetics.nist.gov>.
- (16) Micallef, L. et al. *PLoS ONE* **2014**, *9* (7), e101717. <https://doi.org/10.1371/journal.pone.0101717>.
- (17) Aumont, B. et al. *Atmos. Chem. Phys.* **2005**, *5* (9), 2497–2517. <https://doi.org/10.5194/acp-5-2497-2005>.