



**HAL**  
open science

## The SFiNx Detector System

A.V. Isaev, A.V. Yerebin, N.I. Zamyatin, I.N. Izosimov, A.A. Kuznetsova,  
O.N. Malyshev, R.S. Mukhin, A.G. Popeko, Yu. A. Popov, B. Sailaubekov, et  
al.

► **To cite this version:**

A.V. Isaev, A.V. Yerebin, N.I. Zamyatin, I.N. Izosimov, A.A. Kuznetsova, et al.. The SFiNx Detector System. Phys.Part.Nucl.Lett., 2022, 19 (1), pp.37-45. 10.1134/S154747712201006X . hal-03584718

**HAL Id: hal-03584718**

**<https://hal.science/hal-03584718>**

Submitted on 7 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Study of Spontaneous Fission Using the SFiNx System

A.V. ISAEV<sup>a</sup>, A.V. YEREMIN<sup>a,b</sup>, N.I. ZAMYATIN<sup>a</sup>, I.N. IZOSIMOV<sup>a</sup>,  
A.A. KUZNETSOVA<sup>a</sup>, O.N. MALYSHEV<sup>a,b</sup>, R.S. MUKHIN<sup>a</sup>,  
A.G. POPEKO<sup>a,b</sup>, YU.A. POPOV<sup>a,b</sup>, B. SAILAUBEKOV<sup>a,c</sup>,  
A.I. SVIRIKHIN<sup>a,b</sup>, E.A. SOKOL<sup>a</sup>, M.S. TEZEKBAYEVA<sup>a,c</sup>, D.A. TESTOV<sup>a</sup>,  
M.L. CHELNOKOV<sup>a</sup>, V.I. CHEPIGIN<sup>a</sup>, S. ANTALIC<sup>d</sup>, P. MOSAT<sup>d</sup>,  
P. BRIONNET<sup>e</sup>, B. GALL<sup>e</sup>, O. DORVAUX<sup>e</sup>, K. KESSACI<sup>e</sup>, A. SELLAM<sup>e</sup>,  
A. LOPEZ-MARTENS<sup>f</sup>, K. HAUSCHILD<sup>f</sup>

<sup>a</sup> Joint Institute for Nuclear Research, Dubna, Russia

<sup>b</sup> Dubna State University, Dubna, Russia

<sup>c</sup> Institute of Nuclear Physics, Almaty, Republic of Kazakhstan

<sup>d</sup> Comenius University, Bratislava, Slovakia

<sup>e</sup> IPHC, IN2P3-CNRS, Strasbourg, France

<sup>f</sup> IJCLab, IN2P3-CNRS, Université Paris-Saclay, Orsay, France

A new detector system consisting of 116 <sup>3</sup>He neutron counters and DSSSD array was created. This system can be used to study spontaneous fission properties of short-lived heavy isotopes. The single neutron registration efficiency of the counter assembly was measured as 54.7±0.1% with <sup>248</sup>Cm-source. Using the setup, new data were obtained for prompt neutron yields of <sup>252</sup>No isotope.

PACS numbers: 25.85.Ca; 25.70.-z; 27.90.+b; 29.40.Cs; 29.40.Gx.

### 1. Introduction

The spontaneous fission process defines the boundaries of the existence of heavy elements. At present, there is no complete nuclear fission model, and new experimental data are needed more than ever to improve the theoretical approaches.

Our group has accumulated significant experience in the study of the decay properties of heavy isotopes using VASSILISA [1] and SHELS [2] recoil separators. Important data were obtained for neutron deficient isotopes of Fm [3, 4], No [5, 6] and Rf [7, 8] using a detector [9].

The detector consisted of 54 <sup>3</sup>He neutron counters and a silicon detector array with a 60×60 mm<sup>2</sup> focal plane detector. To increase the collection of

recoil nuclei by a factor of 3, we decided to install a new DSSSD assembly [10] inside the neutron detector with a large  $100 \times 100 \text{ mm}^2$  focal plane detector. It was necessary to completely change the counter's geometry.

## 2. Detector




Fig. 1. Photograph of the SFiNx system. A schematic front view of the assembly is shown in the inset image.

Table 1. The main properties and number of the different types of counters used in the detector. Layers are numbered relative to the central axis of the detector.

Layer	Number	Diameter, mm	Active length, mm	Operating voltage, V	$^3\text{He}$ pressure, atm
1	20	32	530	1400	7
2	24	32	460	1775	
3	28	30	585	1500	
4	44				

The new detection system was named SFiNx (**S**pontaneous **F**ission, **N**eutrons and **x**-rays, fig. 1). The system consists of 116  $^3\text{He}$ -counters of

neutrons that surround the vacuum chamber with DSSSD. The characteristics and number of  $^3\text{He}$  counters are given in the table 1.

The counters are placed in a polyethylene moderator in the form of a regular octagonal prism (height 630 mm, circumscribed circle diameter 762 mm). The outer side of the detector is covered with a 5%-borated polyethylene shield (thickness from 30 to 50 mm) to protect against background neutrons.

The single neutron registration efficiency measured with  $^{248}\text{Cm}$ -source is  $54.7\pm 0.1\%$  (nonlinear least-squares method error). The average neutron lifetime in the setup is  $18.4\pm 0.2\ \mu\text{s}$  (exponent fitting error).

An array of DSSSD detectors inside the vacuum chamber is represented by a focal plane  $128\times 128$ -strip detector ( $100\times 100\ \text{mm}^2$  area,  $500\ \mu\text{m}$  thickness) and 8 side  $16\times 16$ -strip detectors ( $50\times 60\ \text{mm}^2$  area,  $700\ \mu\text{m}$  thickness) [10]. Silicon detectors allow registering  $\alpha$ -,  $\beta$ -particles and fission fragments. Energy resolutions were measured using 8 MeV  $\alpha$ -particles as 20 keV and 30 keV for the focal and side silicon detectors respectively.

### 3. Experiment

An experiment to study  $^{252}\text{No}$  spontaneous fission was carried out on a velocity filter SHELS [2]. The complete fusion reaction of accelerated  $^{48}\text{Ca}^{18+}$  ions with  $^{206}\text{Pb}$  target nuclei was used. The thickness of the  $^{206}\text{PbS}$  target was  $350\ \mu\text{g}/\text{cm}^2$  (on  $1.5\ \mu\text{m}$  Ti backing) and the  $^{206}\text{Pb}$  enrichment was about 97%. The beam energy in the middle of the target was about  $215\pm 2\ \text{MeV}$  (2n-evaporation channel maximum  $\sim 800\ \text{nb}$ ).

After passing through the separator, the recoil nuclei were implanted into the focal plane large DSSSD. The spontaneous fission branching fraction for  $^{252}\text{No}$  is about 32%. When a fission fragment was detected in the focal DSSSD, the coincidence with the recoil nucleus that fell into the same pixel of the detector in the time window 1 ms – 25 s ( $\sim 10\ T_{1/2}$ ) was checked. After that, a search for neutrons was carried out in the time window of 128  $\mu\text{s}$ . A total of 3260 reliable events of  $^{252}\text{No}$  spontaneous fission were found.

The half-life of  $^{252}\text{No}$  was determined as  $2.44\pm 0.05\ \text{s}$  (exponential fitting error) based on «recoil nucleus – fission fragment» correlations. The received half-life is in good agreement with the already measured values  $2.44\pm 0.04\ \text{s}$  [11] and  $2.30\pm 0.22\ \text{s}$  [12].

The prompt neutron multiplicity distribution of  $^{252}\text{No}$  was measured and then true distribution was restored by using the Tikhonov statistical regularisation method [13, 14, 15] (fig. 3). The dispersion of the measured neutron multiplicity distribution is  $\sigma_{\bar{\nu}}^2 = 2.1$ , whereas the dispersion of restored distribution is  $\sigma_{\bar{\nu}r}^2 = 2.2$ . The average number of neutrons per spontaneous fission act of  $^{252}\text{No}$  was obtained as  $\bar{\nu} = 4.25\pm 0.09$  (contains the statistical

error and the efficiency error of the neutron detector) and is in good agreement with previously measured values  $4.15 \pm 0.30$  [16],  $4.43 \pm 0.45$  [17], and  $4.51 \pm 0.25$  [6]. The restored average number of neutrons per fission act is  $\bar{\nu}_r = 4.27 \pm 0.15$ .




Fig. 2. Measured (squares) and restored (circles) prompt neutron multiplicity distributions of  $^{252}\text{No}$  and background neutron spectra (triangles). Lines that connect the points have been added for clarity.

The background neutron multiplicity was estimated during the search for random coincidences of neutrons with  $^{252}\text{No}$   $\alpha$ -particles in a time window of  $128 \mu\text{s}$ .

#### 4. Conclusion

The SFiNx setup contains 116  $^3\text{He}$ -filled proportional counters of neutrons and a DSSSD array has been created.

The new large focal plane DSSSD ( $100 \times 100 \text{ mm}^2$ ) allowed to significantly increase the speed of statistics collection in comparison with our old neutron detector [9]. At the same time, the higher neutrons registration efficiency ( $54.7 \pm 0.1\%$ ) of counter assembly allows us to better reconstruct the neutron multiplicity distributions by the statistical regularisation method.

As a result of the first experiment, new data were obtained on the prompt neutron yields from  $^{252}\text{No}$  spontaneous fission.

The system currently has no analogs in the study of the spontaneous fission properties of short-lived heavy nuclei.

## 5. Acknowledgments

This work was supported by the Russian Foundation for Basic Research (project №18-52-15004).

## References

- [1] A.V. Yereimin et al. “The electrostatic separator VASSILISSA Performance and experimental results”. In: *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 126.1 (1997). International Conference on Electromagnetic Isotope Separators and Techniques Related to Their Applications, pp. 329–333. ISSN: 0168-583X. DOI: [https://doi.org/10.1016/S0168-583X\(96\)01002-6](https://doi.org/10.1016/S0168-583X(96)01002-6).
- [2] A.G. Popeko et al. “Separator for Heavy Element Spectroscopy – velocity filter SHELS”. In: *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 376 (2016). Proceedings of the XVIIth International Conference on Electromagnetic Isotope Separators and Related Topics (EMIS2015), Grand Rapids, MI, U.S.A., 11-15 May 2015, pp. 140–143. ISSN: 0168-583X. DOI: <https://doi.org/10.1016/j.nimb.2016.03.045>.
- [3] A.I. Svirikhin et al. “The emission of prompt neutrons from the spontaneous fission of  $^{252}\text{No}$  and  $^{244}\text{Fm}$ .” In: *The European Physical Journal A* 48.9 (2012), pp. 121–127. DOI: <https://doi.org/10.1140/epja/i2012-12121-7>.
- [4] A.I. Svirikhin et al. “Neutron multiplicity at spontaneous fission of  $^{246}\text{Fm}$ .” In: *The European Physical Journal A* 44.3 (2010), pp. 393–396. DOI: <https://doi.org/10.1140/epja/i2010-10968-0>.
- [5] A.I. Svirikhin et al. “Characteristics of spontaneous fission of  $^{250}\text{No}$ ”. In: *Physics of Particles and Nuclei Letters* 14.4 (2017), pp. 571–575. DOI: <https://doi.org/10.1134/S1547477117040161>.
- [6] A.V. Isaev et al. “Comparative Study of Spontaneous-Fission Characteristics of  $^{252}\text{No}$  and  $^{254}\text{No}$  Isotopes”. In: *Physics of Particles and Nuclei Letters* 18.4 (2021), pp. 449–456. DOI: <https://doi.org/10.1134/S1547477121040087>.
- [7] A.I. Svirikhin et al. “Prompt neutrons from spontaneous  $^{254}\text{Rf}$  fission.” In: *Physics of Particles and Nuclei Letters* 16.6 (2019), pp. 768–771. DOI: <https://doi.org/10.1134/S1547477119060311>.
- [8] A.I. Svirikhin et al. “Spontaneous fission of  $^{256}\text{Rf}$ , new data.” In: *Physics of Particles and Nuclei Letters* 13.4 (2016), pp. 480–482. DOI: <https://doi.org/10.1134/S1547477116040129>.

- [9] A.I. Svirikhin et al. “A detector for studying the characteristics of spontaneous fission of short-lived heavy nuclei.” In: *Instruments and Experimental Techniques* 54.5 (2011), pp. 644–648. DOI: <https://doi.org/10.1134/S0020441211040154>.
- [10] R. Chakma et al. “Gamma and conversion electron spectroscopy using GABRIELA”. In: *The European Physical Journal A* 56.10 (2020), p. 245. DOI: <https://doi.org/10.1140/epja/s10050-020-00242-5>.
- [11] Yu.Ts. Oganessian et al. “Measurements of cross sections for the fusion-evaporation reactions  $^{204,206,207,208}\text{Pb}+^{48}\text{Ca}$  and  $^{207}\text{Pb}+^{34}\text{S}$  : Decay properties of the even-even nuclides  $^{238}\text{Cf}$  and  $^{250}\text{No}$ ”. In: *Physical Review C* 64 (5 2001), p. 054606. DOI: [10.1103/PhysRevC.64.054606](https://doi.org/10.1103/PhysRevC.64.054606).
- [12] C.E. Bemis et al. “Fragment-mass and kinetic-energy distributions from the spontaneous fission of  $^{252}\text{No}$ ”. In: *Physical Review C* 15 (2 1977), pp. 705–712. DOI: [10.1103/PhysRevC.15.705](https://doi.org/10.1103/PhysRevC.15.705).
- [13] V.F. Turchin. “Solution of the Fredholm equation of the first kind in a statistical ensemble of smooth functions”. In: *USSR Computational Mathematics and Mathematical Physics* 7.6 (1967), pp. 79–96. ISSN: 0041-5553. DOI: [https://doi.org/10.1016/0041-5553\(67\)90117-6](https://doi.org/10.1016/0041-5553(67)90117-6).
- [14] V.F. Turchin, V.P. Kozlov, and M.S. Malkevich. “THE USE OF MATHEMATICAL-STATISTICS METHODS IN THE SOLUTION OF INCORRECTLY POSED PROBLEMS”. In: *Soviet Physics Uspekhi* 13.6 (1971), pp. 681–703. DOI: [10.1070/pu1971v013n06abeh004273](https://doi.org/10.1070/pu1971v013n06abeh004273).
- [15] R.S. Mukhin et al. “Reconstruction of Spontaneous Fission Neutron Multiplicity Distribution Spectra by the Statistical Regularization Method”. In: *Physics of Particles and Nuclei Letters* 18.4 (2021), pp. 439–444. DOI: <https://doi.org/10.1134/S1547477121040130>.
- [16] Yu.A. Lazarev et al. “Multiplicity of prompt neutrons from spontaneous fission of the isotope  $^{252}\text{102}$ ”. In: *Physics Letters B* 52.3 (1974), pp. 321–324. ISSN: 0370-2693. DOI: [https://doi.org/10.1016/0370-2693\(74\)90052-5](https://doi.org/10.1016/0370-2693(74)90052-5).
- [17] A.V. Yeremin et al. “Neutron detector at the focal plane of the set up VASSILISSA”. In: *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 539.1 (2005), pp. 441–444. ISSN: 0168-9002. DOI: <https://doi.org/10.1016/j.nima.2004.09.043>.