



HAL
open science

PREDOMOS study, impact of a social intervention program for socially isolated elderly cancer patients: update to the study protocol for a randomized controlled trial

Anne-Laure Couderc, Emilie Nouguerède, Karine Baumstarck, Sandrine Loubière, Hervé Le Caer, Olivier Guillem, Frédérique Rousseau, Laurent Greillier, Emmanuelle Norguet-Monnereau, Maud Cecile, et al.

► **To cite this version:**

Anne-Laure Couderc, Emilie Nouguerède, Karine Baumstarck, Sandrine Loubière, Hervé Le Caer, et al.. PREDOMOS study, impact of a social intervention program for socially isolated elderly cancer patients: update to the study protocol for a randomized controlled trial. *Trials*, 2019, 20 (1), pp.54. 10.1186/s13063-018-3127-0 . hal-03585231

HAL Id: hal-03585231

<https://hal.science/hal-03585231v1>

Submitted on 18 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UPDATE

Open Access



PREDOMOS study, impact of a social intervention program for socially isolated elderly cancer patients: update to the study protocol for a randomized controlled trial

Anne-Laure Couderc^{1,2}, Emilie Nouguerède^{1,3*} , Karine Baumstarck⁴, Sandrine Loubière⁴, Hervé Le Caer⁵, Olivier Guillem^{6,2}, Frédérique Rousseau^{7,2}, Laurent Greillier^{8,2}, Emmanuelle Norguet-Monnereau^{3,2}, Maud Cecile^{7,2}, Rabia Boulahssass⁹, Françoise Le Caer^{10,2}, Sandrine Tournier^{11,2}, Chantal Butaud^{12,2}, Pierre Guillet^{12,2}, Sophie Nahon^{13,2}, Sylvie Kirscher^{14,2}, Nadine Diaz¹⁵, Claire Morando¹⁶, Patrick Villani¹, Pascal Auquier⁴ and Aurélie Daumas¹

Abstract

Background: Social isolation potentiates the risk of death by cancer in the older cancer patient population. The PREDOMOS study investigates the impact of establishing a Program of Social intervention associated with techniques of Domotic and Remote assistance on the improvement of quality of life of older isolated patients, treated for locally advanced or metastatic cancer. This paper updates the pilot trial protocol.

Methods/design: The original protocol was published in *Trials*, accessible at <https://trialsjournal.biomedcentral.com/articles/10.1186/s13063-017-1894-7>.

This update reports on the eligibility criteria expansion and on the adjunction of a cost-utility analysis. We widened the eligible population to patients with locally advanced or metastatic cancer including malignant hemopathies (except acute myeloid leukemia) and to patients in the first and second lines of oncologic treatment. We restricted the inclusion to patients with a Mini Mental State Examination score strictly over 24. In addition to the secondary outcomes outlined in the protocol, a medico-economic analysis has been added to evaluate both the health benefits and costs of the two strategies and calculate the incremental cost-utility ratio of the innovative program assessed, compared to the standard practice.

Trial registration: ClinicalTrials.gov, [NCT02829762](https://clinicaltrials.gov/ct2/show/study/NCT02829762). Registered on 29 June 2016.

Keywords: Social intervention program, Techniques of domotic and remote assistance, Older patient, cancer, Oncogeriatrics, Randomized controlled trial, Cost effectiveness

Background

PREDOMOS (PREcarious elderly patient supported for cancer: impact on quality of life of a DOMotic and remote assistance approach for Socially isolated elderly patients supported for locally advanced or metastatic cancer) is a

randomized controlled trial aiming to evaluate the impact of establishing a Program of Social intervention associated with techniques of Domotic and Remote assistance (PS-DR) on the improvement of quality of life (QoL) at 3 months (primary endpoint) of older isolated patients, treated for locally advanced or metastatic cancer. This study will also assess the effect of the PS-DR on treatment compliance, autonomy maintenance, nutrition, mobility, toxicity, time to treatment failure, survival, social isolation, unexpected hospitalizations, and QoL at 6 months (secondary endpoints). In addition, PREDOMOS aims to

* Correspondence: emilie.nouguerede@ap-hm.fr

¹Service de Médecine Interne Gériatrie et Thérapeutique, CHU Sainte Marguerite, Assistance Publique des Hôpitaux de Marseille (AP-HM), 270 Boulevard de Sainte Marguerite Dromel, 13274 Marseille, cedex 09, France
³Service d'Oncologie Digestive, CHU Timone, AP-HM, 264 Rue Saint Pierre, 13385 Marseille, cedex 05, France

Full list of author information is available at the end of the article



identify individual geriatric characteristics (such as functional, nutritional, cognitive, and mobility status) that would be predictive of clinical outcomes.

As a reminder, 320 individuals are required to obtain 90% power to detect a 10-point difference (standard deviation 25) in QoL score between the two groups (20% loss to follow-up patients expected). Participants are randomized in a 1:1 allocation ratio, either in the control group, receiving usual care, or in the experimental group, receiving usual care associated to PS-DR.

The PREDOMOS study, supported by a grant from the French Ministry of Health, is coordinated by the Assistance Publique des Hôpitaux de Marseille (AP-HM, France) and includes multiple partners in oncologic and geriatric units across France, such as the South Alpes Intercommunal Hospital Center in Gap, the Saint Joseph Hospital and the Paoli Calmettes Institute in Marseille, the Nice University Hospital, the Saint Brieuc, Toulon, and Aix-en-Provence Hospital Centers, and the Saint Catherine Institute in Avignon.

The original study protocol was published by Crétel-Durand et al. [1]. Following the publication of the study protocol, amendments were made to reflect the changes to initial screening and patient recruitment and to add a cost-utility analysis, funded by the Fondation de France. These two changes are outlined below. The amendments of the protocol were approved by the Sud Méditerranée 1 Ethics Committee in June 2018.

Methods/design

Changes in eligibility criteria

To improve patient recruitment, which was difficult from the beginning of the study, the steering committee decided to modify the inclusion criteria. Tumor type restrictions were lifted, allowing patients with locally advanced or metastatic cancer from all organs, including malignant hemopathies, to enter the study. Only patients with acute leukemia will be excluded, in the hope of including patients with a lifespan of more than 6 months. Furthermore, the treatment conditions selected were expanded to include patients cared for in a second line of oncologic treatment. Due to the inclusion of patients with various cancers, we also specified allowed oncological treatments: new generation hormone therapy, targeted therapy, or immunotherapy with or without, concomitant or not, radiotherapy.

Otherwise, to simplify the exclusion criteria, the steering committee added a new inclusion criterion to the protocol, “Mini Mental State Examination (MMSE) score strictly over 24,” to automatically exclude patients unable to complete the QoL questionnaire without help and patients with dementia. We also decided to delete the exclusion criterion “patient with psychiatric troubles”

because in the absence of cognitive impairment these patients are potential candidates like the others.

The inclusion criteria now include the following: (1) patient aged 70 years or older; (2) socially isolated or at risk of social isolation (modified Medical Outcomes Study Social Support Survey (m-MOS-SS) under 80% and/or isolated patient (caregiver living more than 50 km from the patient) and/or primary caregiver to his/her spouse (for a spouse who has a limited autonomy, neurodegenerative disease, cancer, etc., implying regular medical care for at least 3 months); (3) Scale for detection of geriatric frailties in cancer patients over 70 years old (G8 ONCODAGE) score ≤ 14 ; (4) Eastern Cooperative Oncology Group Performance Status (ECOG-PS) ≤ 2 ; (5) Activities of daily living (ADLs) score ≥ 4 ; (6) MMSE > 24 ; (7) lifespan > 6 months; (8) with locally advanced or metastatic cancer, including malignant hemopathies (except acute leukemia); (9) in first or second line of oncologic treatment by chemotherapy, new generation hormone therapy, targeted therapy, or immunotherapy with or without, concomitant or not, radiotherapy; (10) signed informed consent; and (11) affiliation to the French social security system.

The exclusion criteria include: (1) patients with a second cancer; (2) protected adults under guardianship or curatorship; and (3) patient to be immediately directed into a rehabilitation and recuperative care service to benefit from treatment.

Secondary endpoints: adjunction of a cost-utility analysis (CUA)

Home medical care for older cancer patients can be a relevant response to the issue of increasing expenditures related to aging. Indeed, studies have shown that these approaches could be cost-effective; namely, they were effective on a set of clinical criteria at an acceptable cost to the community or the health care system [2–5]. For example, following an initial cost of implementation and equipment, home care interventions are accompanied by avoided costs of medical care (e.g., non-programmed re-hospitalizations, visits, examinations), as well as costs related to the financial burden for the patient. Consequently, cost analyses and cost-effectiveness analyses are crucial for planning proper distribution of health care resources and social/medical interventions in elderly cancer patient populations.

The aim of the economic analysis in the present study is to evaluate both the health benefits and costs of the two strategies and to calculate the incremental cost-effectiveness ratio of the innovative program assessed compared to the standard practice. As recommended by the French Health institution (HAS) [6], the clinical endpoint will be the number of quality-adjusted life years (QALYs) saved. These QALYs are determined by

multiplying life years gained by a value of the utility associated (corresponding to patient's quality of life) during the period under consideration [7]. As the scores in Quality of Life Questionnaire-Cancer (QLQ)-C30 are not utility-based, these scores will be mapped to the EuroQoL Five-levels (EQ-5D) [8, 9]. The QLQ-C30 at T3 will be used to cover the first 3 months (T0 to T3) and the QLQ-C30 at T6 to cover the period between T3 and T6. The time horizon will cover the period during which patients will be exposed to the intervention (i.e., 6 months). This will allow researchers to observe and collect prospectively all health and economic outcomes during the exposed period between the two groups. Therefore, no discounting will be necessary.

The perspective will be that of the society, and the following type of resources will be included in the cost analysis:

- Techniques of domotic and remote assistance (including the cost of equipment installation, renting, technical support, and uninstallation)
- Monthly telephone follow-up by the social worker
- Non-programmed hospitalizations in both groups (including emergency department visits)
- Prescribed exams, laboratory tests, and non-programmed medicines
- Unscheduled visits to the oncologist in both groups (including travel allowances for the patient)
- Costs incurred by the patients (in relation to out-of-pocket charges, extra travel costs). As patients are isolated, we hypothesized that caregiver costs would be negligible.

All resources will be observed, collected through the case report forms (CRFs), and valorized over the period between baseline and the 6 months follow-up. Several sources of unit costs will be investigated for this economic study based on HAS recommendations. Production costs through diagnostic-related groups from the Technical Agency for hospital information (ATIH) will be used for inpatient costs (sources: National Health system and hospital databases). For ambulatory care and examinations, national tariffs are recommended (sources: National databases for medical and paramedical acts, National official list prices for drugs and registers of pharmaceutical specialties, National Table for Biology).

Differences in health and costs outcomes' means will be tested by using the Student *t* test (if normally distributed) and differences between distributions tested with the nonparametric Mann-Whitney *U* test (if means are not normally distributed). Data will be analyzed according to the intention-to-treat principle. Statistical methods will be investigated for dealing with missing data, and their potential impact (according to assumptions made to cope with missing data)

on the CUA findings will be addressed in the sensitivity analysis. To address variability and to assess the generalizability of our results, univariate sensitivity analyses will be performed by varying health and cost variables one by one (including but not limited to incidence of sepsis, recurrence rates, or unit costs). The statistical uncertainty surrounding the incremental cost-utility ratio (due to sampling fluctuations for example) will be captured by a multivariate sensitivity analysis, using, among others, simulation methods such as non-parametric bootstrap methods [10]. In addition, cost-effectiveness acceptability curves will be constructed to represent decision uncertainty surrounding cost-effectiveness estimates [11].

Conclusion

The changes to the PREDOMOS study do not influence the statistical analysis plan; the addition of a CUA results from additional funding, and the new patient inclusion/exclusion criteria will facilitate ongoing recruitment. The results of this study are expected to confirm that a PS-DR may be an interesting care management strategy for isolated older patients with cancer.

Trial status

At the time of manuscript submission, the status of the trial is "recruiting."

Abbreviations

ADL: Activity of daily living; ANSM: French drug and device regulation agency (Agence Nationale de Sécurité du Médicament); ATIH: Technical Agency for hospital information (Agence technique de l'information sur l'hospitalisation); CEA: Cost-effectiveness analysis; CPPSM1: French ethics committee (Comité de Protection des Personnes Sud Méditerranée I); ECOG-PS: Eastern Cooperative Oncology Group Performance Status; EQ-5D: EuroQoL Five-levels; G8 ONCODAGE: Scale for detection of geriatric frailties in cancer patients over 70 years old; HAS: French Health institution (Haute Autorité de Santé); m-MOS-SS: Modified Medical Outcomes Study Social Support Survey; MMSE: Mini Mental State Examination; PS-DR: Program of Social intervention associated with Domotic and Remote assistance techniques; QALY: Quality-adjusted life year; QoL: Quality of life

Acknowledgements

The sponsor was represented by Assistance Publique des Hôpitaux de Marseille, France; and its role was to control the appropriateness of ethical and legal considerations for all centers. The authors are grateful to all the investigators for their participation in the study.

Funding

This work is supported by French ministry of health PREPS-2015-0098 « (Programme de Recherche sur la Performance du Système de Soins), by the Marseilles' integrated cancer center (SIRIC-Marseille): Grant INCa-DGOS-Inserm 6038 and by the Fondation de France ("Soigner, Soulager, Accompagner" Call for proposal).

Availability of data and materials

Not applicable.

Authors' contributions

ALC and AD participated in the protocol design, the regulatory files drafting, and the manuscript drafting; they also carried out the feasibility analysis and coordinated the study. EN participated in the study conception, the protocol design, the regulatory files drafting, and the manuscript drafting and submission, and coordinates the study. HLC participated in the study conception and carried

out the feasibility analysis at his center. OG, FR, LG, ENM, MC, RB, FLC, ST, CB, PG, SN, and SK carried out the feasibility analyses for each of their centers. SL participated in the design of the protocol. ND participated in the design of the social case report forms. CM participated in the regulatory files drafting, submitted the files to the administrative institutions (French regional ethical committee, French drug and device regulation agency), and manages the study. PA and PV participated in the study conception and the protocol design. KB participated in the study conception, designed the protocol methodology, and participated in the manuscript drafting. All authors read and approved the final manuscript.

Authors' information

Not applicable.

Ethics approval and consent to participate

The study is conducted in accordance with the Helsinki declaration and the French laws and regulations (Code de la Santé Publique, article L.1121-1 / Loi de Santé Publique n°2004-806 du 9 août 2004 relative à la politique de santé publique et ses décrets d'application du 27 août 2006) and the International Conference on Harmonization (ICH) E6 Guideline for Good Clinical Practice. Regulatory monitoring will be performed by the sponsor. The sponsor needed the approval of the French authorities, including the French ethics committee (Comité de Protection des Personnes Sud Méditerranée I (CPPSM1)), version 2 of the PREDOMOS protocol submitted on March 23rd 2016 (reference number 16 26 was approved on the 26th May 2016 for all 10 participating centers, the French drug and device regulation agency (Agence Nationale de Sécurité du Médicament (ANSM), registry number 2015-A01976-43) approved on the 29th June 2016, before beginning the study. Substantial changes to the protocol (revised version 4) was approved on May 29th 2018 by the ANSM and on June 25th 2018 by the CPPSM1. The ClinicalTrials.gov identifier is NCT02829762. Informed consent was obtained from all subjects.

Consent for publication

Informed consent was obtained from all subjects.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Service de Médecine Interne Gériatrie et Thérapeutique, CHU Sainte Marguerite, Assistance Publique des Hôpitaux de Marseille (AP-HM), 270 Boulevard de Sainte Marguerite Dromel, 13274 Marseille, cedex 09, France. ²Unit of Coordination in Onco-Geriatrics (UCOG), PACA-west, France. ³Service d'Oncologie Digestive, CHU Timone, AP-HM, 264 Rue Saint Pierre, 13385 Marseille, cedex 05, France. ⁴EA3279, Self-perceived Health Assessment Research Unit, Aix-Marseille University, 27 Boulevard Jean Moulin, 13385 Marseille, cedex 05, France. ⁵Service de Pneumologie, CH Yves le Foll, 10 rue Marcel Proust, Saint-Brieuc, France. ⁶Service d'Onco-Gériatrie, CH Intercommunal des Alpes du Sud Site de Gap (CHICAS), 1 Place Auguste Muret, 05000 Gap, France. ⁷Service d'Oncologie Médicale, Institut Paoli Calmettes, 232 Boulevard de Sainte Marguerite Dromel, 13009 Marseille, France. ⁸Oncologie Multidisciplinaire et Innovation Thérapeutique, CHU NORD, AP-HM, Chemin des Bourrelly, 13915 Marseille, cedex 20, France. ⁹Service de Gérontologie, Hôpital de Cimiez, 4 Avenue Reine Victoria, CS 91179, 06003 Nice, France. ¹⁰Service de Gériatrie, CH Yves le Foll, 10 rue Marcel Proust, Saint-Brieuc, France. ¹¹Service de Gériatrie, Hôpital Saint Joseph, 26 Boulevard Louvain, 13285 Marseille, cedex 08, France. ¹²Unité Mobile de Gériatrie, Hôpital Saint Musse, CH Intercommunal Toulon-La Seyne sur Mer (CHITS), 54 Rue Henri Claire Deville, 83000 Toulon, France. ¹³Service d'Hémo-Oncologie, CH du Pays d'Aix, Avenue les Tamaris, 13616 Aix-en-Provence, France. ¹⁴Service d'Oncologie Médicale, Institut Sainte Catherine (ISC), 250 Chemin de Baigne Pieds, 84918 Avignon, cedex 09, France. ¹⁵Service Social, Hôpital Sainte Marguerite, AP-HM, 270 Boulevard de Sainte Marguerite Dromel, 13274 Marseille, cedex 09, France. ¹⁶Direction de la Recherche Clinique et de l'Innovation (DRCI), AP-HM, 80 Rue Brochier, 13354 Marseille, cedex 05, France.

Received: 4 October 2018 Accepted: 11 December 2018

Published online: 15 January 2019

References

- Crétel-Durand E, Nouguerède E, Le Caer H, Rousseau F, Retornaz F, Guillemin O, Couderc A-L, Greillier L, Norguet E, Cécile M, Boulahssass R, Le Caer F, Tournier S, Butaud C, Guillet P, Nahon S, Poudens L, Kirscher S, Loubière S, Diaz N, Dhone J, Auquier P, Baumstarck K. PREDOMOS study, impact of a social intervention program for socially isolated elderly cancer patients: study protocol for a randomized controlled trial. *Trials*. 2017;18(1):174.
- Byrnes J, Carrington M, Chan Y-K, Pollicino C, Dubrowin N, Stewart S, Scuffham PA. Cost-effectiveness of a home based intervention for secondary prevention of readmission with chronic heart disease. *PLOS One*. 2015;10(12):e0144545.
- Hirani SP, Beynon M, Cartwright M, Rixon L, Doll H, Henderson C, Bardsley M, Steventon A, Knapp M, Rogers A, Bower P, Sanders C, Fitzpatrick R, Hendy J, Newman SP. The effect of telecare on the quality of life and psychological well-being of elderly recipients of social care over a 12-month period: the Whole Systems Demonstrator cluster randomised trial. *Age Ageing*. 2014;43(3):334–41.
- Parker SG, Oliver P, Pennington M, Bond J, Jagger C, Enderby PM, Curless R, Chater T, Vanoli A, Fryer K, Cooper C, Julious S, Donaldson C, Dyer C, Wynn T, John A, Ross D. Rehabilitation of older patients: day hospital compared with rehabilitation at home. A randomised controlled trial. *Health Technol Assess*. 2009;13(39):1–143 iii–iv.
- Tappenden P, Campbell F, Rawdin A, Wong R, Kalita N. The clinical effectiveness and cost-effectiveness of home-based, nurse-led health promotion for older people: a systematic review. *Health Technol Assess*. 2012;16(20):1–72.
- HAS. Choices in Methods for Economic Evaluation [Internet]. Haute Autorité de Santé; 2012. https://www.has-sante.fr/portail/upload/docs/application/pdf/2012-10/choices_in_methods_for_economic_evaluation.pdf. Accessed 10 Mar 2018.
- Van Reenen, Mandy, Janssen, Bas. EQ-5D-5L User Guide Basic information on how to use the EQ-5D-5L instrument [Internet]. EuroQol Research Foundation; 2015. https://euroqol.org/wp-content/uploads/2016/09/EQ-5D-5L_UserGuide_2015.pdf. Accessed 10 Mar 2018.
- Doble B, Lorgelly P. Mapping the EORTC QLQ-C30 onto the EQ-5D-3L: assessing the external validity of existing mapping algorithms. *Qual Life Res*. 2016;25(4):891–911.
- Versteegh MM, Leunis A, Luime JJ, Boggild M, Uyl-de Groot CA, Stolk EA. Mapping QLQ-C30, HAQ, and MSIS-29 on EQ-5D. *Med Decis Making*. 2012; 32(4):554–68.
- Briggs AH. Statistical approaches to handling uncertainty in health economic evaluation. *Eur J Gastroenterol Hepatol*. 2004;16(6):551–61.
- Fenwick E, Claxton K, Sculpher M. Representing uncertainty: the role of cost-effectiveness acceptability curves. *Health Econ*. 2001;10(8):779–87.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

