



HAL
open science

Advances and limits of two model species for ecotoxicological assessment of carbamazepine, two by-products and their mixture at environmental level in freshwater

Fanny Desbiolles, Xavier Moreau, Laetitia de Jong, Laure Malleret, Quentin Grandet-Marchant, Pascal Wong-Wah-Chung, Isabelle Laffont-Schwob

► To cite this version:

Fanny Desbiolles, Xavier Moreau, Laetitia de Jong, Laure Malleret, Quentin Grandet-Marchant, et al.. Advances and limits of two model species for ecotoxicological assessment of carbamazepine, two by-products and their mixture at environmental level in freshwater. *Water Research*, 2020, 169, pp.115267. 10.1016/j.watres.2019.115267 . hal-03587360

HAL Id: hal-03587360

<https://hal.science/hal-03587360>

Submitted on 24 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 **Advances and limits of two model species for ecotoxicological assessment of**
2 **carbamazepine, two by-products and their mixture at environmental level in**
3 **freshwater**

4

5 Desbiolles Fanny^{1,2}, Moreau Xavier^{2*}, de Jong Laetitia², Malleret Laure¹, Grandet-Marchant
6 Quentin², Wong-Wah-Chung Pascal¹, Laffont-Schwob Isabelle^{2,3}

7

8 ¹ Aix Marseille Univ, CNRS, LCE, UMR 7376, ECCOREV FR 3098, Marseille, France
9 (fanny.desbiolles@univ-amu.fr;laure.malleret@univ-amu.fr;pascal.wong-wah-chung@univ-
10 amu.fr).

11 ² Aix Marseille Univ, CNRS, IRD, Avignon Université, IMBE UMR 7263, Marseille, France
12 (xavier.moreau@imbe.fr;laetitia.dejong-
13 moreau@imbe.fr;quentingrandetmarchant@gmail.com).

14 ³ Aix Marseille Univ, IRD, LPED, UMR 151, Marseille, France (isabelle.schwob@univ-
15 amu.fr).

16

17 * Corresponding author. Isabelle Laffont-Schwob (isabelle.schwob@univ-amu.fr)

18 Aix Marseille Univ, Laboratoire Population Environnement Développement, case 10, 3 place
19 Victor Hugo 13331 Marseille cedex 3, France

20

21 **Abstract**

22 In order to evaluate the environmental impacts associated with the presence of low-
23 level of pharmaceuticals in waters, chronic ecotoxicity of carbamazepine (CBZ),
24 oxcarbazepine (OxCBZ) and acridine 9-carboxylic acid (9-CAA) and their mixture was
25 investigated using two species from different trophic levels. Innovative approaches were
26 developed by monitoring: (i) phytometabolites in the duckweed *Lemna minor* L. and, (ii)
27 alterations at the population, cellular and molecular levels on the cnidarian *Hydra*
28 *circumcincta* Schulze 1914, to assess the effects of 14-day chronic exposure. On *H.*
29 *circumcincta*, the approach was conducted at 19.5°C, the optimal growth temperature, and
30 at 23°C, to simulate global warming impact.

31 In *L. minor*, results showed alterations of the nitrogen balance and the chlorophyll
32 indices for OxCBZ and 9-CAA separately at environmental concentrations and for the mixture
33 of the three studied products. However, phenolic compound index deeply varied depending on
34 pharmaceuticals and time of exposure with no specific trend. In *H. circumcincta*, effects on
35 morphology, reproduction and lipid peroxidation were observed at low level of CBZ. OxCBZ
36 impacted the total antioxidant capacity of *H. circumcincta*, whereas exposure to 9-CAA
37 affected all parameters, except the reproduction rate. Finally, 14-day exposure to the mixture
38 of the three products also slowed the reproduction rate and altered the morphology of this
39 cnidarian. Higher damages were detected when experiments were conducted at a temperature
40 3.5°C above the optimum for *Hydra* growth, suggesting the relevance of taking into account
41 possible ecotoxicological consequences of global warming. Our results confirmed the
42 necessity of selecting sensitive organisms at different trophic levels to better assess ecotoxicity
43 of pharmaceuticals and their mixture, especially at environmental levels and facing global
44 changes.

45

46 **Keywords (6):** chronic exposure, sensitive aquatic organisms, low concentrations, cocktail
47 effect, pharmaceuticals, biochemical responses

48

49 **1. Introduction**

50 Aquatic biota is chronically exposed to multiple exogenous substances. Many of these
51 compounds are suspected to impact living organisms and human health, and their
52 combination in complex mixtures could exacerbate their harmful effects (Delfosse et al.,
53 2015). Among them, pharmaceuticals have recently emerged as pollutants of high concern
54 in the aquatic environment, where they are commonly introduced as complex mixtures via
55 wastewater treatment plant (WWTP) effluents (Desbiolles et al., 2018; Gros et al., 2007).
56 Many of the past studies on aquatic organisms have focused on toxicity assessment of single
57 pharmaceuticals (for review see Blaise et al., 2006; Crane et al., 2006; Fent et al., 2006;
58 Godoy et al., 2015). Most of the bioassays currently used investigate short time exposure (72
59 h or 96 h) at concentrations in the mg/L range, i.e. well above those found in the aquatic
60 environment (Desbiolles et al., 2018). Since the concentrations of each pharmaceutical in
61 receiving waters and effluents are low, their potential toxic effects on aquatic organisms and
62 even on humans have been often neglected until recent years. Nonetheless, it has been
63 demonstrated that combined low concentrations of pharmaceuticals having similar biological
64 activities could lead to additive or synergistic effects, that could be ecotoxicologically
65 relevant (Brain et al., 2004; Gros et al., 2007; Kolpin et al., 2002). However, data about
66 chronic ecotoxicity of mixture of compounds at low concentrations are still poorly
67 illustrated.

68 Pharmaceuticals are degraded with highly variable efficiencies at the outlet of WWTP
69 ranging from upper than 80% abatement for ibuprofen to less than 20 % for carbamazepine
70 (Miège et al., 2009). Owing to its low abatement in conventional WWTP, the present study

71 focuses on the anticonvulsant carbamazepine (CBZ) together with two of its by-products,
72 namely oxcarbazepine (OxCBZ) and acridine 9-carboxylic acid (9-CAA). Typical
73 concentrations of CBZ, OxCBZ and 9-CAA range from tens to several hundreds of ng/L in
74 receiving natural waters (Ferrari et al., 2003; Gros et al., 2012; Kaiser et al., 2014).

75 The ecotoxicity of CBZ is poorly documented, especially its possible effects on
76 aquatic macrophytes such as *Lemna minor* (Cleuvers, 2003), while some interesting data are
77 available on the freshwater cnidarian, *Hydra attenuata*, a synonym species of *H.*
78 *circumcincta* (Quinn et al., 2004, 2008a, 2008b, 2009). In this latter species, little effect was
79 observed after short exposure (96 h) to a mixture of 11 pharmaceuticals, including CBZ,
80 present at concentrations levels found in a treated municipal effluent.

81 As commonly admitted in ecotoxicological researches, a battery of several
82 representative aquatic organisms belonging to different biological groups and to different
83 trophic levels is recommended, referred as base-set (Arkhipchuk et al., 2000; Côté et al., 1998;
84 Desbiolles et al., 2018; Vacchi et al., 2013). Thus, in the present study, we have selected a
85 primary producer, the freshwater fast-growing macrophyte *Lemna minor* L. 1753 (Ziegler et
86 al., 2016) and the freshwater carnivorous cnidarian *Hydra circumcincta* Schulze, 1914
87 (Quinn et al., 2008a).

88 To address the issue of toxic effects of chemicals at low levels in mixtures, we aimed
89 at testing effects of chronic (exposure time higher than 7 days) exposures to CBZ, OxCBZ,
90 9-CAA and their mixture at environmentally relevant concentrations on various biological
91 parameters of the two selected model organisms. Photosynthetic pigments and flavonoids
92 are good indicators of plant health (Croft and Chen, 2018). However, sampling for
93 phytometabolites bioassays are usually done at the end of exposure experiments with
94 destructive techniques. We hypothesized that a daily-monitoring of those metabolites will
95 be more accurate for early-detection of effect of pharmaceutical exposure using a non-

96 destructive analytical equipment. Moreover, since in animal models, pharmaceutical
97 exposure may act on physiological and behavioral features, an integrative analysis of
98 pharmaceutical exposure may need to include the study of many endpoints from biochemical
99 to populational levels. Bearing in mind to fit tightly to environmental realistic conditions,
100 our work aimed to investigate how far a panel of tests covering molecular to macroscopic
101 level using two biological models could improve the capability to detect a potential effect of
102 selected pharmaceuticals. Furthermore, crossed-effects between chemical pollutions and
103 climate change were noticed on community structure and ecosystem functioning (Dossena
104 et al., 2012). Thus, to question potential cross-effects of pollution and climate change
105 experiments have been carried out at the classic breeding temperature (19.5°C) and at a
106 higher temperature of 23°C. Indeed, it could be hypothesized that temperature rising affects
107 the sensitiveness of organisms to pollutants and reversely chemical exposure may decrease
108 the thermal tolerance of organisms. Therefore, the crossed effects of these stressors must be
109 investigated as it could entrain a decreased in the fitness of wild populations.

110

111 **2. Materials and methods**

112 **2.1. Pharmaceuticals**

113 Carbamazepine (CBZ) and acridine 9-carboxylic acid (9-CAA) were provided by
114 Sigma-Aldrich (Saint Quentin-Fallavier, France) and oxcarbazepine (OxCBZ), by VWR-
115 International-France (Fontenay-sous-Bois, France) and were available as powders (purity >
116 97 %). The stock solutions of each compound were prepared in ultrapure water (Direct-Q
117 5UV, Millipore®) at concentrations of 100, 40 and 20 mg/L for CBZ, OxCBZ and 9-CAA,
118 respectively, after stirring 2 day-long in the dark and at 25 °C. The solutions were stored in
119 the dark at 4 °C for 2 months, thus enabling the ecotoxicological tests on chronic exposure.
120 The stability of the compounds under these conditions was checked by HPLC monitoring.

121 Test solutions of 600, 27 and 900 ng/L were prepared for CBZ, OxCBZ and 9-CAA
122 respectively and a mixture of them at the same latter concentrations was also prepared. This
123 range of concentrations was chosen based on concentrations found in treated municipal
124 effluents (Ferrari et al., 2003; Gros et al., 2012; Kaiser et al., 2014).

125

126 **2.2. Plant bioassay using phytometabolites**

127 Duckweeds (*Lemna minor* L.) were collected from a freshwater basin of the botanic
128 garden in Saint-Jérôme campus (Aix Marseille University, France). The population was
129 cultivated in a modified 10 % Hoagland solution (Sigma-Aldrich) in glass beakers. Beakers
130 with duckweeds at 100 % confluence were then put in a phytotron (MLR-351H, Sanyo) with
131 a 11-13 h day-night cycle with 60 % of humidity and light intensity of 20 $\mu\text{mol}/\text{m}^2/\text{s}$ at 15 °C
132 to lower algal proliferation for 17 days. Each condition of exposure to pharmaceuticals and
133 their mixture was done in 5 replicates in 40 mL-glass dishes and compared with a control
134 containing only the culture solution. Duckweeds were maintained in the same solution during
135 all the experiment time. Phytometabolites were monitored with a non-destructive
136 measurement equipment (Multiplex®, FORCE-A, Orsay, France) using fluorescence
137 technology with multiple excitations to measure constitutive and induced flavonols,
138 anthocyanins, chlorophylls and a chlorophyll-to-flavonoid ratio referred to as nitrogen
139 balance index (NBI), as previously described by Ellili et al. (2017). For each condition, the
140 duckweed fronds were in contact with the measurement cell of the multiplex to ensure that
141 the focal distance was the same for all samples and measurements were done in triplicate per
142 40 mL-glass dishes.

143

144 **2.3. Animal bioassays**

145 The clonal population of *Hydra circumcincta* arises from a polyp given by the
146 “Laboratoire Interdisciplinaire des Environnements Continentaux” (Lorraine University, Metz,
147 France). Culture maintenance procedure was adapted from that developed by Trottier et al.
148 (1997). Briefly, TES buffer (pH 7) (N-[Tris(hydroxymethyl)-methyl]-2-aminoEthaneSulfonic
149 acid, Merck Chimie SAS, Fontenay-sous-Bois, France) was used as culture media and the
150 polyps were fed every three days with 24 h hatched *Artemia sp.* nauplii. The polyp culture was
151 raised at a constant temperature of $19.5 \pm 0.1^\circ\text{C}$ with a 12-12 h light-dark cycle.

152

153 **2.3.1. Chronic toxicity test**

154 All pharmaceutical solutions were diluted in TES buffer. Experiments were carried out
155 on six sterile multi-well polystyrene microplates. Healthy non-budding *H. circumcincta* polyps
156 of similar size were randomly collected among a population of more than 1000 polyps under a
157 stereomicroscope. For each condition, 12 polyps per well were filled with 8 mL of test solutions
158 and tests were done in triplicate. The multi-well microplates were covered to prevent
159 evaporation and placed into a thermo-regulated incubator at $19.5 \pm 0.1^\circ\text{C}$ or $23 \pm 0.1^\circ\text{C}$ under
160 a 12-12 h light-dark cycle for two weeks. Animals were fed the day before the beginning of the
161 experiment and then each three days (day 3, 6, 9 and 12) with about 10 *Artemia sp.* larvae per
162 polyp. Capacity to catch and ingest prey was recorded. For each well, the media was renewed
163 after a 4-hours feeding period. Daily, morphology was monitored, and the number of buds and
164 new polyps was counted to determine the asexual reproduction. At the end of the two-week
165 exposure period, the media was carefully removed, and the living polyps were rapidly frozen
166 and kept at -80°C until oxidative stress measurements. Criteria used to classify the intensity of
167 toxic effects based on experimental endpoints (i.e. morphological changes, budding and
168 reproductive *Hydra* rates) were the following: (1) non-toxic: 0-20 % effect, (2) moderately

169 toxic: 21-40 % effect, (3) toxic: 41-60 % effect, (4) highly toxic: 61-80 % effect, and (5)
170 extremely toxic: 81-100 % (Arkhipchuk et al., 2006; Wang, 1990).

171

172 **2.3.2. Asexual reproduction: evaluation of budding and reproductive hydra** 173 **rates**

174 The budding rate (K_{tot}) considering the number of attached buds and new polyps (i.e.
175 separate buds) was calculated according to the following formula:

$$176 \quad K_{tot} = (\ln(n_{tot}) - \ln(n)) / d$$

177 in which n is the number of polyps at the beginning of the experiment and n_{tot} the sum of the
178 number of attached buds and the number of polyps (detached buds and initial polyps) after d
179 days of exposure (according to de Jong et al. (2016)).

180 The reproductive *Hydra* rate (RHR) during the two-week exposure to pharmaceuticals
181 was calculated according to the following formula:

$$182 \quad RHR = ((N - n) / n) \times 100$$

183 with N number of *Hydra* polyps after d days of exposure and n number of *Hydra* polyps at the
184 beginning of the experiment (according to Arkhipchuk et al. (2006)).

185

186 **2.3.3. Morphological changes reflecting increasing toxicity**

187 The hereby morphological stage nomenclature used was previously described (de Jong
188 et al., 2016; Pachura-Bouchet et al., 2006). Classically, five morphological stages are reported:
189 *a stage*, normal, *Hydra sp.* polyps exhibit an extended body and tentacles; *b stage*, first sign of
190 intoxication, appearance of bulbs at tips of tentacles; *c stage*, second sign of intoxication,
191 shortened tentacles and body (early phase); *d stage*, shortened tentacles and body (late phase);
192 *e stage* or tulip stage, tentacles and body drastically shortened. This last intoxication stage leads

193 to the animal death by disintegration (*f* stage). Stages *b* to *d* are reversible and are considered
194 as sub-lethal endpoints, while *e* and *f* stages are irreversible and considered as lethal endpoints.

195

196 **2.3.4. Biomarkers of oxidative stress**

197 The disturbance of the oxidative balance equilibrium was evaluated in *H. circumcincta*
198 polyps exposed to chronic (14 days) pharmaceutical concentrations by measuring the total
199 antioxidant capacity (AOC kit, Cayman Chemical, Bertin Pharma, France) and the lipid
200 peroxidation by means of the thiobarbituric acid reactive substance assay (TBARS assay kit,
201 Cayman Chemical, Bertin Pharma). The AOC with the pharmaceuticals was compared with that
202 of Trolox, a water-soluble tocopherol analogue, and was quantified as molar Trolox equivalents.
203 In the TBARS assay, the quantification of the malondialdehyde (MDA) was performed by
204 colorimetry following its controlled reaction with thiobarbituric acid.

205

206 **2.4. Data analysis**

207 Data from phytometabolite monitoring were expressed as ratios. All values of each
208 group on *H. circumcincta* were expressed as mean \pm S.E.M (standard error of the mean). At the
209 end of the exposures, the time, pharmaceutical, and temperature interactions for the hydra
210 bioassays were analyzed by the three-way ANOVA followed by Tukey test for paired
211 comparisons. When the use of parametric tests was not possible, the comparisons between
212 conditions for the duckweed and hydra bioassays were done with the Kruskal-Wallis test
213 followed, if needed, by the *post-hoc* Dunn test.

214

215 **3. Results**

216 **3.1. Plant bioassay using phytometabolites**

217 Three of the monitored indices were considered: one for chlorophylls i.e. part of the
218 primary metabolites, one for phenolic compounds i.e. part of the secondary metabolites and
219 finally, one corresponding to a nitrogen balance index (NBI), proxy of the allocation of N
220 and C resources in the plant. Each index was monitored on *L. minor* for 17 days of exposure
221 to the pharmaceuticals.

222

223 **3.1.1. Index of chlorophylls**

224 Variations of chlorophyll indexes during the exposure to each pharmaceutical
225 compound tested separately at environmental concentration as well as to the mixture are
226 presented in figure 1A. No significant effects of CBZ or 9-CAA were observed on
227 chlorophyll contents in *L. minor* compared to control. However, duckweeds exposed to
228 OxCBZ at 27 ng/L had a chlorophyll index constantly significantly higher than the control
229 along the experiment. Finally, no significant effects were detected on the chlorophyll index
230 in *L. minor* exposed to the mixture as compared to control.

231

232 **3.1.2. Index of phenolic compounds**

233 No significant differences were detected for the index of phenolic compounds of
234 duckweeds when exposed to each pharmaceutical (CBZ, OxCBZ, 9-CAA) separately and to
235 the mixture as compared to control ones (figure 1B), but considerable data fluctuations were
236 noticed.

237

238 **3.1.3. NBI ratio**

239 No significant effects on the NBI ratio were observed over time at the environmental
240 concentration of CBZ (figure 1C). Exposure to OxCBZ significantly increased the NBI ratios
241 as compared to control ones along the exposure period except at days 11 and 17 (figure 1C).

242 These results showed an increase in the pool of nitrogen compounds compared to the pool
243 of carbon compounds when *L. minor* was exposed to OxCBZ. For 9-CAA, the NBI ratios
244 were significantly higher compared to the control at days 2, 5, 9, 11, 14 and 17. For *L. minor*
245 exposed to the pharmaceutical mixture, the NBI ratios were always significantly higher than
246 control ones from the first day of experiment (figure 1C). Consequently, the ecotoxicity on
247 NBI ratio was expressed continuously when the 3 compounds were combined.

248

249 **3.2 Animal bioassay using multi-scale analysis**

250 **3.2.1 Asexual reproduction and feeding behavior**

251 A significant decrease of the budding rate (K_{tot}) was recorded at 19.5 °C in polyps
252 exposed to CBZ at day 6 and in those exposed to the mixture at days 6 and 9 (figure 2A). At
253 23 °C, K_{tot} was significantly lower from day 9 in polyps exposed to CBZ and OxCBZ. A
254 significantly lower K_{tot} was also observed in polyps exposed to the mixture at days 7, 12
255 and 14 (figure 2B). Moreover, a significant effect of the temperature on the K_{tot} values was
256 observed at day 3 ($F_{9,59} = 9.839$, $P < 0.0001$): the K_{tot} values of each group of exposed polyps
257 at 23 °C were significantly increased as compared with those obtained at 19.5 °C (at 19.5 °C,
258 K_{tot} mean is 0.006 and, at 23 °C, K_{tot} mean is 0.066) (figure 2A and B). At the end of the
259 experiment, on day 14, a significant effect of the temperature on the budding rate was
260 observed: the K_{tot} values obtained at 23 °C were lower than those obtained at 19.5 °C ($F_{1,1} =$
261 77.002 , $P < 0.0001$), and a significant effect of the exposure to pharmaceuticals was also
262 noticed ($F_{1,4} = 9.415$, $P < 0.0001$). However, statistical analyses did not show any interactions
263 between the effects of temperature and pharmaceuticals on K_{tot} ($F_{1,4} = 1.623$, $P = 0.169$).

264

265 At 19.5 °C, the reproductive rate (RHR) was significantly lower in polyps exposed to
266 CBZ than controls from day 9 to day 14. Moreover, a significantly lower RHR was also

267 observed in polyps exposed to the mixture from day 12 (figure 2C). At 23 °C, a significantly
268 lower RHR was observed in polyps exposed to CBZ, OxCBZ and the mixture from day 12
269 to the end of the exposure (figure 2D). Significant interactions were detected between
270 temperature and pharmaceutical on the effects on RHR ($F_{1,4} = 6,946$, $P = 0,0001$).

271

272 Whatever the conditions, no alteration of the feeding behavior was observed during
273 the 2-week exposure as all living polyps were still able to catch and ingest their prey.

274

275 **3.2.2 Morphological changes reflecting intoxication**

276 At 19.5 °C, significant morphological changes appeared at the very first day of
277 mixture exposure: only 34.7 % of exposed polyps showed normal morphology (*a* stage)
278 while *b* and *c* stages represented 48.6 % and 16.7 % respectively (figure 3A). Significant
279 morphological changes were still observable at days 2, 3 and 4 in polyps exposed to the
280 mixture. At day 4, *d* stages appeared in the population exposed to the mixture (4.6 %) and in
281 polyps exposed to CBZ (2.8 %), while *a* stages represented 46.2 % and 56.3 %, respectively.
282 From day 5 to 8, all the polyps exhibited normal morphology. Then, at day 9, some polyps
283 exhibited *b* stages, but the percentage of injured polyps were significantly higher only in
284 those exposed to OxCBZ, where *b* stages represented 17.6 %. Finally, from day 10 to the
285 end of the two-week exposure, none of the exposed polyps exhibited signs of impaired
286 morphology.

287

288 At 23 °C, significant morphological changes were recorded only from day 12, where
289 the percentage of *b* stages was significantly higher than in controls for all exposure
290 conditions and represented 10.8 % to 15.9 % of the exposed population (figure 3B). At day
291 14, *c* stages appeared in OxCBZ and mixture exposed polyps and their percentage was

292 significantly higher than in controls; *b* stages were still present for all exposure conditions
293 but as *b* stages also appeared in control group, no significant morphological changes were
294 lasting compared to controls.

295

296 **3.2.3. Biomarkers of oxidative stress**

297 At 19.5 °C, an overall significant difference ($F_{9,29}= 87.04$, $p<0.0001$, two-way
298 ANOVA) in the total antioxidant capacity (AOC) was observed: a 2-week exposure to
299 OxCBZ induced a significant two-fold increase in the AOC compared to controls
300 (figure 4A). Concerning the lipid peroxidation (LPO), an overall significant difference
301 ($F_{9,29}= 68.70$, $P<0.0001$, two-way ANOVA) was detected at 19.5 °C. Indeed, compared to
302 controls, the LPO was significantly higher in polyps exposed to CBZ and to 9-CAA, and
303 significantly lower in those exposed to the mixture, whereas no significant differences were
304 observed in those exposed to OxCBZ (figure 4B).

305 At 23 °C, there were no significant differences in the AOC values between control
306 and exposed groups (figure 4A). However, a significant increase in LPO level was observed
307 in polyps exposed to CBZ and 9-CAA and a significant decrease in polyps exposed to OxCBZ
308 compared with controls (figure 4B). The data showed a significant effect of the temperature
309 since differences were observed between control groups ($F_{1,4}= 540.55$, $P<0.0001$ for AOC
310 and $F_{1,4}= 124.55$, $P<0.0001$ for LPO). Indeed, in control groups, it was observed a significant
311 lower AOC levels at 23 °C than 19.5 °C, while the LPO level was higher at 23 °C than at
312 19.5 °C. Same significant variations according to temperature were observed in exposed
313 polyps: AOC values were lower at 23 °C than at 19.5 °C while, except for OxCBZ, LPO
314 values were higher at 23 °C than at 19.5 °C. (figures 4A and 4B). Moreover, three-way
315 ANOVA results have shown an interaction between the effects of both temperature and

316 pharmaceuticals on AOC ($F_{1,4} = 31.445$, $P < 0.0001$) and on LPO level ($F_{1,4} = 7.147$, $P =$
317 0.001).

318

319 **4. Discussion**

320 The aim of this study was to determine the potential ecotoxicological effects of CBZ,
321 OxCBZ, 9-CAA and their mixture at environmental levels and for exposure times above 7
322 days as required by chronic experiments. No general trends in model organism responses to
323 pharmaceutical exposure were identified, since biochemical responses were diverse.
324 However, mixture effect was mostly different than the effects of pharmaceuticals taken
325 individually, thus confirming research needs on mixture of pharmaceuticals and not only on
326 single pharmaceutical exposure, which is far away from environmental conditions
327 (Desbiolles et al., 2018). Moreover, time of exposure is a key element in the detection of
328 chronic ecotoxicological effect. In this study thanks to the model organisms used, to the
329 multi-parameters monitored and to the non-destructive approach, experimenting long time
330 of exposure was feasible. In this attempt, reduced acute toxicity but significant chronic
331 effects were evidenced for exposure to pharmaceuticals added at environmental
332 concentrations levels.

333

334 **4.1. Chronic effects of CBZ, OxCBZ and 9-CAA on *Lemna minor***

335 A stimulating effect on the chlorophyll index of duckweeds exposed to OxCBZ at
336 27 ng/L was observed with constantly significantly higher values than the control ones.
337 Large variations of the index of phenolic compounds in duckweeds exposed to each
338 pharmaceutical were obtained and precluded any general conclusion. By comparing the
339 exposure to the mixture and to each pharmaceutical taken separately, the effects resulting
340 from the mixture on the NBI ratio were lower than the sum of the effects generated by each

341 compound. Indeed, the mixture exposure resulted in an infra-additive effect or an
342 antagonism. Therefore, the NBI ratio was the most sensitive indicator in duckweeds to
343 mixture exposure showing a cocktail effect of the pharmaceuticals.

344 Literature about ecotoxicity of CBZ on *L. minor* is scarce. However, an EC₅₀ of
345 25.5 mg/L for CBZ was already determined by Cleuvers (2003) after 7 days. In 2015, the
346 concentration of chlorophylls in *L. minor* exposed to various synthetic chemical compounds,
347 including some pharmaceutical products, have been monitored (Fekete-Kertész et al., 2015).
348 This approach was destructive by harvesting duckweeds after 7 days of exposure and
349 performing spectrophotometric assays from the extracts. The authors demonstrated that
350 duckweeds are less sensitive to a pharmaceutical i.e. diclofenac than to pesticides. In our
351 study, the use of a non-destructive measuring equipment to determine indices of
352 phytometabolites allowed to put forward early effects on the metabolism of *L. minor* from
353 the lowest concentrations used. However, the great variability of responses between
354 replicates and over time did not allow to identify a particular trend in the plant response to
355 pharmaceutical exposure.

356 Otherwise, *L. minor* is recognized as a pollutant-tolerant plant species used in
357 phytoremediation to metabolize some pharmaceuticals (Farrell, 2012; Matamoros et al.,
358 2012; Reinhold et al., 2010). Consequently, more research is needed due to the limited
359 number of studies on the effects of CBZ on this plant species and their discordant results.
360 Matamoros and colleagues (2012) hypothesize that the difference in results between studies
361 could be due to different initial concentrations of pharmaceuticals and in the density of plant
362 biomass. *L. minor* may also have a defined photoperiodism dependent on its circadian clock
363 as it has been reported for another duckweed species (Hillman, 1970). Some xenobiotics may
364 also cause a phase shift in the circadian rhythm as demonstrated in *Lemna gibba* (Kondo,
365 1989). Phenolic compound synthesis in plants is controlled by the circadian rhythm

366 (Lattanzio, 2013), which results in a regulatory effect of the biosynthetic pathway via its key
367 enzymes. Among the many functions of phenolic compounds, they play a preponderant role
368 in the response to plant stress (Cheynier et al., 2013). The variability of the metabolic
369 responses in *L. minor* in our work may correspond to multiple responses at the
370 phenylpropanoid pathway leading to the absence of a clear signal related to CBZ and by-
371 product exposure. Given that these phytometabolites cannot be easily used as early
372 biomarkers of the pharmaceutical effects, further studies including more complex mixtures
373 containing chemicals having different usage purpose (pharmaceuticals, pesticides, industrial
374 detergent...) should be explored before renouncing to this nondestructive approach for
375 ecotoxicity assessment.

376

377 **4.2. Chronic effects of CBZ, OxCBZ and 9-CAA on *Hydra circumcincta***

378 According to Maruyama and Sugiyama (1979), two factors determine the growth rate
379 of a hydra population: how frequently mature polyps produce buds and how rapidly newly
380 formed buds mature to start producing their own buds. The time for bud to form a fully
381 developed new hydra is dependent of both feeding rate and temperature (Stiven, 1962). In
382 our study, the feeding rate was constant (each three days), the polyps were equally fed and,
383 whatever the exposure conditions in the present experiments, all polyps were still able to
384 catch and ingest their prey. Therefore, significant differences in the budding rates between
385 control group and exposed ones could only be attributed to exposure conditions. Hence, at
386 day 3, the velocity at which polyps produced new buds at 23 °C, as compared to those
387 maintained at 19.5 °C, can be attributed to the increase of temperature leading to a higher
388 metabolism. At 19.5 °C, the lower budding rate observed at day 6 in polyps exposed to CBZ
389 and at days 6 and 9 in polyps exposed to the mixture could explain in part the lower RHR
390 observed at the following days for the same pharmaceutical exposure. However, as after day

391 9, no more differences in budding rate were observed, it could be hypothesized that the
392 decrease of RHR could also be explained by the action of CBZ and pharmaceutical mixture
393 on the highly molecular conserved signaling pathways involved in the development. For
394 example, pharmacological approaches have demonstrated the importance of the FGFR
395 (“*fibroblast growth factor receptor*”) /Notch (membrane receptor) signaling in bud
396 development because when the FGFR or Notch pathways are inhibited, buds develop well
397 but never detach from the parent (see Galliot, 2012, for review). Therefore, if CBZ and the
398 pharmaceutical mixture acted on such pathways, it could retard the bud detachment (i.e.
399 lower RHR, observed at days 12 and 14) but not necessarily slow the budding development
400 (K_{tot}). At 23 °C and from day 7, significant differences in budding rates of exposed polyps
401 were more frequently observed than at 19.5 °C. Indeed, in polyps exposed to CBZ, OxCBZ
402 but also the mixture, the budding rate (K_{tot}) was lower. This latter result is consistent with
403 a lower observed RHR from day 12 at this temperature. Thus, it seems that the increase of
404 temperature affected the sensitiveness of exposed organisms to CBZ, OxCBZ and the
405 mixture leading to decreases of both K_{tot} and RHR at the end of the exposure. Finally, maybe
406 the decrease of reproductive rates (RHR) at both temperatures and after long term exposure
407 to low concentration of pharmaceuticals, especially to the mixture, could lead to possible
408 deleterious effects on the fitness of chronically exposed wild populations.

409 At 19.5 °C, from the first to the fourth day of mixture exposure, morphological
410 changes revealed effects that can be ranged from moderately toxic to highly toxic according
411 to the nomenclature of Wang (1990) and Arkhipchuk et al. (2006). Then, from day 5 to the
412 end of exposure time (two weeks), a return to a normal morphology was observed. These
413 observations suggest an early setting up of efficient defense mechanisms. At 23 °C, it seemed
414 that the increase in metabolism activity, suggested by an elevated K_{tot} at day 3, may protect
415 from intoxication at the beginning of exposure. Interestingly, at 23 °C, morphological signs

416 of intoxication appeared at day 12 and persisted at day 14, in polyps exposed to OxCBZ and
417 the mixture. It could be hypothesized that the energetic allocation to survive to thermal stress
418 may protect the polyp from intoxication for a while. However, the set-up defense
419 mechanisms could not be maintained to preserve homeostasis for a long time. This
420 hypothesis supports the apparition of *b* stages in controls and an increasing sensitiveness of
421 polyps exposed to OxCBZ and the mixture at day 14.

422 At the end of the two-week exposure, morphological changes did not allow to classify
423 the effects according a toxicity scale (Arkhipchuk et al., 2006; Wang, 1990), as less than
424 20 % of the exposed population showed toxicity signs. Interestingly, at the end of the 2 weeks
425 exposure at 23 °C, even if morphological alterations did not reach 20 % and did not allow to
426 classify the effects according a toxicity scale (Arkhipchuk et al., 2006; Wang, 1990), they
427 were greater than 10 % in polyps exposed to the low pharmaceutical concentrations used in
428 this study. These results showed that *H. circumcincta* is a sensitive species that could be used
429 for detection of early intoxication signs and therefore be a good candidate for utilization in
430 regulatory framework for the determination of classical ecotoxicological endpoints (No
431 Observable Effect Concentration= NOEC, Lowest Observed Effect Concentration= LOEC,
432 Maximum Acceptable Threshold Concentration= MATC= $(\text{NOEC} \times \text{LOEC})^{1/2}$) and
433 especially EC₁₀, an alternative endpoint recommended by several authors instead of the three
434 other above mentioned (Warne and van Dam, 2008). The sensitiveness of *H. circumcincta*
435 to pharmaceuticals was already supported by Quinn et al. (2009).

436 Regarding the biomarkers of the oxidative stress in *H. attenuata*, it has been
437 previously shown that after a 48 h exposure to CBZ at 20 °C, the inducible threshold
438 concentrations (MATC) for heme oxidase and glutathione S-transferase activities are
439 0.03 μM and 0.2 μM CBZ respectively (equivalent to 7 and 47 mg/L respectively) (Quinn et
440 al., 2004). However, these concentrations are considerably higher than the environmental

441 levels of tens to hundreds ng/L tested here separately and in mixture. Our results indicated
442 that, at 19.5 °C, the exposition at realistic pharmaceuticals environmental concentrations of
443 CBZ and 9-CAA did not induce the antioxidant defense systems which in turns lead to a
444 membrane lipid peroxidation. In contrast, as the antioxidant defense systems were induced
445 in polyyps exposed to OxCBZ, no lipid peroxidation was measured in those polyyps. This can
446 be explained by the fact that low doses of CBZ and 9-CAA induce membrane lipid
447 peroxidation that is not counteracted by the antioxidative systems. In polyyps exposed to
448 OxCBZ, the induction of antioxidative system seemed to have prevented lipid peroxidation.
449 The exposure to mixture seemed to have no impact on the oxidative balance parameters (no
450 induced oxidative defense mechanisms and low lipid peroxidation). It can be hypothesized
451 an unknown (agonist/antagonist/synergistic) effect of the combination of the three molecules
452 on the oxidative balance. These results indicated that *Hydra* species is a sensitive model to
453 detect the effects of low concentration of pharmaceuticals on cellular membrane lipid
454 peroxidation. At higher concentrations, it has been reported that hydra may reduce the toxic
455 effect of pharmaceuticals mixture by metabolizing the toxicants (Quinn et al., 2009). Hence,
456 hydra were found to possess glutathione S-transferase activity (Stenersen et al., 1987), an
457 enzyme that plays an important role during phase II biotransformation in the detoxification
458 and metabolism of xenobiotics and endogen compounds (Hoarau et al., 2004). The fact that
459 the increase of temperature led to significantly lower levels of AOC in all groups of *H.*
460 *circumcincta* is consistent with the observations of Dash and Phillips (2012), which have
461 reported that *H. vulgaris* responds to thermal stress by regulating its catalase mRNA
462 transcription, another enzyme involved in the regulation of the oxidative balance. In their
463 study, these authors have investigated the effects of acute thermal stresses (heat shocks) on
464 levels of HvCatalase mRNA. They have observed a decrease in HvCatalase mRNA level in
465 hydra groups exposed to 30 °C or 37 °C for 6 h, as compared to control groups maintained

466 at 18 °C. Our thermal exposure condition (+ 3.5 °C for 14 days) led to similar results, as we
467 observed a lower level of AOC in controls and groups exposed to pharmaceuticals.
468 Moreover, the ANOVA results, showing significant interactions between pharmaceutical
469 and temperature, argue in favor of an increased hydra sensitiveness to pharmaceuticals when
470 temperature rises. Thus, it could be hypothesized that *Hydra* species are able to regulate
471 mRNA antioxidative enzyme levels according to surrounding temperature. As proposed by
472 Dash and Phillips (2012), it would be expected that exposure to increased temperatures for
473 long duration would impair the transcriptional machinery of the cell thereby negatively
474 affecting transcription of mRNA of antioxidative enzymes, and thus, decreasing the total
475 antioxidative capacity. This decrease in AOC could, in turns, lead to an increase in cellular
476 membrane lipid peroxidation.

477 In summary, at the end of the two-week pharmaceutical exposure, none of the
478 biological endpoints allowed to classify the effects according to a toxicity scale (Arkhipchuk
479 et al., 2006; Wang, 1990), as less than 20 % of the exposed population showed toxicity signs.
480 However, altogether, the significant variations of biological parameters studied led to the
481 same conclusions: (1) long term exposure to low pharmaceuticals in mixture had deleterious
482 effects on *H. circumcincta* polyps, and especially decreased reproductive rates suggesting
483 possible harmful effects on the fitness of wild populations chronically exposed; (2)
484 temperature rising also seemed to have harmful effects on the studied biological endpoints.
485 When the temperature rising was combined with exposure to very low doses of
486 pharmaceuticals in mixture, the effects seemed to be more pronounced on some of the
487 biological endpoints (i.e. altered morphology). In a context of global warming climate
488 change, it is possible that the toxic effects of mixtures in non-target aquatic organisms will
489 be higher.

490

491 **5. Conclusion**

492 Our results confirmed the necessity of identifying sensitive organisms belonging to
493 different trophic levels, to better assess ecotoxicity of pharmaceuticals and their mixture
494 especially at environmental levels and facing global changes.

495 Although some variations of phytometabolite composition were noticed under
496 exposure with the selected pharmaceuticals, no specific trend could be identified. The
497 involvement of complex interactions between the pharmaceuticals and the plant metabolic
498 pathways may induce time-dependent effects, which may also be connected with the
499 circadian cycle of the duckweed. Further investigations are needed, especially considering
500 the activities of key-enzymes, which may give more contrasting results. *Hydra circumcincta*
501 appears as an appropriate sensitive model for ecotoxicological studies aiming at
502 investigating long term effect of complex environmental mixtures. Future steps could
503 involve a screening of other biological endpoints and the use of experimental designs,
504 allowing to study more complex environmental mixtures and crossed effects between
505 chemical pollutions and climate change.

506

507 **Acknowledgements**

508 The authors thank the Provence-Alpes-Côte d'Azur regional council and the company
509 Seakalia (Marseille, France) for funding this research through a PhD grant (DEB
510 n°141198/2014). The ECCOREV federation is also acknowledged for funding the project
511 entitled POPMETABOTOX. We also thank the students who participated at several
512 preliminary experiments with *Hydra circumcincta*: L. Bosco, J. Dujardin, M. Caceres and F.
513 Tournier. Many thanks to Labex OT-MED scientific network for inspiring our reflection
514 about warming climate change potential impact.

515

516 **References**

- 517 Arkhipchuk, V.V., Blaise, C., Malinovskaya, M.V., 2006. Use of hydra for chronic toxicity
518 assessment of waters intended for human consumption. *Environ. Pollut.* 142, 200–211.
519 <https://doi.org/10.1016/j.envpol.2005.10.012>
- 520 Arkhipchuk, V.V., Malinovskaya, M.V., Garanko, N.N., 2000. Cytogenetic study of organic
521 and inorganic toxic substances on *Allium cepa*, *Lactuca sativa*, and *Hydra attenuata* cells.
522 *Environ. Toxicol.* 15, 338–344. [https://doi.org/10.1002/1522-7278\(2000\)15:4<338::AID-](https://doi.org/10.1002/1522-7278(2000)15:4<338::AID-)
523 [TOX10>3.0.CO;2-R](https://doi.org/10.1002/1522-7278(2000)15:4<338::AID-TOX10>3.0.CO;2-R)
- 524 Blaise, C., Gagné, F., Eullaffroy, P., Férard, J.-F., 2006. Ecotoxicity of selected pharmaceuticals
525 of urban origin discharged to the Saint-Lawrence river (Québec, Canada): a review. Ecotoxicity
526 of selected pharmaceuticals of urban origin discharged to the Saint-Lawrence river (Québec,
527 Canada): a review. *Braz. J. Aquat. Sci. Technol.* <http://dx.doi.org/10.14210/bjast.v10n2.p29-51>
- 528 Brain, R.A., Johnson, D.J., Richards, S.M., Hanson, M.L., Sanderson, H., Lam, M.W., Young,
529 C., Mabury, S.A., Sibley, P.K., Solomon, K.R., 2004. Microcosm evaluation of the effects of
530 an eight pharmaceutical mixture to the aquatic macrophytes *Lemna gibba* and *Myriophyllum*
531 *sibiricum*. *Aquat. Toxicol.* 70, 23–40. <https://doi.org/10.1016/j.aquatox.2004.06.011>
- 532 Cheynier, V., Comte, G., Davies, K.M., Lattanzio, V., Martens, S., 2013. Plant phenolics:
533 Recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiol. Biochem.*,
534 *Plant Phenolics: biosynthesis, genetics, and ecophysiology* 72, 1–20.
535 <https://doi.org/10.1016/j.plaphy.2013.05.009>
- 536 Clevers, M., 2003. Aquatic ecotoxicity of pharmaceuticals including the assessment of
537 combination effects. *Toxicol. Lett.* 142, 185–194. <https://doi.org/10.1016/S0378->
538 [4274\(03\)00068-7](https://doi.org/10.1016/S0378-4274(03)00068-7)
- 539 Côté, C., Blaise, C., Michaud, J.-R., Ménard, L., Trottier, S., Gagné, F., Lifshitz, R., 1998.
540 Comparisons between microscale and whole-sediment assays for freshwater sediment toxicity

541 assessment. Environ. Toxicol. Water Qual. 13, 93–110. [https://doi.org/10.1002/\(SICI\)1098-](https://doi.org/10.1002/(SICI)1098-)
542 2256(1998)13:1<93::AID-TOX8>3.0.CO;2-2

543 Crane, M., Watts, C., Boucard, T., 2006. Chronic aquatic environmental risks from exposure to
544 human pharmaceuticals. Sci. Total Environ. 367, 23–41.
545 <https://doi.org/10.1016/j.scitotenv.2006.04.010>

546 Croft, H., Chen, J.M., 2018. Leaf Pigment Content. in: Comprehensive Remote Sensing.
547 Elsevier, pp. 117–142. <https://doi.org/10.1016/B978-0-12-409548-9.10547-0>

548 Dash, B., Phillips, T.D., 2012. Molecular characterization of a catalase from *Hydra vulgaris*.
549 Gene 501, 144–152. <https://doi.org/10.1016/j.gene.2012.04.015>

550 de Jong, L., Pech, N., de Aragão Umbuzeiro, G., Moreau, X., 2016. Multi-scale biomarker
551 evaluation of the toxicity of a commercial azo dye (Disperse Red 1) in an animal model, the
552 freshwater cnidarian *Hydra attenuata*. Water Res. 96, 62–73.
553 <https://doi.org/10.1016/j.watres.2016.03.043>

554 Delfosse, V., Dendele, B., Huet, T., Grimaldi, M., Boulahtouf, A., Gerbal-Chaloin, S., Beucher,
555 B., Roecklin, D., Muller, C., Rahmani, R., Cavallès, V., Daujat-Chavanieu, M., Vivat, V.,
556 Pascussi, J.-M., Balaguer, P., Bourguet, W., 2015. Synergistic activation of human pregnane X
557 receptor by binary cocktails of pharmaceutical and environmental compounds. Nat. Commun.
558 6, 8089. <https://doi.org/10.1038/ncomms9089>

559 Desbiolles, F., Malleret, L., Tiliacos, C., Wong-Wah-Chung, P., Laffont-Schwob, I., 2018.
560 Occurrence and ecotoxicological assessment of pharmaceuticals: Is there a risk for the
561 Mediterranean aquatic environment? Sci. Total Environ. 639, 1334–1348.
562 <https://doi.org/10.1016/j.scitotenv.2018.04.351>

563 Dossena, M., Yvon-Durocher, G., Grey, J., Montoya, J.M., Perkins, D.M., Trimmer, M.,
564 Woodward, G., 2012. Warming alters community size structure and ecosystem functioning.
565 Proc. R. Soc. B Biol. Sci. 279, 3011–3019. <https://doi.org/10.1098/rspb.2012.0394>

566 Ellili, A., Rabier, J., Prudent, P., Salducci, M.-D., Heckenroth, A., Lachaâl, M., Laffont-
567 Schwob, I., 2017. Decision-making criteria for plant-species selection for phytostabilization:
568 Issues of biodiversity and functionality. *J. Environ. Manage.* 201, 215–226.
569 <https://doi.org/10.1016/j.jenvman.2017.06.041>

570 Farrell, J., 2012. Duckweed uptake of phosphorus and five pharmaceuticals: Microcosm and
571 wastewater lagoon studies. Thesis of Master Science in Civil and Environmental Engineering.
572 Utah State University, USA.

573 Fekete-Kertész, I., Kunglné-Nagy, Z., Gruiz, K., Magyar, Á., Farkas, É., Molnár, M., 2015.
574 Assessing toxicity of organic aquatic micropollutants based on the total chlorophyll content of
575 *Lemna minor* as a sensitive endpoint. *Period. Polytech. Chem. Eng.* 59, 262–271.
576 <https://doi.org/10.3311/PPch.8077>

577 Fent, K., Escher, C., Caminada, D., 2006. Estrogenic activity of pharmaceuticals and
578 pharmaceutical mixtures in a yeast reporter gene system. *Reprod. Toxicol.* 22, 175–185.
579 <https://doi.org/10.1016/j.reprotox.2006.04.010>

580 Ferrari, B., Paxeus, N., Giudice, R.L., Pollio, A., Garric, J., 2003. Ecotoxicological impact of
581 pharmaceuticals found in treated wastewaters: study of carbamazepine, clofibrac acid, and
582 diclofenac. *Ecotoxicol. Environ. Saf.* 55, 359–370. <https://doi.org/10.1016/S0147->
583 [6513\(02\)00082-9](https://doi.org/10.1016/S0147-6513(02)00082-9)

584 Galliot, B., 2012. Hydra, a fruitful model system for 270 years. *Int. J. Dev. Biol.* 56, 411–423.
585 <https://doi.org/10.1387/ijdb.120086bg>

586 Godoy, A.A., Kummrow, F., Pamplin, P.A.Z., 2015. Ecotoxicological evaluation of
587 propranolol hydrochloride and losartan potassium to *Lemna minor* L. (1753) individually and
588 in binary mixtures. *Ecotoxicology* 24, 1112–1123. <https://doi.org/10.1007/s10646-015-1455-3>

589 Gros, M., Petrović, M., Barceló, D., 2007. Wastewater treatment plants as a pathway for aquatic
590 contamination by pharmaceuticals in the Ebro river basin (Northeast Spain). *Environ. Toxicol.*

591 Chem. 26, 1553–1562. <https://doi.org/10.1897/06-495R.1>

592 Gros, M., Rodríguez-Mozaz, S., Barceló, D., 2012. Fast and comprehensive multi-residue
593 analysis of a broad range of human and veterinary pharmaceuticals and some of their
594 metabolites in surface and treated waters by ultra-high-performance liquid chromatography
595 coupled to quadrupole-linear ion trap tandem mass spectrometry. *J. Chromatogr. A* 1248, 104–
596 121. <https://doi.org/10.1016/j.chroma.2012.05.084>

597 Hillman, W.S., 1970. Carbon dioxide output as an index of circadian timing in *Lemna*
598 photoperiodism. *Plant Physiol.* 45, 273–279.

599 Hoarau, P., Garello, G., Gnassia-Barelli, M., Roméo, M., Girard, J.-P., 2004. Effect of three
600 xenobiotic compounds on Glutathione S-Transferase in the clam *Ruditapes decussatus*. *Aquat.*
601 *Toxicol.* 68, 87–94. <https://doi.org/10.1016/j.aquatox.2004.03.001>

602 Kaiser, E., Prasse, C., Wagner, M., Bröder, K., Ternes, T.A., 2014. Transformation of
603 oxcarbazepine and human metabolites of carbamazepine and oxcarbazepine in wastewater
604 treatment and sand filters. *Environ. Sci. Technol.* 48, 10208–10216.
605 <https://doi.org/10.1021/es5024493>

606 Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B., Buxton,
607 H.T., 2002. Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S.
608 streams, 1999–2000: A national reconnaissance. *Environ. Sci. Technol.* 36, 1202–1211.
609 <https://doi.org/10.1021/es011055j>

610 Kondo, T., 1989. Comparison of phase shifts of the circadian rhythm of K⁺ uptake in *Lemna*
611 *gibba* G3 by various amino acid analogs. *Plant Physiol.* 90, 1600–1608.
612 <https://doi.org/10.1104/pp.90.4.1600>

613 Lattanzio, V., 2013. Phenolic Compounds: Introduction, in: Ramawat, K.G., Mérillon, J.-M.
614 (Eds.), *Natural Products*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1543–1580.
615 https://doi.org/10.1007/978-3-642-22144-6_57

616 Maruyama, T., Sugiyama, T., 1979. A mathematical model for hydra multiplication by asexual
617 budding appendix to “genetic analysis of developmental mechanisms in hydra vii. Statistical
618 analyses of developmental-morphological characters and cellular compositions.” Dev. Growth
619 Differ. 21, 377–381. <https://doi.org/10.1111/j.1440-169X.1979.00377.x>

620 Matamoros, V., Nguyen, L.X., Arias, C.A., Salvadó, V., Brix, H., 2012. Evaluation of aquatic
621 plants for removing polar microcontaminants: A microcosm experiment. Chemosphere 88,
622 1257–1264. <https://doi.org/10.1016/j.chemosphere.2012.04.004>

623 Miège, C., Choubert, J.M., Ribeiro, L., Eusèbe, M., Coquery, M., 2009. Fate of pharmaceuticals
624 and personal care products in wastewater treatment plants – Conception of a database and first
625 results. Environ. Pollut. 157, 1721–1726. <https://doi.org/10.1016/j.envpol.2008.11.045>

626 Pachura-Bouchet, S., Blaise, C., Vasseur, P., 2006. Toxicity of nonylphenol on the cnidarian
627 *Hydra attenuata* and environmental risk assessment. Environ. Toxicol. 21, 388–394.
628 <https://doi.org/10.1002/tox.20201>

629 Quinn, B., Gagné, F., Blaise, C., 2004. Oxidative metabolism activity in *Hydra attenuata*
630 exposed to carbamazepine. Fresenius Environ. Bull. 13, 783–788.

631 Quinn, B., Gagne, F., Blaise, C., 2009. Evaluation of the acute, chronic and teratogenic effects
632 of a mixture of eleven pharmaceuticals on the cnidarian, *Hydra attenuata*. Sci. Total Environ.
633 407, 1072–1079. <https://doi.org/10.1016/j.scitotenv.2008.10.022>

634 Quinn, B., Gagné, F., Blaise, C., 2008a. An investigation into the acute and chronic toxicity of
635 eleven pharmaceuticals (and their solvents) found in wastewater effluent on the cnidarian,
636 *Hydra attenuata*. Sci. Total Environ. 389, 306–314.
637 <https://doi.org/10.1016/j.scitotenv.2007.08.038>

638 Quinn, B., Gagné, F., Blaise, C., 2008b. The effects of pharmaceuticals on the regeneration of
639 the cnidarian, *Hydra attenuata*. Sci. Total Environ. 402, 62–69.
640 <https://doi.org/10.1016/j.scitotenv.2008.04.039>

641 Reinhold, D., Vishwanathan, S., Park, J.J., Oh, D., Michael Saunders, F., 2010. Assessment of
642 plant-driven removal of emerging organic pollutants by duckweed. *Chemosphere* 80, 687–692.
643 <https://doi.org/10.1016/j.chemosphere.2010.05.045>

644 Stenersen, J., Kobro, S., Bjerke, M., Arend, U., 1987. Glutathione transferases in aquatic and
645 terrestrial animals from nine phyla. *Comp. Biochem. Physiol. Part C Comp. Pharmacol.* 86, 73–
646 82. [https://doi.org/10.1016/0742-8413\(87\)90147-2](https://doi.org/10.1016/0742-8413(87)90147-2)

647 Stiven, A.E., 1962. The effect of temperature and feeding on the intrinsic rate of increase of
648 three species of hydra. *Ecology* 43, 325–328. <https://doi.org/10.2307/1931991>

649 Trottier, S., Blaise, C., Kusui, T., Johnson, E.M., 1997. Acute toxicity assessment of aqueous
650 samples using a microplate-based *Hydra attenuata* assay. *Environ. Toxicol. Water Qual.* 12,
651 265–271. [https://doi.org/10.1002/\(SICI\)1098-2256\(1997\)12:3<265::AID-TOX10>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1098-2256(1997)12:3<265::AID-TOX10>3.0.CO;2-9)


652 Vacchi, F.I., Albuquerque, A.F., Vendemiatti, J.A., Morales, D.A., Ormond, A.B., Freeman,
653 H.S., Zocolo, G.J., Zanoni, M.V.B., Umbuzeiro, G., 2013. Chlorine disinfection of dye
654 wastewater: Implications for a commercial azo dye mixture. *Sci. Total Environ.* 442, 302–309.
655 <https://doi.org/10.1016/j.scitotenv.2012.10.019>

656 Wang, W., 1990. Toxicity assessment of pretreated industrial wastewaters using higher plants.
657 *Res. J. Water Pollut. Control Fed.*, 62, 853-860.





658 Warne, M.S.J., van Dam, R., 2008. NOEC and LOEC data should no longer be generated or
659 used. *Australas. J. Ecotoxicol.* 14, 1.

660 Ziegler, P., Sree, K.S., Appenroth, K.-J., 2016. Duckweeds for water remediation and toxicity
661 testing. *Toxicol. Environ. Chem.* 98, 1127–1154.
662 <https://doi.org/10.1080/02772248.2015.1094701>

663



666 Figure 1: Boxplots of chlorophyll index **(A)**, phenolic compound index **(B)**, and nitrogen
667 balance index (NBI) **(C)** measured during 17 days in *Lemna minor* after exposure to single
668 pharmaceuticals and the low-dose mixture (CONT: control; CBZ: Carbamazepine (600 ng/L);
669 OxCBZ: Oxcarbazepine (27 ng/L); 9-CAA: acridine 9-carboxylic acid (900 ng/L), MIX:
670 mixture of CBZ, OxCBZ and 9-CAA at respectively 600, 27 and 900 ng/L. Boxplots,
671 considered per day of measure, those marked with asterisks * are significantly different from
672 controls, at $p \leq 0.05$.




Control
 CBZ
 OxCBZ
 *** 9CAA
 Mixture

674 **Figure 2:** Asexual reproduction expressed as mean budding rate (Ktot) at 19.5 °C (A) and at
 675 23 °C (B) in *Hydra circumcincta* exposed 14 days to pharmaceuticals. Asexual reproduction
 676 expressed as mean reproductive *Hydra* rate (RHR) at 19.5 °C (C) and at 23 °C (D) in *H.*
 677 *circumcincta* exposed 14 days to pharmaceuticals.

678 CONT: control; CBZ: Carbamazepine (600 ng/L); OxCBZ: Oxcarbazepine (27 ng/L); 9-
 679 CAA: acridine 9-carboxylic acid (900 ng/L); Mix: Mixture of CBZ, OxCBZ and 9-CAA at
 680 600, 27 and 900 ng/L respectively.


681 *: $P \leq 0.05$, **: $P \leq 0.01$, ***: $P \leq 0.001$ significantly different from controls at the corresponding
 682 day.



683

684 **Figure 3:** Cumulative histograms of morphological stage occurrence (in percent) in *Hydra*
 685 *circumcincta* reflecting intoxication according to time (from day 0 to day 14) during
 686 pharmaceutical exposures at 19.5 °C (A) and at 23 °C (B). For better readability, standard
 687 errors and days where morphological changes were not observed (98-100 % of *a* stages) are
 688 omitted.

689 CONT: Control; CBZ: Carbamazepine (600 ng/L); OxCBZ: Oxcarbazepine (27 ng/L); 9-
 690 CAA: acridine 9-carboxylic acid (900 ng/L); Mix: Mixture of CBZ, OxCBZ and 9-CAA at
 691 600, 27 and 900 ng/L, respectively. For stage description see 2.3.3. section. *: $P \leq 0.05$, **:
 692 $P \leq 0.01$, ***: $P \leq 0.001$ significantly different from controls at the corresponding day.



693
 694 **Figure 4:** Effects of 14-day exposure to pharmaceuticals on biomarkers of oxidative stress
 695 in *Hydra circumcincta* at two different temperatures. **(A)** Antioxidant capacity (AOC)
 696 expressed as mM Trolox/mg proteins and, **(B)** Lipid peroxidation (LPO) measured by means
 697 of the TBARS assay (expressed in malondialdehyde-MDA- concentration in μmole/mg
 698 proteins).

699 CONT: Controls; CBZ: Carbamazepine (600 ng/L); OxCBZ: Oxcarbazepine (27 ng/L); 9-
 700 CAA: acridine 9-carboxylic acid (900 ng/L); Mix: Mixture of CBZ, OxCBZ and 9-CAA at
 701 respectively 600, 27 and 900 ng/L. *: $P \leq 0.05$, ***: $P < 0.001$, significantly different from
 702 corresponding control at 19.5 °C; \$\$\$: $P \leq 0.001$, significantly different from corresponding
 703 control at 23 °C; #: $P \leq 0.05$, ####: $P \leq 0.001$, significantly different between 19.5 °C and 23 °C.

704