



**HAL**  
open science

# Stabilization and approximate null-controllability of a large class of diffusive equations from thick control supports

Paul Alphonse, Jérémy Martin

► **To cite this version:**

Paul Alphonse, Jérémy Martin. Stabilization and approximate null-controllability of a large class of diffusive equations from thick control supports. 2021. hal-03588500v1

**HAL Id: hal-03588500**

**<https://hal.science/hal-03588500v1>**

Preprint submitted on 8 Jan 2021 (v1), last revised 24 Feb 2022 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# STABILIZATION AND APPROXIMATE NULL-CONTROLLABILITY OF A LARGE CLASS OF DIFFUSIVE EQUATIONS FROM THICK CONTROL SUPPORTS

PAUL ALPHONSE AND JÉRÉMY MARTIN

ABSTRACT. We prove that the thickness is a necessary and sufficient geometric condition that ensures the (rapid) stabilization or the approximate null-controllability with uniform cost of a large class of evolution equations posed on the whole space  $\mathbb{R}^n$  and associated with operators of the form  $F(|D_x|)$ , the function  $F : [0, +\infty) \rightarrow \mathbb{R}$  being bounded below and continuous. We also provide explicit feedbacks and constants associated with these stabilization properties. Our results apply in particular for the half heat equation associated with the function  $F(t) = t$ , for which null-controllability is known to fail from thick control supports. More generally, the notion of thickness was known to be a necessary and sufficient condition for the null-controllability of the fractional heat equations associated with the functions  $F(t) = t^{2s}$  in the case  $s > 1$ , and that this null-controllability property from thick control supports does not hold when  $0 < s \leq 1$ .

## 1. INTRODUCTION

This paper is devoted to investigate stabilization and approximate null-controllability issues for control systems of the following form

$$(E_F) \quad \begin{cases} \partial_t f(t, x) + F(|D_x|)f(t, x) = h(t, x)1_\omega(x), & t > 0, x \in \mathbb{R}^n, \\ f(0, \cdot) = f_0 \in L^2(\mathbb{R}^n), \end{cases}$$

where the operator  $F(|D_x|)$  is the Fourier multiplier associated with the symbol  $F(|\xi|)$ , the function  $F : [0, +\infty) \rightarrow \mathbb{R}$  being bounded below and continuous, and  $\omega \subset \mathbb{R}^n$  is a Borel set with positive Lebesgue measure.

The study of the (rapid) stabilization and the (approximate) null-controllability of evolution equations of the form  $(E_F)$  has been much addressed recently [2, 3, 9, 12, 16, 17, 19, 21]. The Schrödinger counterparts of these equations [18] or the same equations posed on bounded domains [15, 23] have also been studied. In this work, we consider supports  $\omega \subset \mathbb{R}^n$  which are thick.

**Definition 1.1.** Given  $\gamma \in (0, 1)$  and  $L > 0$ , the set  $\omega \subset \mathbb{R}^n$  is said to be  $\gamma$ -thick at scale  $L$  when it is measurable and satisfies

$$\forall x \in \mathbb{R}^n, \quad \text{Leb}(\omega \cap (x + [0, L]^n)) \geq \gamma L^n,$$

where  $\text{Leb}$  denotes the Lebesgue measure.

This notion of thickness has appeared to play a key role in the null-controllability theory since the works [7, 22], where the authors established that this is a necessary and sufficient geometric condition that ensures the null-controllability of the heat equation posed on  $\mathbb{R}^n$ , which is the equation  $(E_F)$  associated with the function  $F(t) = t^2$ . The same phenomena holds true more generally for the evolution equations associated with fractional Laplacians  $(-\Delta)^s$  (case where  $F(t) = t^{2s}$ ) under the same setting and when  $s > 1/2$ , as proven in [3], and also quite surprisingly for the Schrödinger counterparts of these equations in the one dimensional setting and when  $s \geq 1/2$ , see [18] (Corollary 2.8). It is also known from the

---

2020 *Mathematics Subject Classification.* 93D15, 93B05, 35R11, 26E10.

*Key words and phrases.* Stabilization, Approximate null-controllability, Thick sets, Quasi-analytic sequences, Diffusive equations.

works [11, 12] that in the case  $0 < s \leq 1/2$ , the fractional heat equation  $(E_{t^{2s}})$  is not null-controllable from thick control supports anymore in general. Other classes of degenerate parabolic equations of hypoelliptic type, as evolution equations associated with accretive quadratic operators or (non-autonomous) Ornstein-Uhlenbeck operators, were also proven to be null-controllable from thick control supports, see e.g. [2, 4, 5]. In this work, we prove that a very general class of equations of the form  $(E_F)$  are approximately null-controllable with uniform cost from the support control  $\omega \subset \mathbb{R}^n$  if and only if  $\omega$  is a thick set. Our result holds in particular for the half heat equation  $(E_t)$  which is not null-controllable from thick control supports, see e.g. [12] (Theorem 2.3) or [16] (Theorem 1.1).

The study of the (rapid) stabilization of the control system  $(E_F)$ , as for it, has been addressed very recently in the works [9, 17, 21]. It has been proven in [9] (Theorem 1.1) that for all  $s > 0$ , the fractional heat equation  $(E_{t^{2s}})$  is exponentially stabilizable from the support  $\omega$  if and only if  $\omega$  is thick. It is also known from [17] (Example 1) that the very same equation  $(E_{t^{2s}})$  is rapidly stabilizable from complementaries of Euclidean balls in  $\mathbb{R}^n$  when  $0 < s < 1$ . In this paper, we establish that the control system  $(E_F)$  is exponentially stabilizable from  $\omega$  if and only if  $\omega$  is a thick set when  $\inf F \leq 0$  (in the case  $\inf F > 0$ , the control system  $(E_F)$  is stable) and  $\liminf_{+\infty} F > 0$ . Moreover, we provide explicit formulas for the feedbacks  $K$  and the constants associated with this stabilization, which allows us to prove that when  $\lim_{+\infty} F = +\infty$ , the control system  $(E_F)$  is rapidly stabilizable from  $\omega$  if and only if  $\omega$  is thick. In particular, we recover [9] (Theorem 1.1) (with new explicit feedbacks) and we generalize [17] (Example 1). We also prove that when  $\omega$  is not dense in  $\mathbb{R}^n$ , the equation  $(E_F)$  is never rapidly stabilizable in the particular case where  $F$  admits a finite limit at  $+\infty$ .

In a nutshell, our results highlight the importance of the notion of thickness not only in the null-controllability theory, since this is also a necessary and sufficient geometric condition that ensures the stabilization or the approximate null-controllability with uniform cost of a large class of diffusive equations  $(E_F)$ .

*Outline of the work.* In Section 2, we present in details the main results contained in this work. Section 3 is devoted to the proofs of the results concerning the stabilization and the rapid stabilization of the control system  $(E_F)$ . Basic properties of quasi-analytic sequences are presented in Section 4, which allow to establish the results concerning the approximate null-controllability of the evolution equation  $(E_F)$  in Section 5. Finally, Section 6 is an Appendix concerning the proof of an observability result used in Section 5.

*Notations.* The following notations and conventions will be used all over the work:

1. The canonical Euclidean scalar product of  $\mathbb{R}^n$  is denoted by  $\cdot$  and  $|\cdot|$  stands for the associated canonical Euclidean norm.
2. For all measurable subset  $\omega \subset \mathbb{R}^n$ , the inner product of  $L^2(\omega)$  is defined by

$$\langle u, v \rangle_{L^2(\omega)} = \int_{\omega} u(x) \overline{v(x)} \, dx, \quad u, v \in L^2(\omega),$$

while  $\|\cdot\|_{L^2(\omega)}$  stands for the associated norm. Moreover,  $\mathcal{L}(L^2(\omega))$  stands for the set of bounded operators on  $L^2(\omega)$ .

3. For all function  $u \in \mathcal{S}(\mathbb{R}^n)$ , the Fourier transform of  $u$  is denoted  $\widehat{u}$  or  $\mathcal{F}u$  and defined by

$$\widehat{u}(\xi) = (\mathcal{F}u)(\xi) = \int_{\mathbb{R}^n} e^{-ix \cdot \xi} u(x) \, dx, \quad \xi \in \mathbb{R}^n.$$

With this convention, Plancherel's theorem states that

$$\forall u \in L^2(\mathbb{R}^n), \quad \|\widehat{u}\|_{L^2(\mathbb{R}^n)} = (2\pi)^{n/2} \|u\|_{L^2(\mathbb{R}^n)}.$$

4. We denote by  $\nabla_x$  the gradient and we set  $D_x = -i\nabla_x$ . Moreover,  $F(|D_x|)$  stands for the Fourier multiplier associated with the symbol  $F(|\xi|)$  for all continuous function  $F : [0, +\infty) \rightarrow \mathbb{R}$ .

5. For all measurable subset  $\omega \subset \mathbb{R}^n$ ,  $1_\omega$  stands for the characteristic function of  $\omega$ .

## 2. STATEMENT OF THE MAIN RESULTS

This section is devoted to present in details the main results contained in this work. Let us begin by defining precisely the different concepts related to the control system  $(E_F)$  we are interested in:

- (i) The control system  $(E_F)$  is said to be *null-controllable* from the support control  $\omega$  in time  $T > 0$  when for all  $f_0 \in L^2(\mathbb{R}^n)$ , there exists a control  $h \in L^2((0, T) \times \omega)$  such that the mild solution of  $(E_F)$  satisfies  $f(T, \cdot) = 0$ .
- (ii) The control system  $(E_F)$  is said to be *approximately null-controllable* from the support control  $\omega$  in time  $T > 0$  if for all  $\varepsilon > 0$  and  $f_0 \in L^2(\mathbb{R}^n)$ , there exists a control  $h \in L^2((0, T) \times \omega)$  such that the mild solution of  $(E_F)$  satisfies  $\|f(T, \cdot)\|_{L^2(\mathbb{R}^n)} \leq \varepsilon$ .
- (iii) The control system  $(E_F)$  is said to be *approximately null-controllable with uniform cost* from the support control  $\omega$  in time  $T > 0$  if for all  $\varepsilon > 0$ , there exists a positive constant  $C_{\varepsilon, T} > 0$  such that for all  $f_0 \in L^2(\mathbb{R}^n)$ , there exists a control  $h \in L^2((0, T) \times \omega)$  such that the mild solution of  $(E_F)$  satisfies  $\|f(T, \cdot)\|_{L^2(\mathbb{R}^n)} \leq \varepsilon$  with moreover

$$\int_0^T \|h(t, \cdot)\|_{L^2(\omega)}^2 dt \leq C_{\varepsilon, T} \|f_0\|_{L^2(\mathbb{R}^n)}^2.$$

- (iv) The control system  $(E_F)$  is said to be *exponentially stabilizable* from the support control  $\omega$  at rate  $\alpha > 0$  if there exist a positive constant  $M_\alpha \geq 1$  and a feedback  $K_\alpha \in \mathcal{L}(L^2(\mathbb{R}^n))$  such that for all  $t \geq 0$ ,

$$\|e^{-t(F(|D_x|) + 1_\omega K_\alpha)}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq M_\alpha e^{-\alpha t},$$

where  $\mathcal{L}(L^2(\mathbb{R}^n))$  denotes the set of bounded operators on  $L^2(\mathbb{R}^n)$ . When the feedback  $K$  can be chosen equal to zero, the control system  $(E_F)$  is said to be *stable*. The existence of the semigroup generated by the operator  $F(|D_x|) + 1_\omega K_\alpha$  is ensured by the theory of bounded perturbation of semigroups, see e.g. [8] (Theorem III.1.3).

- (v) The control system  $(E_F)$  is said to be *rapidly stabilizable* from the support control  $\omega$  if it is exponentially stabilizable from  $\omega$  at any rate  $\alpha > 0$ .

**2.1. Stabilization.** First of all, we are interesting in tackling stabilization issues for the evolution system  $(E_F)$ . Let us begin by noticing that when  $\inf F > 0$ , we get from Plancherel's theorem that

$$\forall t \geq 0, \quad \|e^{-tF(|D_x|)}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq e^{-(\inf F)t},$$

so the control system  $(E_F)$  is stable. The interesting case is therefore when  $\inf F \leq 0$ . In this case, we prove that the thickness of the support  $\omega \subset \mathbb{R}^n$  is a necessary geometric condition that ensures the stabilization of the equation  $(E_F)$ , and a sufficient one when assuming in addition that  $\liminf_{+\infty} F > 0$ . We also provide explicit feedbacks and quantitative estimates associated with this stabilization.

**Theorem 2.1.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  be a measurable set.*

- (i) *When  $\inf F \leq 0$  and the evolution system  $(E_F)$  is exponentially stabilizable from  $\omega$ , then the set  $\omega$  is thick.*
- (ii) *When  $\liminf_{+\infty} F > 0$  and  $\omega$  is a thick set, then there exist some positive constants  $C = C(\omega) \geq 1$  and  $R_0 = R_0(F) > 0$  such that for all  $R \geq R_0$  and  $t \geq 0$ ,*

$$(2.1) \quad \|e^{-t(F(|D_x|) + C e^{CR} \alpha_R 1_\omega K_R)}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq C e^{CR} e^{-\alpha_R t/2},$$

where we set  $\alpha_R = \inf_{r \geq R} F(r)$  and where  $K_R$  stands for the following orthogonal projection

$$K_R : L^2(\mathbb{R}^n) \rightarrow \{f \in L^2(\mathbb{R}^n) : \text{Supp } \widehat{f} \subset B(0, R)\}.$$

The evolution system  $(E_F)$  is therefore exponentially stabilizable from  $\omega$  under these two assumptions.

**Corollary 2.2.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function satisfying  $\inf F \leq 0$  and  $\liminf_{+\infty} F > 0$ , and  $\omega \subset \mathbb{R}^n$  be a measurable set. The evolution system  $(E_F)$  is exponentially stabilizable from  $\omega$  if and only if  $\omega$  is thick.*

It is a very interesting issue to know if the control system  $(E_F)$  is exponentially stabilizable when  $\inf F \leq 0$  and  $\liminf_{+\infty} F \leq 0$ . We will not tackle such a question in this work.

As a consequence of the quantitative stabilization estimates (2.1), we directly obtain the following result concerning the rapid stabilization of the evolution system  $(E_F)$  under the assumption  $\lim_{+\infty} F = +\infty$ , by applying Theorem 2.1 to the function  $F - \inf F$ .

**Corollary 2.3.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function satisfying  $\lim_{+\infty} F = +\infty$  and  $\omega \subset \mathbb{R}^n$  be a measurable set. The evolution system  $(E_F)$  is rapidly stabilizable from  $\omega$  if and only if  $\omega$  is thick.*

**Example 2.4.** For all real number  $s > 0$ , let us consider the function  $F_s : [0, +\infty) \rightarrow [0, +\infty)$  defined for all  $t \geq 0$  by  $F_s(t) = t^s$ . We also consider  $\omega \subset \mathbb{R}^n$  a measurable set with positive Lebesgue measure. It follows from Corollaries 2.2 and 2.3 that the associated control system  $(E_{F_s})$  is exponentially stabilizable from the support  $\omega$  if and only if it is rapidly stabilizable from  $\omega$  if and only if  $\omega$  is a thick set. Moreover, we deduce from Theorem 2.1 that when  $\omega$  is thick, there exists a positive constant  $C > 0$  such that for all  $R > 0$  and  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-t((-\Delta)^s + Ce^{CR}R^s 1_{\omega K_R})}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq Ce^{CR}e^{-R^s t/2},$$

where  $K_R$  stands for the following orthogonal projection

$$K_R : L^2(\mathbb{R}^n) \rightarrow \{f \in L^2(\mathbb{R}^n) : \text{Supp } \hat{f} \subset B(0, R)\}.$$

These explicit stabilization estimates allow to recover [9] (Theorem 1.1) and also to generalize [17] (Example 1).

In the case where  $\lim_{+\infty} F < +\infty$ , we only provide a necessary condition for the control system  $(E_F)$  to be rapidly stabilizable. The following result implies in particular that when the function  $F$  has a finite limit in  $+\infty$  and when the support  $\omega \subset \mathbb{R}^n$  is not dense in  $\mathbb{R}^n$ , then the equation  $(E_F)$  is not rapidly stabilizable from  $\omega$ .

**Proposition 2.5.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  be a measurable set which is not dense in  $\mathbb{R}^n$ . We assume that  $\lim_{t \rightarrow +\infty} F(t)$  exists and is a non-negative real number (the function  $F$  is therefore bounded). Then, if  $\alpha > \sup F$ , the equation  $(E_F)$  is not stabilizable from  $\omega$  at rate  $\alpha$ .*

**2.2. Cost-uniformly approximate null-controllability.** In the second part of this work, we study the cost-uniformly approximate null-controllability of the equations  $(E_F)$ . We will not address this question for general bounded below continuous functions  $F$ , but only for the ones generating a *quasi-analytic sequence*. Let us precisely define this class of functions. Associated with the function  $F$  is the following log-convex sequence  $\mathcal{M}^F$  whose elements  $M_k^F$ , assumed to be positive real numbers, are defined by

$$(2.2) \quad M_k^F = \sup_{r \geq 0} r^k e^{-F(r)}, \quad k \geq 0.$$

We say that the sequence  $\mathcal{M}^F$  is *quasi-analytic* when, for all real numbers  $a < b$ , the associated Denjoy-Carleman class

$$\mathcal{C}_{\mathcal{M}^F}([a, b]) = \{f \in C^\infty([a, b]) : \forall k \geq 0, \forall x \in [a, b], |f^{(k)}(x)| \leq M_k^F\},$$

is quasi-analytic, meaning that any function  $f$  in this class satisfying

$$\exists x_0 \in [a, b], \forall k \geq 0, \quad f^{(k)}(x_0) = 0,$$

is identically equal to zero. We refer to Section 4 where the notion of quasi-analytic sequence is discussed, and where Denjoy-Carleman's theorem is presented, giving a useful characterization of such sequences.

When the function  $F$  generates a quasi-analytic sequence  $\mathcal{M}^F$  of positive real numbers, the solutions of the homogeneous counterpart of the equation  $(E_F)$  belong to quasi-analytic classes of functions (see Subsection 5.1 for more details). By exploiting this quasi-analytic regularity, we prove that the notion of thickness is a necessary and sufficient geometric condition that ensures the cost-uniformly approximate null-controllability of the evolution equations  $(E_F)$  in any positive time.

**Theorem 2.6.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  be a measurable set. We assume that the sequence  $\mathcal{M}^F$  associated with the function  $F$  defined in (2.2) is a quasi-analytic sequence of positive real numbers. Then, for all positive time  $T > 0$ , the diffusive equation  $(E_F)$  is cost-uniformly approximately null-controllable from the support control  $\omega$  in time  $T$  if and only if  $\omega$  is thick.*

The necessary part of Theorem 2.6 comes to the fact that the cost-uniformly approximate null-controllability implies the rapid stabilization, see Proposition 5.3, and is therefore a consequence of Corollary 2.3 (notice that the assumption on  $F$  in Theorem 2.6 implies in particular that  $\lim_{+\infty} F = +\infty$ .)

Let us now present explicit examples of functions  $F$  generating quasi-analytic sequences  $\mathcal{M}^F$  and for which Theorem 2.6 therefore applies.

**Example 2.7.** Let us assume that the non-negative continuous function  $F : [0, +\infty) \rightarrow [0, +\infty)$  is such that  $\Theta \leq F$ , where the weight  $\Theta : [0, +\infty) \rightarrow [0, +\infty)$  satisfies the following properties

- (i)  $\Theta(0) = 0$  and  $\Theta$  is non-decreasing with  $\lim_{t \rightarrow +\infty} \Theta(t) = +\infty$ ,
- (ii)  $\Theta$  is lower-semicontinuous and convex,
- (iii)

$$\int_0^{+\infty} \frac{\Theta(t)}{1+t^2} dt = +\infty.$$

It follows from the work [10] (see also Proposition 4.6 in the present work) that the sequence  $\mathcal{M}^\Theta$  associated with the weight  $\Theta$ , defined in (2.2), is quasi-analytic. Moreover, we have  $\mathcal{M}^F \leq \mathcal{M}^\Theta$  since  $\Theta \leq F$  and Lemma 4.4 implies that the sequence  $\mathcal{M}^F$  is also quasi-analytic. We deduce that for all Borel set  $\omega \subset \mathbb{R}^n$  with positive Lebesgue measure and all positive time  $T > 0$ , the equation  $(E_F)$  is cost-uniformly approximate null-controllable from the set  $\omega$  in time  $T$  if and only if  $\omega$  is thick. A relevant particular example is when  $F(t) = \Theta(t) = t$ , since the associated evolution equation is the half heat equation  $(E_t)$  posed on the whole space associated with the operator  $\sqrt{-\Delta}$ , known to be not null-controllable from any non dense control support  $\omega$ , see [12] (Theorem 2.3) or [16] (Theorem 1.1). This evolution equation is then a relevant example where the thick condition fails to be sufficient for the (strong) null-controllability but appears to be necessary and sufficient for the cost-uniformly approximate null-controllability.

Actually, we are able to derive cost-uniformly approximate null-controllability results for much less diffusive equations than the half heat equation  $(E_t)$ , as illustrated in the two following examples.

**Example 2.8.** Let  $s \geq 1$ ,  $0 \leq \delta \leq 1$  be non-negative real numbers and  $F_{s,\delta} : [0, +\infty) \rightarrow [0, +\infty)$  be the non-negative continuous function defined for all  $t \geq 0$  by

$$F_{s,\delta}(t) = \frac{t^s}{\log^\delta(e+t)}.$$

We check in Corollary 4.8 that the associated sequence  $\mathcal{M}^{F_{s,\delta}}$  defined in (2.2) is a quasi-analytic sequence of positive real numbers. Therefore, for all Borel set  $\omega \subset \mathbb{R}^n$  with positive

Lebesgue measure and all positive time  $T > 0$ , the equation  $(E_{F_{s,\delta}})$  is cost-uniformly approximate null-controllable from the set  $\omega$  in time  $T$  if and only if  $\omega$  is thick.

**Example 2.9.** Let  $p \geq 1$  be a positive integer and  $F_p : [0, +\infty) \rightarrow [0, +\infty)$  be the non-negative continuous function defined for all  $t \geq 0$  by

$$F_p(t) = \frac{t}{g(t)(g \circ g)(t)\dots g^{\circ p}(t)}, \quad \text{where } g(t) = \log(e + t),$$

with  $g^{\circ p} = g \circ \dots \circ g$  ( $p$  compositions). We check in Proposition 4.7 that the associated sequence  $\mathcal{M}^{F_p}$  defined in (2.2) is quasi-analytic. As a consequence, for all Borel set  $\omega \subset \mathbb{R}^n$  with positive Lebesgue measure and all positive time  $T > 0$ , the equation  $(E_{F_p})$  is cost-uniformly approximate null-controllable from the set  $\omega$  in time  $T$  if and only if  $\omega$  is thick.

Regarding the weaker notion of approximate null-controllability presented in the beginning of this section, the geometry of the control support is much simpler. Indeed, the following proposition ensures that the control system  $(E_F)$  is approximate null-controllable in any positive time  $T > 0$  and from any measurable set  $\omega \subset \mathbb{R}^n$  with positive Lebesgue measure when  $F$  generates a log-convex quasi-analytic sequence  $\mathcal{M}^F$ .

**Proposition 2.10.** *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  be a measurable set with positive Lebesgue measure. If the sequence  $\mathcal{M}^F$  associated with the function  $F$  defined in (2.2) is a quasi-analytic sequence of positive real numbers, then for all positive time  $T > 0$ , the diffusive equation  $(E_F)$  is approximate null-controllable from the support control  $\omega$  in time  $T$ .*

In particular, diffusive equations discussed in Examples 2.7, 2.8 and 2.9 are approximate null-controllable in any positive time  $T > 0$  from any measurable subset  $\omega \subset \mathbb{R}^n$  satisfying  $\text{Leb}(\omega) > 0$ .

### 3. (RAPID) STABILIZATION OF DIFFUSIVE EQUATIONS

The aim of this section is to prove Theorem 2.1 and Proposition 2.5 concerning the stabilization and the rapid stabilization properties of the following general control system

$$(E_F) \quad \begin{cases} \partial_t f(t, x) + F(|D_x|)f(t, x) = h(t, x)1_\omega(x), & t > 0, x \in \mathbb{R}^n, \\ f(0, \cdot) = f_0 \in L^2(\mathbb{R}^n). \end{cases}$$

where  $F : [0, +\infty) \rightarrow \mathbb{R}$  is a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  is a measurable set with positive Lebesgue measure.

**3.1. Proof of Theorem 2.1: necessary part.** First of all, let us assume that the equation  $(E_F)$  is stabilizable from the set  $\omega$ , with the extra assumption that  $\inf F \leq 0$ . We aim at proving that the support control  $\omega$  is then thick. To that end, we will use the following nice characterization of the stabilization in terms of observability estimates:

**Theorem 3.1** (Theorem 1 in [21]). *The following assertions are equivalent:*

- (i) *The evolution system  $(E_F)$  is stabilizable from  $\omega$ .*
- (ii) *For all  $\varepsilon \in (0, 1)$ , there exists  $T > 0$  and  $C > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,*

$$\|e^{-TF(|D_x|)}g\|_{L^2(\mathbb{R}^n)}^2 \leq C \int_0^T \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

- (iii) *There exists  $\varepsilon \in (0, 1)$ ,  $T > 0$  and  $C > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,*

$$\|e^{-TF(|D_x|)}g\|_{L^2(\mathbb{R}^n)}^2 \leq C \int_0^T \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

According to the above theorem, assuming that the equation  $(E_F)$  is stabilizable from  $\omega$  is equivalent in assuming that there exist  $\varepsilon \in (0, 1)$ ,  $T > 0$  and  $C > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$(3.1) \quad \|e^{-TF(|D_x|)}g\|_{L^2(\mathbb{R}^n)}^2 \leq C \int_0^T \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

The strategy consists in applying this observability estimate for well-chosen functions  $g \in L^2(\mathbb{R}^n)$ . This approach has especially been used in the works [2, 3, 4, 9] in which stabilization or null-controllability issues are studied for fractional heat equations or evolution equations associated with (non)-autonomous Ornstein-Uhlenbeck operators posed on the whole space  $\mathbb{R}^n$ .

Fixing  $x_0 \in \mathbb{R}^n$  and considering  $\xi_0 \in \mathbb{R}^n$  and  $l \gg 1$  whose values will be adjusted later, we consider the Gaussian function  $g_{l,\xi_0}$  defined by

$$\forall x \in \mathbb{R}^n, \quad g_{l,\xi_0}(x) = \frac{1}{l^n} \exp\left(ix \cdot \xi_0 - \frac{|x - x_0|^2}{2l^2}\right).$$

Classical results concerning Fourier transform of Gaussian functions show that

$$(3.2) \quad \forall \xi \in \mathbb{R}^n, \quad \widehat{g}_{l,\xi_0}(\xi) = (2\pi)^{n/2} \exp\left(-ix_0 \cdot (\xi - \xi_0) - \frac{l^2|\xi - \xi_0|^2}{2}\right).$$

On the one hand, it follows from Plancherel's theorem that the left-hand side of the inequality (3.1) applied to the functions  $g_{l,\xi_0}$  is a positive constant independent of the point  $x_0$ , denoted  $\delta_{l,\xi_0} > 0$  in the following and given by

$$(3.3) \quad \begin{aligned} \delta_{l,\xi_0} &= \|e^{-TF(|D_x|)}g_{l,\xi_0}\|_{L^2(\mathbb{R}^n)}^2 = \int_{\mathbb{R}^n} |e^{-ix_0 \cdot (\xi - \xi_0)} e^{-TF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2}|^2 d\xi \\ &= \frac{1}{l^n} \int_{\mathbb{R}^n} |e^{-TF(|\xi/l + \xi_0|)} e^{-|\xi|^2/2}|^2 d\xi > 0. \end{aligned}$$

On the other hand, we get that the  $L^2$ -norm of the function  $g_{l,\xi_0}$  does not depend on the point  $x_0 \in \mathbb{R}^n$  also and is given by the following Gaussian integral

$$\|g_{l,\xi_0}\|_{L^2(\mathbb{R}^n)}^2 = \frac{1}{l^{2n}} \int_{\mathbb{R}^n} e^{-|x|^2/l^2} dx = \left(\frac{\pi}{l^2}\right)^{n/2}.$$

Let us check that the point  $\xi_0 \in \mathbb{R}^n$  and the large positive parameter  $l \gg 1$  can be adjusted so that  $\delta_{l,\xi_0} - \varepsilon \|g_{l,\xi_0}\|_{L^2(\mathbb{R}^n)}^2 > 0$ , that is,

$$(3.4) \quad \int_{\mathbb{R}^n} |e^{-TF(|\xi/l + \xi_0|)} e^{-|\xi|^2/2}|^2 dx > \varepsilon \pi^{n/2}.$$

Since  $\varepsilon \in (0, 1)$  and the function  $F$  satisfies  $\inf F \leq 0$ , we can assume that the point  $\xi_0 \in \mathbb{R}^n$  is chosen in order to satisfy  $e^{-TF(|\xi_0|)} > \varepsilon$ . The dominated convergence theorem then implies that

$$\begin{aligned} \lim_{l \rightarrow +\infty} \int_{\mathbb{R}^n} |e^{-TF(|\xi/l + \xi_0|)} e^{-|\xi|^2/2}|^2 dx &= e^{-TF(|\xi_0|)} \int_{\mathbb{R}^n} |e^{-|\xi|^2/2}|^2 dx \\ &= e^{-TF(|\xi_0|)} \pi^{n/2} > \varepsilon \pi^{n/2}. \end{aligned}$$

The parameter  $l \gg 1$  can therefore be adjusted so that (3.4) holds. The values of  $\xi_0 \in \mathbb{R}^n$  and  $l \gg 1$  are now fixed. We therefore deduce from (3.1) and (3.4) that

$$(3.5) \quad M_{l,\xi_0} \leq C \int_0^T \|e^{-tF(|D_x|)}g_{l,\xi_0}\|_{L^2(\omega)}^2 dt \quad \text{with} \quad M_{l,\xi_0} = \delta_{l,\xi_0} - \varepsilon \|g_{l,\xi_0}\|_{L^2(\mathbb{R}^n)}^2.$$



Moreover, by introducing  $\mathcal{F}_\xi^{-1}$  the partial inverse Fourier transform with respect to the variable  $\xi \in \mathbb{R}^n$  and using (3.2), the right-hand side of this inequality writes as

$$\begin{aligned} \int_0^T \|e^{-tF(|D_x|)} g_{l,\xi_0}\|_{L^2(\omega)}^2 dt &= (2\pi)^n \int_0^T \int_\omega |\mathcal{F}_\xi^{-1}(e^{-ix_0 \cdot (\xi - \xi_0)} e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt \\ &= (2\pi)^n \int_0^T \int_\omega |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x - x_0)|^2 dx dt \\ &= (2\pi)^n \int_0^T \int_{\omega - x_0} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt. \end{aligned}$$

Given  $r > 0$  a positive radius whose value will be chosen later, we split the previous integral in two parts and obtain the following estimate

$$\begin{aligned} (3.6) \quad \int_0^T \|e^{-tF(|D_x|)} g_{l,\xi_0}\|_{L^2(\omega)}^2 dt &\leq (2\pi)^n \int_0^T \int_{(\omega - x_0) \cap B(0,r)} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt \\ &\quad + (2\pi)^n \int_0^T \int_{|x|>r} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt. \end{aligned}$$

Now, we study one by one the two integrals appearing in the right-hand side of (3.6). First, notice that for all  $0 \leq t \leq T$ ,

$$\begin{aligned} \|\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})\|_{L^\infty(\mathbb{R}^n)} &\leq \frac{1}{(2\pi)^n} \|e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2}\|_{L^1(\mathbb{R}^n)} \\ &\leq \frac{1}{(2\pi)^n} \|e^{-l^2|\xi - \xi_0|^2/2}\|_{L^1(\mathbb{R}^n)} = \frac{1}{(2\pi)^n} \left(\frac{\pi}{l^2}\right)^{n/2}. \end{aligned}$$

It therefore follows from the invariance by translation of the Lebesgue measure that

$$\begin{aligned} (3.7) \quad (2\pi)^n \int_0^T \int_{(\omega - x_0) \cap B(0,r)} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt &\leq \left(\frac{\pi}{l^2}\right)^{n/2} \int_0^T \text{Leb}((\omega - x_0) \cap B(0,r)) dt = T \left(\frac{\pi}{l^2}\right)^{n/2} \text{Leb}(\omega \cap B(x_0, r)). \end{aligned}$$

In order to control the second integral, we use the dominated convergence theorem which justifies the following convergence

$$\int_0^T \int_{|x|>r} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt \xrightarrow{r \rightarrow +\infty} 0,$$

since

$$\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2}) \in L^2([0, T] \times \mathbb{R}^n).$$

Thus, we can choose the radius  $r \gg 1$  large enough so that

$$(3.8) \quad C(2\pi)^n \int_0^T \int_{|x|>r} |\mathcal{F}_\xi^{-1}(e^{-tF(|\xi|)} e^{-l^2|\xi - \xi_0|^2/2})(x)|^2 dx dt \leq \frac{M_{l,\xi_0}}{2}.$$

Gathering (3.5), (3.6), (3.7) and (3.8), we obtain the following estimate

$$\forall x_0 \in \mathbb{R}^n, \quad \frac{M_{l,\xi_0}}{2} \leq TC \left(\frac{\pi}{l^2}\right)^{n/2} \text{Leb}(\omega \cap B(x_0, r)).$$

This proves that the support control  $\omega$  is actually a thick set.

**3.2. Proof of Theorem 2.1: sufficient part.** In this second subsection, we check that the evolution equation  $(E_F)$  is actually stabilizable when  $\omega$  is a thick set, assuming this time that  $\liminf_{+\infty} F > 0$ . The strategy consists in constructing an adapted Lyapunov function, inspired by the proof of [23] (Theorem 1.1) in which the author studies the stabilization properties of the heat equation posed on bounded domains of  $\mathbb{R}^n$ .

Since  $\liminf_{+\infty} F > 0$  by assumption, there exists a positive constant  $R_0 > 0$  such that

$$\forall R \geq R_0, \quad \alpha_R := \inf_{r \geq R} F(r) > 0.$$

Let us fix  $R \geq R_0$  and consider two positive real numbers  $\lambda_R, \mu_R > 0$  to be chosen later. For all initial datum  $f_0 \in L^2(\mathbb{R}^n)$  and all time  $t \geq 0$ , we define  $f(t)$  as the mild solution of the control system  $(E_F)$  with feedback  $\lambda_R K_R$  at time  $t$ , that is

$$\forall t \geq 0, \quad f(t) = e^{-t(F(|D_x|) + \lambda_R 1_\omega K_R)} f_0,$$

where  $K_R$  stands for the following orthogonal projection

$$(3.9) \quad K_R : L^2(\mathbb{R}^n) \rightarrow \{g \in L^2(\mathbb{R}^n) : \text{Supp } \hat{g} \subset B(0, R)\}.$$

We want to prove that the two constants  $\lambda_R$  and  $\mu_R$  can be adjusted so that the following estimate holds for all  $t \geq 0$ ,

$$(3.10) \quad \|f(t)\|_{L^2(\mathbb{R}^n)}^2 \leq \mu_R e^{-\alpha_R t} \|f_0\|_{L^2(\mathbb{R}^n)}^2.$$

To that end, we consider the following Lyapunov function

$$(3.11) \quad V(y) = \mu_R \|K_R y\|_{L^2(\mathbb{R}^n)}^2 + \|(1 - K_R)y\|_{L^2(\mathbb{R}^n)}^2, \quad y \in L^2(\mathbb{R}^n).$$

Notice that the function  $V \circ f$  is differentiable on  $(0, +\infty)$ , with

$$\forall t > 0, \quad \frac{d}{dt} V(f(t)) = \mu_R \frac{d}{dt} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 + \frac{d}{dt} \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2.$$

We shall need to estimate each term of the right-hand side of this equality. On the one hand, we have that for all  $t > 0$ ,

$$\begin{aligned} \mu_R \frac{d}{dt} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 &= 2 \langle K_R f(t), K_R f'(t) \rangle_{L^2(\mathbb{R}^n)} \\ &= -2\mu_R \langle K_R f(t), F(|D_x|) K_R f(t) \rangle_{L^2(\mathbb{R}^n)} - 2\lambda_R \mu_R \|K_R f(t)\|_{L^2(\omega)}^2. \end{aligned}$$

The operator  $F(|D_x|)$  being accretive, we get that for all  $t > 0$ ,

$$\langle K_R f(t), F(|D_x|) K_R f(t) \rangle_{L^2(\mathbb{R}^n)} \geq 0,$$

and as a consequence,

$$\mu_R \frac{d}{dt} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 \leq -2\lambda_R \mu_R \|K_R f(t)\|_{L^2(\omega)}^2.$$

Moreover, O. Kovrijkine established in [14] (Theorem 3) a quantitative version of the Logvinenko-Sereda theorem for thick sets which will allow us to control the right-hand side of the above estimate. Precisely, this result is the following:

**Theorem 3.2** (Theorem 3 in [14]). *There exists a universal positive constant  $C_n \geq e$  depending only on the dimension  $n \geq 1$  such that for all  $\gamma$ -thick at scale  $L > 0$  subset  $\omega \subset \mathbb{R}^n$ ,*

$$\forall R > 0, \forall f \in L^2(\mathbb{R}^n), \text{Supp } \hat{f} \subset [-R, R]^n, \quad \|f\|_{L^2(\mathbb{R}^n)} \leq \left(\frac{C_n}{\gamma}\right)^{C_n(1+LR)} \|f\|_{L^2(\omega)}.$$

We therefore deduce from this theorem and the definition (3.9) of the orthogonal projection  $K_R$  that there exists a positive constant  $C = C(\omega) \geq 1$  only depending on the thick set  $\omega$  (and not on the positive real number  $R$ ) such that for all  $t > 0$ ,

$$(3.12) \quad \mu_R \frac{d}{dt} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 \leq -2\lambda_R \mu_R C^{-1} e^{-CR} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2.$$

On the other hand, the second term we aim at controlling is given for all  $t > 0$  by

$$(3.13) \quad \begin{aligned} \frac{d}{dt} \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2 &= 2\langle (1 - K_R)f(t), (1 - K_R)f'(t) \rangle_{L^2(\mathbb{R}^n)} \\ &= -2\langle (1 - K_R)f(t), F(|D_x|)(1 - K_R)f(t) \rangle_{L^2(\mathbb{R}^n)} - 2\lambda_R \langle (1 - K_R)f(t), (1 - K_R)1_\omega K_R f(t) \rangle_{L^2(\mathbb{R}^n)}. \end{aligned}$$

We notice that by definition of the orthogonal projection  $K_R$ , the Fourier transforms of the functions  $(1 - K_R)f(t)$  are supported in  $\mathbb{R}^n \setminus B(0, R)$ , which implies that for all  $t > 0$

$$(3.14) \quad 2\langle (1 - K_R)f(t), F(|D_x|)(1 - K_R)f(t) \rangle_{L^2(\mathbb{R}^n)} \geq 2\alpha_R \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2.$$

By using in addition Cauchy-Schwarz' and Young's inequalities, we obtain

$$(3.15) \quad \begin{aligned} -2\lambda_R \langle (1 - K_R)f(t), (1 - K_R)1_\omega K_R f(t) \rangle_{L^2(\mathbb{R}^n)} &\leq 2\lambda_R \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)} \|K_R f(t)\|_{L^2(\mathbb{R}^n)} \\ &\leq \frac{\lambda_R^2}{\alpha_R} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 + \alpha_R \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2. \end{aligned}$$

Combining the estimates (3.13), (3.14) and (3.15), we obtain that for all  $t > 0$ ,

$$\frac{d}{dt} \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2 \leq \frac{\lambda_R^2}{\alpha_R} \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 - \alpha_R \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2.$$

This inequality and (3.12) then imply that for all  $t > 0$ ,

$$\frac{d}{dt} V(f(t)) \leq -2 \left( \lambda_R \mu_R C^{-1} e^{-CR} - \frac{\lambda_R^2}{\alpha_R} \right) \|K_R f(t)\|_{L^2(\mathbb{R}^n)}^2 - \alpha_R \|(1 - K_R)f(t)\|_{L^2(\mathbb{R}^n)}^2.$$

By making the following choices for the constants  $\mu_R$  and  $\lambda_R$ ,

$$\mu_R = 2C^2 e^{2CR}, \quad \lambda_R = C e^{CR} \alpha_R,$$

we get that for all  $t > 0$ ,

$$\frac{d}{dt} V(f(t)) \leq -\alpha_R V(f(t)).$$

This latest estimate and Grönwall's inequality readily imply that for all  $t \geq 0$ ,

$$V(f(t)) \leq e^{-\alpha_R t} V(f(0)),$$

and then, since  $\mu_R \geq 1$ , we obtain

$$\|f(t)\|_{L^2(\mathbb{R}^n)}^2 \leq \mu_R e^{-\alpha_R t} \|f_0\|_{L^2(\mathbb{R}^n)}^2.$$

This ends the proof of the estimate (3.10), and therefore the one of Theorem 2.1.

**3.3. Proof of Proposition 2.5.** In this last subsection, we prove Proposition 2.5 which provides a negative result for the rapid stabilization of the evolution equation ( $E_F$ ). We assume that  $\omega$  is not dense in  $\mathbb{R}^n$  and also that  $\lim_{t \rightarrow +\infty} F(t)$  exists and is a non-negative real number  $L \geq 0$ . Since the function  $F$  is continuous, this implies that  $F$  is bounded  $\sup F < +\infty$ . We aim at proving that if  $\alpha > \sup F$ , then the equation ( $E_F$ ) is not stabilizable from  $\omega$  at rate  $\alpha$ . To that end, we will use the following interpretation of stabilization at rate  $\alpha > 0$  in terms of observability.

**Proposition 3.3** (Theorem 1.1 in [17]). *If the evolution system ( $E_F$ ) is stabilizable from  $\omega$  at rate  $\alpha > 0$ , then there exists a positive constant  $A_\alpha > 0$  such that for all  $T > 0$ , there exists a positive constant  $C_{\alpha, T} > 0$  satisfying that for all  $g \in L^2(\mathbb{R}^n)$ ,*

$$\|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\alpha, T} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + A_\alpha e^{-2\alpha T} \|g\|_{L^2(\mathbb{R}^n)}^2.$$

The proof of Proposition 3.3 is contained in the proof of Theorem 1.1 in [17], although [17] (Theorem 1.1) only states characterizations of complete stabilization. For the sake of completeness, we recall the arguments given by the authors of [17] in Section 6.

Proceeding by contradiction, we consider  $\alpha > \sup F$  and assume that the equation ( $E_F$ ) is stabilizable at rate  $\alpha$  from  $\omega$ . According to Proposition 3.3, there exists a positive

constant  $A_\alpha > 0$  such that for all  $T > 0$ , there exists a positive constant  $C_{\alpha,T} > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$(3.16) \quad \|e^{-TF(|D_x|)}g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\alpha,T} \int_0^T \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt + A_\alpha e^{-2\alpha T} \|g\|_{L^2(\mathbb{R}^n)}^2.$$

Since we get that for all  $T > 0$  and  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-TF(|D_x|)}g\|_{L^2(\mathbb{R}^n)} \geq e^{-T \sup F} \|g\|_{L^2(\mathbb{R}^n)},$$

it follows from (3.16) that for all  $T > 0$  and  $g \in L^2(\mathbb{R}^n)$ ,

$$(3.17) \quad \|g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\alpha,T} e^{2T \sup F} \int_0^T \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt + A_\alpha e^{2(\sup F - \alpha)T} \|g\|_{L^2(\mathbb{R}^n)}^2.$$

Notice that since  $\sup F - \alpha < 0$ , we have  $\lim_{T \rightarrow +\infty} e^{2(\sup F - \alpha)T} = 0$ . This, together with (3.17), implies that there exists a positive constant  $T_0 > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$(3.18) \quad \|g\|_{L^2(\mathbb{R}^n)}^2 \leq 2C_{\alpha,T_0} e^{2T_0 \sup F} \int_0^{T_0} \|e^{-tF(|D_x|)}g\|_{L^2(\omega)}^2 dt.$$

Since  $\omega$  is not dense in  $\mathbb{R}^n$  and that the evolution equation ( $E_F$ ) is invariant under translations, we can assume that there exists a positive radius  $r > 0$  such that  $B(0,r) \subset \mathbb{R}^n \setminus \omega$ . Let us fix a non-zero  $L^2$ -function  $\psi$  and define the function  $g_h$  for all  $h > 0$  by

$$\forall x \in \mathbb{R}^n, \quad g_h(x) = \psi\left(\frac{x}{h}\right).$$

On the one hand, we have that for all  $h > 0$ ,

$$(3.19) \quad \|g_h\|_{L^2(\mathbb{R}^n)}^2 = h^n \|\psi\|_{L^2(\mathbb{R}^n)}^2.$$

On the other hand, we get that for all  $h > 0$ ,

$$(3.20) \quad \begin{aligned} \int_0^{T_0} \|e^{-tF(|D_x|)}g_h\|_{L^2(\omega)}^2 dt &\leq \int_0^{T_0} \|e^{-tF(|D_x|)}g_h\|_{L^2(\mathbb{R}^n \setminus B(0,r))}^2 dt \\ &= h^n \int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 dt. \end{aligned}$$

It follows from (3.18), (3.19) and (3.20) that for all  $h > 0$ ,

$$\|\psi\|_{L^2(\mathbb{R}^n)}^2 \leq 2C_{\alpha,T_0} e^{2T_0 \sup F} \int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 dt.$$

Moreover, since  $L \geq 0$ , we get

$$\begin{aligned} &\int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 dt \\ &\leq 2 \int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi - e^{-tL}\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 dt + 2 \int_0^{T_0} \|e^{-tL}\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 dt \\ &\leq 2 \int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi - e^{-tL}\psi\|_{L^2(\mathbb{R}^n)}^2 dt + 2T_0 \|\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2, \end{aligned}$$

and this implies that

$$\begin{aligned} \|\psi\|_{L^2(\mathbb{R}^n)}^2 &\leq 4C_{\alpha,T_0} e^{2T_0 \sup F} \int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi - e^{-tL}\psi\|_{L^2(\mathbb{R}^n)}^2 dt \\ &\quad + 4C_{\alpha,T_0} T_0 e^{2T_0 \sup F} \|\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2. \end{aligned}$$

To obtain a contradiction, it remains to check that each term of the right-hand side of the above inequality converges to 0 as  $h \rightarrow 0^+$ . For the first term, we deduce from Plancherel's theorem and the dominated convergence theorem that

$$\int_0^{T_0} \|e^{-tF(|D_x|/h)}\psi - e^{-tL}\psi\|_{L^2(\mathbb{R}^n)}^2 dt = \frac{1}{(2\pi)^n} \int_0^{T_0} \|e^{-tF(|\xi|/h)}\widehat{\psi} - e^{-tL}\widehat{\psi}\|_{L^2(\mathbb{R}^n)}^2 dt \xrightarrow{h \rightarrow 0} 0.$$

For the second one, the dominated convergence theorem readily implies that

$$\|\psi\|_{L^2(\mathbb{R}^n \setminus B(0,r/h))}^2 \xrightarrow{h \rightarrow 0} 0.$$

This concludes the proof of Proposition 2.5.

#### 4. QUASI-ANALYTIC SEQUENCES

This section is devoted to recall some properties of quasi-analytic sequences.

**4.1. Notion of quasi-analytic sequence.** Let us first recall that a sequence  $(M_k)_k$  of non-negative real numbers is said to be *log-convex* when it satisfies

$$(4.1) \quad \forall k \geq 1, \quad M_k^2 \leq M_{k+1}M_{k-1}.$$

A relevant example of log-convex sequence is the sequence  $\mathcal{M}^F$  associated with any bounded below continuous function  $F : [0, +\infty) \rightarrow \mathbb{R}$  whose elements  $M_k^F$  are defined for all  $k \geq 0$  by

$$(4.2) \quad M_k^F = \sup_{r \geq 0} r^k e^{-F(r)}.$$

Let us also recall that a sequence  $(M_k)_k$  of positive real numbers is called quasi-analytic when for all real numbers  $a < b$ , the associated Denjoy-Carleman class

$$\mathcal{C}_{\mathcal{M}}([a, b]) = \{f \in \mathcal{C}^\infty([a, b]) : \forall k \geq 0, \forall x \in [a, b], |f^{(k)}(x)| \leq M_k\},$$

is quasi-analytic, meaning that any function belonging to this class is identically equal to zero when satisfying

$$\exists x_0 \in [a, b], \forall k \geq 0, \quad f^{(k)}(x_0) = 0.$$

There exist several necessary and sufficient equivalent conditions that ensure a log-convex sequence to be quasi-analytic. In this work, we will make use of the following one, which is a particular case of Denjoy-Carleman's theorem, see e.g. [13].

**Theorem 4.1** (Denjoy-Carleman). *A log-convex sequence  $(M_k)_k$  of positive real numbers is quasi-analytic if and only if*

$$\sum_{k=0}^{+\infty} \frac{M_k}{M_{k+1}} = +\infty.$$

For all connected open set  $U \subset \mathbb{R}^n$ , let us consider the more general Denjoy-Carleman class associated with a sequence  $\mathcal{M}$  of positive real numbers

$$\mathcal{C}_{\mathcal{M}}(U) = \{f \in \mathcal{C}^\infty(U) : \forall \beta \in \mathbb{N}^n, \|\partial_x^\beta f\|_{L^\infty(U)} \leq M_{|\beta|}\}.$$

We check that when the sequence  $\mathcal{M}$  is quasi-analytic, then all the associated classes  $\mathcal{C}_{\mathcal{M}}(U)$  are also quasi-analytic.

**Proposition 4.2.** *Let  $U \subset \mathbb{R}^n$  be a connected open set and  $\mathcal{M}$  be a quasi-analytic sequence of positive real numbers. Then, the associated class  $\mathcal{C}_{\mathcal{M}}(U)$  is quasi-analytic, meaning that any function  $f$  belonging to this class that vanishes at infinite order in a point of the set  $U$  is identically equal to zero.*

*Proof.* This proof is based on elementary topology arguments. Let  $f \in \mathcal{C}_{\mathcal{M}}(U)$  vanishing at infinite order in a point of the set  $U$ , that is,

$$(4.3) \quad \exists x_0 \in U, \forall \beta \in \mathbb{N}^n, \quad \partial_x^\beta f(x_0) = 0.$$

We set

$$\Omega = \{x \in U : \forall \beta \in \mathbb{N}^n, \partial_x^\beta f(x) = 0\}.$$

The set  $\Omega$  is non empty and closed in  $U$ , since  $f$  is a smooth function. Let us check that this is an open subset of  $U$ . Since  $U$  is connected this will imply that  $\Omega = U$  and therefore that the function  $f$  is identically equal to zero. Let  $x \in \Omega$ . Since  $U$  is open, there

exists a radius  $r > 0$  such that  $\overline{B(x, r)} \subset U$ . Fixing  $y \in B(x, r)$ , we consider the function  $g_y : [-r, r] \rightarrow \mathbb{R}$  defined by

$$g_y(t) = f\left(x + \frac{t(y-x)}{r}\right), \quad t \in [-r, r].$$

Since  $f \in C_{\mathcal{M}}(U)$ , we get that  $g \in C_{\mathcal{M}}([-r, r])$ . Moreover, we deduce from (4.3) that  $g^{(k)}(0) = 0$  for all  $k \geq 0$ . The sequence  $\mathcal{M}$  being quasi-analytic by assumption, this implies that  $g$  is identically equal to zero on  $[-r, r]$ , and so  $f(y) = g(r) = 0$ . The set  $\Omega$  is therefore actually open on  $U$ .  $\square$

We are now in position to state the main proposition of this subsection, which is instrumental in the proof of Theorem 2.6. This result is established in the work [10].

**Proposition 4.3** (Corollary 2.8 in [10]). *Let  $\mathcal{M}$  be a log-convex quasi-analytic sequence,  $t > 0$  be a positive real number and  $\omega \subset (0, 1)^n$  be a measurable set. If  $\text{Leb}(\omega) \geq \lambda$  with  $0 < \lambda \leq 1$ , then there exists a positive constant  $C(\lambda, t, \mathcal{M}, n) > 0$  such that for all  $f \in C_{\mathcal{M}}((0, 1)^n)$  satisfying  $\|f\|_{L^\infty((0, 1)^n)} \geq t$ ,*

$$\|f\|_{L^2((0, 1)^n)} \leq C(\lambda, t, \mathcal{M}, n) \|f\|_{L^2(\omega)}.$$

A quantitative version of this result (with  $L^\infty$  norms) is stated by Nazarov, Sodin and Volberg in [20] and this stronger result provides an explicit dependence of  $C(\lambda, t, \mathcal{M}, n)$  with respect to  $\lambda$ ,  $t$  and  $\mathcal{M}$  when  $n = 1$ .

**4.2. Examples.** Now, we establish useful results to construct quasi-analytic sequences and we also provide examples of such sequences. Let us begin with a straightforward general lemma.

**Lemma 4.4.** *Let  $\mathcal{M}$  and  $\mathcal{M}'$  be two sequences of positive real numbers satisfying  $\mathcal{M} \leq \mathcal{M}'$ . Then, if  $\mathcal{M}'$  is a quasi-analytic sequence, so is the sequence  $\mathcal{M}$ .*

*Proof.* Since  $\mathcal{M} \leq \mathcal{M}'$  by assumption, we get that  $C_{\mathcal{M}}([a, b]) \subset C_{\mathcal{M}'}([a, b])$  for all real numbers  $a < b$  and the result follows.  $\square$

We now focus on the log-convex sequences of the form  $\mathcal{M}^F$  defined in (4.2), which are the ones considered in this work. First, we study the stability of the quasi-analyticity of the sequence  $\mathcal{M}^F$  when modifying the function  $F$ .

**Lemma 4.5.** *If the sequence  $\mathcal{M}^F$  associated with a bounded below continuous function  $F : [0, +\infty) \rightarrow \mathbb{R}$  is quasi-analytic, so is  $\mathcal{M}^{TF+c}$  for all  $c \in \mathbb{R}$  and  $T > 0$ .*

*Proof.* By considering the function  $G = F + c/T$ , whose associated sequence  $\mathcal{M}^G$  is immediately quasi-analytic, we can assume that  $c = 0$ . Let us consider a positive integer  $p \geq 1$  so that  $T \geq 1/p > 0$ . We therefore have that for all  $k \geq 0$ ,

$$M_k^{TF} = \sup_{r \geq 0} r^k e^{-TF(r)} \leq \sup_{r \geq 0} r^k e^{-\frac{1}{p}F(r)} = \left( \sup_{r \geq 0} r^{kp} e^{-F(r)} \right)^{\frac{1}{p}} = (M_{kp}^F)^{\frac{1}{p}} =: M_k.$$

From Lemma 4.4, it is sufficient to check that the sequence  $(M_k)_k$  is quasi-analytic to end the proof of Lemma 4.5. To that end, we will use Denjoy-Carleman's theorem, that is Theorem 4.1, and prove that

$$\sum_{k=0}^{+\infty} \frac{M_k}{M_{k+1}} = +\infty.$$

Notice that since the sequence  $\mathcal{M}^F$  is log-convex, the sequence  $(M_k^F/M_{k+1}^F)_k$  is non-increasing. As a consequence, we get that

$$\sum_{k=0}^{+\infty} \frac{M_k}{M_{k+1}} = \sum_{k=0}^{+\infty} \left( \frac{M_{kp}^F}{M_{k(p+1)}^F} \right)^{\frac{1}{p}} = \sum_{k=0}^{+\infty} \left( \prod_{j=0}^{p-1} \frac{M_{kp+j}^F}{M_{k(p+j)+1}^F} \right)^{\frac{1}{p}} \geq \sum_{k=0}^{+\infty} \frac{M_{kp+p-1}^F}{M_{kp+p}^F}.$$

We therefore need to prove that

$$(4.4) \quad \sum_{k=0}^{+\infty} \frac{M_{kp+p-1}^F}{M_{kp+p}^F} = +\infty.$$

Still using that the sequence  $(M_k^F/M_{k+1}^F)_k$  is non-increasing, we obtain by also applying a Euclidean division by the positive integer  $p \geq 1$  that

$$\sum_{k=0}^{+\infty} \frac{M_k^F}{M_{k+1}^F} = \sum_{r=0}^{p-1} \sum_{k=0}^{+\infty} \frac{M_{kp+r}^F}{M_{kp+r+1}^F} \leq p \sum_{k=0}^{+\infty} \frac{M_{kp}^F}{M_{kp+1}^F}.$$

Yet, the log-convex sequence  $\mathcal{M}^F$  is quasi-analytic by assumption, and Theorem 4.1 implies that

$$(4.5) \quad \sum_{k=0}^{+\infty} \frac{M_k^F}{M_{k+1}^F} = +\infty \quad \text{and so} \quad \sum_{k=0}^{+\infty} \frac{M_{kp}^F}{M_{kp+1}^F} = +\infty.$$

Finally, exploiting a last time the non-increasing property of the sequence  $(M_k^F/M_{k+1}^F)_k$ , we obtain that

$$\sum_{k=0}^{+\infty} \frac{M_{(k+1)p}^F}{M_{(k+1)p+1}^F} \leq \sum_{k=0}^{+\infty} \frac{M_{(k+1)p-1}^F}{M_{(k+1)p}^F}.$$

This inequality and (4.5) show that (4.4) holds. This ends the proof of Lemma 4.5.  $\square$

It is a very interesting problem to characterize the functions  $F$  that generate quasi-analytic sequences  $\mathcal{M}^F$ . This question has been addressed by B. Jaye and M. Mitkovski in [10], where these authors provided a necessary and sufficient condition on some functions  $F$  that ensures the associated sequence  $\mathcal{M}^F$  to be quasi-analytic, by exploiting Denjoy-Carleman's theorem.

**Proposition 4.6** ([10]). *Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a function satisfying*

- (i)  $F(0) = 0$  and  $F$  is non-decreasing with  $\lim_{t \rightarrow +\infty} F(t) = +\infty$ ,
- (ii)  $F$  is lower-semicontinuous and convex.

*Then, the sequence  $\mathcal{M}^F$  associated with the function  $F$  defined in (4.2) is quasi-analytic if and only if*

$$\int_0^{+\infty} \frac{F(t)}{1+t^2} dt = +\infty.$$

The following lemma provides some examples of functions  $F$  generating quasi-analytic sequences  $\mathcal{M}^F$  which are not covered by Proposition 4.6.

**Proposition 4.7.** *Let  $p \geq 1$  be a positive integer and  $F_p : [0, +\infty) \rightarrow [0, +\infty)$  be the non-negative function defined for all  $t \geq 0$  by*

$$F_p(t) = \frac{t}{g(t)(g \circ g)(t) \dots g^{\circ p}(t)}, \quad \text{where } g(t) = \log(e+t),$$

*with  $g^{\circ p} = g \circ \dots \circ g$  ( $p$  compositions). The associated sequence  $\mathcal{M}^{F_p}$  defined in (4.2) is a quasi-analytic sequence of positive real numbers.*

*Proof.* We first notice that the sequence  $\mathcal{M}^{F_p}$  is well-defined by (4.2) as the supremum of a continuous function on  $[0, +\infty)$  with finite limit when  $t \rightarrow +\infty$ . Define  $\phi_p : [0, +\infty) \rightarrow [0, +\infty)$  by

$$\phi_p(t) = g(t)(g \circ g)(t) \dots g^{\circ p}(t), \quad t \geq 0.$$

The function  $\phi_p$  is positive, increasing and it can be readily checked that

$$(4.6) \quad \frac{\phi_p(k)}{\phi_p(2k\phi_p(k))} \xrightarrow{k \rightarrow +\infty} 1.$$

Moreover, by direct computations, we have that for all  $t \geq 0$ ,

$$\begin{aligned} \frac{\phi_p'(t)}{\phi_p(t)} &= \sum_{i=1}^p \frac{(g^{\circ i})'(t)}{g^{\circ i}(t)} = \sum_{i=1}^p \frac{1}{g^{\circ i}(t)} \prod_{j=1}^i g'(g^{\circ(j-1)}(t)) \\ &= \sum_{i=1}^p \frac{1}{g^{\circ i}(t)} \prod_{j=1}^i \frac{1}{e + g^{\circ(j-1)}(t)} \\ &= \frac{1}{e+t} \sum_{i=1}^p \frac{1}{g^{\circ i}(t)} \prod_{j=2}^i \frac{1}{e + g^{\circ(j-1)}(t)} \underset{t \rightarrow +\infty}{=} o\left(\frac{1}{t}\right), \end{aligned}$$

that is

$$(4.7) \quad \phi_p'(t) \underset{t \rightarrow +\infty}{=} o\left(\frac{\phi_p(t)}{t}\right).$$

For all positive integer  $k \geq 1$ , the supremum  $M_k^{F_p} = \sup_{r \geq 0} r^k e^{-F_p(r)}$  is reached at a point  $t_k > 0$  satisfying the following equation

$$(4.8) \quad t_k F_p'(t_k) = k,$$

which is equivalent to

$$\frac{t_k}{\phi_p(t_k)} \left(1 - t_k \frac{\phi_p'(t_k)}{\phi_p(t_k)}\right) = k.$$

Notice that the application  $t \rightarrow t F_p'(t)$  satisfies  $\lim_{t \rightarrow +\infty} t F_p'(t) = +\infty$  since  $\phi_p(t) \underset{t \rightarrow +\infty}{=} o(t)$  and there exists  $T > 0$  such that this application is increasing on  $[T, +\infty)$ . It follows that for all  $k \gg 1$  sufficiently large, there exists a unique  $t_k \in [0, +\infty)$  satisfying (4.8). By defining  $u_k = 2k\phi_p(k)$ , it follows from (4.6) and (4.7) that

$$\frac{u_k}{\phi_p(u_k)} \left(1 - u_k \frac{\phi_p'(u_k)}{\phi_p(u_k)}\right) = 2k \frac{\phi_p(k)}{\phi_p(2k\phi_p(k))} \left(1 - u_k \frac{\phi_p'(u_k)}{\phi_p(u_k)}\right) \underset{k \rightarrow +\infty}{\sim} 2k.$$

As a consequence, we get that for all  $k \gg 1$  sufficiently large,

$$t_k \leq u_k = 2k\phi(k).$$

We deduce that for all  $k \gg 1$ ,

$$\frac{M_{k-1}^{F_p}}{M_k^{F_p}} \geq \frac{t_k^{k-1} e^{-F(t_k)}}{t_k^k e^{-F(t_k)}} = \frac{1}{t_k} \geq \frac{1}{2k\phi(k)}.$$

It is well-known that the series  $\sum \frac{1}{2k\phi(k)}$  is divergent and then, we obtain

$$\sum_{k=0}^{+\infty} \frac{M_k^{F_p}}{M_{k+1}^{F_p}} = +\infty.$$

Since the sequence  $\mathcal{M}^{F_p}$  is log-convex by construction, we deduce from Theorem 4.1 that the sequence  $\mathcal{M}^{F_p}$  is quasi-analytic. This ends the proof of Proposition 4.7.  $\square$

**Corollary 4.8.** *Let  $s \geq 1$ ,  $0 \leq \delta \leq 1$  be non-negative real numbers and  $F_{s,\delta} : [0, +\infty) \rightarrow [0, +\infty)$  be the non-negative continuous function defined by*

$$F_{s,\delta}(t) = \frac{t^s}{\log^\delta(e+t)}, \quad t \geq 0.$$

*The associated sequence  $\mathcal{M}^{F_{s,\delta}}$  defined in (4.2) is a quasi-analytic sequence of positive real numbers.*

*Proof.* Notice that there exists a positive constant  $C_{s,\delta} > 0$  such that for all  $t \geq 0$ ,

$$\frac{t^s}{\log^\delta(e+t)} \geq C_{s,\delta} \left( \frac{t}{\log(e+t)} - 1 \right).$$

Corollary 4.8 is then a consequence of Lemma 4.4, Lemma 4.5 and Proposition 4.7.  $\square$



## 5. PROOF OF THE APPROXIMATE NULL-CONTROLLABILITY RESULTS

This section is devoted to the proofs of the results stated in Section 2.2, dealing with the (cost-uniformly) approximate null-controllability of the evolution equation

$$(E_F) \quad \begin{cases} \partial_t f(t, x) + F(|D_x|)f(t, x) = h(t, x)1_\omega(x), & t > 0, x \in \mathbb{R}^n, \\ f(0, \cdot) = f_0 \in L^2(\mathbb{R}^n), \end{cases}$$

where the bounded below continuous function  $F : [0, +\infty) \rightarrow \mathbb{R}$  is assumed to be associated with a quasi-analytic sequence  $\mathcal{M}^F$  of positive real numbers, defined in (2.2), and  $\omega \subset \mathbb{R}^n$  is a measurable set with positive Lebesgue measure.

**5.1. Proof of Proposition 2.10.** The purpose of this first subsection is to prove that the evolution equation  $(E_F)$  is approximate null-controllable from the support  $\omega$  in any positive time  $T > 0$  (with no extra assumption on  $\omega$ ). It is well-known, see e.g. [6] (Theorem 2.43, page 56) or [21] (Remark 16), that the approximate null-controllability (without uniform cost) of the system  $(E_F)$  is equivalent to a unique continuation property of the adjoint system. More precisely, the adjoint system

$$\begin{cases} \partial_t g(t, x) + F(|D_x|)g(t, x) = 0, & t > 0, x \in \mathbb{R}^n, \\ g(0, \cdot) = g_0 \in L^2(\mathbb{R}^n), \end{cases}$$

is said to satisfy the unique continuation property from  $\omega \subset \mathbb{R}^n$  at some time  $T > 0$  if and only if for all initial datum  $g_0 \in L^2(\mathbb{R}^n)$ , we have

$$(\forall t \in [0, T], 1_\omega e^{-tF(|D_x|)}g_0 = 0) \Rightarrow (e^{-TF(|D_x|)}g_0 = 0).$$

Let  $T > 0$  be a fixed positive time. We aim at proving that the unique continuation property holds from the control support  $\omega$ .

Let us first establish that for all  $g \in L^2(\mathbb{R}^n)$ , the function  $e^{-TF(|D_x|)}g$  belongs to a class of quasi-analytic functions. We consider  $g \in L^2(\mathbb{R}^n)$  not being identically equal to zero (the result is straightforward when  $g = 0$ ). Notice that for all  $\beta \in \mathbb{N}^n$ ,

$$\xi \rightarrow \xi^\beta \mathcal{F}(e^{-TF(|D_x|)}g) \in L^1(\mathbb{R}^n),$$

since we have

(5.1)

$$\begin{aligned} \int_{\mathbb{R}^n} |\xi^\beta \mathcal{F}(e^{-TF(|D_x|)}g)(\xi)| \, d\xi &= \int_{\mathbb{R}^n} |\xi^\beta e^{-TF(|\xi|)}\widehat{g}(\xi)| \, d\xi \\ &\leq (2\pi)^{n/2} \left( \int_{\mathbb{R}^n} |\xi|^{2|\beta|} e^{-2TF(|\xi|)} \, d\xi \right)^{\frac{1}{2}} \|g\|_{L^2(\mathbb{R}^n)} \\ &= (2\pi)^{n/2} \left( \int_{\mathbb{R}^n} \frac{(|\xi|^{2|\beta|} + |\xi|^{2|\beta|+2n})e^{-2TF(|\xi|)}}{1 + |\xi|^{2n}} \, d\xi \right)^{\frac{1}{2}} \|g\|_{L^2(\mathbb{R}^n)} \\ &\leq C_n ((M_{|\beta|}^{TF})^2 + (M_{|\beta|+n}^{TF})^2)^{\frac{1}{2}} \|g\|_{L^2(\mathbb{R}^n)} < +\infty, \end{aligned}$$

the positive constant  $C_n > 0$  being given by

$$C_n = (2\pi)^{n/2} \left( \int_{\mathbb{R}^n} \frac{1}{1 + |\xi|^{2n}} \, d\xi \right)^{\frac{1}{2}} < +\infty.$$

Since the sequence  $\mathcal{M}^{TF}$  is log-convex, there exists a positive constant  $A_{TF} > 0$  such that

$$\forall k \geq 0, \quad M_k^{TF} \leq A_{TF} M_{k+1}^{TF},$$

and we get from (5.1) that for all  $\beta \in \mathbb{N}^n$ ,

$$\int_{\mathbb{R}^n} |\xi^\beta \mathcal{F}(e^{-TF(|D_x|)}g)(\xi)| \, d\xi \leq C_{n,TF} M_{|\beta|+n}^{TF} \|g\|_{L^2(\mathbb{R}^n)},$$

with  $C_{n,TF} = C_n(1 + A_{TF}^{2n})^{\frac{1}{2}}$ . It then follows from the Fourier inverse formula that

$$\forall \beta \in \mathbb{N}^n, \quad \partial_x^\beta (e^{-TF(|D_x|)} g) \in L^\infty(\mathbb{R}^n),$$

and that there exists a positive constant  $C'_{n,TF} > 0$  such that

$$(5.2) \quad \forall \beta \in \mathbb{N}^n, \quad \|\partial_x^\beta (e^{-TF(|D_x|)} g)\|_{L^\infty(\mathbb{R}^n)} \leq C'_{n,TF} M_{|\beta|+n}^{TF} \|g\|_{L^2(\mathbb{R}^n)}.$$

Notice that for all positive real number  $\lambda > 0$ , the sequence  $\mathcal{M}_\lambda$  whose elements  $M_{\lambda,k}$  are given for all  $k \geq 0$  by

$$(5.3) \quad M_{\lambda,k} = C'_{n,TF} \lambda^k M_{k+n}^{TF} \|g\|_{L^2(\mathbb{R}^n)},$$

also defines a log-convex sequence which satisfies

$$\sum_{k=0}^{+\infty} \frac{M_{\lambda,k}}{M_{\lambda,k+1}} = \frac{1}{\lambda} \sum_{k=n}^{+\infty} \frac{M_k^{TF}}{M_{k+1}^{TF}}.$$

Since the sequence  $\mathcal{M}^F$  is quasi-analytic, so is the sequence  $\mathcal{M}^{TF}$  according to Lemma 4.5 and therefore, Theorem 4.1 implies that

$$\sum_{k=0}^{+\infty} \frac{M_k^{TF}}{M_{k+1}^{TF}} = +\infty.$$

A second application of this theorem shows that the log-convex sequence  $\mathcal{M}_\lambda$  is also quasi-analytic. This fact, combined with the estimate (5.2) and Proposition 4.2, implies that for all  $g \in L^2(\mathbb{R}^n)$ , the function  $e^{-TF(|D_x|)} g$  belongs to the quasi-analytic class  $\mathcal{C}_{\mathcal{M}_1}(\mathbb{R}^n)$ .

Now, let us check that the unique continuation property holds. To that end, we make a weaker assumption by considering a function  $g \in L^2(\mathbb{R}^n)$  not identically equal to zero such that

$$(5.4) \quad 1_\omega e^{-TF(|D_x|)} g = 0.$$

Since  $\text{Leb}(\omega) > 0$ , if we consider the cube  $Q = (0, 1)^n$ , then

$$\exists \lambda_0 > 0, \forall \lambda \geq \lambda_0, \quad \text{Leb}(\omega \cap \lambda Q) > 0.$$

Let us consider  $\lambda \geq \lambda_0$  and define the function  $g_\lambda \in L^2(\mathbb{R}^n)$  by

$$g_\lambda(x) = (e^{-TF(|D_x|)} g)(\lambda x), \quad x \in \mathbb{R}^n.$$

It readily follows from the estimate (5.2) that for all  $\beta \in \mathbb{N}^n$ ,

$$\|\partial_x^\beta g_\lambda\|_{L^\infty(Q)} \leq \lambda^{|\beta|} \|\partial_x^\beta (e^{-TF(|D_x|)} g)\|_{L^\infty(\mathbb{R}^n)} \leq C'_{n,TF} \lambda^{|\beta|} M_{|\beta|+n}^{TF} \|g\|_{L^2(\mathbb{R}^n)}.$$

Since the sequences  $\mathcal{M}_\lambda$  defined in (5.3) are quasi-analytic and that  $g_\lambda \in \mathcal{C}_{\mathcal{M}_\lambda}(\mathbb{R}^n)$  from the above estimate, Proposition 4.3 implies that for all  $\lambda \geq \lambda_0$ , there exists a positive constant  $C > 0$  such that

$$\|g_\lambda\|_{L^2(Q)} \leq C \|g_\lambda\|_{L^2((\lambda^{-1}\omega) \cap Q)}.$$

As a consequence, we get from the assumption (5.4) that for all  $\lambda \geq \lambda_0$ ,

$$\|e^{-TF(|D_x|)} g\|_{L^2(\lambda Q)} \leq C \lambda^{-n} \|e^{-TF(|D_x|)} g\|_{L^2(\omega \cap \lambda Q)} = 0.$$

It follows that  $1_{\lambda Q} e^{-TF(|D_x|)} g = 0$  for all  $\lambda \geq \lambda_0$ , and therefore  $e^{-TF(|D_x|)} g = 0$ . Then, the unique continuation property holds and the evolution equation  $(E_F)$  is approximate null-controllable from  $\omega$  at time  $T > 0$ .

**5.2. Proof of Theorem 2.6: sufficient part.** In this subsection, we aim at proving that when the support control  $\omega$  is thick, then the equation  $(E_F)$  is cost-uniformly approximate null-controllable from  $\omega$  in any positive time  $T > 0$ . To that end, we shall use the following observability characterization of the cost-uniformly approximate null-controllability:

**Theorem 5.1** (Proposition 6 in [21]). *The following assertions are equivalent:*

- (i) *The evolution system  $(E_F)$  is cost-uniformly approximately null-controllable from  $\omega$ .*
- (ii) *For all  $\varepsilon \in (0, 1)$  and  $T > 0$ , there exists a positive constant  $C_{\varepsilon, T} > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,*

$$(5.5) \quad \left\| e^{-TF(|D_x|)} g \right\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\varepsilon, T} \int_0^T \left\| e^{-tF(|D_x|)} g \right\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

According to Theorem 5.1, we need to prove that the observability estimate (5.5) holds. The first step consists in establishing the following uncertainty principle:

**Proposition 5.2.** *For all time  $T > 0$  and for all  $\varepsilon > 0$ , there exists a positive constant  $C_{\varepsilon, T} > 0$  such that for all  $t \geq T/2$  and  $g \in L^2(\mathbb{R}^n)$ ,*

$$\left\| e^{-tF(|D_x|)} g \right\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\varepsilon, T} \left\| e^{-tF(|D_x|)} g \right\|_{L^2(\omega)}^2 + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

*Proof.* This proof is adapted from the one of [10] (Theorem 1.3). Since  $\omega$  is a thick set, there exists  $\gamma \in (0, 1]$  and  $L > 0$  by definition such that

$$\forall x \in \mathbb{R}^n, \quad \text{Leb}(\omega \cap (x + [0, L]^n)) \geq \gamma L^n,$$

where  $\text{Leb}$  denotes the Lebesgue measure on  $\mathbb{R}^n$ . For  $\alpha \in LZ^n$ , let us define  $Q(\alpha)$  as the cube centered in  $\alpha$  with sides of length  $L$ , that is,

$$Q(\alpha) = \alpha + [0, L]^n.$$

Notice that the family of cubes  $(Q(\alpha))_{\alpha \in LZ^n}$  covers the space  $\mathbb{R}^n$ :

$$(5.6) \quad \mathbb{R}^n = \bigcup_{\alpha \in LZ^n} Q(\alpha).$$

Let  $g \in L^2(\mathbb{R}^n)$ ,  $T > 0$ ,  $\varepsilon > 0$  and  $t \geq \frac{T}{2}$ . A cube  $Q(\alpha)$  is said to be good if and only if it satisfies the property

$$(5.7) \quad \forall \beta \in \mathbb{N}^n, \quad \left\| \partial_x^\beta e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2 \leq \frac{2^{2|\beta|+n}}{\varepsilon} (M_{|\beta|}^{TF})^2 \left\| e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2,$$

where the positive real numbers  $M_k^{TF}$  are the elements of the sequence  $\mathcal{M}^{TF}$  generated by the function  $TF$ , and given for all  $k \geq 0$  by

$$M_k^{TF} = \sup_{r \geq 0} r^k e^{-TF(r)}.$$

Naturally, a cube  $Q(\alpha)$  is said to be bad if it is not good, that is, when

$$(5.8) \quad \exists \beta_0 \in \mathbb{N}^n, \quad \left\| \partial_x^{\beta_0} e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2 > \frac{2^{2|\beta_0|+n}}{\varepsilon} (M_{|\beta_0|}^{TF})^2 \left\| e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2.$$

Notice from the covering property (5.6) that

$$(5.9) \quad \left\| e^{-tF(|D_x|)} g \right\|_{L^2(\mathbb{R}^n)}^2 = \sum_{\text{good cubes}} \left\| e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2 + \sum_{\text{bad cubes}} \left\| e^{-tF(|D_x|)} g \right\|_{L^2(Q(\alpha))}^2.$$

We will estimate independently the two terms of the right-hand side of this equality. Let us begin with the second one. From Parseval's theorem, we have that for all  $\beta \in \mathbb{N}^n$ ,

$$\begin{aligned}
 (5.10) \quad \|\partial_x^\beta e^{-tF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{-2tF(|\xi|)} |\xi^\beta|^2 |\widehat{g}(\xi)|^2 d\xi \\
 &\leq \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{-2TF(|\xi|)} |\xi|^{2|\beta|} |\widehat{g}(\xi)|^2 d\xi \\
 &\leq \left( \sup_{r \geq 0} r^{|\beta|} e^{-TF(r)} \right)^2 \|g\|_{L^2(\mathbb{R}^n)}^2 = (M_{|\beta|}^{TF})^2 \|g\|_{L^2(\mathbb{R}^n)}^2.
 \end{aligned}$$

It follows from its definition (5.8) that if  $Q(\alpha)$  is a bad cube, there exists  $\beta_0 \in \mathbb{N}^n$  such that

$$\begin{aligned}
 (5.11) \quad \|e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2 &\leq \frac{\varepsilon}{2^{2|\beta_0|+n} (M_{|\beta_0|}^{TF})^2} \|\partial_x^{\beta_0} e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2 \\
 &\leq \sum_{\beta \in \mathbb{N}^n} \frac{\varepsilon}{2^{2|\beta|+n} (M_{|\beta|}^{TF})^2} \|\partial_x^\beta e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2.
 \end{aligned}$$

By summing over all the bad cubes, we obtain from (5.10) and (5.11) that

$$\begin{aligned}
 (5.12) \quad \int_{\bigcup \text{bad cubes}} |e^{-tF(|D_x|)} g(x)|^2 dx &\leq \varepsilon \sum_{\text{bad cubes}} \sum_{\beta \in \mathbb{N}^n} \frac{1}{2^{|\beta|+1} (M_{|\beta|}^{TF})^2} \|\partial_x^\beta e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2 \\
 &\leq \varepsilon \sum_{\beta \in \mathbb{N}^n} \frac{1}{2^{2|\beta|+n} (M_{|\beta|}^{TF})^2} \|\partial_x^\beta e^{-tF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \\
 &\leq \varepsilon \sum_{\beta \in \mathbb{N}^n} \frac{1}{2^{2|\beta|+n}} \|g\|_{L^2(\mathbb{R}^n)}^2 \leq \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2,
 \end{aligned}$$

since

$$\sum_{\beta \in \mathbb{N}^n} \frac{1}{2^{2|\beta|+n}} = \sum_{k=0}^{+\infty} \binom{k+n-1}{n} \frac{1}{2^{2k+n}} \leq \sum_{k=0}^{+\infty} 2^{k+n-1} \frac{1}{2^{2k+n}} = 1.$$

It remains to estimate the first term of the right-hand side of the equality (5.9). Let  $Q(\alpha)$  be a good cube. We define the following multivariable function  $\varphi : [0, 1]^n \rightarrow [0, 1]^n$  by

$$(5.13) \quad \varphi(z) = \frac{L^{\frac{n}{2}} (e^{-tF(|D_x|)} g)(\alpha + Lz)}{\|e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}}, \quad z \in [0, 1]^n.$$

Notice that the function  $\varphi$  satisfies

$$(5.14) \quad \|\varphi\|_{L^\infty([0,1]^n)} = \frac{L^{\frac{n}{2}} \|e^{-tF(|D_x|)} g\|_{L^\infty(Q(\alpha))}}{\|e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}} \geq 1.$$

As the cube  $[0, 1]^n$  satisfies the cone condition, the following Sobolev embedding holds

$$W^{n,2}([0, 1]^n) \hookrightarrow L^\infty([0, 1]^n),$$

see e.g. [1] (Theorem 4.12). This implies that there exists a positive constant  $C_n > 0$ , only depending on the dimension  $n \geq 1$ , such that

$$(5.15) \quad \forall u \in W^{n,2}([0, 1]^n), \quad \|u\|_{L^\infty([0,1]^n)} \leq C_n \|u\|_{W^{n,2}([0,1]^n)}.$$

From the definition (5.7) of good cube and the estimate (5.15), it follows that for all  $\beta \in \mathbb{N}^n$ ,

$$\begin{aligned}
(5.16) \quad \|\partial_x^\beta \varphi\|_{L^\infty([0,1]^n)}^2 &\leq C_n^2 \sum_{\tilde{\beta} \in \mathbb{N}^n, |\tilde{\beta}| \leq n} \|\partial_x^{\beta+\tilde{\beta}} \varphi\|_{L^2([0,1]^n)}^2 \\
&= \frac{C_n^2}{\|e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2} L^{2|\beta|} \sum_{\tilde{\beta} \in \mathbb{N}^n, |\tilde{\beta}| \leq n} L^{2|\tilde{\beta}|} \|\partial_x^{\beta+\tilde{\beta}} e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))}^2 \\
&\leq C_n^2 (2L)^{2|\beta|} 2^n \varepsilon^{-1} \sum_{\tilde{\beta} \in \mathbb{N}^n, |\tilde{\beta}| \leq n} (2L)^{2|\tilde{\beta}|} (M_{|\beta|+|\tilde{\beta}|}^{TF})^2.
\end{aligned}$$

Notice that since the sequence  $\mathcal{M}^{TF}$  is log-convex, there exists a positive constant  $A_{TF} > 0$  such that

$$\forall k \geq 0, \quad M_k^{TF} \leq A_{TF} M_{k+1}^{TF}.$$

Therefore, there exists a new positive constant  $D_{n,L} > 0$  depending on  $n$  and  $L$ , such that for all  $\beta \in \mathbb{N}^n$ ,

$$(5.17) \quad \|\partial_x^\beta \varphi\|_{L^\infty([0,1]^n)}^2 \leq \varepsilon^{-1} A_{TF}^{2n} D_{n,L}^2 (2L)^{2|\beta|} (M_{|\beta|+n}^{TF})^2.$$

Let us recall that by assumption, the sequence  $\mathcal{M}^F$  associated with the function  $F$  defines a quasi-analytic sequence of positive real numbers, and so is  $\mathcal{M}^{TF}$  from Lemma 4.5. Notice that the slightly modified sequence  $\mathcal{M}$  whose elements  $M_k$  are given for all  $k \geq 0$  by

$$M_k = \varepsilon^{-1/2} A_{TF}^n D_{n,L} (2L)^k M_{k+n}^{TF}, \quad k \geq 0,$$

also defines a log-convex sequence that satisfies

$$\sum_{k=0}^{+\infty} \frac{M_k}{M_{k+1}} = \frac{1}{2L} \sum_{k=n}^{+\infty} \frac{M_k^{TF}}{M_{k+1}^{TF}}.$$

Since the sequence  $\mathcal{M}^{TF}$  is quasi-analytic, Theorem 4.1 implies that

$$\sum_{k=0}^{+\infty} \frac{M_k^{TF}}{M_{k+1}^{TF}} = +\infty,$$

and a second application of this theorem provides that the log-convex sequence  $\mathcal{M}$  is also quasi-analytic. Since  $\varphi \in \mathcal{C}_{\mathcal{M}}((0,1)^n)$ , we therefore deduce from (5.14), (5.17) and Proposition 4.3 that there exists a positive constant  $C_{\varepsilon,\gamma,n,T,F} > 0$  independent on  $g$  and  $\alpha$  such that

$$(5.18) \quad \|\varphi\|_{L^2([0,1]^n)} \leq C_{\varepsilon,\gamma,n,T,F} \|\varphi\|_{L^2(E)},$$

where  $E = \frac{\omega-\alpha}{L} \cap [0,1]^n \subset [0,1]^n$  satisfies  $\text{Leb}(E) \geq \gamma > 0$ . It follows directly from (5.13) and (5.18) that

$$(5.19) \quad \|e^{-tF(|D_x|)} g\|_{L^2(Q(\alpha))} \leq C_{\varepsilon,\gamma,n,T,F} \|e^{-tF(|D_x|)} g\|_{L^2(\omega \cap Q(\alpha))}.$$

By summing over all the good cubes, we deduce from (5.6) and (5.19) that

$$\int_{\bigcup_{\text{good cubes}} Q(\alpha)} |e^{-tF(|D_x|)} g(x)|^2 dx \leq C_{\varepsilon,\gamma,n,T,F}^2 \int_{\omega} |e^{-tF(|D_x|)} g(x)|^2 dx.$$

This ends the proof of Proposition 5.2.  $\square$

We can now tackle the proof of the observability estimate (5.5). Let  $\varepsilon > 0$  and  $T > 0$ . It follows from Proposition 5.2 that there exists a positive constant  $C_{\varepsilon,T} > 0$  such that for all  $t \geq T/2$  and  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-tF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\varepsilon,T} \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

By using that for all  $s_1 \geq s_2 \geq 0$  and  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-s_1 F(|D_x|)} g\|_{L^2(\mathbb{R}^n)} \leq \|e^{-s_2 F(|D_x|)} g\|_{L^2(\mathbb{R}^n)},$$

it follows that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$\begin{aligned} \|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 &\leq \frac{2}{T} \int_{\frac{T}{2}}^T \|e^{-tF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 dt \\ &\leq \frac{2C_{\varepsilon,T}}{T} \int_{\frac{T}{2}}^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2 \\ &\leq \frac{2C_{\varepsilon,T}}{T} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2. \end{aligned}$$

This ends the proof of the estimate (5.5).

**5.3. Cost-uniformly approximate null-controllability vs rapid stabilization.** To end this section, let us quickly check that cost-uniformly approximate null-controllability implies rapid stabilization:

**Proposition 5.3.** *If the control system  $(E_F)$  is cost-uniformly approximately null-controllable from  $\omega$  at some positive time  $T > 0$ , then it is rapidly stabilizable from  $\omega$ .*

*Proof.* Assume that the system  $(E_F)$  is cost-uniformly approximately null-controllable from  $\omega$  at some positive time  $T > 0$ . According to Theorem 5.1, the following observability estimate holds: for all  $\varepsilon > 0$ , there exists a positive constant  $C_{\varepsilon,T} > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$(5.20) \quad \|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\varepsilon,T} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + \varepsilon \|g\|_{L^2(\mathbb{R}^n)}^2.$$

In order to prove that  $(E_F)$  is rapidly stabilizable, it is sufficient to show that for all  $\mu > 0$ , the system  $(E_{F_\mu})$  is stabilizable, where  $F_\mu := F - \mu$ . To that end, we shall use Theorem 3.1, already exploited in Section 3.1. From (5.20), by multiplying by  $e^{2T\mu}$ , we get that for all  $\varepsilon > 0$ , there exists a positive constant  $C_{\varepsilon,T} > 0$  such that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-TF_\mu(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\varepsilon,T} \int_0^T \|e^{-tF_\mu(|D_x|)} g\|_{L^2(\omega)}^2 dt + \varepsilon e^{2T\mu} \|g\|_{L^2(\mathbb{R}^n)}^2.$$

By taking  $\varepsilon = \frac{e^{-2T\mu}}{2}$  in the above estimate, it follows from the characterization (iii) of Theorem 3.1 that the system  $(E_{F_\mu})$  is stabilizable from  $\omega$ . Since  $\mu > 0$  can be chosen arbitrary large, we deduce that the system  $(E_F)$  is rapidly stabilizable from  $\omega$ .  $\square$

## 6. APPENDIX

This appendix is devoted to the proof of Proposition 3.3. The following arguments are due to H. Liu, G. Wang, Y. Xu and H. Yu and are originally presented in [17] (Section 2). Let  $F : [0, +\infty) \rightarrow \mathbb{R}$  be a bounded below continuous function and  $\omega \subset \mathbb{R}^n$  be a Borel set with positive Lebesgue measure. Assume that the control system  $(E_F)$  is stabilizable from  $\omega$  at rate  $\alpha > 0$ . We aim at proving that there exists a positive constant  $A_\alpha > 0$  such that for all  $T > 0$ , there exists a positive constant  $C_{\alpha,T} > 0$  satisfying that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq C_{\alpha,T} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + A_\alpha e^{-2\alpha T} \|g\|_{L^2(\mathbb{R}^n)}^2.$$

By definition of stabilization, there exists a positive constant  $M_\alpha > 0$  and a linear feedback  $K_\alpha \in \mathcal{L}(L^2(\mathbb{R}^n))$  such that

$$\forall t \geq 0, \quad \|e^{-t(F(|D_x|) + 1_\omega K_\alpha)}\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \leq M_\alpha e^{-\alpha t}.$$

Let  $T > 0$  fixed. First notice from Duhamel's-type formula for bounded perturbations of semigroups, see e.g. [8] (Corollary III.1.7), that for all  $f \in L^2(\mathbb{R}^n)$ ,

$$e^{-T(F(|D_x|)+1_\omega K_\alpha)} f = e^{-TF(|D_x|)} f - \int_0^T e^{-(T-t)F(|D_x|)} 1_\omega K_\alpha e^{-t(F(|D_x|)+1_\omega K_\alpha)} f dt.$$

By using the fact that the evolution operators  $e^{-tF(|D_x|)}$  are selfadjoint, the above formula implies that for all  $f, g \in L^2(\mathbb{R}^n)$ ,

$$\begin{aligned} & \langle e^{-TF(|D_x|)} g, f \rangle_{L^2(\mathbb{R}^n)} = \langle g, e^{-TF(|D_x|)} f \rangle_{L^2(\mathbb{R}^n)} \\ &= \langle g, e^{-T(F(|D_x|)+1_\omega K_\alpha)} f \rangle_{L^2(\mathbb{R}^n)} + \int_0^T \langle g, e^{-(T-t)F(|D_x|)} 1_\omega K_\alpha e^{-t(F(|D_x|)+1_\omega K_\alpha)} f \rangle_{L^2(\mathbb{R}^n)} dt \\ &= \langle g, e^{-T(F(|D_x|)+1_\omega K_\alpha)} f \rangle_{L^2(\mathbb{R}^n)} + \int_0^T \langle 1_\omega e^{-(T-t)F(|D_x|)} g, K_\alpha e^{-t(F(|D_x|)+1_\omega K_\alpha)} f \rangle_{L^2(\mathbb{R}^n)} dt \\ &\leq M_\alpha e^{-\alpha T} \|f\|_{L^2(\mathbb{R}^n)} \|g\|_{L^2(\mathbb{R}^n)} + \|K_\alpha\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \|f\|_{L^2(\mathbb{R}^n)} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)} dt. \end{aligned}$$

By applying the above inequality to  $f = e^{-TF(|D_x|)} g$ , it follows that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)} \leq \|K_\alpha\|_{\mathcal{L}(L^2(\mathbb{R}^n))} \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)} dt + M_\alpha e^{-\alpha T} \|g\|_{L^2(\mathbb{R}^n)}.$$

Finally, we deduce from Hölder's inequality and the classical convexity inequality

$$\forall a, b > 0, \quad (a + b)^2 \leq 2(a^2 + b^2),$$

that for all  $g \in L^2(\mathbb{R}^n)$ ,

$$\|e^{-TF(|D_x|)} g\|_{L^2(\mathbb{R}^n)}^2 \leq 2\|K_\alpha\|_{\mathcal{L}(L^2(\mathbb{R}^n))}^2 T \int_0^T \|e^{-tF(|D_x|)} g\|_{L^2(\omega)}^2 dt + 2M_\alpha^2 e^{-2\alpha T} \|g\|_{L^2(\mathbb{R}^n)}^2,$$

and this concludes the proof of Proposition 3.3.

## REFERENCES

- [1] R.A. ADAMS, J.J.F. FOURNIER, *Sobolev spaces*, Second edition, Pure and Applied Mathematics (Amsterdam), 140. Elsevier/Academic Press, Amsterdam (2003).
- [2] P. ALPHONSE, *Régularité des solutions et contrôlabilité d'équations d'évolution associées à des opérateurs non-autoadjoints*, PhD Thesis, Université de Rennes 1 (2020).
- [3] P. ALPHONSE, J. BERNIER, *Smoothing properties of fractional Ornstein-Uhlenbeck semigroups and null-controllability*, Bull. Sci. Math. 165 (2020), p. 102914.
- [4] K. BEAUCHARD, M. EGIDI, K. PRAVDA-STAROV, *Geometric conditions for the null-controllability of hypoelliptic quadratic parabolic equations with moving control supports*, C. R. Math. Acad. Sci. Paris 358 (2020), no. 6, pp. 651-700.
- [5] K. BEAUCHARD, P. JAMING, K. PRAVDA-STAROV, *Spectral inequality for finite combinations of Hermite functions and null-controllability of hypoelliptic quadratic equations*, to appear in Studia Math. (2020).
- [6] J.-M. CORON, *Control and nonlinearity*, Mathematical Surveys and Monographs, Vol. 136, American Mathematical Society, Providence, RI (2007).
- [7] M. EGIDI, I. VESELIĆ, *Sharp geometric condition for null-controllability of the heat equation on  $\mathbb{R}^d$  and consistent estimates on the control cost*, Arch. Math. (Basel) 111 (2018), no. 1, pp. 85-99.
- [8] K.-J. ENGEL, R. NAGEL, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, Vol 194, Springer-Verlag, New York (2000).
- [9] S. HUANG, G. WANG, M. WANG, *Characterizations of stabilizable sets for some parabolic equations in  $\mathbb{R}^n$* , J. Differential Equations 272 (2021), pp. 255-288.
- [10] B. JAYE, M. MITKOVSKI, *Quantitative uniqueness property for  $L^2$  functions with fast decaying, or sparsely supported, Fourier transform*, preprint (2018), [arXiv:1808.02149](https://arxiv.org/abs/1808.02149).
- [11] A. KOENIG, *Lack of null-Controllability for the fractional heat equation and related equations*, SIAM J. Control Optim. 58 (2020), no. 6, pp. 3130-3160.
- [12] A. KOENIG, *Contrôlabilité de quelques équations aux dérivées partielles paraboliques peu diffusives*, Ph.D thesis, Université Côte d'Azur (2019).
- [13] P. KOOSIS, *The Logarithmic Integral I*, Cambridge University Press, 1988.
- [14] O. KOVRIJKINE, *Some results related to the Logvinenko-Sereda Theorem*, Proc. Amer. Math. Soc. 129, (2001), no. 10, pp. 3037-3047.

- [15] G. LEBEAU, L. ROBBIANO, *Contrôle exact de l'équation de la chaleur*, Comm. Partial Differential Equations 20 (1995), no. 1-2, pp. 335-356.
- [16] P. LISSY, *A non-controllability result for the half-heat equation on the whole line based on the prolate spheroidal wave functions and its application to the Grushin equation*, preprint (2020), [hal-02420212](#).
- [17] H. LIU, G. WANG, Y. XU, H. YU, *Characterizations on complete stabilizability*, preprint (2020), [arXiv:2012.07253](#)
- [18] J. MARTIN, K. PRAVDA-STAROV, *Geometric conditions for the exact controllability of fractional free and harmonic Schrödinger equations*, to appear in J. Evol. Equ. (2020), [arXiv:2007.04096](#).
- [19] L. MILLER, *A direct Lebeau-Robbiano strategy for the observability of heat-like semigroups*, Discrete Contin. Dyn. Syst. Ser. B 14 (2010), no. 4, pp. 1465-1485.
- [20] F. NAZAROV, M. SODIN, A. VOLBERG, *Lower bounds for quasianalytic functions, I. How to control smooth functions?*, Math. Scand. 95 (2004), no. 1, pp. 59-79.
- [21] E. TRÉLAT, G. WANG, Y. XU, *Characterization by observability inequalities of controllability and stabilization properties*, Pure Appl. Anal. 2 (2020), no. 1, pp. 93-122.
- [22] G. WANG, M. WANG, C. ZHANG, Y. ZHANG, *Observable set, observability, interpolation inequality and spectral inequality for the heat equation in  $\mathbb{R}^n$* , J. Math. Pures Appl. (9) 126 (2019), pp. 144-194.
- [23] S. XIANG, *Quantitative rapid and finite time stabilization of the heat equation*, preprint (2020), [arXiv:2010.04696](#).

(PAUL ALPHONSE) UNIVERSITÉ DE LYON, ENSL, UMPA - UMR 5669, F-69364 LYON  
Email address: [paul.alphonse@ens-lyon.fr](mailto:paul.alphonse@ens-lyon.fr)

(JÉRÉMY MARTIN) UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES  
Email address: [jeremy.martin@ens-rennes.fr](mailto:jeremy.martin@ens-rennes.fr)