



HAL
open science

APOGEE view of the globular cluster NGC 6544

F Gran, M Zoccali, A Rojas-Arriagada, I Saviane, R Contreras ramos, R Beaton, D Bizyaev, R Cohen, J.G. Fernández-Trincado, D García-Hernández, et al.

► **To cite this version:**

F Gran, M Zoccali, A Rojas-Arriagada, I Saviane, R Contreras ramos, et al.. APOGEE view of the globular cluster NGC 6544. *Monthly Notices of the Royal Astronomical Society*, 2021, 504 (3), pp.3494-3508. 10.1093/mnras/stab1051 . hal-03592365

HAL Id: hal-03592365

<https://hal.science/hal-03592365v1>

Submitted on 20 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

APOGEE view of the globular cluster NGC 6544

F. Gran¹,^{2,3}★ M. Zoccali^{1,2}, A. Rojas-Arriagada^{1,2}, I. Saviane,³ R. Contreras Ramos,^{1,2} R. Beaton,⁴ D. Bizyaev^{5,6}, R. E. Cohen,⁷ J. G. Fernández-Trincado,^{8,9,10} D. A. García-Hernández,^{11,12} D. Geisler,^{13,14,15} R. R. Lane,^{8,10} D. Minniti,^{16,17} C. Moni Bidin,¹⁸ C. Nitschelm,¹⁹ J. Olivares Carvajal,^{1,2} K. Pan,⁵ F. I. Rojas^{1,2} and S. Villanova¹³

¹Instituto de Astrofísica, Av. Vicuña Mackenna 4860, Santiago, Chile

²Instituto Milenio de Astrofísica, Santiago, Chile

³European Southern Observatory, Alonso de Córdova 3107, Santiago, Chile

⁴Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08544, USA

⁵Apache Point Observatory and New Mexico State University, P.O. Box 59, Sunspot, NM 88349-0059, USA

⁶Sternberg Astronomical Institute, Moscow State University, Moscow, Russia

⁷Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁸Instituto de Astronomía y Ciencias Planetarias, Universidad de Atacama, Copayapu 485, Copiapó, Chile

⁹Institut Utinam, CNRS UMR 6213, Université Bourgogne-Franche-Comté, OSU THETA Franche-Comté, Observatoire de Besançon, BP 1615, F-25010 Besançon Cedex, France

¹⁰Centro de Investigación en Astronomía, Universidad Bernardo O'Higgins, Avenida Viel 1497, Santiago, Chile

¹¹Instituto de Astrofísica de Canarias (IAC), E-38205 La Laguna, Tenerife, Spain

¹²Universidad de La Laguna (ULL), Departamento de Astrofísica, E-38206 La Laguna, Tenerife, Spain

¹³Departamento de Astronomía, Casilla 160-C, Universidad de Concepción, Concepción, Chile

¹⁴Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, Universidad de La Serena, Avenida Raúl Bitrán S/N, La Serena, Chile

¹⁵Departamento de Astronomía, Facultad de Ciencias, Universidad de La Serena, Av. Juan Cisternas 1200, La Serena, Chile

¹⁶Depto. de Ciencias Físicas, Facultad de Ciencias Exactas, Universidad Andrés Bello, Av. Fernández Concha 700, Las Condes, Santiago, Chile

¹⁷Vatican Observatory, V-00120 Vatican City State, Italy

¹⁸Instituto de Astronomía, Universidad Católica del Norte, Avenida Angamos 0610, Antofagasta, Chile

¹⁹Centro de Astronomía (CITEVA), Universidad de Antofagasta, Avenida Angamos 601, Antofagasta 1270300, Chile

Accepted 2021 April 2. Received 2021 March 2; in original form 2020 August 20

ABSTRACT

The second phase of the APOGEE survey is providing near-infrared (near-IR), high-resolution, high signal-to-noise spectra of stars in the halo, disc, bar, and bulge of the Milky Way. The near-IR spectral window is especially important in the study of the Galactic bulge, where stars are obscured by the dust and gas of the disc in its line of sight. We present a chemical characterization of the globular cluster NGC 6544 with high-resolution spectroscopy. The characterization of the cluster chemical fingerprint, given its status of ‘interloper’ towards the Galactic bulge and clear signatures of tidal disruption in its core is crucial for future chemical tagging efforts. Cluster members were selected from the DR16 of the APOGEE survey, using chemodynamical criteria of individual stars. A sample of 23 members of the cluster was selected. An analysis considering the intracluster abundance variations, known as anticorrelations is given. According to the red giant branch (RGB) content of the cluster, the iron content and α -enhancement are $[\text{Fe}/\text{H}] = -1.44 \pm 0.04$ dex and $[\alpha/\text{Fe}] = 0.20 \pm 0.04$ dex, respectively. Cluster members show a significant spread in $[\text{Fe}/\text{H}]$ and $[\text{Al}/\text{Fe}]$ that is larger than expected based on measurement errors. An $[\text{Al}/\text{Fe}]$ spread, signal of an Mg–Al anticorrelation is observed and used to constrain the cluster mass budget, along with C, N, Mg, Si, K, Ca, and Ce element variations discussed. Across all the analysed evolutionary stages (RGB and asymptotic giant branch), about $\sim 2/3$ (14 out of 23) show distinct chemical patterns, possibly associated with second-generation stars.

Key words: surveys – proper motions – stars: abundances – stars: evolution – Galaxy: bulge – globular clusters: individual: NGC 6544.

1 INTRODUCTION

As the oldest objects in our Galaxy, Galactic globular clusters (GCs) play a crucial role in the characterization of the early phases of its

formation. While most halo clusters have been extensively studied, GCs close to the plane of the Milky Way (MW) have not been equally explored. The variable amount of gas and dust and the contamination by disc and bulge stars in the disc and bulge line of sights were the main reasons to evade observations in this area. Bulge GCs, in particular, that can be associated with this Galactic component based on their positions, kinematics, and metallicities (e.g. Minniti 1995;

* E-mail: fegran@uc.cl

Table 1. NGC 6544 coordinates (J2000), tidal radius, and distance from the Sun (Cohen et al. 2014), metallicity (Harris 1996, 2010 edition), and PMs (Baumgardt et al. 2019).

RA (hh:mm:ss)	Dec. (dd:mm:ss)	ℓ (deg)	b (deg)	r_t (arcmin)	d_\odot (kpc)	[Fe/H] (dex)	$\mu_\alpha \cos \delta$ (mas yr $^{-1}$)	μ_δ (mas yr $^{-1}$)
18:07:20.12	−24:59:53.6	5.8365	−2.2024	$\simeq 19$	2.46 ± 0.09	−1.4	$−2.34 \pm 0.04$	$−18.66 \pm 0.04$

Pérez-Villegas et al. 2019), have been avoided until recently, when near-infrared (near-IR) detectors and spectrographs allowed us to reduce the effect of extinction (e.g. Valenti, Ferraro & Origlia 2007, 2010; Bica, Ortolani & Barbuy 2016; Cohen et al. 2017, 2018).

NGC 6544 ($\ell = 5.83$, $b = -2.20$; Cohen et al. 2014) is a perfect example of a poorly studied GC, despite being at a distance of only ~ 2.5 kpc in the bulge direction. The first attempt to characterize NGC 6544 come from 160 stars measured with photographic plates, Alcaino (1983) reported a very reddened cluster, with $E(B - V) = 0.70$ mag, at a distance of 2.8 kpc. During the last decade, Valenti et al. (2010) analysed the first near-IR CCD photometry of the cluster and used the slope of the red giant branch (RGB) to derive a metallicity of $[\text{Fe}/\text{H}] = -1.36$ dex. Based on low-resolution spectroscopy, Carretta et al. (2009b) and Saviane et al. (2012) found consistent results of the cluster metallicity with $[\text{Fe}/\text{H}]$ of -1.47 and -1.43 dex, respectively, both slightly lower than the photometric determination. Recently, Cohen et al. (2014) combined near-IR photometry from the VISTA Variable in the Vía Láctea survey (VVV; Minniti et al. 2010; Saito et al. 2012) with optical photometry from the *Hubble Space Telescope*, deriving $d = 2.46$ kpc, $E(B - V) = 0.79$ mag, and constraining the tidal radius with a lower limit of $r_t = 19$ arcmin. They also concluded that the cluster is likely to have undergone a tidal stripping process, under the assumption that NGC 6544 is a halo cluster currently passing close to the Galactic disc, and projected towards the bulge (Bica et al. 2016).

Recently, Contreras Ramos et al. (2017), Pérez-Villegas et al. (2019), and Massari, Koppelman & Helmi (2019) confirmed that the cluster origin was not the bulge based on absolute proper motions (PMs), from which they derive the orbit of NGC 6544. Interestingly, Contreras Ramos et al. (2017) found a remarkable elongation of the shape of the cluster in the direction of the Galactic Centre, unfortunately without any tidal tail signature. Finally, and as a side product of the numerous efforts to characterize the GC content in the *Gaia* Data Release 2 (DR2) catalogue (Gaia Collaboration 2018a,b; Lindegren et al. 2018), Vasiliev (2019), Baumgardt & Hilker (2018), and Baumgardt et al. (2019) derived several structural and photometric properties for NGC 6544. This is how very precise estimates, based solely on cluster members of the distance, PMs, core, tidal and half-light radii, and mass were derived.

This paper presents the first high-resolution spectral analysis of 23 NGC 6544 cluster members, to characterize its internal chemical composition and compare it with other GCs and the bulge field. The structure of this paper is as follows: Section 2 describes the used data, Section 3 describes cluster members' selection process, Section 4 provides the metallicity determination for the cluster, Section 5 refers to the chemical abundances of the individual stars and its variations within the cluster, Section 6 gives a context of our findings within a sample of GCs, and finally, Section 7 presents our concluding remarks.

2 APOGEE SURVEY DATA

The APOGEE survey (Majewski et al. 2017) is one of the projects carried out within the SDSS-III (Eisenstein et al. 2011) and the SDSS-

IV (Blanton et al. 2017) Collaboration. APOGEE uses a multifibre spectrograph (Wilson et al. 2010, 2012, 2019) to obtain high-resolution ($R \sim 22\,000$) high-signal-to-noise ratio (SNR) (> 100) spectra in near-IR wavelengths (H band; $1.51\text{--}1.69$ μm) in order to observe all the components of the MW (halo, disc, bar, and bulge) from the Apache Point Observatory (APO; Gunn et al. 2006) and Las Campanas Observatory (LCO; Bowen & Vaughan 1973). The survey targets mainly giant stars imposing a colour cut in the dereddened near-IR colour–magnitude diagram (CMD; Zasowski et al. 2013, 2017). After the observations, all the data collected are processed by the APOGEE pipeline (Nidever et al. 2015) to extract and co-add multiple visits of the same target and calculate accurate radial velocities to the level of 0.1 km s $^{-1}$ for an SNR > 20 star with at least three observations. Finally, all the individual and co-added spectra are processed by the APOGEE stellar parameter and chemical abundance pipeline (ASPCAP; García Pérez et al. 2016) to derive abundances for up to 26 chemical elements (Shetrone et al. 2015; Smith et al. 2021) in the publicly available Data Release 16 (DR16; Ahumada et al. 2020). The DR16 version of the APOGEE catalogue is presented by Jönsson et al. (2020), and it includes data for more than $\sim 437\,000$ stars across all the components in the MW.

3 NGC 6544 IN THE APOGEE DR16

We searched for stars that might belong to known GCs in the APOGEE public DR16, restricted to the bulge area observed by the VVV survey footprint. Originally, NGC 6544 was not part of the APOGEE-1 calibration set of clusters (observational campaign from APO), and targeting efforts of cluster members started with APOGEE-2 (APO and LCO observations). In total, more than 690 stars lie within 45 arcmin of the cluster centre. The spatial selection cut was performed using the cluster centre derived by Cohen et al. (2014), as listed in Table 1, whose centre is shifted by 7 arcsec with respect to the value provided by the Harris (1996, 2010 edition) catalogue. Most of the stars within this radius are bulge stars, although 26 stars group around a V_{HELIO} and $[\text{Fe}/\text{H}]$ loci far from the main field distribution, as shown in Fig. 1 (left-hand panel). Note that of the 26 stars selected, there are three with repeated observations from both hemispheres (APO and LCO).

From here on, we will merge the entries selecting the ones with lower overall errors (APO). Shown here are the heliocentric velocity and iron abundance derived from the APOGEE reduction pipeline and ASPCAP, respectively. The cluster members tend to agglomerate in a region of the $V_{\text{HELIO}}\text{--}[\text{Fe}/\text{H}]$ space marked with a box with $[\text{Fe}/\text{H}]_{\text{median}} = -1.46$ dex and $V_{\text{HELIO,median}} = -38.17$ km s $^{-1}$. The radial velocity limits of the box have been set as the maximum dispersion found in massive GCs (McLaughlin & van der Marel 2005; Watkins et al. 2015) of $\lesssim 20$ km s $^{-1}$, while for the iron content, we select all the stars with $-2.0 \lesssim [\text{Fe}/\text{H}] \lesssim -1.4$ because there are no more stars in this range to be considered cluster members.

As an additional test, we cross-match the same bulge field stars around NGC 6544 with the *Gaia* DR2 to obtain PMs with the idea of applying a kinematic cut in our sample. For all the cluster candidates, we found a match that gives us the PMs ($\mu_\alpha \cos \delta$ and μ_δ in mas yr $^{-1}$),




Figure 1. Left-hand panel: Heliocentric radial velocity versus $[\text{Fe}/\text{H}]$ for all the APOGEE targets within 45 arcmin from the centre of NGC 6544. A group of stars, candidate cluster members, clearly separate from the bulk of field (bulge) stars. See text for the box limits. Right-hand panel: VPD for the same field around the cluster. A small concentration of stars can be seen as a coherent group isolated from the bulge field stars. We choose the radius of the circle as 1.5 mas yr^{-1} . Note that the RR Lyrae star used to derive the distance to the cluster is marked with a cross.

which we show as the vector-point diagram (VPD) in Fig. 1 (right-hand panel). We select a boundary limit of 1.5 mas yr^{-1} to select cluster members within the VPD. The median PM of the matched APOGEE stars shows a perfect agreement with the value published by Vasiliev (2019) and Baumgardt et al. (2019). We want to perform the cluster selection process as blind as possible therefore we do not put additional parameter constraints.

In total, the same initial 23 stars were selected as probable NGC 6544 members. The final selection of cluster candidates is also shown on their position in the dereddened $(J-K_s)$ versus K_s CMD, using the 2MASS magnitudes (Skrutskie et al. 2006) transformed to the VISTA filters.¹ Fig. 2 shows the CMD of NGC 6544 with the selected APOGEE targets shown with symbols. For stars with $K_s > 12 \text{ mag}$, VVV dereddened magnitudes come from the PM selected cluster members, according to Contreras Ramos et al. (2017). However, for $K_s < 12 \text{ mag}$, stars are saturated in the VVV survey, and therefore we perform a dynamical selection (with 1.5 mas yr^{-1} tolerance) based on the *Gaia* DR2 PMs to select 2MASS members (Skrutskie et al. 2006; Valenti et al. 2007). Three isochrones from the PARSEC library (Bressan et al. 2012; Chen et al. 2014, 2015; Tang et al. 2014; Marigo et al. 2017; Pastorelli et al. 2019), were overlotted to the data. The predefined parameters were used to retrieve the isochrones in the version 1.2S. We choose ages of 10, 12, and 14 Gyr that can be noticed from left to right in the turn-off area. As expected for the given ages, all the isochrones follow an almost identical line in the main sequence, RGB, and asymptotic giant branch (AGB). The same metallicity and α -enhancement derived for the cluster (see Section 5 for the exact values) were used.

The inset in Fig. 2 also shows the $T_{\text{eff}}-\log g$ diagram for the observed stars with the isochrones mentioned above. The isochrones help identify the RGB bump in the inset panel, highlighted with two intersecting shaded areas at $T_{\text{eff}} \sim 4950 \text{ K}$ and $\log g \sim 2.0$, splitting the RGB sample into two. Note that all the stars in the $T_{\text{eff}}-\log g$ diagram appear to be shifted to the respective isochrones towards cooler temperatures or higher gravities. The shift is small, and similar to the total error on the parameters, with values of $\Delta T_{\text{eff}} \sim 100 \text{ K}$




Figure 2. Dereddened $(J-K_s) \times K_s$ CMD of NGC 6544 members from VVV ($K_s \gtrsim 11 \text{ mag}$; Contreras Ramos et al. 2017) and 2MASS ($K_s \lesssim 11 \text{ mag}$; Skrutskie et al. 2006; Valenti et al. 2007). As mentioned in the text, different symbols were used to differentiate within stellar stages. Circles, triangles, and stars were used to highlight different evolutionary stages based on their positions on the $T_{\text{eff}}-\log g$ diagram (see Section 3 for definitions). Subsequent figures will use the same markers. The RR Lyrae star analysed in Section 4.2 is marked with a cross. Inset: $T_{\text{eff}}-\log g$ diagram of the 23 APOGEE cluster candidates together with the same three PARSEC isochrones. All of them for the same metallicity and α -enhancement values of the cluster, for ages of 10, 12, and 14 Gyr. The RGB bump is marked at the intersection of the two shaded areas.

¹<http://casu.ast.cam.ac.uk/surveys-projects/vista/technical/photometric-properties>

and $\Delta \log g \sim 0.2$ dex in the uncorrelated case and half of them if we consider a simultaneous shift on both measurements. Despite this fact and as expected for a single GC, stars follow the evolutionary sequences, confirming that the stellar surface parameters are rather well determined. On the contrary, the targets are more spread out in the observed CMD because of the large differential reddening along the line of sight: up to $\Delta E(J - K_s) \sim 0.4$ mag within arcmin scales around the cluster centre, as already noted by Contreras Ramos et al. (2017). It is for this reason that we only use the position of the stars in the $T_{\text{eff}} - \log g$ diagram to classify the cluster members into three groups: the main two in the RGB evolutionary phase, below and above the bump and the third one, associated with the AGB phase. Take in consideration that the RGB bump might differ whether we select a stellar model or perform an empirical estimate, as shown in Cohen et al. (2017) for NGC 6544. The shift in magnitudes can be as large as $\Delta K_s \sim 0.2 - 0.3$ mag for this cluster, depending on the derived distance, α -enhancement, and to a lesser extent by age (Salaris et al. 2007).

As Fig. 2 suggests, we split the cluster sample into three groups: stars below and above the RGB bump (lower RGB or IRGB as circles, and upper RGB or uRGB as triangles), and the AGB group (as stars). Each group consists of 13, 6, and 4 stars for the IRGB, uRGB, and AGB, respectively, with a total of 23 observed cluster stars. These groups were also marked with different symbols according to APOGEE derived T_{eff} and $\log g$ values.

In Sections 4 and 5, abundance differences will be reviewed, but as expected from an evolutionary point of view, stars below the RGB bump will be carbon enhanced and nitrogen depleted, and the contrary for the stars above the bump.

4 NGC 6544 FUNDAMENTAL PARAMETERS

4.1 Metallicity and α -enhancement values

From the photometric point of view, some of the most uncertain parameters to determine from a CMD are the metallicity, α -enhancement, and age of a stellar population. Even a slight change in the first two affects the main-sequence turn-off point location, and therefore the estimated age of a stellar population (e.g. Oliveira et al. 2020; Souza et al. 2020). Here, the cluster metallicity is known in great detail even across different stages of stellar evolution, and the combination of PMs and radial velocities ensures that all the analysed stars move coherently in space and form part of the cluster.

One of the first striking results that the spectra present is the systematic difference between the DR16 derived metallicity of the RGB (both groups) and that of the AGB. The marked difference between populations of a GC was not present in previous studies, while using APOGEE spectra but applying different reduction methods (García-Hernández et al. 2015; Masseron et al. 2019; Mészáros et al. 2020). In this cluster, we are finding that the median metallicity with its standard error for the entire RGB is $[\text{Fe}/\text{H}] = -1.44 \pm 0.04$ dex, that is composed of a lower and upper part with -1.44 ± 0.05 and -1.46 ± 0.04 dex, respectively. RGB metallicities agree within its errors, nevertheless are systematically higher than those of AGB stars of $[\text{Fe}/\text{H}] = -1.66 \pm 0.10$ dex. Across the paper, we use the standard deviation of the median as a statistically significant measurement of the errors.

The discrepant values of iron content for stars in different evolutionary stages (RGB and AGB in this case) are not a new finding. Indeed, Ivans et al. (2001) found that AGB stars in M5 had $[\text{Fe}/\text{H}]$ lower than RGB stars, by 0.15 dex. They concluded that this finding was in agreement with the prediction of Thévenin & Idiart

(1999), i.e. that in the atmosphere of metal poor, late-type stars, most Fe I lines are formed out of local thermal equilibrium (LTE), and therefore by assuming LTE, derived Fe I abundances are lower than real. This is due to the fact that in thin, transparent atmospheres, some of the Fe is ionized by UV radiation coming from the stellar interior. The effect is larger for less dense atmospheres at a given T_{eff} , and also larger for higher T_{eff} at a given luminosity, therefore it is expected to be larger in AGB stars than in RGB stars. According to Thévenin & Idiart (1999), Fe II lines are not affected by this non-LTE effect. More recently and using optical spectra, Lapenna et al. (2014, 2015) and Mucciarelli et al. (2015a,b) all confirmed that the putative spread in the iron abundance of stars in 47Tuc, M62, M22, NGC 3201, reported in previous studies, was due to the same effect. Namely, Fe I abundances in AGB stars were lower than those of RGB stars. On the contrary, iron abundances derived from Fe II lines showed a very good agreement in both evolutionary phases. However, converging metallicities between the RGB and AGB stages were found by García-Hernández et al. (2015), Masseron et al. (2019), and Mészáros et al. (2020) when characterizing APOGEE near-IR spectra that uses only Fe I lines. With these conflicting evidence in context, it is clear that we have no clear explanation of the phenomena that NGC 6544 AGB stars exhibit, giving that they are dynamically confirmed cluster members. For that reason, we have isolated the behaviour of the different evolutionary stages to compute the median $[\text{Fe}/\text{H}]$.

A max-to-mean difference of 0.23 dex is observed in $[\text{Fe}/\text{H}]$ (or 0.17 dex within each RGB subdivision), a value that exceeds the observational errors in NGC 6544 (typically ± 0.02 dex), and in other GCs (Carretta et al. 2009b). It will be evident from Section 5 that the star-to-star variations exceed the observational error, even among the RGB and AGB stars themselves. We cannot exclude that extra-enrichment processes happened in NGC 6544 (as explained in Renzini 2008), or even that this trend is the result of the interaction with the MW (Contreras Ramos et al. 2018a; Kundu, Minniti & Singh 2019a; Kundu et al. 2019b). Given the absolute V magnitude (M_V) of -6.94 mag as a proxy of the total mass, no other cluster with similar M_V shows an $[\text{Fe}/\text{H}]$ dispersion greater than ~ 0.07 dex (Harris 1996; Bailin 2019). Up to now, we do not have a clear explanation of the origin of the $[\text{Fe}/\text{H}]$ spread of NGC 6544, considering the limited number of stars observed.

Finally, the ASPCAP pipeline returns an estimation for the $[\alpha/\text{Fe}]$ content of each analysed star. Following the description in García Pérez et al. (2016), this contribution is measured using O, Mg, Si, S, Ca, and Ti from the spectra. The median $[\alpha/\text{Fe}]$ value obtained from cluster combined RGB stars is 0.20 ± 0.04 dex, matching canonical values for GCs. If we separate stars in the lower and upper part of the RGB, we find medians of 0.22 ± 0.04 and 0.19 ± 0.03 dex, respectively. The same deviation from the RGB values is found in the AGB stars, with a median of 0.25 ± 0.08 dex. We present median observed quantities, instead of mean, because of a significant spread in the $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$ of cluster stars.

4.2 Distance to NGC 6544

Distance is a crucial parameter that is often derived from CMD fitting; although it degenerates with reddening and metallicity. A CMD-independent measurement of the distance can be sought to confirm that the isochrone parameters were accurate enough. Fig. 2 uses a reddening value of $E(J - K_s) = 0.36$ mag (Contreras Ramos et al. 2017) in the VISTA filters or 0.40 mag in the 2MASS system and a distance modulus of $dm = 11.92$ found in this work, which

Table 2. RR Lyrae star used to verify the NGC 6544 distance modulus. Magnitudes in each filter corresponds to the mean quantity. J - and H -band values were derived from only a few epochs, as described in Saito et al. (2012).

Stellar parameters from <i>Gaia</i> DR2	
Source ID	4065784375110674176
RA (J2000)	271.868215
Dec. (J2000)	-25.031377
ℓ (deg)	5.8226
b (deg)	2.2458
G (mag)	14.5874 ± 0.0311
BP (mag)	15.2310 ± 0.1308
RP (mag)	13.6256 ± 0.0770
$\mu_\alpha \cos \delta$ (mas yr $^{-1}$)	-1.821 ± 0.074
μ_δ (mas yr $^{-1}$)	-18.195 ± 0.061
Stellar parameters from OGLE survey	
OGLE ID	OGLE-BLG-RRLYR-13867
V (mag)	13.584
I (mag)	15.048
P (d)	0.5723310 ± 0.0000001
$I_{\text{amplitude}}$ (mag)	0.777
Stellar parameters from VVV survey	
VVV ID	VVV180728.37-250153.1
J (mag)	12.164 ± 0.02
H (mag)	11.932 ± 0.03
K_s (mag)	11.694 ± 0.01
Derived stellar parameters	
$E(V - I)$	0.98 ± 0.07 mag
$E(J - K_s)$	0.31 ± 0.03 mag
A_{K_s}	0.25 ± 0.02 mag
dm (mag)	11.92 ± 0.18
Distance (kpc)	2.43 ± 0.20

is consistent with the distances derived by Cohen et al. (2014) and Contreras Ramos et al. (2017) of $dm = 11.96$ mag.

Although only one RR Lyrae star has been detected to move along with the cluster (see Fig. 1, right-hand panel), it can be used to verify the assumed distance modulus. The RR Lyrae is listed in the online catalogue of Clement et al. (2001), it was also detected by the Optical Gravitational Lensing Experiment (OGLE) survey (OGLE-BLG-RRLYR-13867;² Soszyński et al. 2014, 2019), and by the VVV survey (Contreras Ramos, private communication). Its main properties are presented in Table 2, yielding a distance modulus of $dm = 11.92 \pm 0.18$ mag (2.43 ± 0.20 kpc) using the Muraveva et al. (2015) period–luminosity relation for the K_s band, an intrinsic ($J-K_s$) RR Lyrae colour of 0.17 ± 0.03 (Navarrete et al. 2015; Contreras Ramos et al. 2018b) and the Cardelli, Clayton & Mathis (1989) extinction law, which is comparable to other distance measurements. We also report the near-IR reddening and extinction values given by the RR Lyrae star in Table 2. Consider that all the near-IR measurements presented here are in the 2MASS filter system. Note that $E(J - K_s)$ and $E(V - I)$ derived are consistent within errors with Contreras Ramos et al. (2017) and Cohen et al. (2014), respectively. To calculate $E(V - I)$, we use the same derivation described by Pietrukowicz et al. (2015), considering the actual metallicity of the cluster derived above and the Catelan, Pritzl & Smith (2004) period–luminosity relations. There is also a discrepancy in the $E(J - K_s)$

²Available at the OGLE website: <http://ogledb.astrouw.edu.pl/~ogle/OCVS/o.php?OGLE-BLG-RRLYR-13867>.

value derived by Cohen et al. (2014) which is ~ 0.1 mag shifted towards higher reddening. This shift will be addressed in the next subsection as a VVV survey calibration issue discovered in Hajdu et al. (2020).

Finally, Vasiliev & Baumgardt (2021) recently derived a parallax-driven value of $d = 2.54 \pm 0.07$ kpc ($\overline{\pi} = 0.394 \pm 0.011$ mas) to NGC 6544. This is an independent distance determination, which is also in agreement with all of the previous measurements showed in this section.

4.3 Reddening law towards NGC 6544

Given the multisurvey synergies that we have presented until this point, the final quantity that we want to constrain is the reddening law towards NGC 6544. Using the spectroscopic data that we are analysing, it is possible to derive isochrone-based or synthetic colours for all the stars in the cluster. We will consider, however, only the RGB stars, given the uncertain nature of the AGB stars as discussed in Section 4.1. We cross-match the star positions to a list of surveys that have observed the area with 1 arcsec tolerance, founding that there are observations from the ultraviolet (UV) to the mid-IR for part of the stars in our sample. Photometric information from *SWIFT* (Roming et al. 2005), PanStarrs DR2 (Chambers et al. 2016; Flewelling et al. 2020), *Gaia* DR2 (Gaia Collaboration 2018a; Lindegren et al. 2018), 2MASS (Skrutskie et al. 2006), Spitzer/GLIMPSE (Benjamin et al. 2003; Churchwell et al. 2009), and AllWISE (Wright et al. 2010; Mainzer et al. 2011) was collected. In total, 20 filters (UVOT/UVW2; g , r , i , z , y ; G , BP , RP ; J , H , K_s ; [3.6], [4.5], [5.8], [8.0]; and $W1$, $W2$, $W3$, $W4$) contain fluxes for our stars. Given the crowded nature of a GC, even with the radius tolerance of 1 arcsec, surveys with large pixel scale will often present incorrect or blended magnitudes for the analysed stars. Note also, that we retrieve *Swift* matches for only five stars in the NGC 6544 sample, in contrast to all the other surveys in which we obtain a high counterpart recovery percentage (~ 95 percent or 18 out of 19 possible matches in the worst case). Considering the relevance of the B and V filters in the extinction characterization, to obtain these magnitudes we apply a colour transformation present in Tonry et al. (2012) to the PanStarrs g and r bandpasses.

To derive synthetic magnitudes, we interpolate the T_{eff} and $\log g$ from the APOGEE DR16 into a PARSEC isochrone of 12 Gyr and the cluster metallicity to obtain the absolute magnitudes in different filters. With the absolute theoretical magnitudes, we are able to calculate the colour excess term just subtracting to the observed colour term. This step is necessary to compare with other results, like the one that we derive from the RR Lyrae colours, or the canonical $E(B - V)$ value. Indeed, it is important to mention that there should be statistical and systematic errors not considered here, like the isochrone election or additional differential reddening effects.

Our results for the $E(B - V) = 0.77 \pm 0.19$ mag coincides with the one derived by Cohen et al. (2014) of 0.79 ± 0.01 mag. However, the $E(g - r) = 1.19 \pm 0.16$ value from our method is greater than previously published values. Extinction maps released³ by Green et al. (2014, 2019) account for a value of $E(g - r) = 0.58 \pm 0.03$ mag at the cluster distance.

In parallel, Cohen et al. (2014), Contreras Ramos et al. (2017), and Surot et al. (2019, 2020) give values of the $E(J - K_s)$ of 0.43 ± 0.03 , 0.40 , and 0.47 ± 0.09 mag, respectively, for which we agree with our method. It is important to note that to produce the

³Available at the website: <http://argonaut.skymaps.info/>.

Table 3. Median extinction coefficients per bandpass derived for the NGC 6544 stars.

Filter	x (μm^{-1})	Wavelength (μm)	A_λ (mag)	A_λ/A_V
UVW2	4.67	0.21	4.51 ± 0.90	1.69
<i>B</i>	2.30	0.44	3.42 ± 0.28	1.28
<i>g</i>	2.08	0.48	3.12 ± 0.19	1.17
BP	1.88	0.53	2.55 ± 0.10	0.96
<i>V</i>	1.83	0.55	2.72 ± 0.14	1.02
<i>r</i>	1.62	0.62	2.35 ± 0.10	0.88
<i>G</i>	1.49	0.67	1.93 ± 0.08	0.72
<i>i</i>	1.33	0.75	1.90 ± 0.13	0.71
RP	1.25	0.80	1.44 ± 0.06	0.54
<i>z</i>	1.15	0.87	1.46 ± 0.12	0.55
<i>y</i>	1.04	0.96	1.16 ± 0.05	0.43
<i>J</i>	0.81	1.24	0.40 ± 0.03	0.15
<i>H</i>	0.60	1.66	0.19 ± 0.03	0.07
K_s	0.46	2.16	0.16 ± 0.03	0.06
W1	0.30	3.32	0.03 ± 0.03	0.01
[3.6]	0.28	3.52	0.06 ± 0.02	0.02
[4.5]	0.22	4.45	0.11 ± 0.03	0.04
W2	0.22	4.56	0.15 ± 0.02	0.05
[5.8]	0.18	5.61	0.06 ± 0.03	0.02
[8.0]	0.13	7.70	0.08 ± 0.18	0.03
W3	0.09	10.79	0.31 ± 0.04	0.12
W4	0.05	21.91	0.00 ± 0.09	0.00

best fit for the Fig. 2 CMD, we use the $E(J - K_s)$ value present in Contreras Ramos et al. (2017) (0.36 mag). This value is provided in the VISTA system of magnitudes, and it is affected by the VISTA-2MASS colour transformations, and the correction to the zero-point calibration published by Hajdu et al. (2020). After transformation to the 2MASS framework, the Contreras Ramos et al. (2017) value of $E(J - K_s) = 0.40$ mag, is consistent with our results. The small discrepancy that still remains of the order of ~ 0.04 with our $E(J - K_s)$ derived value can be explained by the strong differential reddening of the cluster, which have amplitudes of up to $\Delta E(J - K_s) = 0.33$ mag depending on the spatial location across the cluster area (Contreras Ramos, private communication). Another possible explanation is possibly the recalibration process proposed by Hajdu et al. (2020), however, this effect should be an order of magnitude less prominent than the last one. With this analysis, we have settled the near-IR reddening discrepancy, preferring the value given by Cohen et al. (2014) and this work of $E(J - K_s) = 0.44 \pm 0.01$.

Finally, we take the difference within the observed star magnitudes and the theoretical absolute magnitude, shifted to the cluster distance ($dm = 11.92$ from Table 2). The difference corresponds to the total extinction A_λ present in the line of sight per filter, which is presented in Table 3. Giving the positive physical nature of the extinction values, we have not considered some slightly negative results as a product of statistical fluctuations in the data, which were probably caused, as we commented earlier, by the differential reddening not taken into account in this step. As expected, a monotonic decrease trend is observed as the wavelength increases, only interrupted by the reddest filters W2, [8.0], and W3 for which we assume that several stars contribute to the matched magnitude in its corresponding catalogues. Note also that W4 shows a very high dispersion compared to the others near- and mid-IR filters that follow the overall trend. Fig. 3 shows the inverse wavelength and normalized extinction values with respect to the *V* band in the same format as in Cardelli et al. (1989). We use the `DUST_EXTINCTION PYTHON` package to reproduce the MW $R(V)$ extinction curve at different $R_V = A_V/E(B - V)$ values.




Figure 3. Wavelength (as $x \equiv \lambda^{-1} [\text{nm}^{-1}]$ and λ [nm]) against the normalized extinction values with respect to the *V* band $A(x)/A(V)$. The two extinction curves shown represents the total-to-selective ratio of $R_V = 3.5$ (continuous), and $R_V = 5.8$ (dashed) between 0.3 and 5.0 nm. In total, 20 filters from the UV to the mid-IR with its derives extinction values are shown. The *V* band is highlighted with a shaded area because, by definition, all the extinction curves pass through this point.

As it is clear from the figure, total-to-selective extinction ratio of $R_V = 3.5 \pm 0.19$ deduced from the data is enough to reproduce the overall optical trend ($0.4 \lesssim \lambda$ (nm) $\lesssim 1.0$). However, this value present inconsistencies with the UV and mid-IR data. We attribute this discrepancy to the source confusion caused by the combined effect of the crowding level typical of GCs and the lower spatial resolution (i.e. larger pixel size) of these photometric surveys. The figure also shows an extreme case with $R_V = 5.9$ that fits solely the UV observation, not reproducing any other available filters to illustrate the erroneous status of the measurement. Given the similarities in the optical and IR parts of the spectral energy distribution within the $R_V = 3.1$ and $R_V = 3.5$ curves of less than ~ 5 per cent, we are not able to differentiate the preferred model, both being consistent with the current data.

As a final caveat on this procedure, all of the extinction derivation relies on the fact that isochrones are model independent, which is not completely accurate in this case. The PARSEC set of isochrones uses the Cardelli et al. (1989) recipe to apply the interstellar extinction to their data, adopting a canonical value of $R_V = 3.1$. Nevertheless, given the short distance of the cluster this effect should not represent a dramatic change in our results.

5 FINGERPRINT OF AN ELUSIVE CLUSTER

As a result of the earliest star formation in galaxies, GCs are one of the main tools that we have to unveil the early stages and evolution of the MW. Long considered as simple stellar populations, GCs have revealed a more complex composition of their stars, which appears to belong to different generations, each one with star-to-star abundance variations, visible at photometric and spectroscopic scale (Gratton, Carretta & Bragaglia 2012; Bastian & Lardo 2018, and references therein).

In this sense, these multiple populations (MPs) within a GC refer basically to a first (or primordial generation, FG) and at last a second generation (SG) of stars. The FG of stars pollutes the SG with processed material, and in this way different subpopulations arise in the cluster. The composition of the SG stars varies almost at an individual cluster level, manifested through several sequences in




Figure 4. T_{eff} , $\log g$, and $[X/\text{Fe}]$ for surface parameters and abundances in Table A1. It can be noticed from the Figure, for a given metallicity there is a large spread for certain elements ($[\text{N}/\text{Fe}]$, $[\text{C}/\text{Fe}]$, $[\text{Na}/\text{Fe}]$, and $[\text{Al}/\text{Fe}]$). The AGB group (green stars) and one uRGB star (yellow triangle) is completely offset from the global $[\text{Fe}/\text{H}]$ trend. This variety of $[\text{Fe}/\text{H}]$ values was the main reason that led us to consider and publish the median over the mean values for the elemental abundances.

CMDs, as well as light-element variations within the clusters (C, N, O, Na, Mg, Al, Si, among other elements; see Bastian & Lardo 2018, for a recent review).

We want to extend this background to NGC 6544 if possible, trying to search for MPs within the member stars through light-element abundances. In order to interpret NGC 6544 chemical properties, we compare its abundance patterns with other GCs observed by APOGEE, and with nearby bulge field stars. All the abundance comparisons were performed with the APOGEE DR16 values. Cross-matched stars from other GCs were located in the DR16 using the APOGEEID available for all the recent APOGEE publications.

5.1 Abundance analysis and element variations

The present analysis adds a substantial contribution to the chemical characterization of NGC 6544. Our sample of cluster members ascends to 23 and allows us to observe intracluster abundance variations comparatively. In previous publications, Nataf et al. (2019) and Mészáros et al. (2020), only found 2 and 7 stars in the APOGEE DR14 and 16, respectively. This difference is originated because of the growing number of observations produced by the APOGEE survey over time, and the fact that we do not use the GC flag that is present in all the APOGEE DRs (TARGET_FLAGS={APOGEE_SCI_CLUSTER, APOGEE2_SCI_CLUSTER}). Recently, Horta et al. (2020) analysed most of the GCs present in the APOGEE catalogue including NGC 6544, in a broad approach to compare the accreted and *in situ* classification given by Massari et al. (2019). They calculated mean metallicities, radial velocities, and $[\text{Si}/\text{Fe}]$ abundances to derive general GCs properties of the MW, but without detailing any other cluster features. The paper indicates that 21 NGC 6544 stars were recovered from the cluster reaching identical results than the ones presented here. We estimate that the different number of stars analysed originated on the automatic elimination of stars, which

removes two of the most metal-poor AGB stars (see fig. 2 in the appendix of Horta et al. 2020)

Table A1 contains the abundance values, derived by the ASPCAP pipeline, for a subset of all the elements reported in the DR16. Specifically, we select the following elements: $[\text{C}/\text{Fe}]$, $[\text{N}/\text{Fe}]$, $[\text{O}/\text{Fe}]$, $[\text{Na}/\text{Fe}]$, $[\text{Mg}/\text{Fe}]$, $[\text{Al}/\text{Fe}]$, $[\text{Si}/\text{Fe}]$, $[\text{K}/\text{Fe}]$, and $[\text{Ca}/\text{Fe}]$, together with $[\alpha/\text{Fe}]$, and $[\text{Fe}/\text{H}]$, in order to analyse known (anti) correlations for other GCs and to insert NGC 6544 within the scenario proposed by Jönsson et al. (2018), Zasowski et al. (2019), and Masseron et al. (2019) for the bulge and GC samples, respectively.

Fig. 4 shows the abundance of each element as a function of iron, for all the cluster members. Nevertheless, we will base the discussion on RGB stars only, because of the metallicity offset of AGB stars mentioned in Section 4. Fig. 4 shows a slight trend in sections showing the evolution of T_{eff} , $\log g$, and $[\alpha/\text{Fe}]$ with metallicity for IRGB stars, but the trend vanishes as stars cross the RGB bump. A clear dichotomy for stars in the IRGB and uRGB is present in the $[\text{C}/\text{Fe}]$ abundances as expected from stellar evolution with C-enhanced stars below the RGB bump and C-poor stars above the same point (Iben 1968; Lardo et al. 2012; Cassisi & Salaris 2013).

Even if there are only a few Na lines in the APOGEE H -band spectra, only half of the cluster sample has measured abundances from the ASPCAP pipeline. They show a wide spread of Na abundance that resemble the trend presented in Masseron et al. (2019) and Carretta et al. (2009a). A detailed discussion of known (anti) correlations will be given in the next subsections.

5.1.1 The Mg–Al anticorrelation and the non-existent Si-leakage

The Mg–Al chain is a complex set of proton capture nuclear reactions that need high temperatures (50 MK) to trigger (Prantzos, Charbonnel & Iliadis 2007; Carretta et al. 2009a; Cassisi & Salaris 2013). For this reason, the anticorrelation between Mg and Al that is naturally produced through this chain, cannot happen inside low-mass stars. Massive stars are thought to be responsible for producing




Figure 5. Known Mg–Al, C–N, and O–Na anticorrelations, along with other discussed abundance planes in Section 5: Si–Al, Mg–K, and Ca–Al. This Figure uses the same background and symbols described in Fig. 4, and for the background the matched APOGEE stars from the Schiavon et al. (2017), Masseron et al. (2019), and Mészáros et al. (2020) analysed cluster stars. We also apply a Gaussian smoothing to this sample, shown as the coloured background, to emphasize the high-density areas of the distributions.

the observed anticorrelations, with some external mechanism needed to transport processed material to the surface of low-mass RGB stars (Renzini et al. 2015; Bastian & Lardo 2018).

Upper left panel in Fig. 5 shows the comparison of our [Mg/Fe] and [Al/Fe] abundances with stars of other APOGEE observed GCs (Schiavon et al. 2017; Masseron et al. 2019; Mészáros et al. 2020) from the Galactic bulge to the halo. For both catalogues, its stars were matched with the DR16, resulting in more than 680 stars in total. Overall the red giant stars in NGC 6544 follow the same trend that other GCs at the same metallicity as M13, M2, and M3 with [Fe/H] equal to -1.53 , -1.65 , and -1.50 dex, respectively. For a cluster as low mass as NGC 6544, an approximate mono abundance of [Mg/Fe] is expected. Despite that, we found that there is some spread around the median [Mg/Fe] value of 0.24 ± 0.07 dex. Nevertheless, this spread is well within two times the observational error that also depends on the evolutionary stage marked with different symbols on the Figure. A clear sign of the Mg–Al anticorrelation is more evident in the [Al/Fe] abundances, in which we observe a high amplitude of almost $\Delta[\text{Al/Fe}] \sim 0.8$ dex. The same result was reported by Carretta et al. (2009a), also observing a [Mg/Fe]-depletion for stars with [Al/Fe] ≥ 0.8 dex in massive and/or metal-poor GCs. We report a maximum [Al/Fe] abundance of 0.52 dex, which agrees with the results of Carretta et al. (2009a). There appears to be a gap around [Al/Fe] = 0 dex, and even AGB stars appear to split in two groups. Often, a large Al-spread and decrease in the [Mg/Fe] for stars with [Al/Fe] ≤ 0.8 dex, could suggest a possible Si-leakage from the Mg–Al chain, that should be seen as an enhancement in the [Si/Fe] abundance. We do not detect any sign of [Mg/Fe] depletion. Indeed, although the Mg–Al knee is observable in the APOGEE data, none of the cluster members shows an evident Mg-poor, Al-rich behaviour. In

fact, there is only a few clusters observed with APOGEE in which this effect have been documented (see Masseron et al. 2019; Mészáros et al. 2020, for more details). Another observable consequence of the Mg-depletion would be the increase of the [Si/Fe] spread among individual stars in a GC, also known as Si-leakage, which is also not observed. This effect have been observed and described in depth in GCs more metal-poor than NGC 6544 (Mészáros et al. 2020), like NGC 6341/M92 ([Fe/H] = -2.31 dex) and NGC 7078/M15 ([Fe/H] = -2.37 dex). Within the cluster, the [Si/Fe] abundances remain constant, with a median value of 0.22 ± 0.05 dex, for cluster stars showing large differences in [Al/Fe]. Two groups of stars with different [Al/Fe] abundances can be identified, with separation around [Al/Fe] ~ 0 dex. We will assign the labels of FG and SG to the groups depending on the [Al/Fe] abundance lower, and greater than 0.0 dex, respectively. These two groups, FG and SG, median [Al/Fe] values have a total difference of 0.42 dex and a standard deviation of 0.23 dex, without counting the AGB stars. Both values agrees with the results of Mészáros et al. (2020) for other APOGEE observed clusters around the metallicity that we derive for NGC 6544. Similarly, we can separate FG and SG stars in the C–N plane.

5.1.2 The C–N anticorrelation

The C–N anticorrelation is a natural consequence of the CNO cycle, which depletes C and O and enhances N in the H-burning (Cassisi & Salaris 2013). These products may be carried to the upper layers of the star either through some extra mixing phenomenon along the RGB, or because of pollution of the stellar atmosphere by the winds of high-mass AGB stars. The central upper panel in Fig. 5 shows the

anticorrelation for NGC 6544 in comparison with the GCs analysed by Schiavon et al. (2017), Masseron et al. (2019), and Mészáros et al. (2020). Notice that we must be careful with the $[C/Fe]$ values derived by ASPCAP. The value represents the best-fitting value for the whole spectrum, and according to Mészáros et al. (2015), only upper values can be measured below $[Fe/H] = -1.7$ dex. Fortunately, only two AGB stars fall under this metal-poor regime, and only one does not return any value in the DR16. A clear separation between pre and post-RGB bump is seen through the $[C/Fe]$ abundances.

5.1.3 The Na–O anticorrelation

The Na–O anticorrelation is produced by oxygen and sodium, produced by the combined action of the CNO and Ne–Na cycles (Gratton, Sneden & Carretta 2004; Cassisi & Salaris 2013). While the CNO cycle decreases the O abundance (maintaining the $C + N + O$ overall count), the Ne–Na cycle produces Na, creating the anticorrelation. The latter cycle is efficient in intermediate-mass stars during the thermal pulses of the AGB phase, but can also be active at the bottom of the convective envelope in RGB stars, since it requires lower temperatures compared with the Mg–Al cycle (around 40 MK). Observationally, Na is difficult to measure in APOGEE spectra, because its two lines are very weak in the *H* band. That is the reason that all the $[Na/Fe]$ quantities reported (and its associated errors) must be considered with caution, as the most uncertain values are presented in this work. On the same behaviour, O can be difficult to detect for metal-poor stars (see the previous subsection, and Masseron et al. 2019). As a result, only a few cluster stars have both elements measured, and are shown in the lower left panel of Fig. 5. As a broad overview, NGC 6544 stars follow the expected anticorrelation, like other APOGEE clusters, with a large $[Na/Fe]$ spread of $\Delta[Na/Fe] \sim 1.1$ dex. This is in agreement with the results from other clusters (Gratton, Sneden & Carretta 2004), which also shows a larger spread in $[Na/Fe]$ with respect to $[Al/Fe]$.

5.1.4 The odd-Z element K

Finally, we analyse an element that present strange patterns in the NGC 6544 APOGEE DR16 data. Potassium, as an odd-Z element, is produced from argon by a proton capture reaction. It needs a much higher temperature than other cycles reviewed in this section, with 120–180 MK to trigger (Ventura et al. 2012; Iliadis et al. 2016). There are available only a few lines in the APOGEE spectra, but as shown in Masseron et al. (2019), $[K/Fe]$ was successfully recovered and reported a negligible K production over all the GCs analysed.

Contrary to the results for other GCs, we observe a broad $[K/Fe]$ spread, with up to ~ 1.25 dex in range. This abundance variation matches the values reported by Carretta et al. (2013) and Mucciarelli, Merle & Bellazzini (2017) with high $[K/Fe]$ variance. The central lower panel in Fig. 5 shows the Mg–K plane. As seen in the background for other GCs observed by APOGEE (Schiavon et al. 2017; Masseron et al. 2019; Mészáros et al. 2020), a constant value around solar composition was reported, i.e. no K-production was found. K-enhanced stars are indeed expected in GCs if the polluters achieve high temperatures to create potassium. Like K-rich stars, subsequent generations will also show Mg-depleted and Si-enhanced stars due to the lower temperature barrier of the Mg–Al–Si cycle, as shown in Mészáros et al. (2020) for some APOGEE GC stars.

With the present observations, we cannot determine the origin of the peculiar K-enhancement, which is not compatible with a constant

value given the significant star-to-star variations. Optical follow-up is necessary to constraint the discrepancy extent of our measurements.

5.1.5 The α -element Ca

Most of the calcium that we can measure in GCs stars is produced by supernovae, and therefore not affected by the H-burning processes neither of the stars nor of the polluters. Mészáros et al. (2020) reported constant $[Ca/Fe]$ values for APOGEE GCs, though warning about the weakness of the Ca lines in the near-IR at low metallicities. Their results are shown in the lower right panel of Fig. 5 as background, along with the NGC 6544 stars.

In general, cluster stars follow the overall trend of constant calcium abundance, centred at $[Ca/Fe] = 0.21 \pm 0.09$ dex. There is a deviation of this behaviour, but only for stars with $[Al/Fe] \lesssim 0.15$ dex. These four stars, (three IRGB and one AGB), however, also present large $[Ca/Fe]$ errors according to ASPCAP. If we do not take into account these four outliers, the calcium abundance can be described as a constant, with star-to-star deviations within the errors.

5.1.6 Neutron-capture elements: Ce and Nd

Neutron capture elements, like cerium and neodymium, can be formed by a wide variety of channels in different contexts within the Galaxy (Sneden, Cowan & Gallino 2008). However, in a GC, the possible scenarios that can occur can be limited to a few like a burning shell in a massive stars or an evolved AGB star in the thermally pulsating phase (Bisterzo et al. 2011, 2014). Within the APOGEE survey, multiple measurements of Ce and Nd have been reported since its characterization and inclusion in subsequent DRs (Hasselquist et al. 2016; Cunha et al. 2017). Given the limited range of temperatures in which Ce can be reliably fitted and measured ($T_{\text{eff}} < 4400$ K according to Mészáros et al. 2020), all of the ASPCAP $[Ce/Fe]$ abundances reported here and in the DR16 not in this temperature range, can be considered only as upper limits. ASPCAP (García Pérez et al. 2016), as an automatic abundance pipeline, cannot handle correctly the difference between an upper limit and a non-detection in which the line is absent from the spectra. This is the case of all the Ce abundance measured and reported in this section. Moreover, to better constrain the Ce abundances, the BACCHUS (Masseron, Merle & Hawkins 2016) code was used to properly account for the non-detections, even including more Ce lines to the fitting procedure (Masseron et al. 2019; Mészáros et al. 2020).

Given that information, we can explain the behaviour of the $[Ce/Fe]$ within the cluster. Fig. 6 shows the $[Ce/Fe]$ abundance as a function of the metallicity and $[Al/Fe]$ for NGC 6544 and other APOGEE observed clusters. The wide range of $[Ce/Fe]$ of 1.9 dex is solely explained by the fact that ASPCAP is not able to correctly fit the Ce line. As shown in Mészáros et al. (2020), ω Cen presents the most dramatic Ce enrichment of ~ 1 dex across ~ 1 dex of $[Fe/H]$ enrichment, followed by NGC 1851/47Tuc. This marked difference exemplifies how our measurements do not represent the actual state of NGC 6544 $[Ce/Fe]$. Further analysis with cooler stars within the cluster is necessary before reliably establish the Ce status of NGC 6544.

Finally, no $[Nd/Fe]$ have been reported for the selected NGC 6544 stars due to the combination of T_{eff} and $[Fe/H]$ that the spectra of the GC presents. According to Hasselquist et al. (2016), only a fraction close to the ~ 20 per cent of the total APOGEE spectra will show Nd variations that are strong enough to be detected at a high SNR.




Figure 6. Upper limits for the [Ce/Fe] detections versus [Fe/H] (left-hand panel) and [Al/Fe] (right-hand panel) for the APOGEE observed clusters present in the DR16. NGC 6544 stars are colour coded with the same markers as in Figs 4 and 5. A spread of almost ~ 2 dex in [Ce/Fe] can be explained by the inability of the ASPCAP pipeline to treat upper limits in the analysed spectra (see text).

5.2 Al–N correlation and its relation with NGC 6544 mass

Finally, and with the aim to check the consistency of the previous APOGEE results, we perform the same analysis of Nataf et al. (2019) that relates the amplitude of the Al and N spread with the cluster mass. This relation was built in the past DR14 version of the APOGEE data base, however, considering that the calibrations within DRs consists only in shifts on the abundance plane, we can assume that is still valid within errors. To built the relation, we need first to define the $\Delta[X/Fe]$ quantity, which is defined as

$$\Delta[X/Fe]_i = [X/Fe]_{\text{Gen II},i} - ([X/Fe]_{\text{Gen I}}),$$

which relates the difference within the [X/Fe] value of each i -star with SG abundance patterns with the mean value of [X/Fe] but measured for FG stars. As derived in equation (6) of Nataf et al. (2019), we apply the relation within [Fe/H] and $\Delta[N/Fe]$ of the individual SG stars, and the cluster mass $\log M/M_\odot$ to derive the total $\Delta[Al/Fe]$ of the same stars. Given that the only unknown quantity is the total cluster mass, we can apply this relation to constrain the possible NGC 6544 mass values. The procedure is performed by comparing the reason of both $\Delta[Al/Fe]$ quantities, derived from aluminum abundances, and from fixing the cluster mass. After calculating the median of all the stars, the most accurate mass will produce a value of $\Delta Al = \Delta[Al/Fe]_{\text{spectra}}/\Delta[Al/Fe]_{\text{fixed mass}}$ closer to 1. The results shown in Fig. 7 correlate well with the statements of Nataf et al. (2019) in both the [Al/Fe]–[N/Fe] correlation (left-hand panel) and the $\Delta[Al/Fe]$ – $\Delta[N/Fe]$ plane. Note that there are two different NGC 6544 total masses in the literature that differs by a factor of ~ 2 , one listed in the Harris (1996) catalogue and used by Nataf et al. (2019) of $\log M/M_\odot = 5.05$, and the other one derived by Baumgardt & Hilker (2018) of $\log M/M_\odot = 4.80$. Also is important to highlight that at the time of fixing the cluster mass from the relation to calculate the trend line, we also fixed the metallicity of the cluster to [Fe/H] = -1.44 dex, as is stated in Section 4.1. We use both masses and conclude that the Baumgardt & Hilker (2018) fully agrees within the errors the Al–N–mass relation, which is showed as a continuous line in the right-hand panel of Fig. 7. To quantify this statement, both fixed masses produces a value of ΔAl equal to 0.82 and 1.00, for the Nataf et al. (2019) and Baumgardt & Hilker (2018) masses, respectively.




Figure 7. Left-hand panel: [Al/Fe]–[N/Fe] correlation for the cluster stars. A clear separation between FG and SG at [N/Fe] < 0.4 dex and [Al/Fe] < 0 dex for both abundances, respectively. Same symbols were used than in other figures for the different evolutionary stages. The red cross indicate the mean value of the FG stars, used later in the $\Delta[X/Fe]$ values. Right-hand panel: $\Delta[N/Fe]$ – $\Delta[Al/Fe]$ diagram as is shown in Nataf et al. (2019), with the continuous solid line representing the best fit using the lower mass value indicated in the legend, with its corresponding error as shaded area. Additionally, we include the fit as a dashed line (and hatched area for the error) with the mass value used by Nataf et al. (2019) to derive this relation. Note that only SG stars are showed in this panel for which $\Delta[X/Fe]$ can be measured.

6 GALACTIC CONTEXT

NGC 6544 is projected on the sky towards the bulge area in a very crowded region at $(\ell, b) \sim (5.8^\circ, -2.2^\circ)$. The first comprehensive observations describe the cluster as a relatively metal-poor one, unlike most other inner Galactic GCs (Cohen et al. 2014, 2017, 2018, and clusters there in). Indeed, Bica et al. (2016) and Contreras Ramos et al. (2017) classify NGC 6544 as a halo intruder in the bulge region based on metallicity estimates and simple orbital constraints. More recently, Pérez-Villegas et al. (2019) derived a more robust probability-based membership based on state-of-the-art Galactic potentials for GCs in the inner Galaxy. With a very high percentage (within ~ 80 – 97 percent depending on the Galactic bar parameters), NGC 6544 was classified as thick disc GC. Based on its [Fe/H], we can state that the cluster is consistent with the metal-poor tail of the canonical thick disc. In order to contextualize the present results for the cluster, we built a catalogue of GCs observed by APOGEE (Schiavon et al. 2017; Masseron et al. 2019; Mészáros et al. 2020). The median metallicity and total α -enhancement, as measured by ASPCAP in the DR16, were calculated. Fig. 8 shows the chemical neighbourhood of NGC 6544, together with other clusters in the halo (NGC 7078/M 15, NGC 6341/M 92, NGC 5024/M 53, NGC 5466, NGC 7089/M 2, NGC 5272/M 3, NGC 6205/M 13, and NGC 5904/M 5), thick disc (NGC 6171/M 107, Pal 6, and NGC 6838/M 71), and bulge (NGC 6522, Ter 5, NGC 6528, and NGC 6553). We have also added a sample of randomly selected field stars from APOGEE DR16, representing 10 percent of the total catalogue, as comparison.

As a detailed comparison between the abundances of cluster stars and those of its spatial neighbourhood, we show in Fig. 9 the abundances of several elements in the cluster and in a sample of 3731 field stars located within 2° from the cluster centre, which are mostly bulge stars. The normalized abundance distribution is shown as a density estimation for both bulge and cluster population,




Figure 8. Tinsley diagram of the $[\alpha/\text{Fe}]$ enrichment across metallicity for the observed GCs by APOGEE (Schiavon et al. 2017; Masseron et al. 2019; Mészáros et al. 2020). Despite being in a similar position of some halo GCs (NGC 5272/M 3, NGC 7089/M 2, and NGC 6205/M 13 with $[\text{Fe}/\text{H}] = -1.49, -1.56,$ and -1.60 dex, respectively), NGC 6544 (shown as a red star) was classified as thick disc GC by orbital reasons. As a background reference (small points), we place a randomly selected 10 per cent of the whole APOGEE catalogue. The bulge, disc, and halo are visible, even showing the sequence of anomalous stars at $[\alpha/\text{Fe}] \sim 0.2$ dex, described by Zasowski et al. (2019).




Figure 9. $[X/\text{Fe}]$ abundance density estimation comparison between bulge (left grey symbols) and cluster (right orange symbols) stars. Each violin representation was normalized, and indicates with horizontal lines the median and limits of the distribution.

whereby the median, min, and max of the measurements of each element are marked. As it is clear from the figure, cluster stars show a wider range of abundances than field stars, for all the elements except Mg and Si. The effect is especially clear in $[\text{C}/\text{Fe}]$, $[\text{N}/\text{Fe}]$, $[\text{Na}/\text{Fe}]$, $[\text{Al}/\text{Fe}]$ (with hints of the bimodality reported in Section 5), and $[\text{K}/\text{Fe}]$. Compared with previous results for field stars that may belong to GCs (Fernández-Trincado et al. 2016, 2017), we do not find any unusual abundance ratios in the nearby bulge stars that can be directly linked to NGC 6544.

Finally, given the remarkable elongation displayed by NGC 6544 caused by the Galaxy (Contreras Ramos et al. 2017), we also search for extra tidal stars within the APOGEE DR16. We use the derived metallicity and space velocity for the cluster within this work as a first guess. Unfortunately, we did not find additional members within $\sim 10^\circ$ from the cluster centre. The use of more extended and deep data is mandatory to constrain the expected tidal tails of NGC 6544. Upcoming releases of the APOGEE data, and also forthcoming stellar wide-field multi-object spectroscopic surveys like 4MOST (de Jong et al. 2019) and MOONS (Cirasuolo et al. 2011), will help to identify extra tidal members, if they exist, as it was strongly suggested by previous studies (Cohen et al. 2014).

7 CONCLUDING REMARKS

We have performed a near-IR high-resolution spectral analysis for the reddened GC NGC 6544. With 23 confirmed members in the APOGEE DR16 data, this is the most complete study up to date, to characterize the abundance patterns of this cluster. Across the RGB (below and above the bump) and AGB, the abundances derived by the ASPCAP pipeline were presented and discussed in the context of other Galactic GCs. Known anticorrelations (Mg–Al, C–N, Na–O) were used to isolate two generations of stars, an FG with 9 members and an SG with 14 members. For a random targeting, these numbers are in very good agreement with predictions about the number of FG stars that still remain in GCs (close to $\sim 1/3$ of the total population).

A large $[\text{Al}/\text{Fe}]$ spread of 0.78 dex, negligible $[\text{Mg}/\text{Fe}]$ dispersion, a significant variation of CNO, across all the sampled evolutionary phases was a pivotal factor to divide different generations of stars. The metallicity and α -element abundances for RGB stars, from which we have a bonafide abundance determination, are -1.44 ± 0.05 and 0.20 ± 0.04 dex, respectively. Using a correlation within the cluster aluminium, nitrogen, and metallicity, we were able to constraint the NGC 6544 mass, favouring the less massive proposed scenario with $\log M/M_\odot = 4.80$.

Multisurvey synergies were used to derive other cluster properties, such as the distance, reddening law towards the cluster and confirming its Galactic classification. Finally, no extra tidal or chemically peculiar stars were found around the cluster, using data from the APOGEE DR16. Still, this picture could change with future and deeper spectroscopic surveys, which might explain the spatial elongation of NGC 6544.

ACKNOWLEDGEMENTS

Support for some of the authors is provided by the BASAL Center for Astrophysics and Associated Technologies (CATA) through grant PFB-06, and the Ministry for the Economy, Development, and Tourism, Programa Iniciativa Científica Milenio through grant IC120009, awarded to the Millennium Institute of Astrophysics (MAS).

This work is part of the PhD thesis of FG, funded by grant CONICYT-PCHA Doctorado Nacional 2017-21171485. FG acknowledges support by CONICYT-PCHA Doctorado Nacional 2017-21171485. ARA acknowledges support from FONDECYT through grant 3180203. We acknowledge support from FONDECYT Regular grants No. 1150345 (MZ and FG), and No. 1170121 (DM). DAGH acknowledges support from the State Research Agency (AEI) of the Spanish Ministry of Science, Innovation and Universities (MCIU) and the European Regional Development Fund (FEDER) under grant AYA2017-88254-P. SV gratefully acknowledges the support provided by FONDECYT reg. n. 1170518. SV gratefully acknowledges the support from the Chilean BASAL Centro de Excelencia en Astrofísica y Tecnologías Afines (CATA) grant AFB-170002. JGF-T is supported by FONDECYT No. 3180210 and Becas Iberoamérica Investigador 2019, Banco Santander Chile. DG gratefully acknowledges support from the Chilean Centro de Excelencia en Astrofísica y Tecnologías Afines (CATA) BASAL grant AFB-170002. DG also acknowledges financial support from the Dirección de Investigación y Desarrollo de la Universidad de La Serena through the Programa de Incentivo a la Investigación de Académicos (PIA-DIDULS).

We gratefully acknowledge the use of data from the VVV/VVVx ESO Public Survey program ID 179.B-2002/198.B-2004 taken with the VISTA telescope, and data products from the Cambridge Astronomical Survey Unit (CASU). The VVV Survey data are made public at the ESO Archive.

Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS website is www.sdss.org.

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) University of Tokyo, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatory of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional/MCTI,

The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, and NEOWISE, which is a project of the Jet Propulsion Laboratory/California Institute of Technology. WISE and NEOWISE are funded by the National Aeronautics and Space Administration.

This paper uses data from the MW-BULGE-PSFPHOT compilation (Surot et al. 2019, 2020). This research has made use of the services of the ESO Science Archive Facility.

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

This research made use of: TOPCAT (Taylor 2005); PANDAS (Pandas Development Team 2020); IPYTHON (Pérez & Granger 2007); NUMPY (van der Walt, Colbert & Varoquaux 2011); MATPLOTLIB (Hunter 2007); ASTROPY, a community-developed core PYTHON package for Astronomy (Astropy Collaboration 2013, 2018); DUST_EXTINCTION, an Astropy-affiliate package; and SCIKIT-LEARN (Pedregosa et al. 2011). This research has made use of NASA’s Astrophysics Data System.

DATA AVAILABILITY

All APOGEE DR16 data used in this study are publicly available and can be found at: <https://www.sdss.org/dr16/>. However, we prepared Table A1 as a supplemental material in the journal webpage.

REFERENCES

- Ahumada R. et al., 2020, *ApJS*, 249, 3
 Alcaino G., 1983, *A&AS*, 52, 105
 Astropy Collaboration, 2013, *A&A*, 558, A33
 Astropy Collaboration, 2018, *AJ*, 156, 123
 Bailin J., 2019, *ApJS*, 245, 5
 Bastian N., Lardo C., 2018, *ARA&A*, 56, 83
 Baumgardt H., Hilker M., 2018, *MNRAS*, 478, 1520
 Baumgardt H., Hilker M., Sollima A., Bellini A., 2019, *MNRAS*, 482, 5138
 Benjamin R. A. et al., 2003, *PASP*, 115, 953
 Bica E., Ortolani S., Barbay B., 2016, *Publ. Astron. Soc. Aust.*, 33, e028
 Bisterzo S., Gallino R., Straniero O., Cristallo S., Käppeler F., 2011, *MNRAS*, 418, 284
 Bisterzo S., Travaglio C., Gallino R., Wiescher M., Käppeler F., 2014, *ApJ*, 787, 10
 Blanton M. R. et al., 2017, *AJ*, 154, 28

- Bowen I. S., Vaughan A. H. J., 1973, *Appl. Opt.*, 12, 1430
- Bressan A., Marigo P., Girardi L., Salasnich B., Dal Cero C., Rubele S., Nanni A., 2012, *MNRAS*, 427, 127
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
- Carretta E., Bragaglia A., Gratton R., D'Orazi V., Lucatello S., 2009b, *A&A*, 508, 695
- Carretta E., Gratton R. G., Bragaglia A., D'Orazi V., Lucatello S., Sollima A., Sneden C., 2013, *ApJ*, 769, 40
- Carretta E. et al., 2009a, *A&A*, 505, 117
- Cassisi S., Salaris M., 2013, Wiley-VCH, Germany. Old Stellar Populations: How to Study the Fossil Record of Galaxy Formation
- Catelan M., Pritzl B. J., Smith H. A., 2004, *ApJS*, 154, 633
- Chambers K. C. et al., 2016, preprint ([arXiv:1612.05560](https://arxiv.org/abs/1612.05560))
- Chen Y., Bressan A., Girardi L., Marigo P., Kong X., Lanza A., 2015, *MNRAS*, 452, 1068
- Chen Y., Girardi L., Bressan A., Marigo P., Barbieri M., Kong X., 2014, *MNRAS*, 444, 2525
- Churchwell E. et al., 2009, *PASP*, 121, 213
- Cirasuolo M., Afonso J., Bender R., Bonifacio P., Evans C., Kaper L., Oliva E., Vanzi L., 2011, *Messenger*, 145, 11
- Clement C. M. et al., 2001, *AJ*, 122, 2587
- Cohen R. E., Mauro F., Alonso-García J., Hempel M., Sarajedini A., Ordoñez A. J., Geisler D., Kalirai J. S., 2018, *AJ*, 156, 41
- Cohen R. E., Mauro F., Geisler D., Moni Bidin C., Dotter A., Bonatto C., 2014, *AJ*, 148, 18
- Cohen R. E., Moni Bidin C., Mauro F., Bonatto C., Geisler D., 2017, *MNRAS*, 464, 1874
- Contreras Ramos R. et al., 2017, *A&A*, 608, A140
- Contreras Ramos R. et al., 2018a, *ApJ*, 863, 78
- Contreras Ramos R. et al., 2018b, *ApJ*, 863, 79
- Cunha K. et al., 2017, *ApJ*, 844, 145
- de Jong R. S. et al., 2019, *Messenger*, 175, 3
- Eisenstein D. J. et al., 2011, *AJ*, 142, 72
- Fernández-Trincado J. G. et al., 2016, *ApJ*, 833, 132
- Fernández-Trincado J. G. et al., 2017, *ApJ*, 846, L2
- Flewelling H. A. et al., 2020, *ApJS*, 251, 7
- Gaia Collaboration, 2018a, *A&A*, 616, A1
- Gaia Collaboration, 2018b, *A&A*, 616, A12
- García-Hernández D. A., Mészáros S., Monelli M., Cassisi S., Stetson P. B., Zamora O., Shetrone M., Lucatello S., 2015, *ApJ*, 815, L4
- García Pérez A. E. et al., 2016, *AJ*, 151, 144
- Gratton R., Sneden C., Carretta E., 2004, *ARA&A*, 42, 385
- Gratton R. G., Carretta E., Bragaglia A., 2012, *A&AR*, 20, 50
- Green G. M., Schlafly E., Zucker C., Speagle J. S., Finkbeiner D., 2019, *ApJ*, 887, 93
- Green G. M. et al., 2014, *ApJ*, 783, 114
- Gunn J. E. et al., 2006, *AJ*, 131, 2332
- Hajdu G., Dékány I., Catelan M., Grebel E. K., 2020, *Exp. Astron.*, 49, 217
- Harris W. E., 1996, *AJ*, 112, 1487
- Hasselquist S. et al., 2016, *ApJ*, 833, 81
- Horta D. et al., 2020, *MNRAS*, 493, 3363
- Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
- Iben I., 1968, *Nature*, 220, 143
- Iliadis C., Karakas A. I., Prantzos N., Lattanzio J. C., Doherty C. L., 2016, *ApJ*, 818, 98
- Ivans I. I., Kraft R. P., Sneden C., Smith G. H., Rich R. M., Shetrone M., 2001, *AJ*, 122, 1438
- Jönsson H. et al., 2018, *AJ*, 156, 126
- Jönsson H. et al., 2020, *AJ*, 160, 120
- Kundu R., Fernández-Trincado J. G., Minniti D., Singh H. P., Moreno E., Reylé C., Robin A. C., Soto M., 2019b, *MNRAS*, 489, 4565
- Kundu R., Minniti D., Singh H. P., 2019a, *MNRAS*, 483, 1737
- Lapenna E., Mucciarelli A., Ferraro F. R., Origlia L., Lanzoni B., Massari D., Dalessandro E., 2015, *ApJ*, 813, 97
- Lapenna E., Mucciarelli A., Lanzoni B., Ferraro F. R., Dalessandro E., Origlia L., Massari D., 2014, *ApJ*, 797, 124
- Lardo C., Pancino E., Mucciarelli A., Milone A. P., 2012, *A&A*, 548, A107
- Lindgren L. et al., 2018, *A&A*, 616, A2
- Mainzer A. et al., 2011, *ApJ*, 731, 53
- Majewski S. R. et al., 2017, *AJ*, 154, 94
- Marigo P. et al., 2017, *ApJ*, 835, 77
- Massari D., Koppelman H. H., Helmi A., 2019, *A&A*, 630, L4
- Masseron T., Merle T., Hawkins K., 2016, Astrophysics Source Code Library, record ascl:1605.004
- Masseron T. et al., 2019, *A&A*, 622, A191
- McLaughlin D. E., van der Marel R. P., 2005, *ApJS*, 161, 304
- Mészáros S. et al., 2015, *AJ*, 149, 153
- Mészáros S. et al., 2020, *MNRAS*, 492, 1641
- Minniti D., 1995, *AJ*, 109, 1663
- Minniti D. et al., 2010, *New Astron.*, 15, 433
- Mucciarelli A., Lapenna E., Massari D., Ferraro F. R., Lanzoni B., 2015a, *ApJ*, 801, 69
- Mucciarelli A., Lapenna E., Massari D., Pancino E., Stetson P. B., Ferraro F. R., Lanzoni B., Lardo C., 2015b, *ApJ*, 809, 128
- Mucciarelli A., Merle T., Bellazzini M., 2017, *A&A*, 600, A104
- Muraveva T. et al., 2015, *ApJ*, 807, 127
- Nataf D. M. et al., 2019, *AJ*, 158, 14
- Navarrete C. et al., 2015, *A&A*, 577, A99
- Nidever D. L. et al., 2015, *AJ*, 150, 173
- Oliveira R. A. P. et al., 2020, *ApJ*, 891, 37
- Pandas Development Team, 2020, pandas-dev/pandas: Pandas
- Pastorelli G. et al., 2019, *MNRAS*, 485, 5666
- Pedregosa F. et al., 2011, *J. Mach. Learn. Res.*, 12, 2825
- Pérez-Villegas A., Barbuy B., Kerber L. O., Ortolani S., Souza S. O., Bica E., 2019, *MNRAS*, 491, 3251
- Pérez F., Granger B. E., 2007, *Comput. Sci. Eng.*, 9, 21
- Pietrukowicz P. et al., 2015, *ApJ*, 811, 113
- Prantzos N., Charbonnel C., Iliadis C., 2007, *A&A*, 470, 179
- Renzini A., 2008, *MNRAS*, 391, 354
- Renzini A. et al., 2015, *MNRAS*, 454, 4197
- Roming P. W. A. et al., 2005, *Space Sci. Rev.*, 120, 95
- Saito R. K. et al., 2012, *A&A*, 537, A107
- Salaris M., Held E. V., Ortolani S., Gullieuszik M., Momany Y., 2007, *A&A*, 476, 243
- Saviane I., da Costa G. S., Held E. V., Sommariva V., Gullieuszik M., Barbuy B., Ortolani S., 2012, *A&A*, 540, A27
- Schiavon R. P. et al., 2017, *MNRAS*, 466, 1010
- Shetrone M. et al., 2015, *ApJS*, 221, 24
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Smith V. V., et al., 2021, preprint ([arXiv:2103.10112](https://arxiv.org/abs/2103.10112))
- Sneden C., Cowan J. J., Gallino R., 2008, *ARA&A*, 46, 241
- Soszyński I. et al., 2014, *Acta Astron.*, 64, 177
- Soszyński I. et al., 2019, *Acta Astron.*, 69, 321
- Souza S. O., Kerber L. O., Barbuy B., Pérez-Villegas A., Oliveira R. A. P., Nardiello D., 2020, *ApJ*, 890, 38
- Surot F., Valenti E., Gonzalez O. A., Zoccali M., Sökmen E., Hidalgo S. L., Minniti D., 2020, *A&A*, 644, A140
- Surot F. et al., 2019, *A&A*, 623, A168
- Tang J., Bressan A., Rosenfield P., Slemmer A., Marigo P., Girardi L., Bianchi L., 2014, *MNRAS*, 445, 4287
- Taylor M. B., 2005, in Shopbell P., Britton M., Ebert R., eds, ASP Conf. Ser. Vol. 347, Astronomical Data Analysis Software and Systems XIV. Astron. Soc. Pac., San Francisco, p. 29
- Thévenin F., Idiart T. P., 1999, *ApJ*, 521, 753
- Tony J. L. et al., 2012, *ApJ*, 750, 99
- Valenti E., Ferraro F. R., Origlia L., 2007, *AJ*, 133, 1287
- Valenti E., Ferraro F. R., Origlia L., 2010, *MNRAS*, 402, 1729
- van der Walt S., Colbert S. C., Varoquaux G., 2011, *Comput. Sci. Eng.*, 13, 22
- Vasiliev E., 2019, *MNRAS*, 484, 2832
- Vasiliev E., Baumgardt H., 2021, preprint ([arXiv:2102.09568](https://arxiv.org/abs/2102.09568))
- Ventura P., D'Antona F., Di Criscienzo M., Carini R., D'Ercole A., vesperini E., 2012, *ApJ*, 761, L30

- Watkins L. L., van der Marel R. P., Bellini A., Anderson J., 2015, *ApJ*, 803, 29
- Wilson J. C. et al., 2010, in McLean I. S., Ramsay S. K., Takami H., eds, Proc. SPIE Conf. Ser. Vol. 7735, Ground-Based and Airborne Instrumentation for Astronomy III. SPIE, Bellingham, p. 77351C
- Wilson J. C. et al., 2012, in McLean I. S., Ramsay S. K., Takami H., eds, Proc. SPIE Conf. Ser. Vol. 8446, Ground-Based and Airborne Instrumentation for Astronomy IV. SPIE, Bellingham, p. 84460H
- Wilson J. C. et al., 2019, *PASP*, 131, 055001
- Wright E. L. et al., 2010, *AJ*, 140, 1868
- Zasowski G. et al., 2013, *AJ*, 146, 81
- Zasowski G. et al., 2017, *AJ*, 154, 198
- Zasowski G. et al., 2019, *ApJ*, 870, 138

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

NGC6544_sup_material.dat

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: APPENDIX

Table A1. Identifiers and stellar parameters, $[\alpha/\text{Fe}]$, $[\text{Fe}/\text{H}]$, and elemental abundances from the NGC 6544 members. Median values, standard deviation, and mean ASPCAP errors are also derived.

APOGEE ID	Type	T_{eff} (K)	$\log g$ (dex)	V_{HELIO} (km s^{-1})	SNR	$[\text{Fe}/\text{H}]$ (dex)	$[\alpha/\text{Fe}]$ (dex)	$[\text{C}/\text{Fe}]$ (dex)	$[\text{N}/\text{Fe}]$ (dex)	$[\text{O}/\text{Fe}]$ (dex)	$[\text{Na}/\text{Fe}]$ (dex)	$[\text{Mg}/\text{Fe}]$ (dex)	$[\text{Al}/\text{Fe}]$ (dex)	$[\text{Si}/\text{Fe}]$ (dex)	$[\text{K}/\text{Fe}]$ (dex)	$[\text{Ca}/\text{Fe}]$ (dex)	$[\text{Ce}/\text{Fe}]$ (dex)
2M18070312-2501429	lrgb	4949.61	2.42	-41.9	119.35	-1.47	0.16	0.23	0.47	0.23	0.11	0.17	0.30	0.22	0.75	0.23	-0.25
2M18070657-2500417	agb	4797.31	1.49	-30.9	291.71	-1.55	0.11	-0.14	0.53	0.21	-	0.12	0.25	0.16	0.42	0.24	0.28
2M18071190-2458586	lrgb	4929.79	2.27	-39.2	70.09	-1.41	0.23	0.42	0.12	0.52	0.34	0.27	0.23	0.20	-0.05	0.53	-
2M18071347-2458525	urgb	4782.61	1.95	-39.7	169.30	-1.48	0.18	-0.15	0.55	0.21	0.50	0.24	0.11	0.24	0.16	0.18	-0.17
2M18071474-2454595	lrgb	4943.43	2.17	-37.9	205.98	-1.54	0.26	0.10	0.17	0.57	0.44	0.23	-0.20	0.22	0.24	0.19	-0.79
2M18071492-2457397	agb	4819.18	1.71	-34.2	152.02	-1.62	0.29	-0.23	-0.23	0.28	-	0.31	-0.26	0.29	0.39	0.18	-0.13
2M18071622-2457231	urgb	4688.18	1.76	-51.4	206.36	-1.60	0.27	-0.27	0.21	0.37	-	0.40	-0.20	0.26	0.80	0.28	0.06
2M18071727-2504387	lrgb	5018.44	2.42	-36.6	94.82	-1.44	0.14	0.14	0.55	0.33	0.57	0.17	0.07	0.13	-0.47	0.21	0.14
2M18071947-2500517	lrgb	4986.75	2.27	-50.6	224.29	-1.46	0.23	0.10	-0.08	0.53	-0.59	0.26	-0.25	0.19	0.19	0.24	0.22
2M18072003-2502407	lrgb	5074.18	2.50	-38.5	114.17	-1.52	0.24	0.16	0.33	0.50	-	0.37	0.06	0.20	0.63	0.21	-0.84
2M18072114-2458359	lrgb	4861.69	2.23	-37.7	115.97	-1.39	0.20	0.08	0.49	0.19	0.62	0.26	0.52	0.33	0.39	0.30	-0.16
2M18072123-2502436	lrgb	5112.84	2.65	-30.5	117.47	-1.37	0.16	0.17	0.92	-	-	0.26	0.39	0.20	-0.04	-0.01	-
2M18072204-2501100	urgb	4674.79	1.73	-37.5	159.89	-1.44	0.21	-0.20	0.09	0.30	-0.13	0.22	-0.26	0.22	0.07	0.21	-0.17
2M18072226-2500325	urgb	4470.25	1.39	-38.2	492.25	-1.44	0.13	-0.36	0.83	0.19	-	0.13	0.22	0.16	0.42	0.22	0.39
2M18072230-2458509	agb	4896.84	1.74	-33.4	323.80	-1.76	0.31	-0.03	-	0.39	0.11	0.30	-0.21	0.28	0.42	0.10	-
2M18072402-2458088	lrgb	5099.30	2.51	-37.2	79.06	-1.37	0.20	0.25	0.71	-	-	0.22	0.25	0.27	0.39	0.11	-
2M18072492-2458004	agb	5062.54	1.90	-36.8	82.36	-1.71	0.21	0.06	0.03	0.21	-	0.32	0.23	0.29	-	-0.01	0.61
2M18072612-2457497	lrgb	4972.73	2.32	-42.4	99.79	-1.45	0.18	0.20	0.41	0.33	-	0.27	0.10	0.19	-0.34	0.27	-0.39
2M18072927-2502396	lrgb	4912.17	2.20	-40.2	109.95	-1.44	0.13	0.02	0.62	0.13	-	0.19	0.35	0.21	0.67	0.32	-0.93
2M18072970-2456359	urgb	4316.64	1.19	-47.1	650.68	-1.43	0.19	-0.40	0.71	0.21	0.16	0.16	-0.06	0.20	0.29	0.14	-0.53
2M18073271-2500281	lrgb	4885.97	2.16	-40.1	72.44	-1.43	0.23	-0.07	0.65	0.23	0.47	0.25	0.22	0.29	0.07	0.06	-0.97
2M18073312-2457396	lrgb	5029.56	2.35	-40.7	129.57	-1.49	0.24	0.09	-0.09	0.58	-	0.20	-0.21	0.24	0.30	0.22	0.96
2M18074298-2452101	urgb	4756.59	1.85	-36.7	226.44	-1.47	0.20	-0.09	0.09	0.31	-0.46	0.19	-0.11	0.20	0.37	0.10	0.36
Cluster median	-	-	-	-38.17	129.57	-1.46	0.20	0.06	0.44	0.30	0.25	0.24	0.10	0.22	0.33	0.21	-0.16
Cluster 1σ	-	-	-	5.10	138.04	0.10	0.05	0.20	0.31	0.14	0.38	0.07	0.23	0.05	0.31	0.11	0.52
ASPCAP mean error	-	113.57	0.09	0.06	-	0.02	0.02	0.05	0.07	0.05	0.13	0.03	0.03	0.02	0.10	0.06	0.11

This paper has been typeset from a \LaTeX file prepared by the author.