



SUSTAINED FLUVIAL DEPOSITION RECORDED IN MARS' NOACHIAN STRATIGRAPHIC RECORD

Francesco Salese, William McMahon, Matthew Balme, V. Ansan, Joel Davis,
Maarten Kleinhans

► To cite this version:

Francesco Salese, William McMahon, Matthew Balme, V. Ansan, Joel Davis, et al.. SUSTAINED FLUVIAL DEPOSITION RECORDED IN MARS' NOACHIAN STRATIGRAPHIC RECORD. Nature Communications, 2020, 11 (1), pp.2067. <10.1038/s41467-020-15622-0>. <hal-03595032>

HAL Id: hal-03595032

<https://hal.science/hal-03595032v1>

Submitted on 3 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

SUSTAINED FLUVIAL DEPOSITION RECORDED IN MARS' NOACHIAN STRATI- GRAPHIC RECORD

**Francesco Salese^{1,2,†,*}, William J. McMahon^{1,†}, Matthew R. Balme³, Veronique Ansan⁴,
Joel M. Davis⁵ and Maarten G. Kleinhans¹**

¹Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands

²International Research School of Planetary Sciences, Università Gabriele D'Annunzio, Pescara, Italy

³Planetary Environments Group, Open University, Walton Hall, Milton Keynes, UK

⁴LPG Nantes, UMR6112, CNRS-Université de Nantes, 2 rue de la Houssinière, BP 92208, 44322, Nantes cedex 3, France

⁵Department of Earth Sciences, Natural History Museum, Cromwell Road, Kensington, London, SW7 5BD, UK.

* Corresponding author e-mail f.salese@uu.nl.

† These authors contributed equally to this work.

ABSTRACT

Orbital observation has revealed a rich record of fluvial landforms on Mars, with much of this record dating 3.6-3.0 Ga. Despite widespread geomorphic evidence, few analyses of Mars' alluvial sedimentary-stratigraphic record exist, with detailed studies of alluvium largely limited to smaller sand-bodies amenable to study in-situ by rovers. These typically metre-scale outcrop dimensions have prevented interpretation of larger scale channel-morphology and long-term basin evolution, vital for understanding the past martian climate. Here we give an interpretation of a large sedimentary succession at Izola mensa within the NW Hellas Basin rim. The succession comprises channel and barform packages which together demonstrate that river deposition was already well established >3.7 Ga. The deposits mirror terrestrial analogues subject to low-peak discharge variation, implying that river deposition at Izola was subject to sustained, potentially perennial, fluvial flow. Such conditions would require an environment capable of maintaining large volumes of water for extensive time-periods, necessitating a precipitation-driven hydrological cycle.

INTRODUCTION

While the present-day martian surface is generally dry and cold, its geomorphic record contains compelling evidence for the former presence of liquid water¹⁻⁷. In addition to this rich geomorphic archive, Mars' increasingly accessible sedimentary rock record provides a repository of information from which to study how planet-wide patterns in deposition have changed over time. From orbit, martian sedimentary rocks have been observed for more than twenty years⁸⁻¹³, but detailed descriptions of large Noachian alluvial successions have so far been lacking. In fact, unequivocal sedimentary-stratigraphic evidence of alluvium has only been identified in-situ by rover-led studies¹⁴, and whilst rovers provide unprecedented direct access to extraterrestrial strata¹⁵, methodologies are typically limited by accessible outcrop dimensions. For example, the Shaler outcrop, an interpreted Hesperian fluvial deposit identified by Curiosity¹⁶, comprises a single 70-centimetre thick, 20-metre wide sandstone-body. This is a crucial scale difference when compared with orbital investigation: the physical dimensions of most rock outcrops studied on the ground to date are smaller than the dimensions of many geomorphic components of even moderate-sized extant river systems^{17,18}.

In this study we use High Resolution Imaging Science Experiment (HiRISE) image (25 cm/pixel) and topographic (1 m/pixel) data to describe the sedimentary-stratigraphic architecture of a far-larger, 1500-metre-wide, 190-metre-thick sedimentary succession. The recently identified Izola outcrop is located in the northwestern rim of the Hellas basin (Fig. 1A), a ~ 2000 km diameter impact structure containing a variety of 3.7 Ga Noachian Fe/Mg phyllosilicate-rich sedimentary intercrater plains, overlain by Hesperian-aged (~3.3 Ga) lava flows¹⁹.

These intercrater plains offer erosional windows which expose stratigraphic sections with well-preserved channel-forms, and which must be older than the ~3.7 Ga overlying plains¹⁹. The channel-forms and associated sedimentary packages are interpreted as the product of an actively depositing fluvial system, with the final sedimentary architecture suggesting that these Noachian-aged rivers were not typified by high-energy episodic floods, but rather perennial or semi-perennial fluvial flow. The scale and completeness of the sedimentary succession offers a so far unique opportunity to assess the larger-scale morphology of an evolving Noachian-aged fluvial system.

RESULTS AND DISCUSSION

The Sedimentary Succession

The outcrop exposes layered sedimentary strata which display a variety of large-scale stratal architectures consistent with an alluvial interpretation. Alluvial sedimentary strata can be subdivided into genetically related three-dimensional packages²⁰⁻²². The aggregate of these packages is referred to as the succession's sedimentary architecture, and is the product of the scale and behavior of the fluvial system over time. The outcrop appears to have undergone little post-depositional deformation, has a gentle dip and large-dimensions, so is suitable for the analysis of sedimentary architecture (Supplementary Figure 5). A hierarchy of bounding surfaces is applied to divide stratigraphy, hierarchically ordered to reflect river processes at varying scales (Table 1, Supplementary Figure 1). Two distinct sedimentary packages were recognized in the studied outcrop: 1) channelized packages; and 2) inclined accretionary surfaces. As details of sedimentary facies (usually discriminated by grain size and centimetre to decimetre-scale bedding and sedimentary structure)

are unattainable without in-situ investigation on the ground, only the three-dimensional geometry of the sedimentary packages are described, with no details of internal structure attempted.

Channelized Packages

In the observed stratigraphy, packages bound by lower erosional, channel-shaped (fourth-order) surfaces and truncated by flat, erosional (fourth- and fifth-order) surfaces are attributed to channel-fill deposition (Fig. 2, Fig. 3B, Supplementary Figure 2A). They are 5 to 15 metres thick, with observable lateral extents of up to 210 metres (although outcrop limitations restrict observation of the full lateral extent of many channel-forms: Table 2, Supplementary Figure 2). Internally, packages appear succession-dominated²³, comprising multiple aggrading (third-order) surfaces (Fig. 3B). Final channel banks and former channel margins coalesce, indicating that the original channels laterally migrated (Fig. 3B, 3D). Some channelized packages have a distinct channel wing (yellow asterisks in Fig. 2B; Supplementary Figure 3-4), which may archive a genetically associated levee or crevasse and thus strengthen the alluvial interpretation. Particular areas of the outcrop show high concentrations of discrete, but partly amalgamated, channelized packages (Fig. 3B). In these areas, smaller channelized packages may be nested inside larger examples (Fig. 3B), implying that periods of net-erosion locally occurred within a dominantly depositional regime. Areas containing clusters of channels are bottomed and topped by laterally extensive (up to 640 metre), low relief, fifth-order surfaces, which suggest a change in the type or location of the dominant depositional process. These surfaces probably reflect channel avulsion, in which the location of the active channel changes abruptly. The sediments enclosed between these fifth-order surfaces were therefore laid down between avulsion events and are thus defined as channel-belts²⁴.

Inclined Accretionary Surfaces

In the observed stratigraphy, wedge-shaped packages comprise gently inclined depositional surfaces (third and fourth order; Fig. 3D). They are topped and bottomed by flat, erosional surfaces (both fourth order), except on occasions where they can be traced laterally into an associated channelized package (Fig. 3B). Most frequently only erosional remnants are preserved, with deposits often passing laterally into areas of non-exposure (Fig. 3D). Thicknesses range from 1.5 to 14 metres, similar to the associated channels. Geometry and internal stacking patterns are suggestive of shifting fluvial barforms. For example, in some instances, bottomsets of fourth order surfaces are downlapped by third order foresets (Fig. 3D), demonstrating discrete intervals of bar-building. The outcrop orientation with respect to paleoflow is not well known, meaning it is not possible to build any unequivocal consensus on the direction of barform accretion with respect to the flow of the original channels (e.g., downstream accretion, lateral accretion). However, the identification of channel-forms in conjunction with the outcrops alignment with the regional slope direction suggests a broadly depositional-strike succession.

Alternative Explanations

Without in-situ validation of fluvial deposition other alternative possibilities must additionally be considered. In this section we examine two alternative origins for the studied sedimentary succession: aeolian deposition, and deposition within submarine channels.

Aeolian: if these deposits are have an aeolian origin, then the working hypothesis is that the channel-forms are in fact localised scour fills formed by migrating aeolian strata, and that this succession records periods of aeolian accumulation punctuated by erosion or stagnation. However, there are disconnects between the stratigraphic architecture described here, and that of large-scale cross-bedding typical of aeolian bedforms described on both Earth^{25,26} and Mars²⁷. No aeolian interpre-

tation can explain the observable lateral migration surfaces associated with the discrete channel-ized packages (Fig. 3B). Multiple generations of aeolian dunes would partially erode preceding examples, perceived at outcrop by marked downlapping surfaces not apparent anywhere across the succession. Also, the fill geometry of aeolian dune deposits is often symmetrical and concordant with the trough base, unless the final outcrop is cut obliquely to the original flow direction, in which case planes are perceived to fill asymmetrically and downlap onto the trough base²⁸ (again, an architectural style not recognised here). Finally, multiple possible examples of channel-wings, evident from a distinct inflexion point, are found across the outcrop (Fig. 2B, Supplementary Figure 3,4). Such elements can be variably interpreted as the topmost story of an individual channel or a genetically associated levee or crevasse²³, but do not concur with any aeolian model.

Submarine channels: certain depositional structures are found in both submarine and alluvial environments (e.g., channel wings), but several architectural characteristics in these deposits favour alluvial interpretation and suggest against a submarine origin. Evidence of lateral migration is widespread across the outcrop (i.e., inclined accretionary surfaces). Such an architectural style is more typical of alluvial settings, with submarine slope architectures more regularly dominated by vertical accretion²⁹. Some inclined accretion surfaces additionally have distinct foreset and bottomset elements (Fig. 3D), consistent with those of fluvial barforms which scale to bankfull water-depth. Furthermore, most described submarine slope channels on Earth are at least an order of magnitude greater in size than the channels identified in this study³⁰, which have dimensions consistent with many other geomorphic fluvial channels identified across the martian surface³¹. Finally, regional observations of paleolakes occupying the Hellas basin at various stages in its evolution³² are compatible with an alluvial interpretation for these deposits.

Although a degree of ambiguity exists in any interpretation of ancient sedimentary strata, whether on Earth or Mars, and disputes between subaqueous and subaerial interpretations are commonplace on both planets^{10,33,34}, considering the discussion herein, we argue that outcrop evidence strongly favours a fluvial origin for the described Izola architecture.

Depositional Environment

From outcrop measurements, the channel and barform thicknesses are up to 13 and 14 metres respectively. This provides minimum constraints for bankfull water-depths at the time of deposition. Numerical models demonstrate that preserved sand-body thickness is on average 30% of the original channel depth³⁵, so this approximation is almost certainly an underestimate. These hypothesized water depths would be comparable with those of modern, mature fluvial systems on Earth draining extensive basins³⁶, suggesting these Noachian-aged rivers had similar drainage capacities.

Channels and barforms account for the entire observable stratigraphy. The evidence for channels being succession dominated (Fig. 3B) implies that entrenchment was quickly followed by bed aggradation²³. The occurrence of distinct channel clusters is consistent with channel belts with relatively stable banks and limited lateral mobility³⁷. The internal structure of the identified barforms comprises remnants of numerous phases of bar-building activity (Fig. 3D). On Earth, deposits of rivers with low peak discharge variability preserve macroform structure, enabling the reconstruction of barform morphology^{35,38}. The similar preserved architectural style seen here, of relatively intact barforms and channels, implies that the depositing Izola rivers were also characterized by low peak discharge variability. Episodic flooding events may not have been responsible for deposition, but rather long-term, potentially perennial fluvial flow. This conclusion also indicates that the Izola rivers were not the product of meltwater flowing from a glacial front, as these

environments are more widely associated with fluctuating, high discharges, and so would be unlikely to leave a stratigraphic record comprising relatively intact channels and bars.

Siliciclastic Deposition On An Unvegetated Planet

Depositional models are largely based on sedimentary environments on Earth, where physical form and process is near ubiquitously influenced by biology³⁹. Comparisons between the terrestrial and martian sedimentary record therefore require careful consideration before application. For instance, copious observations demonstrate the various ways extant vegetation modifies fluvial processes and landforms^{40,41}. Most important from an astrobiological standpoint is the likelihood of alluvial mud becoming preserved on an unvegetated planet. In addition to increasing mud production through chemical weathering⁴², terrestrial vegetation promotes mud retention on the continents through above-ground baffling and below-ground stabilization^{41,43,44}. Through these combined processes, an upsurge in mudstone abundance within alluvial stratigraphy is observed in stratigraphic alignment with evolving land plants⁴⁵. Before this time terrestrial alluvium is predominantly sand-grade or coarser with few preserved muddy floodplain facies^{45,46}. Direct comparison between the pre-vegetation Earth and unvegetated martian record might therefore imply that the studied Izola alluvium might also be lacking in preserved mudstone. Such tangible characteristics of the terrestrial pre-vegetation alluvial record have been previously used as supportive evidence for environmental interpretations elsewhere on Mars: for example, the scarcity of pre-vegetation terrestrial floodplain mudstones was used to recommend a lacustrine, not floodplain, origin of the Sheepbed mudstone at Gale crater¹⁶. The abundance of mudstone in the studied outcrop here is not known or speculated upon, as their intrinsic fine-grained components require rover-based observations. However, the preserved channel-belt architecture, comprising relatively intact channels and barforms, suggests some degree of original channel belt stability, with naturally shear-

resistant sediment such as mud a possible candidate^{47,48}. With regard to channel and barform facies, evidence of relatively stable, deep-channeled drainage on pre-vegetation Earth is increasingly reported³⁷. Such findings are helping to dispel notions that pre-vegetation rivers were ubiquitously wide and shallow⁴⁹, an observation that can now be extended to Mars (Fig. 2).

Preservation Of Time And Implications For Early Mars Climate

Disentangling the total duration recorded from any sedimentary outcrop is difficult, as stratigraphic records are highly fragmentary, incomplete chronicles of time^{50,51}. Despite uncertainty, broad approximations from the studied outcrop can be made which can inform ongoing debates about the early martian climate⁵². The outcrop comprises at least four possible channel belts (Fig. 2), discrete packages of strata bound by laterally extensive (fifth order) surfaces and, on Earth, channel belts of this scale can require up to $\sim 10^4$ - 10^5 terrestrial years to deposit⁵² (though we recognize that this approximation may differ under Mars' distinct boundary conditions). However, the amount of additional time hidden within the bounding fifth order surfaces between individual channel-belts is unknown (and unknowable)⁵³. If a channel-belt interpretation is correct, the laterally extensive surfaces developed after the active channel-belt was transposed through avulsion. During this time, deposition was likely occurring elsewhere in the basin, and the region covered by our sedimentary outcrop was undergoing intervals of net erosion or stasis⁵⁴. In other words, hiatuses in deposition in the studied outcrop were accompanied by deposition elsewhere in the basin. This implies that the time recorded in the 190-metre thick succession represents only a fraction of the total time fluvial deposition was ongoing in this region. The majority of strata and time will have either been lost to erosion, or preserved in outcrops as of yet undiscovered, or currently buried and not amenable to study. While we are only beginning to understand the chronostratigraphic exactness of

sedimentary rock outcrops on Earth^{53,55,56}, let alone Mars, it appears likely that the period of deposition in the northwestern rim of the Hellas basin exceeded 10^5 terrestrial years. Furthermore, the preservation of relatively intact channel-margins and barforms advocates that throughout this protracted period fluvial deposition was a relatively constant phenomena³⁸.

The architectural interpretation of this so far unique sedimentary succession feeds into ongoing debates about the early martian climate. Our interpretation of long-lived, deep, perennial or semi-perennial rivers necessitates a climate in which active water-conduits were maintained for 10^5 years or longer. For the first time, orbital data has allowed us to examine, through detailed high-resolution architectural analysis, a large (1500 metre by 190 metre) pre-late Noachian outcrop, and draw reliable paleoenvironmental interpretations based on sedimentary-stratigraphic evidence. Our observations and analysis favour steady water discharges that are most consistent with a precipitation-driven hydrological cycle. This conclusion aligns with previous arguments for the prolonged presence of water on the early martian surface drawn from alternative geomorphological^{2,6,14,19} and mineralogical^{57,58} observations.

Acknowledgements This research was funded by F.S. Marie Curie Individual Postdoctoral Fellowship (WET_MARS, Grant Agreement 795192). W.M and M.K were funded by the European Research Council (ERC Consolidator grant 647570 to M.K). The authors wish to thank the Mars Reconnaissance Orbiter and HiRISE teams and USGS/IAU for having named the studied outcrop and the nearby crater in honour of the Slovenian town of Izola.

Author contributions F.S. conceptualized the study; F.S. and W.J.M. devised the methodology, analyzed the data and wrote the paper; M.R.B., V.A., J.M.D., M.G.K. revised the paper and contributed to the discussion; J.M.D. processed the HiRISE stereo pairs and built the digital elevation model; F.S. and M.K. acquired funding.

Competing interests The authors declare no competing interests.

Data Availability Statement The HiRISE data that support the findings of this study were obtained freely from the Planetary Data System (PDS) and are publicly available online at <https://pds.nasa.gov/index.shtml>. Satellite imagery and the Extended Data were generated with ISIS 3 (Integrated Software for Imagers and Spectrometers) available online at <https://isis.astrogeology.usgs.gov>. All these data were integrated into ArcMap 10.6 project. The DTM was produced using the USGS Integrated Software for Imagers and Spectrometers (ISIS 3) software and the BAE photogrammetric package SOCET SET with a post spacing of 1 m/pixel.

Correspondence and requests for materials should be addressed to Francesco Salese (f.salese@uu.nl)

245

246 **METHODS**

247 **Dtm Construction**

248 High Resolution Imaging Science Experiment⁵⁹ image ESP_055357_1540 acquired in May 2018
249 along with HiRISE image PSP_003799_1540 (acquired in 2007) enabled the construction of a
250 centimetre scale DTM. Both images have 25.6 cm/pixel (with 1 x 1 binning) resolution so objects
251 down to 77 cm can be resolved. A digital elevation model (DEM) was produced from the HiRISE
252 images ESP_055357_1540 and PSP_003799_1540 using the USGS Integrated Software for Im-
253 agers and Spectrometers (ISIS) software and the BAE photogrammetric package SOCET SET
254 according to a previously used methodology⁶⁰. Tie points were automatically populated in SOCET
255 SET between the two images. We ran a series of bundle adjustments, removing erroneous tie points
256 until the remaining points had an RMS pixel matching error of ≤ 0.6 pixels. The resultant DEM
257 was then tied to Mars Orbital Laser Altimeter⁶¹ topography and exported with a horizontal post
258 spacing of 1 m/pixel and a vertical precision of ~ 1 m.

259 **Calculation Of Sedimentary Package Dimensions**

260 The acquired HiRISE DTM of the Hellas outcrop was of sufficient resolution to enable accurate
261 tracing of beds and plotting of architectural elements (Figure 2). Channels depths were obtained
262 measuring the exact elevation of the channel top and the exact elevation of the channel base using
263 ArcMap 10.6 elevation tools. Channel widths were measured by tracing an edge-to-edge topo-
264 graphic channel profile using the HiRISE DEM. This allowed calculation of true thickness, given
265 that they are bound by laterally-extensive fifth-order surfaces, which are almost flat lying (Sup-
266 plementary Figure 5).

267

268 **Architectural Analysis**

269 Line drawings were only attempted at areas where stratigraphy is clearly visible. Outcrop orienta-
270 tion with respect to paleoflow is not known, so no architectural elements with distinct directional
271 components (e.g., downstream accretion, lateral accretion) were assigned. Sediment grain size is
272 also unknown, thereby prohibiting the distinction between active and abandoned channel-fill de-
273 posits. The completed architectural panel enabled the various sediment stacking patterns and lat-
274 eral relationships assessed.

275 **Bounding Surface Hierarchy**

276 A hierarchy of bounding surfaces was used to describe partitions of fluvial strata at outcrop, with
277 different order surfaces reflecting river processes at varying scales⁶² (Table 1, Supplementary Fig-
278 ure 1). The principles of the hierarchical division applied here follows that detailed in previous
279 papers^{22,62}. Succinctly, zeroth, first, and second order surfaces relate to foreset, set and coset
280 boundaries respectively, and are not observable from HiRISE imagery. Third and fourth order
281 surfaces indicate the presence of macroforms (e.g., a barform deposit) or a channel. Fourth order
282 surfaces represent the upper and lower boundaries of the macroform or channel, whereas third
283 order surfaces relate to internal growth increments (indicating flow fluctuation, but no significant
284 changes in predominant fluvial style). Fifth order surfaces are the highest order observed at the
285 studied outcrop and bind major depositional packages (e.g., channel-belts). More fifth order sur-
286 faces are likely present in the studied outcrop than highlighted on Figure 2. This is simply because
287 their confident recognition depends on an understanding of their relationship with lower order

surfaces, and in some instances the vagaries of outcrop exposure (particularly towards the bottom of the section) prevent this.

Estimation Of Outcrop Age

The age of the intercrater plains, which form the planform cover of the studied sedimentary-stratigraphic succession, has been estimated in a previous work¹⁹ using various crater count techniques. They date $3.70 \pm 0.03/-0.04$ Ga (Noachian). The latter represents the age of the surface at the top of the outcrop studied in this work, which lies in the S1 unit (Figure 18¹⁹). Crater counts were performed on Context Camera (CTX; 5–6m/pixel) data using Crater Tools. Crater statistics and crater model ages were analyzed with Craterstats2 software. For more details and references see a previous study¹⁹.

REFERENCES

- 1 Carr, M. H. The Martian drainage system and the origin of valley networks and fretted channels. *Journal of Geophysical Research: Planets* **100**, 7479-7507 (1995).
- 2 Davis, J. M., Balme, M., Grindrod, P. M., Williams, R. M. E. & Gupta, S. Extensive Noachian fluvial systems in Arabia Terra: Implications for early Martian climate. *Geology* **44**, 847-850 (2016).
- 3 Di Achille, G. & Hynek, B. M. Ancient ocean on Mars supported by global distribution of deltas and valleys. *Nature Geoscience* **3**, 459 (2010).
- 4 Howard, A. D., Moore, J. M. & Irwin III, R. P. An intense terminal epoch of widespread fluvial activity on early Mars: 1. Valley network incision and associated deposits. *Journal of Geophysical Research: Planets* **110** (2005).
- 5 Malin, M. C. & Edgett, K. S. Evidence for persistent flow and aqueous sedimentation on early Mars. *Science* **302**, 1931-1934 (2003).
- 6 Mangold, N., Quantin, C., Ansan, V., Delacourt, C. & Allemand, P. Evidence for precipitation on Mars from dendritic valleys in the Valles Marineris area. *Science* **305**, 78-81 (2004).
- 7 Salese, F., Pondrelli, M., Neeseman, A., Schmidt, G. & Ori, G. G. Geological Evidence of Planet-Wide Groundwater System on Mars. *Journal of Geophysical Research: Planets* **124**, 374-395 (2019).
- 8 Ansan, V. *et al.* Stratigraphy, mineralogy, and origin of layered deposits inside Terby crater, Mars. *Icarus* **211**, 273-304, doi:10.1016/j.icarus.2010.09.011 (2011).
- 9 Cardenas, B. T., Mohrig, D. & Goudge, T. A. Fluvial stratigraphy of valley fills at Aeolis Dorsa, Mars: Evidence for base-level fluctuations controlled by a downstream water body. *GSA Bulletin* **130**, 484-498 (2018).
- 10 Dromart, G., Quantin, C. & Broucke, O. Stratigraphic architectures spotted in southern Melas Chasma, Valles Marineris, Mars. *Geology* **35**, 363-366 (2007).

322 11 Goudge, T. A., Mohrig, D., Cardenas, B. T., Hughes, C. M. & Fassett, C. I. Stratigraphy and
323 paleohydrology of delta channel deposits, Jezero crater, Mars. *Icarus* **301**, 58-75 (2018).
324 12 Lucchitta, B. K., Ferguson, H. M. & Summers, C. Sedimentary Deposits in the Northern Lowland
325 Plains, Mars. *J Geophys Res-Solid* **91**, E166-E174, doi:DOI 10.1029/JB091iB13p0E166 (1986).
326 13 Malin, M. C. & Edgett, K. S. Sedimentary rocks of early Mars. *Science* **290**, 1927-1937 (2000).
327 14 Williams, R. M. E. *et al.* Martian fluvial conglomerates at Gale crater. *science* **340**, 1068-1072
328 (2013).
329 15 Edgar, L. A. *et al.* Shaler: in situ analysis of a fluvial sedimentary deposit on Mars. *Sedimentology*
330 **65**, 96-122 (2018).
331 16 Grotzinger, J. P. *et al.* A habitable fluvio-lacustrine environment at Yellowknife Bay, Gale Crater,
332 Mars. *Science* **343**, 1242777 (2014).
333 17 McMahon, W. J. & Davies, N. S. The shortage of geological evidence for pre-vegetation
334 meandering rivers. *Fluvial Meanders and Their Sedimentary Products in the Rock Record*, 119-148
335 (2018).
336 18 Miall, A. D. How do we identify big rivers? And how big is big? *Sedimentary Geology* **186**, 39-50
337 (2006).
338 19 Salese, F. *et al.* A sedimentary origin for intercrater plains north of the Hellas basin: Implications
339 for climate conditions and erosion rates on early Mars. *Journal of Geophysical Research: Planets*
340 **121**, 2239-2267 (2016).
341 20 Allen, J. R. L. Studies in fluvial sedimentation: bars, bar-complexes and sandstone sheets (low-
342 sinuosity braided streams) in the Brownstones (L. Devonian), Welsh Borders. *Sedimentary*
343 *Geology* **33**, 237-293 (1983).
344 21 Friend, P. F. Towards the field classification of alluvial architecture or sequence. *Modern and*
345 *ancient fluvial systems*, 345-354 (1983).
346 22 Miall, A. D. Architectural-element analysis: a new method of facies analysis applied to fluvial
347 deposits. *Earth-Science Reviews* **22**, 261-308 (1985).
348 23 Gibling, M. R. Width and thickness of fluvial channel bodies and valley fills in the geological record:
349 a literature compilation and classification. *Journal of sedimentary Research* **76**, 731-770 (2006).
350 24 McLaurin, B. T. & Steel, R. J. Architecture and origin of an amalgamated fluvial sheet sand, lower
351 Castlegate Formation, Book Cliffs, Utah. *Sedimentary Geology* **197**, 291-311 (2007).
352 25 Kocurek, G. & Day, M. What is preserved in the aeolian rock record? A Jurassic Entrada Sandstone
353 case study at the Utah–Arizona border. *Sedimentology* **65**, 1301-1321 (2018).
354 26 Mountney, N. P., Posamentier, H. W. & Walker, R. G. Eolian facies models. *SPECIAL PUBLICATION-*
355 *SEPM* **84**, 19 (2006).
356 27 Banham, S. G. *et al.* Ancient Martian aeolian processes and palaeomorphology reconstructed from
357 the Stimson formation on the lower slope of Aeolis Mons, Gale crater, Mars. *Sedimentology* **65**,
358 993-1042 (2018).
359 28 DeCelles, P. G., Langford, R. P. & Schwartz, R. K. Two new methods of paleocurrent determination
360 from trough cross-stratification. *Journal of Sedimentary Research* **53**, 629-642 (1983).
361 29 Jobe, Z. R., Howes, N. C. & Auchter, N. C. Comparing submarine and fluvial channel kinematics:
362 Implications for stratigraphic architecture. *Geology* **44**, 931-934 (2016).
363 30 Kolla, V., Posamentier, H. W. & Wood, L. J. Deep-water and fluvial sinuous channels—
364 Characteristics, similarities and dissimilarities, and modes of formation. *Marine and Petroleum*
365 *Geology* **24**, 388-405 (2007).
366 31 Kite, E. S. *et al.* Persistence of intense, climate-driven runoff late in Mars history. *Science Advances*
367 **5**, eaav7710 (2019).
368 32 Wilson, S. A., Moore, J. M., Howard, A. D. & Wilhelms, D. E. Evidence for ancient lakes in the Hellas
369 region. *Lakes on Mars*, 195-222 (2010).

370 33 Bourgeois, J. & Som, S. Stratigraphic architectures spotted in southern Melas Chasma, Valles
371 Marineris, Mars: COMMENT: COMMENT. *Geology* **35**, e145-e145 (2007).

372 34 Wizevich, M. C., Ahern, J. & Meyer, C. A. The Triassic of southwestern Switzerland–Marine or non-
373 marine, that is the question! *Palaeogeography, Palaeoclimatology, Palaeoecology* **514**, 577-592
374 (2019).

375 35 van de Lageweg, W. I., van Dijk, W. M., Box, D. & Kleinhans, M. G. Archimetrics: a quantitative tool
376 to predict three-dimensional meander belt sandbody heterogeneity. *The Depositional Record* **2**,
377 22-46 (2016).

378 36 Ashworth, P. J. & Lewin, J. How do big rivers come to be different? *Earth-Science Reviews* **114**, 84-
379 107 (2012).

380 37 Ielpi, A., Ventra, D. & Ghinassi, M. Deeply channelled Precambrian rivers: Remote sensing and
381 outcrop evidence from the 1.2 Ga Stoer Group of NW Scotland. *Precambrian Research* **281**, 291-
382 311 (2016).

383 38 Fielding, C. R., Alexander, J. & Allen, J. P. The role of discharge variability in the formation and
384 preservation of alluvial sediment bodies. *Sedimentary Geology* **365**, 1-20 (2018).

385 39 Davies, N. S., Shillito, A. P., Slater, B. J., Liu, A. G. & McMahon, W. J. Evolutionary synchrony of
386 Earth's biosphere and sedimentary-stratigraphic record. *Earth-Science Reviews*, 102979 (2019).

387 40 Corenblit, D., Davies, N. S., Steiger, J., Gibling, M. R. & Bornette, G. Considering river structure and
388 stability in the light of evolution: feedbacks between riparian vegetation and
389 hydrogeomorphology. *Earth Surface Processes and Landforms* **40**, 189-207 (2015).

390 41 Kleinhans, M. G., de Vries, B., Braat, L. & van Oorschot, M. Living landscapes: Muddy and
391 vegetated floodplain effects on fluvial pattern in an incised river. *Earth surface processes and*
392 *landforms* **43**, 2948-2963 (2018).

393 42 Hazen, R. M. *et al.* Clay mineral evolution. *American Mineralogist* **98**, 2007-2029 (2013).

394 43 Gurnell, A. Plants as river system engineers. *Earth Surface Processes and Landforms* **39**, 4-25
395 (2014).

396 44 Mitchell, R. L. *et al.* Mineral weathering and soil development in the earliest land plant
397 ecosystems. *Geology* **44**, 1007-1010 (2016).

398 45 McMahon, W. J. & Davies, N. S. Evolution of alluvial mudrock forced by early land plants. *Science*
399 **359**, 1022-1024 (2018).

400 46 Gibling, M. R. & Davies, N. S. Palaeozoic landscapes shaped by plant evolution. *Nature Geoscience*
401 **5**, 99 (2012).

402 47 Lapôtre, M. G. A., Ielpi, A., Lamb, M. P., Williams, R. M. E. & Knoll, A. H. Model for the formation
403 of single-thread rivers in barren landscapes and implications for pre-Silurian and martian fluvial
404 deposits. *Journal of Geophysical Research: Earth Surface* (2019).

405 48 Matsubara, Y. *et al.* River meandering on Earth and Mars: A comparative study of Aeolis Dorsa
406 meanders, Mars and possible terrestrial analogs of the Usuktuk River, AK, and the Quinn River,
407 NV. *Geomorphology* **240**, 102-120 (2015).

408 49 McMahon, W. J. & Davies, N. S. Physical and biological functioning in Proterozoic rivers: evidence
409 from the archetypal pre-vegetation alluvium of the Torridon Group, NW Scotland. *Scottish Journal*
410 *of Geology* (2019).

411 50 Barrell, J. Rhythms and the measurements of geologic time. *Bulletin of the Geological Society of*
412 *America* **28**, 745-904 (1917).

413 51 Miall, A. D. Updating uniformitarianism: stratigraphy as just a set of 'frozen accidents'. *Geological*
414 *Society, London, Special Publications* **404**, 11-36 (2015).

415 52 Wordsworth, R. D., Kerber, L., Pierrehumbert, R. T., Forget, F. & Head, J. W. Comparison of "warm
416 and wet" and "cold and icy" scenarios for early Mars in a 3-D climate model. *Journal of*
417 *Geophysical Research: Planets* **120**, 1201-1219 (2015).

- 53 Davies, N. S., Shillito, A. P. & McMahon, W. J. Where does the time go? Assessing the
418 chronostratigraphic fidelity of sedimentary rock outcrops in the Pliocene-Pleistocene Red Crag
419 Formation, eastern England. *Journal of the Geological Society*, jgs2019-2056 (2019).
420
- 54 Tipper, J. C. The importance of doing nothing: stasis in sedimentation systems and its stratigraphic
421 effects. *Geological Society, London, Special Publications* **404**, 105-122 (2015).
422
- 55 Holbrook, J. & Miall, A. D. Time in the Rock: A field guide to interpreting past events and processes
423 from a fragmentary siliciclastic archive. *Earth-Science Reviews*, 103121 (2020).
424
- 56 Paola, C., Ganti, V., Mohrig, D., Runkel, A. C. & Straub, K. M. Time not our time: Physical controls
425 on the preservation and measurement of geologic time. *Annual Review of Earth and Planetary*
426 *Sciences* **46**, 409-438 (2018).
427
- 57 Bibring, J. P. *et al.* Global mineralogical and aqueous Mars history derived from OMEGA/Mars
428 Express data. *science* **312**, 400-404 (2006).
429
- 58 Ehlmann, B. L. & Edwards, C. S. Mineralogy of the Martian surface. *Annual Review of Earth and*
430 *Planetary Sciences* **42**, 291-315 (2014).
431
- 59 McEwen, A. S. *et al.* Mars Reconnaissance Orbiter's High Resolution Imaging Science Experiment
432 (HiRISE). *J Geophys Res-Planet* **112**, doi:Artn E05s02
433 10.1029/2005je002605 (2007).
434
- 60 Kirk, R. L. *et al.* Ultrahigh resolution topographic mapping of Mars with MRO HiRISE stereo images:
435 Meter-scale slopes of candidate Phoenix landing sites. *Journal of Geophysical Research: Planets*
436 **113** (2008).
437
- 61 Smith, D. E. *et al.* Mars Orbiter Laser Altimeter: Experiment summary after the first year of global
438 mapping of Mars. *J Geophys Res-Planet* **106**, 23689-23722, doi:Doi 10.1029/2000je001364 (2001).
439
- 62 Holbrook, J. Origin, genetic interrelationships, and stratigraphy over the continuum of fluvial
440 channel-form bounding surfaces: an illustration from middle Cretaceous strata, southeastern
441 Colorado. *Sedimentary Geology* **144**, 179-222 (2001).
442
443
444
445




Fig. 1 General context of study location at Izola mensa. a Mars Orbiter Laser Altimeter (MOLA) topographic map centred on the NW rim of the Hellas basin, showing a depression (blue/violet) in the centre-lower right corner, the cratered highlands (brown-orange) in the upper left corner and the location of the studied outcrop within the red box. The magenta arrow indicates the slope direction of the modern surface. The identified macroforms suggest that the outcrop is broadly cut in the depositional-strike direction, in alignment with regional slope. The topographic elevation spans from 1500 m (brown) to -1350 m (white). Bottom right inset shows the regional context. Mars Orbiter Laser Altimeter (MOLA) topographic map centred on the northwestern shoulder of the Hellas basin. Study area indicated by white box. The topographic elevation spans from 4319 m (brown) to 8194 m (white). b High-resolution imaging science experiment (HiRISE) contour lines from HiRISE digital elevation model (1 m/pixel) (HiRISE stereo pairs ESP_055357_1540; PSP_003799_1540) over HiRISE visible image (25 cm/pixel). The Izola outcrop is located at 25.88°S and 54.29°E and faces almost N-S. c 3D view without vertical exaggeration of the Izola outcrop shown in b. HiRISE ESP_055357_1540 (25 cm/pixel) draped on HiRISE digital elevation model (1 m/pixel).

446

447

448




Fig. 2 Architectural analysis of Martian channel forms and inclined accretionary surfaces at the Izola outcrop. a High-resolution imaging science experiment (HiRISE) image of the studied outcrop (ESP_055357_1540; 25 cm/pixel resolution). Objects down to 77 cm can be resolved. The outcrop is 1500-m-wide and 190-m-thick. Note the different scale between the vertical and horizontal axis (vertically exaggerated) b Line drawing of a to illustrate an architectural interpretation, displayed inc. Line drawings were only attempted in areas where stratigraphy was clearly visible (no lines were joined across areas on non-exposure). The yellow asterisk and hashtag link to Supplementary Figs. S3 and S4, both possible examples of channel wings. Identified fifth-order surfaces are labelled with letters and surface dimensions and listed in Table 2. The well-exposed channels are labelled with numbers and their parameters reported in Table 2.




Fig. 3 Example channelised packages and inclined accretionary surfaces. a Close up of white box indicated in Fig 2b. b Architectural interpretation of a displaying a number of channels (blue lines). Some examples preserve former channel margins, strong evidence for original channel lateral migration. Channelised packages have associated inclined accretion surfaces (orange lines), with all packages topped and floored by more extensive fifth-order surfaces (red lines). Fifth-order surfaces possibly archive avulsion events. A previous channel margin (1+ 3) laterally coalesces with the final channel margin (2+ 4) indicating that the original active channel migrated laterally. A nested channel-cut (5) is present within a larger channelised package. c Close up of white box indicated in Fig 2b. d Examples of inclined accretion surfaces with distinct foreset and bottomset elements (topsets truncated by overlying strata). Downlapping of internal, third-order surfaces suggest multiple accretion phases. Note that line drawings are not attempted in areas where exposure is poor. A third-order surface (7) downlaps the bottomset of an underlying fourth-order surface (6) indicating bar migration. A subsequent fourth-order surface (8) truncates the previous fourth-order surface (7) indicating multiple phases of bar-building activity.

Bounding surface order	Maximum lateral extent	Bounded units	Sedimentary process
Fifth	640 metres	Channel belts	Switch between dominant depositional process (e.g., channel-belt avulsion)
Fourth	210 metres	Channel fills, fluvial barforms	Termination of accretion or reworking of a discrete macroform
Third	180 metres	Macroforms	Accretion of a discrete macroform
Second	Not observable (rover only)	Cosets	Accretion and reworking of mesoforms
First	Not observable (rover only)	Sets	Ripple or dune migration
Zeroth	Not observable (rover only)	Laminae	Burst-sweep cycle

Table 1 Hierarchical division of bounding surfaces applied in this study. Surface ranking and process interpretations adopted from previous studies^{22,62}.

4 th order ID	Depth (m)	Width (m)	Width/Depth Ratio
1	8	150	18
2	11	110	10
3	15	210	14
4	7	110	15
5	15	170	11

5th order ID

A-A'	undefined	570	n/a
B-B'	21	345	n/a
C-C'	30	640	n/a

Table 2. 4th and 5th order observed bounding surface dimensions. The ID locations of these surfaces are reported in Figure 2C.