

Multistage stochastic programs with the entropic risk measure

Bernardo Pagnoncelli, Oscar Dowson, David Morton

► To cite this version:

Bernardo Pagnoncelli, Oscar Dowson, David Morton. Multistage stochastic programs with the entropic risk measure. 23ème congrès annuel de la Société Française de Recherche Opérationnelle et d'Aide à la Décision, INSA Lyon, Feb 2022, Villeurbanne - Lyon, France. hal-03596198

HAL Id: hal-03596198 https://hal.science/hal-03596198v1

Submitted on 3 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Multistage stochastic programs with the entropic risk measure

Bernardo K. Pagnoncelli¹, Oscar Dowson², David Morton³

¹ SKEMA Business School, Université Côte d'Azur, F-59800, Lille, France

bernardo.pagnoncelli@skema.edu

² Department of Industrial Engineering and Management Sciences, Northwestern University o.dowson@gmail.com

³ Department of Industrial Engineering and Management Sciences, Northwestern University david.morton@northwestern.edu

Mots-clés : stochastic programming; convex risk measures; entropic risk measure;

1 Introduction

In order to make sequential decisions under uncertainty, it is convenient to consider *risk* measures. Most works use dynamic programming, and consider the so-called *nested* risk measures. [2, 3]. When we care about the outcome of our decision at the last stage, *end-of-horizon* formulations are more appropriate. In this work we study the entropic risk measure, for which the two approaches coincide.

2 Conditional consistency and the entropic risk measure

Consider a sequence of correlated real-valued random variables, $Z = \{Z_t\}_{t=1}^T$. We have

$$End - of - Horizon - Risk(Z) = \mathbb{F}[Z_1 + Z_2 + \dots + Z_T],$$

$$Nested - Risk(Z) = \mathbb{F}\left[Z_1 + \mathbb{F}_{Z_2|Z_1}[Z_2 + \mathbb{F}_{Z_3|Z_2}[\dots + \mathbb{F}_{Z_T|Z_{T-1}}[Z_T]]\right].$$

Définition 1 Let (X_1, X_2) and (Y_1, Y_2) be two-dimensional vectors for which requisite expectations are finite. A risk measure \mathbb{F} is said to be conditionally consistent if :

$$\mathbb{F}[X_1 + X_2] \le \mathbb{F}[Y_1 + Y_2] \iff \mathbb{F}[X_1 + \mathbb{F}_{X_2|X_1}[X_2]] \le \mathbb{F}[Y_1 + \mathbb{F}_{Y_2|Y_1}[Y_2]].$$

An example of a convex risk measure that is conditionally consistent is the entropic $(\gamma > 0)$:

$$\mathbb{ENT}_{\gamma}[Z] = \frac{1}{\gamma} \log\left(\mathbb{E}[e^{\gamma Z}]\right).$$
(1)

2.1 Risk-averse multistage Stochastic Programming (MSSP)

We consider a T-stage MSSP given by

$$V_{t}(x_{t-1}, \omega_{t}) = \min_{\bar{x}_{t}, x_{t} \ge 0} \quad c_{t}^{\top} x_{t} + \mathbb{F}_{\omega_{t+1} \in \Omega_{t+1}} [V_{t+1}(x_{t}, \omega_{t+1})] \\ \bar{x}_{t} = x_{t-1} \quad [\lambda] \\ A_{t} x_{t} + B_{t} \bar{x}_{t} = b_{t}.$$
(2)

We can form single-cut master problems that approximate the stage-wise problems as follows :

$$V_{t}^{K}(x_{t-1},\omega_{t}) = \min_{\bar{x}_{t},x_{t}\geq0,\theta_{t+1}\geq-M_{t+1}} c_{t}^{\top}x_{t} + \theta_{t+1} \bar{x}_{t} = x_{t-1} [\lambda] A_{t}x_{t} + B_{t}\bar{x}_{t} = b_{t} \theta_{t+1}\geq\alpha_{t+1,k} + \beta_{t+1,k}^{\top}x_{t}, \quad k = 1,\dots, K-1,$$
(3)

where M_{t+1} is a lower bound on $\mathbb{F}[V_{t+1}(\cdot, \omega_{t+1})]$. Explicit values of α_{t+1} and β_{t+1} can be computed for the entropic risk measure, and such formulation is amenable to SDDP.

3 Numerical illustration

Consider a road network with three arcs (see [1]) as shown in Figure 1. Travel times are independent, and we must choose between $\mathbb{F}[X + Z]$ and $\mathbb{F}[Y + Z]$.




FIG. 1 – Road network with three independent travel times, X, Y, and Z.

The results are shown in Figure 2 : for the $\mathbb{CV}@\mathbb{R}$, when $\gamma \in (0.865, 0.97)$, the optimal decision to the end-of-horizon model is to switch back to X. In Figure 2b, since the end-of-horizon and nested formulations are equivalent, there is only one line visible on the graph.




FIG. 2 – x = 0 (road Y), x = 1 (road X) against γ using $\mathbb{CV}@\mathbb{R}$ (a) and entropic (b)

4 Conclusions

We studied the entropic risk measure in MSSP. We showed that it can be embedded within decomposition algorithms such as the SDPP, and exemplified its use in a transportation problem.

Références

- R. Cominetti and A. Torrico. Additive consistency of risk measures and its application to risk-averse routing in networks. *Math. of Oper. Res.*, 41(4):1510–1521, 2016.
- [2] Andy Philpott, Vitor de Matos, and Erlon Finardi. On solving multistage stochastic programs with coherent risk measures. *Operations Research*, 61(4):957–970, 2013.
- [3] A. Shapiro, W. Tekaya, J.P. da Costa, and M. P. Soares. Risk neutral and risk averse stochastic dual dynamic programming method. *EJOR*, 224(2) :375–391, 2013.