

Animal social networks: Towards an integrative framework embedding social interactions, space and time

Sebastian Sosa, David Jacoby, Mathieu Lihoreau, Cédric Sueur

▶ To cite this version:

Sebastian Sosa, David Jacoby, Mathieu Lihoreau, Cédric Sueur. Animal social networks: Towards an integrative framework embedding social interactions, space and time. Methods in Ecology and Evolution, 2021, 12 (1), pp.4-9. 10.1111/2041-210x.13539. hal-03608063

HAL Id: hal-03608063 https://hal.science/hal-03608063

Submitted on 7 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. 1 Animal social networks: towards an integrative framework embedding social

2 interactions, space and time

- 3 Sebastian Sosa^{1a}, David M.P. Jacoby², Mathieu Lihoreau³, Cédric Sueur^{1,4}
- ⁴ ¹ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

⁵ ² Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY,

6 UK

⁷ ³ Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI);

- 8 CNRS, University Paul Sabatier Toulouse III
- 9⁴ Institut Universitaire de France, 75007 Paris, France
- 10 ^a Corresponding author
- 11 All authors contributed equally
- 12

13 Fifty years of social network analysis

Social groups take a myriad of forms, reflecting the countless different ways in which 14 15 animals can interact and associate (Wilson 2000). This diversity calls for a broad and 16 dynamic toolkit that is both robust in allowing rigorous quantification of different 17 societies but also flexible in its ability to account for and control the nuances associated with each ecological system. Since Sade (1972) first used social network analysis to 18 study heterogeneity in the affiliative social interactions of primates and their 19 relationship with dominance rank, this approach considerably developed for the 20 exploration and hypothesis testing of different aspects of animal social interactions 21 22 (Hasenjager, Leadbeater & Hoppitt; Hinde 1976; Brent, Lehmann & Ramos-Fernández 2011). 23

Over the last 50 years, social network analysis has grown into a diverse toolkit 24 that transcends animal behaviour and ecology allowing biologists to understand the 25 many facets of sociality, from mechanistic processes to ecological and evolutionary 26 27 functions (Cantor et al.; Hasenjager, Leadbeater & Hoppitt; Sosa, Sueur & Puga-28 Gonzalez). For example, Bejder, Fletcher and Bräger (1998) proposed advanced permutation techniques to examine spatial associations between individuals, and 29 30 Croft, James and Krause (2008) together with Whitehead (2008) aggregated most of 31 our knowledge on the study of animal sociality in their seminal books. The growth in 32 the popularity of social network analysis in animal behaviour and ecology has been favoured by methodological advances (Whitehead 2008), and automated monitoring 33 34 techniques (Smith & Pinter-Wollman) have also played a significant role by scaling up 35 research questions to new and previously intractable species and systems. Using 36 these tools for data collection and analysis has further engaged a lively community of 37 researchers that, together, have contributed a near constant refinement and evolution 38 of social network analysis methods and its application to animals.

The Joint Special Feature in *Methods in Ecology and Evolution* and the *Journal* 39 of Animal Ecology is a celebration of research by animal social network scientists, 40 41 introducing novel methods and questions pertaining to Animal Social Network Analysis 42 (ASNA). It brings together research that highlights developments in computational methods, novel considerations about bias in ASNA, advances in the study of intrinsic 43 and extrinsic factors shaping individuals and group social structure and how animal 44 networks vary under different social environments. In doing so, we believe this Special 45 Feature offers exciting directions for future research to better cope with the complexity 46 47 of animal social structures.

48

49 **Controlling for biases in animal social network analysis**

Few wild biological data and analytical methods are immune to biases inflicted by 50 51 specifics of sampling protocols or study organism. Consequently, an important challenge for ASNA studies is the need to consider the robustness of current 52 methodological approaches. In this joint Special Feature, Sosa, Sueur and Puga-53 54 Gonzalez (consider network measures and their variants, and highlight the necessity for future research to state the variant used as this may skew the interpretation of the 55 results. Once a network measure is selected, taking care to avoid multiple hypothesis 56 testing (Webber, Schneider & Vander Wal 2020), it is essential to ensure the reliability 57 of statistical tests. Puga-Gonzalez, Sueur and Sosa outline how limitations of current 58 59 permutation approaches extend to different data sampling scenarios, and Franks et al. offer solutions to avoid spurious results that call for greater emphasis on effect sizes 60 as a means of making reliable statistical inference. Together, these three 61 62 methodological studies underline the continued need for caution when developing analytical solutions and interpreting ASNA results. ASNA is continuously being refined 63 64 and improved (Franks et al.; Weiss et al.), but it will take continued efforts to keep improving existing solutions and to address the multitude of remaining issues, such as 65 the appropriateness of different network measures, the rates of type I and type II errors 66 in standard null hypothesis-testing procedures, and finding robust solutions to 67 68 measuring effect sizes.

With these considerations in mind, new methods continue to be presented, 69 70 allowing researchers to address novel challenges in ASNA such as accurate estimates of social trait heritability (Radersma), social drivers of animal movement (Milner, 71 72 Blackwell & Niu) and to apply techniques developed for data collected using markrelease-recapture data (Silk et al.) and bio-logging methods (Gilbertson, White & Craft; 73 Gomes, Boogert & Cardoso). The miniaturization of bio-logging devices now enable 74 the study of a wider variety of organisms, from insects to cetaceans (Börger et al. 75 76 2020). Bio-logging provides access to new data sources, using less invasive methods 77 and continuous collection, considerably expanding our knowledge of how animals and groups behave in the wild (Smith & Pinter-Wollman). These advances, however, have 78 79 led to a substantial methodological shift in network construction and analysis (Godfrey et al. 2014; Spiegel et al. 2016). One specific challenge that Gomes et al. tackle is how 80 to determine cut-offs while delimiting social associations and, thereby, providing a 81 standard procedure for building social networks from proximity-based association data 82 collected through radio frequency identification detection. 83

84

85 Intrinsic and extrinsic factors shaping animal networks

Research in ASNA has focused on how intrinsic and extrinsic factors shape individual 86 social traits and their consequences on fitness (Silk et al. 2009; Silk et al. 2010; Brent, 87 Ruiz-Lambides & Platt 2017), and species social diversity (Balasubramaniam et al. 88 2017). Studies have highlighted a broad range of intrinsic factors that shape 89 heterogeneity in individual social traits such as age (Almeling et al. 2016), sex 90 (Borgeaud et al. 2017), kinship (Hirsch, Stanton & Maldonado 2012), personality 91 (Krause, James & Croft 2010), pathogens (Romano et al. 2016) etc. (see Cantor et al. 92 & Sosa, Sueur and Puga-Gonzalez for an overview). Here, Brandl et al. provide a good 93 example of how these intrinsic factors might interlink by showing how synchrony in 94 95 reproductive timing can shape present and future individual social bonds.

Although ecologists and evolutionary biologists have extensively explored the effects of variation in the physical environment on the development and fitness of individuals and their consequences on population dynamics, these aspects remain overlooked in the study of animal societies (Spiegel *et al.* 2017; Webber & Vander Wal 2018), but see (Moscovice, Sueur & Aureli 2020). Space use is often implicated as strongly influencing social dynamics, as this constrains the frequency or time with which individuals can interact or be associated (Webber & Vander Wal 2018). As a result, space use represents an important dimension for many social processes, such
 as disease spread (Albery *et al.*). A particular focus of this joint Special Feature is on
 the study of how habitat constraints or spatial ecology can shape group social structure
 through individual movements (Albery *et al.;* Milner, Blackwell & Niu; Pasquaretta *et al.*).

In addition, few studies have investigated the multivariable aspects of individual 108 sociality (Brandell et al.) and movements (Milner, Blackwell & Niu). Here, Brandell et 109 al. determine the relative influence of environmental factors, biotic interactions, 110 infectious disease and group composition on group spatial networks in two social 111 species of carnivores. Such a multivariable approach is of considerable interest to 112 understand the relative influence of habitat structure and/or social factors in shaping 113 social structure and to better understand their dynamics, host pathogen dynamics, or 114 115 species-species assemblages (Massol et al.). The new methods presented in this joint Special Feature (Albery et al.; Massol et al.; Milner, Blackwell & Niu; Pasquaretta et 116 117 al.) will help better understand such complex related processes.

118

119 Animal networks under different environments

Social structure represents the most plastic aspect of animal societies as individuals 120 can, through social interactions, regulate conflicts (Aureli & de Waal 2000), create 121 affiliative bonds (De Waal & Roosmalen 1979), cooperate (Seyfarth & Cheney 2012), 122 transmit information and learn (Hoppitt & Laland 2013). This allows them to cope with 123 ecological constraints specific to their living environment. There is growing evidence 124 125 for a dynamic eco-evolutionary feedback between the (social) environment and social structure (Cantor et al.; Smolla & Akçay 2019; Sueur et al. 2019; Romano, Macintosh 126 & Sueur 2020; Udiani & Fefferman 2020) as individuals that better adjust their 127 behaviour in response to the challenges, both external to and inherent in social 128 relationships, within the context of their own dynamic social networks, might increase 129 130 their own fitness (Romano, Macintosh & Sueur 2020).

131

Research in ASNA is increasingly integrating the concept of social plasticity, the degree to which an individual varies its social behaviour dependent upon external factors. This is important not only for exploring the mechanisms driving individual heterogeneity in sociality but also for understanding how plasticity manifests at both the individual and the group level (Ilany & Akçay 2016; Montiglio, McGlothlin & Farine

2018). The study of social dynamics, such as how individuals sociality changes in 137 response to demographic changes (Borgeaud *et al.* 2017), is made possible thanks in 138 part to the application of time-aggregated network analysis (Hobson, Avery & Wright 139 2013) for which specialised analysis packages exist now (Bonnell & Vilette; Sosa et al. 140 2020). Several studies in this joint Special Feature explore these aspects by 141 addressing, for example, how mechanistic factors allow animals to cope with 142 demographic changes (Farine), how networks are shaped by group phenotypic 143 composition (Dakin et al.) and environmental conditions (Burns et al.), and how inter-144 145 group encounters shape overall network structure (Preston et al.). Together, these studies highlight that individual sociality is not the only plastic trait but that there are 146 147 also numerous extrinsic and intrinsic constraints that drive group social structure dynamism. 148

Similarly, while past research has established that sociality affects fitness (Silk *et al.* 2009; Silk *et al.* 2010; Brent, Ruiz-Lambides & Platt 2017), Formica *et al.* show that this effect may be condition-dependent whilst Turner *et al.* show that different types of social interactions have different effects on individual fitness according to ontogenetic stages. Such dynamics in individual sociality, group structure as well as condition-dependent effects of sociality on fitness may help better explain evolutionary processes such as population structure and gene flow dynamics (Zonana *et al.*).

156

157 Anthropogenic impacts on animal societies.

At the heart of understanding the intrinsic link between habitat, movement and social 158 159 behaviour is the pressing acknowledgement that animals are increasingly inhabiting a 160 world impacted by anthropogenic-driven disturbances. As so, it is important that we 161 understand the response of social animals to such change (Bond et al.). The major question of how social structure is shaped by the broader environment represents a 162 substantial challenge to address. Do networks change in response to environmental 163 164 changes in a consistent way? For example, Bond et al. test how the social structure of communities of giraffes are impacted by human disturbance, revealing a signature of 165 166 network structure that matches those detected in small captive groups of birds exposed 167 to social disturbances (Maldonado-Chaparro et al. 2018).

As humans and wildlife increasingly share space, new methods that integrate biologging technologies (Gilbertson, White & Craft; Gomes, Boogert & Cardoso) and/or that use citizen scientists to collect and record data on grouping animals (Aplin *et al.*), will no doubt prove invaluable for anticipating how and to what extent animal groups and populations will be affected by future environmental changes during the Anthropocene. In doing so researchers will also need to consider carefully how these approaches can be optimised to reduce our impact on animal welfare. Thus, future adaptations in ASNA methods must also be mindful to align with a changing ethical landscape (Soulsbury *et al.* 2020).

177

178 Networks beyond social interactions: cascading effects across levels of179 organisation

180 We now have access to unprecedented amounts of data including, for example, 181 bio-logged individual physiological measurements, remotely sensed environmental variables or heterospecific space use and behaviour. Importantly, these data span 182 183 intrinsic (e.g. physiological, ontogenesis) and extrinsic (e.g. habitat, climate, parasites, other species and/or groups) factors, and their variation over time. Contributions in this 184 185 Joint Special Feature provide insight into these developments that are sharpening the picture of the various evolutionary processes involved in shaping social heterogeneity 186 and plasticity. The Figure 1 provides a comprehensive view of the different levels of 187 organisation embedded in the study of animal social networks, i.e. individuals, social 188 structure, habitat, environment, and their dynamics. 189

We hope that this will serve as a basis for novel questions and for identifying 190 the methodological challenges to come in order to determine the relative influence of 191 each of these factors by integrating interaction networks across levels of organisation 192 (Brandell et al.; Jacoby & Freeman 2016; Sueur et al. 2019). A particularly promising 193 194 line of research is the development of multi-layered networks (Silk et al. 2018; Mourier, Ledee & Jacoby 2019; Fisher & Pinter-Wollman 2020), hierarchically embedded 195 196 interaction networks (Montiglio et al. 2020), or bipartite networks (Massol et al.) that can expand exploration of interactions beyond social groups, spanning from cells to 197 198 whole ecosystems, and their dynamics. For instance, Massol et al. used bipartite networks to analyse the structure of host-microbiota interaction networks. The method 199 200 Massol et al. considers and assesses multiple drivers of network structures across 201 species. Importantly, this approach can be adapted to study other aspects of 202 connectivity in networks. For example, it could be used to study the multifactorial drivers embedding social and spatial networks of pollinators, spatial patterns of pollen 203 204 flow, and reproductive networks of plants, potentially revolutionizing research in pollination ecology through the application of predictive models (Pasquaretta *et al.*206 2019).

207

208 Concluding remarks

Considerable progress has been made since the first application of social network 209 analysis to animals (Sade 1972) and the first specific ASNA approaches were 210 developed (Bejder, Fletcher & Bräger 1998). This joint Special Feature is reflective of 211 the milestones reached in the past few years in ASNA, presenting new methods that 212 213 redefine analytical standards and covering topics as diverse as social transmission, 214 epidemiology, quantitative genetics, social structure plasticity, fitness consequences, 215 habitat use and applied conservation. Although social network analysis enables to 216 examine many facets of social phenomena, there is still much to be done to build 217 bridges across disciplines. As the research tools and questions developed for animal social network analysis strengthen, they will also have the potential inform pressing 218 219 global events (Firth et al. 2020). Opening up a dialog across disciplines will be particularly important in the development and application of holistic frameworks that 220 221 embed social interactions, space and time, to address topics such as the diversity (Sah, Méndez & Bansal 2019) and complexity (Kappeler et al. 2019) of animal societies 222 and heterospecific assemblages (Farine, Garroway & Sheldon 2012). 223

224

225 Acknowledgements

We are grateful to all editors of MEE and JAE for managing manuscripts submitted to this special feature. We also appreciate the many suggestions by Damien Farine and Eric Vander Wal. We finally thank all the authors who participated to this call.




Figure 1. ASNA synthetic framework for integrating social, spatial and temporal network features in the study of animal social networks. Below we highlight some questions that could be addressed by this approach.

1. Which, how and why extrinsic biotic (e.g. resource dispersion, pathogens) and abiotic factors (e.g. habitat heterogeneity, entropic

disturbances) affect social behaviour, group-level dynamics and population-level social structure? For example, climatic factors that

shape habitats through plant phenology and resource dispersion may affect animal social structure and habitat use (Albery et al.;

Brandell *et al.;* Burns *et al.;* Gilbertson, White & Craft; Milner, Blackwell & Niu; Pasquaretta *et al.*).

2. Which, how and why intrinsic factors (*e.g.* social organization, individuals' development, physiological markers) shape individualgroup- and/or population-level social structure? Variation in biotic and abiotic factors can affect physiological markers, propensity to cooperate, propensity to disperse in individuals, which may, in turn, affect sociality and group structure (Brandl *et al.;* Dakin *et al.;* Turner *et al.*).

3. How and why extrinsic and intrinsic factors can drive interactions between groups (*e.g.* group size, group encounters), species (*e.g.* host-pathogen, plant-pollinator), and ecosystem assemblages? The impact of biotic and abiotic factors on sociality may affect group-group encounters and competition for resources, and shape species-species assemblages and ecosystem structures (Massol *et al.;* Preston *et al.*).

4. Which, how and why extrinsic and intrinsic factors drive individual- and group-level social plasticity, inter-group social structure, and ecosystem dynamics (*e.g.* spatial network variation across seasons)? Considering the hypothesis in points 1, 2, 3, we may expect inter- and intra-group differences according to seasonality or social organisation of species [e.g. (*Brandell et al.*)].

5. Which, how and why are the consequences of such social plasticity on developmental, life history traits, ecological genetics, evolutionary biology, populations and community ecology? Considering influences determining sociality and plasticity, as well as condition-dependent effects of sociality on fitness (Formica *et al.;* Turner *et al.*), facilitates a better understanding of evolution processes such as group/population structures, sexual selection and gene flow (Zonana *et al.*). 6. Points 1 to 5 focus on the drivers shaping sociality in the same time point and space of the observed sociality. However, a temporal
analysis of how past individuals' sociality, extrinsic and intrinsic factors may shape current individuals' sociality (Farine), space use
Albery *et al.* (), groups and populations social structures, and species-species assemblages is needed to better understand global
system dynamics.

255 **REFERENCES**

- Albery, G.F., Kirkpatrick, L., Firth, J.A. & Bansal, S. Unifying spatial and social network analysis
 in disease ecology. *Journal of Animal Ecology*, n/a.
- Almeling, L., Hammerschmidt, K., Sennhenn-Reulen, H., Freund, A.M. & Fischer, J. (2016)
 Motivational shifts in aging monkeys and the origins of social selectivity. *Current Biology*, 26, 1744-1749.
- Aplin, L.M., Major, R.E., Davis, A. & Martin, J.M. A citizen science approach reveals long-term
 social network structure in an urban parrot, Cacatua galerita. *Journal of Animal Ecology*, n/a.
 Aureli, F. & de Waal, F.B.M. (2000) *Natural conflict resolution*. University of California Press.
- Aureli, F. & de Waal, F.B.M. (2000) *Natural conflict resolution*. University of California Press.
- Balasubramaniam, K.N., Beisner, B.A., Berman, C.M., De Marco, A., Duboscq, J., Koirala, S.,
 Majolo, B., MacIntosh, A.J., McFarland, R. & Molesti, S. (2017) The influence of
 phylogeny, social style, and sociodemographic factors on macaque social network structure.
 American journal of primatology.
- Bejder, L., Fletcher, D. & Bräger, S. (1998) A method for testing association patterns of social
 animals. *Animal Behaviour*, 56, 719-725.
- Bonnell, T.R. & Vilette, C. Constructing and analysing time-aggregated networks: The role of
 bootstrapping, permutation and simulation. *Methods in Ecology and Evolution*, n/a.
- Borgeaud, C., Sosa, S., Sueur, C. & Bshary, R. (2017) The influence of demographic variation on
 social network stability in wild vervet monkeys. *Animal Behaviour*, **134**, 155-165.
- Börger, L., Bijleveld, A.I., Fayet, A.L., Machovsky-Capuska, G.E., Patrick, S.C., Street, G.M. &
 Vander Wal, E. (2020) Biologging special feature. *Journal of Animal Ecology*, 89, 6-15.
- Brandell, E.E., Fountain-Jones, N.M., Gilbertson, M.L.J., Cross, P.C., Hudson, P.J., Smith, D.W.,
 Stahler, D.R., Packer, C. & Craft, M.E. Group density, disease, and season shape territory
 size and overlap of social carnivores. *Journal of Animal Ecology*, n/a.
- Brandl, H.B., Griffith, S.C., Farine, D.R. & Schuett, W. Wild zebra finches that nest synchronously
 have long-term stable social ties. *Journal of Animal Ecology*, n/a.
- Brent, L.J., Ruiz-Lambides, A. & Platt, M.L. (2017) Family network size and survival across the
 lifespan of female macaques. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20170515.
- Brent, L.J.N., Lehmann, J. & Ramos-Fernández, G. (2011) Social network analysis in the study of
 nonhuman primates: A historical perspective. *American journal of primatology*, 73, 720-730.
- Burns, D.D.R., Franks, D.W., Parr, C. & Robinson, E.J.H. Ant colony nest networks adapt to
 resource disruption. *Journal of Animal Ecology*, n/a.
- Cantor, M., Maldonado-Chaparro, A.A., Beck, K.B., Brandl, H.B., Carter, G.G., He, P., Hillemann,
 F., Klarevas-Irby, J.A., Ogino, M., Papageorgiou, D., Prox, L. & Farine, D.R. The
 importance of individual-to-society feedbacks in animal ecology and evolution. *Journal of Animal Ecology*, n/a.
- 291 Croft, D., James, R. & Krause, J. (2008) Exploring animal social networks. Princeton University Press.
- Dakin, R., Moore, I.T., Horton, B.M., Vernasco, B.J. & Ryder, T.B. Testosterone-mediated
 behaviour shapes the emergent properties of social networks. *Journal of Animal Ecology*, n/a.
- De Waal, F.M. & Roosmalen, A. (1979) Reconciliation and consolation among chimpanzees.
 Behavioral Ecology and Sociobiology, 5, 55-66.
- Farine, D.R. Structural trade-offs can predict rewiring in shrinking social networks. *Journal of Animal Ecology*, n/a.
- Farine, D.R., Garroway, C.J. & Sheldon, B.C. (2012) Social network analysis of mixed-species
 flocks: exploring the structure and evolution of interspecific social behaviour. *Animal Behaviour*, 84, 1271-1277.
- Firth, J.A., Hellewell, J., Klepac, P., Kissler, S., Kucharski, A.J. & Spurgin, L.G. (2020) Using a real world network to model localized COVID-19 control strategies. *Nature medicine*, 1-7.
- Fisher, D.N. & Pinter-Wollman, N. (2020) Using multilayer network analysis to explore the
 temporal dynamics of collective behaviour. *Current zoology*.

- Formica, V., Donald, H., Marti, H., Irgebay, Z. & Brodie III, E. Social network position
 experiences more variable selection than weaponry in wild subpopulations of forked fungus
 beetles. *Journal of Animal Ecology*, n/a.
- Franks, D.W., Weiss, M.N., Silk, M.J., Perryman, R.J.Y. & Croft, D.P. Calculating effect sizes in animal social network analysis. *Methods in Ecology and Evolution*, n/a.
- Gilbertson, M.L.J., White, L.A. & Craft, M.E. Trade-offs with telemetry-derived contact networks
 for infectious disease studies in wildlife. *Methods in Ecology and Evolution*, n/a.
- Godfrey, S.S., Ansari, T.H., Gardner, M.G., Farine, D.R. & Bull, C.M. (2014) A contact-based
 social network of lizards is defined by low genetic relatedness among strongly connected
 individuals. *Animal Behaviour*, 97, 35-43.
- Gomes, A.C.R., Boogert, N.J. & Cardoso, G.C. Network structure and the optimization of
 proximity-based association criteria. *Methods in Ecology and Evolution*, n/a.
- Hasenjager, M.J., Leadbeater, E. & Hoppitt, W. Detecting and quantifying social transmission using
 network-based diffusion analysis. *Journal of Animal Ecology*, n/a.
- 319 Hinde, R.A. (1976) Interactions, relationships and social structure. Man, 1-17.
- Hirsch, B.T., Stanton, M.A. & Maldonado, J.E. (2012) Kinship shapes affiliative social networks
 but not aggression in ring-tailed coatis. *PloS one*, 7.
- Hobson, E.A., Avery, M.L. & Wright, T.F. (2013) An analytical framework for quantifying and
 testing patterns of temporal dynamics in social networks. *Animal Behaviour*, 85, 83-96.
- Hoppitt, W. & Laland, K.N. (2013) Social learning: an introduction to mechanisms, methods, and models.
 Princeton University Press.
- Ilany, A. & Akçay, E. (2016) Personality and social networks: A generative model approach.
 Integrative and comparative biology, 56, 1197-1205.
- Jacoby, D.M. & Freeman, R. (2016) Emerging network-based tools in movement ecology. *Trends in ecology & evolution*, **31**, 301-314.
- Kappeler, P.M., Clutton-Brock, T., Shultz, S. & Lukas, D. (2019) Social complexity: patterns,
 processes, and evolution. Springer.
- Krause, J., James, R. & Croft, D. (2010) Personality in the context of social networks. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 4099-4106.
- Maldonado-Chaparro, A.A., Alarcón-Nieto, G., Klarevas-Irby, J.A. & Farine, D.R. (2018)
 Experimental disturbances reveal group-level costs of social instability. *Proceedings of the Royal Society B*, 285, 20181577.
- Massol, F., Macke, E., Callens, M. & Decaestecker, E. A methodological framework to analyse
 determinants of host-microbiota networks, with an application to the relationships
 between Daphnia magna's gut microbiota and bacterioplankton. *Journal of Animal Ecology*,
 n/a.
- Milner, J.E., Blackwell, P.G. & Niu, M. Modelling and inference for the movement of interacting
 animals. *Methods in Ecology and Evolution*, n/a.
- Montiglio, P.-O., Gotanda, K.M., Kratochwil, C.F., Laskowski, K.L. & Farine, D.R. (2020)
 Hierarchically embedded interaction networks represent a missing link in the study of
 behavioral and community ecology. *Behavioral Ecology*, **31**, 279-286.
- Montiglio, P.O., McGlothlin, J.W. & Farine, D.R. (2018) Social structure modulates the
 evolutionary consequences of social plasticity: A social network perspective on interacting
 phenotypes. *Ecology and evolution*, 8, 1451-1464.
- Moscovice, L.R., Sueur, C. & Aureli, F. (2020) How socio-ecological factors influence the
 differentiation of social relationships: an integrated conceptual framework. *Biology Letters*,
 16, 20200384.
- Mourier, J., Ledee, E.J. & Jacoby, D.M. (2019) A multilayer perspective for inferring spatial and
 social functioning in animal movement networks. *bioRxiv*, 749085.

- Pasquaretta, C., Dubois, T., Gomez-Moracho, T., Delepoulle, V.P., Le Loc'h, G., Heeb, P. &
 Lihoreau, M. Analysis of temporal patterns in animal movement networks. *Methods in Ecology and Evolution*, n/a.
- Pasquaretta, C., Jeanson, R., Pansanel, J., Raine, N.E., Chittka, L. & Lihoreau, M. (2019) A spatial
 network analysis of resource partitioning between bumblebees foraging on artificial flowers
 in a flight cage. *Movement ecology*, 7, 1-10.
- Preston, E.F.R., Thompson, F.J., Ellis, S., Kyambulima, S., Croft, D.P. & Cant, M.A. Network level consequences of outgroup threats in banded mongooses: Grooming and aggression
 between the sexes. *Journal of Animal Ecology*, n/a.
- Puga-Gonzalez, I., Sueur, C. & Sosa, S. Null models for animal social network analysis and data
 collected via focal sampling: Pre-network or node network permutation? *Methods in Ecology and Evolution*, n/a.
- Radersma, R. Estimating heritability of social phenotypes from social networks. *Methods in Ecology and Evolution*, n/a.
- Romano, V., Duboscq, J., Sarabian, C., Thomas, E., Sueur, C. & MacIntosh, A.J. (2016) Modeling
 infection transmission in primate networks to predict centrality-based risk. *American journal* of primatology.
- Romano, V., Macintosh, A.J. & Sueur, C. (2020) Stemming the flow: Information, infection, and
 social evolution. *Trends in Ecology & Evolution*.
- Sade, D. (1972) Sociometrics of *Macaca mulatta* Linkages and cliques in grooming matrices. *Folia Primatologica*, 18, 196-223.
- Sah, P., Méndez, J.D. & Bansal, S. (2019) A multi-species repository of social networks. *Scientific data*, 6, 1-6.
- Seyfarth, R.M. & Cheney, D.L. (2012) The evolutionary origins of friendship. *Annual review of psychology*, 63, 153-177.
- Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig, R.M.,
 Seyfarth, R.M. & Cheney, D.L. (2009) The benefits of social capital: close social bonds
 among female baboons enhance offspring survival. *Proceedings of the Royal Society B: Biological Sciences*, 276, 3099-3104.
- Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig, R.M.,
 Seyfarth, R.M. & Cheney, D.L. (2010) Strong and consistent social bonds enhance the
 longevity of female baboons. *Current Biology*, 20, 1359-1361.
- Silk, M.J., Finn, K.R., Porter, M.A. & Pinter-Wollman, N. (2018) Can multilayer networks advance
 animal behavior research? *Trends in ecology & evolution*, 33, 376-378.
- Silk, M.J., McDonald, R.A., Delahay, R.J., Padfield, D. & Hodgson, D.J. CMRnet: An R package
 to derive networks of social interactions and movement from mark-recapture data. *Methods in Ecology and Evolution*, n/a.
- Smith, J.E. & Pinter-Wollman, N. Observing the unwatchable: Integrating automated sensing,
 naturalistic observations, and animal social network analysis in the age of big data. *Journal* of Animal Ecology.
- Smolla, M. & Akçay, E. (2019) Cultural selection shapes network structure. *Science advances*, 5,
 eaaw0609.
- Sosa, S., Puga-Gonzalez, I., Hu, F., Pansanel, J., Xie, X. & Sueur, C. (2020) A multilevel statistical
 toolkit to study animal social networks: the Animal Network Toolkit Software (ANTs) R
 package. *Scientific Reports*, 10, 1-8.
- Sosa, S., Sueur, C. & Puga-Gonzalez, I. Network measures in animal social network analysis: Their
 strengths, limits, interpretations and uses. *Methods in Ecology and Evolution*, n/a.
- Soulsbury, C., Gray, H., Smith, L., Braithwaite, V., Cotter, S., Elwood, R.W., Wilkinson, A. &
 Collins, L.M. (2020) The welfare and ethics of research involving wild animals: A primer. *Methods in Ecology and Evolution.*

- 404 Spiegel, O., Leu, S.T., Bull, C.M. & Sih, A. (2017) What's your move? Movement as a link between 405 personality and spatial dynamics in animal populations. *Ecology Letters*, **20**, 3-18.
- Spiegel, O., Leu, S.T., Sih, A. & Bull, C.M. (2016) Socially interacting or indifferent neighbours?
 Randomization of movement paths to tease apart social preference and spatial constraints. *Methods in Ecology and Evolution*, 7, 971-979.
- Sueur, C., Romano, V., Sosa, S. & Puga-Gonzalez, I. (2019) Mechanisms of network evolution: a
 focus on socioecological factors, intermediary mechanisms, and selection pressures.
 Primates, 60, 167-181.
- Turner, J.W., Robitaille, A.L., Bills, P.S. & Holekamp, K.E. Early-life relationships matter: Social
 position during early life predicts fitness among female spotted hyenas. *Journal of Animal Ecology*, n/a.
- Udiani, O. & Fefferman, N.H. (2020) How disease constrains the evolution of social systems.
 Proceedings of the Royal Society B, 287, 20201284.
- Webber, Q.M., Schneider, D.C. & Vander Wal, E. (2020) Is less more? A commentary on the
 practice of 'metric hacking'in animal social network analysis. *Animal Behaviour*, 168, 109120.
- 420 Webber, Q.M. & Vander Wal, E. (2018) An evolutionary framework outlining the integration of 421 individual social and spatial ecology. *Journal of Animal Ecology*, **87**, 113-127.
- Weiss, M.N., Franks, D.W., Brent, L.J.N., Ellis, S., Silk, M.J. & Croft, D.P. Common datastream
 permutations of animal social network data are not appropriate for hypothesis testing using
 regression models. *Methods in Ecology and Evolution*, n/a.
- Whitehead, H. (2008) Analyzing animal societies: quantitative methods for vertebrate social analysis. University
 of Chicago Press.
- 427 Wilson, E.O. (2000) Sociobiology: The new synthesis. Harvard University Press.
- Zonana, D.M., Gee, J.M., Breed, M.D. & Doak, D.F. Dynamic shifts in social network structure
 and composition within a breeding hybrid population. *Journal of Animal Ecology*, n/a.

430