



HAL
open science

On the impact of grassland management on soil carbon stocks: a worldwide meta-analysis

K. Phukubye, M. Mutema, N. Buthelezi, P. Muchaonyerwa, C. Cerri, Vincent Chaplot

► **To cite this version:**

K. Phukubye, M. Mutema, N. Buthelezi, P. Muchaonyerwa, C. Cerri, et al.. On the impact of grassland management on soil carbon stocks: a worldwide meta-analysis. *Geoderma Régional*, 2022, 28, pp.e00479. 10.1016/j.geodrs.2021.e00479 . hal-03608182

HAL Id: hal-03608182

<https://hal.science/hal-03608182>

Submitted on 8 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **On the impact of grassland management on soil carbon stocks: a worldwide meta-**
2 **analysis**

3 K Phukubye¹, M Mutema², N Buthelezi¹, P Muchaonyerwa¹, C Cerri³, V Chaplot^{1, 4*}

4

5 ¹School of Agricultural, Earth and Environmental Sciences, University of KwaZulu-Natal,
6 Private Bag X01, Scottsville 3209, Pietermaritzburg, Republic of South Africa;

7 ²Agricultural Research Council-Agricultural Engineering, Private Bag X519 Silverton 0127,
8 Pretoria, Republic of South Africa;

9 ³Unversidade de São Paulo, Piracicaba, Brazil;

10 ⁴Institut de Recherche pour le Développement (IRD), Laboratoire d'Océanographie et du
11 Climat: Expérimentations et approches numériques, UMR IRD-CNRS-MNHN-UPMC 7159,
12 5 place Jussieu, 75252 PARIS Cedex 05, France.

13 *Corresponding author: vincent.chaplot@ird.fr

14

15 **Abstract**

16 Grasslands occupy 70% of whole agricultural land and hold significant amounts of carbon, a
17 key element in the regulation of Earth's soils fertility, biomass production and climate.
18 Previous work has shown that carbon stocks of grassland soils have been largely depleted
19 worldwide due to missuse or mismanagement but that shifts in management could also
20 potentially increase soil carbon stocks and mitigate against the degradation of natural
21 ecosystems. However, the existing literature points to large discrepancies in the impact of
22 grassland management practices on soil carbon, which the present study investigated. Here we
23 considered 235 experimental sites in 18 countries across the world where shifts in grassland
24 management involved different grazing strategies (free, F vs controlled, C; high, H vs low, L
25 density grazers), grazers exclusion (E), mowing (M) and burning (B). The best performing
26 practice was controlled grazing with high density of grazers (CHG) with an average soil
27 organic carbon content (SOC_C) increase of 21% and with 100 % of the studies pointing to a
28 SOC_C increase. This was followed by E (14.9%; 60%) and FLG (13.3%; 80%). On average,
29 burning grasslands, decreases SOC_C by 9.3% but 31% of the studies pointed to an increase,
30 thus indicating discrepancies in the impact of grassland management. CLG and mowing did
31 not significantly impact SOC_C. These results also indicated that B decreased SOC_C the most
32 under moist to humid climates (-10.9% vs -1.7% under arid to semi-arid), while that E was
33 only beneficial in arid to semi-arid grasslands. Adoption of rotational high-intensity grazing in
34 place of free grazing grasslands, should be seriously considered by policy and decision
35 makers to mitigate against climate change while fostering economic and social development.

36

37 **Key words:** soil organic carbon, rehabilitation, grazing, multiple soil classes, meta-analysis

38

39 **Abbreviations**

40	B	Burning
41	BD	Soil bulk density
42	C	Carbon
43	C/N	carbon to nitrogen ratio
44	CHG	Controlled high grazing
45	CLG	Controlled low grazing
46	CV	Coefficient of variation
47	N	Sample size
48	E	Grazer exclusion
49	FLG	Free grazing with low intensity
50	FHG	Free grazing with high intensity
51	LAT	Latitude (°)
52	LONG	Longitude (°)
53	M	Mowing with residues retained
54	MAP	Mean annual precipitation (mm year ⁻¹)
55	MAT	Mean annual air temperature (°C year ⁻¹)
56	Max	Maximum
57	Min	Minimum
58	PC	Principal Component
59	PCA	Principal Component Analyses
60	Quart	Quartile
61	SEM	Standard error of mean
62	SOC _C	Change in soil organic carbon content (%)
63	SOC _S	Soil organic carbon stocks (kg C m ²)
64	SON _C	Soil organic nitrogen content (%)
65	Z	Altitude (masl)
66	ΔSOC _C	Change in soil organic carbon content (%)
67	ΔSOC _C >0	Number of studies with positive change in soil organic carbon content (%)
68	ΔSON _C	Change in soil organic nitrogen content (%)
69	ΔBD	Change in bulk density (%)
70	ΔC/N	Change in carbon to nitrogen ratio (%)
71		

72 **1 Introduction**

73 In 2015, 192 countries ratified the COP21 in Paris to keep global temperature rise this century
74 well below 2 degrees Celsius above pre-industrial levels and to pursue efforts to limit further
75 increase of the temperature. At that occasion, transfer of atmospheric carbon (C) to soils, to
76 increase the soil organic C stocks, was suggested as an effective and low cost means to meet
77 the COP21 objective (Minasny et al. 2017). Spill over benefits such as improved soil health,
78 soil fertility and reduced soil erosion will accrue to potentiate food production (Lal 2016).

79 Grasslands, which are natural ecosystems with grass as the dominant vegetation, occupy
80 about 40% of the earth's land surface area and 70% of all agricultural land (FAO, 2015).
81 Despite being mostly found under the arid to semi-arid climates of Africa, Asia and America
82 where net primary production is significantly lower than the worldwide average (Abdalla et
83 al., 2018), grassland soils store about 10% of the global soil organic C (SOC) stocks, which is
84 nearly 50% more than is stored in forests worldwide (FAO 2007). Previous work established
85 that grassland soil C stocks are sensitive to changes in land use and management. Guo and
86 Gifford (2002) reported that converting grassland to cropland is detrimental to SOC stocks
87 with a worldwide average loss of 59 %. A comprehensive meta-analysis by Dlamini et al.
88 (2016), using 628 soil profiles, revealed that grassland degradation further declines SOC
89 stocks by a worldwide average of 9 % and that this decline was associated with a loss of basal
90 grass cover. Higher average SOC stock decline (-16%) was found under dry climates (<600
91 mm of rainfall per annum) than wet climates (-8% for >1000 mm yr⁻¹ zone).

92 It was also learned from Abdalla et al. (2018) that when grazing intensity rises above carrying
93 capacity, SOC stocks decrease at an average rate of -0.9% per year, which can be attributed to
94 a net loss of grass and/or inability of frequently grazed plants to build root C reserves, thus
95 unable to sustain atmospheric C allocation to soils (Savory and Parsons, 1980). On the other
96 hand, Conant et al. (2017) reported that only 49% of worldwide studies on improved

97 grassland management practices such as low stocking rates, exclusion of grazing livestock
98 and planned rotational grazing enhanced SOC stocks, thus pointing to the absence of
99 consensus on the grassland management strategies to promote for increasing SOC stocks.
100 There is also no consensus on the impact of practices that do not involve livestock such as
101 burning, which is a common practice for increasing fodder production and quality, whilst
102 avoiding bush encroachments (Tainton, 1999). For instance, Abdalla et al. (2016) reported
103 that SOC stocks increased from 1.4 to 1.6 kg C m⁻² (i.e. by 14%) in the 0-0.2 m soil depth of a
104 long-term (70 years) annual burning trial in sub-tropical humid South Africa. However,
105 Granged et al. (2011) reported a decrease of 35% for a site in Spain, and Nardoto and
106 Bustamante (2003) reported a 13% decline for another site in Brasil. Discrepancies on the
107 impact of burning also exist within single ecosystems. For example, Snayman (2002) reported
108 a decrease of -33% while Manson et al. (2000) reported an increase of +22% in SOC stocks of
109 a grasslands subjected to burning in savannah climate of South Africa.

110 Previous studies that quantified consequences of grassland degradation (Dlamini et al., 2016),
111 conversion of forest and/or cropland to grassland, burning, fertilization and reclamation of
112 grass (Conant et al., 2001; 2018), grazing intensity (Abdalla, 2018) on grassland SOC
113 demonstrated that SOC is very sensitive to changes in management practices. The studies also
114 showed large discrepancies in the impacts of these practices on SOC changes. For instance,
115 Conant et al. (2017) reported an average SOC increase of 0.28 Mg C ha⁻¹ yr⁻¹ following
116 adoption of improved grazing but with changes in SOC stocks varying between -30% at one
117 study site and 70% at another, which points to a need to understand the reasons behind such
118 discrepancies before suggesting adoption of any given practice at large scale. Investigation of
119 inconsistencies on the effects of a given practice to enhance grassland SOC stocks thus
120 becomes an important objective. Therefore, the main purpose of the current paper was to
121 investigate discrepancies in the effects of selected grassland management practices; burning,

122 low and high intensity grazing (free or controlled), mowing and livestock exclusion, on SOC
123 stocks.

124

125 **2 Materials and methods**

126 **2.1 Database preparation**

127 A comprehensive search was performed on the Web of Science to harvest published research
128 papers on grassland management practices and their impacts on soil carbon stocks and other
129 selected soil properties such as nitrogen content, bulk density and soil water content, between
130 January and July 2017. The key words used for the search from title, abstracts and keywords
131 included “grass” AND/OR “grassland” AND/OR “veld” AND/OR “burning of grass”,
132 AND/OR “livestock enclosure” AND/OR “livestock exclusion” AND/OR “grazing, intensity”
133 AND/OR “grazing density” AND/OR “soil carbon, nitrogen stocks”. All references in the
134 papers found were also assessed in order to increase the search. The search process identified
135 342 papers. Only studies reporting on results based on grassland management trials longer
136 than three years and including soil carbon concentration and stocks in their analyses were
137 selected. Sixty-two papers met the acceptance criteria. The 62 papers reported on trials
138 performed on 235 experimental sites in 18 countries across the world (Figure 1). Table 1
139 shows a summarised version of the database. The studies had to compare management
140 practices against free grazing as reference. Different management practices were considered
141 from free grazing at high intensity (FHG) and controlled grazing at low or high intensity
142 (CLG, CHG), burning (Burn), mowing (M) and cattle exclusion (E). The intervals between
143 burning incidences varied with study site and were recorded in the same way as the duration
144 of livestock enclosure, which also varied with experimental sites.

145

146 **2.2 Data extraction**

147 *Variables of interest*

148 Data on soil organic carbon content (SOC_C), soil organic nitrogen content (SOC_N), soil
149 organic carbon stocks (SOC_S) and soil bulk density (BD) were collected from the papers.
150 Here we used data from the shallowest available soil layer where most of soil carbon lies
151 (Dlamini et al., 2015), mostly from the 0-0.05 m layer but when not available to 0.05m the
152 data came to the shallowest depth available to a maximum of 0.3m (n=15). In order to
153 compare SOC_S from different thicknesses of the top-soil layer, the data were converted to soil
154 carbon stocks per m³ of soil using the depth of soil layer considered. Moreover, the data on
155 soil organic matter content were converted to SOC_C using the 1.72 ratio (Pribyl, 2010), a
156 conversion factor that assumes organic matter contains 58% organic carbon. Data for each
157 management were subtracted from reference treatment (free grazing at low density) and
158 expressed in gram or percent difference per year to remove the differences in length of trials.
159 Finally data of carbon stocks were also reported per cubic metre of soil to improve the
160 comparisons between layers of different thicknesses. When not available in the research
161 papers, data of BD were estimated using values of SOC_C and SOC_S. The missing SOC_C or
162 SOC_S values were estimated in a similar way, and when not possible the data points had no
163 information (see table 5 for number of data points).

164 Here we considered the treatment means, the sample sizes (i.e. number of replications), and
165 the standard deviation (SD) between replications. When not directly available these
166 parameters were either estimated from the replicates or recalculated such as for SD that was in
167 some cases obtained from standard error data [SE = SD (n^{-1/2})]. In case of the results were
168 only presented in the format of figures, which occurred for two studies, the data were
169 extracted through graph digitisation (use of “GetData Graph Digitizer 2.26”)

170

171 ***Secondary variables***

172 The environmental factors were related to site location (LONG: longitude, LAT: latitude and
173 Z: altitude), climate (MAP: mean annual precipitation, and MAT: mean annual air
174 temperature), thickness of the entire soil profile and soil texture (clay, silt and sand content) of
175 the upper layer. These variables were stratified into 3 classes (low, medium and high, Table
176 3). The soil order (WRB, 2016) at each data point was considered and the climates were
177 defined following Mathew et al. (2017).

178

179 **2.3 Statistical analysis**

180 The first step of the statistical analysis was to calculate descriptive statistics for the selected
181 variables (Tables 4 and 5). Meta-analysis was then applied to the data to quantify the
182 significance of the impact of the grassland management techniques under study on SOC_s. The
183 MetaEasy software (v1.0.5, Kontopantelis and Reeves 2009) was used to compute effect sizes
184 (a quantitative measure of the magnitude of the relationship between two variables, the larger
185 the effect size the stronger the correlation between the two variables) and standard errors (SE)
186 from each technique. MetaEasy allows the standardisation of effect size according to eight
187 possible methods based on the sample size and available information, and ultimately selects
188 the method that minimises the biases (Kontopantelis and Reeves 2009). Bootstrap resampling
189 was used to assign confidence intervals (CI) at 95%.

190 The multiple correlations between grassland management techniques and soil and
191 environmental conditions were further investigated using principal component analysis (PCA)
192 (Figure 3). PCAs convert non-linearly correlated factors and variables into linear
193 combinations called principal components (Jambu, 1991) and based on the assumption of
194 normality, which was not necessarily true with the study due to the disparate nature of the
195 data.

196

197 **3 Results**

198 **3.1 Variations of soil and environmental conditions across the data set**

199 Table 4 presents the descriptive statistics depicting variability of the environmental conditions
200 across the data set. The mean annual precipitation (MAP) was 711 mm yr⁻¹ with values
201 ranging from 100 mm yr⁻¹ in China (Rong et al., 2014) to 2326 mm yr⁻¹ in Brazil (Conant et
202 al., 2017). Mean MAT was 14.0°C with data varying from -8.5°C in Canada (Donkor et al.,
203 2001) to 32°C in Ethiopia (Kassahun et al., 2012). Altitude (mean of 956 m) ranged from 4 m
204 in Brasil (Conant et al., 2017) to 3500 m in China (Wei et al., 2011). Soil texture also varied
205 greatly across sites with clay content (mean, 24.9%) varying from 3.0% in China (Pei et al.
206 2007) to 74% in Brazil (Nardoto and Bustamante, 2003).

207

208 **3.2 Variations in soil carbon and nitrogen**

209 Soil carbon and nitrogen varied greatly within the data set (Table 5). The average soil organic
210 carbon content (SOC_C) computed from all reference treatments was 2.26% with a standard
211 error mean of ±0.19% (Table 5). Values varied from 0.06% in the USA under Aridisols and
212 6% clay content (Neff et al., 2005) to 20.16% under gleyic andosols in Japan (Shimoda and
213 Takahashi, 2009). The mean soil organic nitrogen (SON_C) at the reference sites was
214 0.19±0.02 with values ranging from a minimum of 0.01% also found by Neff et al., (2005) to
215 1.17% under Mollisols with 10% clay content in Canada (Evans et al 2012). The resultant
216 average carbon to nitrogen (C/N) ratio at these sites was 11.52±1.10.

217 Soil bulk density at reference sites averaged 1.22±0.02 g cm⁻³ and soil organic carbon stocks
218 (SOC_S) was 2.02±0.08 kg C m² with values between 0.27 to 17.80 kg C m² (Table 5).

219 On average, changes in grassland management decreased BD by 0.40±0.15% yr⁻¹, enhanced
220 SOC_C by 0.71±0.52% yr⁻¹, SON_C by 1.17±0.84% yr⁻¹ and SOC_S by 0.24±0.54% yr⁻¹ (Table 5).

221 The change in SOC_C (ΔSOC_C) varied from a $-25.9\% \text{ yr}^{-1}$ under free grazing with high density
222 grazers in Argentina (Abril and Bucher 1999) to $28.9\% \text{ yr}^{-1}$ at a livestock enclosure site in
223 China (Yang et al., 2016).

224

225 **3.2 Effects of grassland management practice on soil carbon**

226 Soil carbon stocks were significantly affected by only one out of the six grassland management
227 techniques (Fig. 2A). Controlled grazing with high density grazers (CHG) significantly
228 enhanced soil carbon stocks (SOC_S) with an effect size of 0.477 and a top-soil SOC_S increase
229 by an average $5.9\% \text{ yr}^{-1}$ or $94.5 \text{ g m}^{-3} \text{ yr}^{-1}$ (Figure 2A). In contrast, under the conditions of the
230 study, the effect on SOC_S was neutral for enclosure (effect size = 0.004), mowing (0.009),
231 high density free grazing (-0.013), burning (-0.002) and controlled grazing with low density
232 grazers (-0.016).

233 Similarly, CHG was the only technique to significantly enhance soil carbon content (SOC_C)
234 with an effect size of 0.70 and a mean percentage change of $6.6\% \text{ yr}^{-1}$ (Fig. 2B). Finally, for
235 top-soil bulk density (BD) the effect of grassland management (Fig. 2C) was significant for
236 burning (effect size of 0.071; $1.08\% \text{ yr}^{-1}$ increase) and controlled grazing at low density
237 (0.074 ; $1.07\% \text{ yr}^{-1}$).

238

239 **3.3 Effect of soil and environmental conditions on the changes in soil carbon**

240 Figure 3 shows the variations in soil carbon stocks (ΔSOC_S) following the changes in
241 management for different soil orders. The statistics for Mollisols were performed using 64
242 data points, Alfisols (60), Oxisols (43), Aridisols (24), Vertisols (17), Entisols (13), Ultisols
243 (2) and with 14 data points with undetermined soil order. The median stock difference was
244 with $-10 \text{ gC m}^{-3} \text{ yr}^{-1}$ the lowest for Oxisols, followed by Aridisols and Mollisols (2.1 and 3.2),

245 Entisols (6.0), Alfisols (8.1), Vertisols (14.0) and Ultisols (85.2), all these differences being
246 not significant at $P < 0.05$ level.

247

248 As pointed out by figure 4, management duration did not have a significant impact ($P < 0.05$)
249 on the changes of soil organic carbon stocks with average carbon gains of $30.43 \text{ gC m}^{-3} \text{ y}^{-1}$ for
250 durations lower than 10 years, $5.59 \text{ gC m}^{-3} \text{ y}^{-1}$ for 10-20 years but -15.51 for 20-30 years and
251 $-0.62 \text{ gC m}^{-3} \text{ y}^{-1}$ for more than 30 years.

252 Figure 5 shows the multiple correlations between the selected variables of interest (ΔSOC_C ,
253 ΔSOC_S , ΔSON_C , $\Delta\text{C/N}$, ΔBD) on one hand, and soil and environmental factors on the other
254 hand. On figure 3A, focusing on all management techniques, the first two axes of the PCA
255 explained 94% of the data variability, with Axis 1 explaining 66% while Axis 2 accounted for
256 28%. In axis 1, ΔSOC_C , ΔSOC_S and ΔSON_C had positive coordinates while ΔBD had negative
257 coordinates, indicating a tendency for soil C and N to increase concomitantly, and to be
258 associated to a decrease in soil compaction. Figure 3A also shows a positive correlation
259 between ΔSOC_C and ΔSON_C on one hand and soil clay content on the other, meaning that
260 changes in soil C and N were the greatest at higher soil clay content ($r=0.67$, Table 6) but also
261 warmer climatic conditions ($r=0.58$ with MAT). Changes in the C/N ratio correlated the most
262 with SOC_C , pointing to an increase in C relative to N as soil carbon content increases ($r=0.62$,
263 Table 6).

264 The PCA generated from the grazing treatments only (Fig. 4B) pointed to a correlation
265 between MAT, MAP and clay content on the one hand and SOC_C , SOC_N and SOC_S in the
266 other hand, thus showing a tendency for soil C and N to increase the most under high clay,
267 temperature and rainfall conditions. From Table 7 we also learn that controlled grazing with
268 high density grazers (CHG) had a singular behaviour with the greatest SOC_S gains under arid
269 to humid and low clay conditions. Controlled high grazing (CHG) enhanced SOC_S the most
270 under moist-humid ($233 \text{ g m}^{-3} \text{ yr}^{-1}$ or $8.9 \% \text{ yr}^{-1}$) and arid to semi-arid ($15.6 \text{ g m}^{-3} \text{ yr}^{-1}$ or 1.9
271 $\% \text{ yr}^{-1}$) and as compared to wet climates ($0.5 \% \text{ yr}^{-1}$) and the increase was greater under low
272 clay conditions ($211.7 \text{ g m}^{-3} \text{ yr}^{-1}$ or $3.15 \% \text{ yr}^{-1}$ at $<20\%$ clay) (Table 7). CHG thus appeared

273 to induce highest SOC recovery under moist-humid and sandy soil conditions than under wet
274 and clayey ones.

275 In contrast, controlled low grazing (CLG) only slightly increased SOC_s under sandy and arid
276 conditions but these differences were not significant. CLG enhanced SOC_s by 3.1% yr⁻¹ under
277 the wet climate (Table 7).

278 Finally, grass burning significantly decreased SOC_s from Arid to humid climates but not for
279 wet ones (Table 8) and lessened SOC_s for all textures. The PCA of Fig. 5D showed that the
280 impact of grazer exclusion (E) on soil C and N did not correlate with climate. However,
281 Table 7 indicated SOC_s slight gains under arid to humid conditions and significant losses
282 under wet climate (-2.1% yr⁻¹).

283

284 **4 Discussion**

285 **4.1 The impact of grassland management on soil C and other selected soil properties**

286 From our dataset, controlled grazing at high density (CHG) was the only grassland
287 management technique that significantly increased top-soil organic carbon content (SOC_C)
288 and stocks (SOC_s) as compared to controlled grazing at low density (CLG), free grazing at
289 high density (FHG), grazer exclusion (E), mowing (M), and grass burning (B). CHG yielded
290 the highest soil C increase in the study by Gao et al. (2009) in China finding a 48% SOC_C rise,
291 Chaplot et al (2016), in South Africa found a 35-45% increase, Li et al (2007) in China (20%)
292 and Manley et al 1995 in the USA found 20%. From these studies we also learn that soil C
293 recovery is likely to be due to increase in soil surface coverage by grass (Chaplot et al. 2016)
294 that potentiates C inputs to the soil by plants. Greater soil coverage by grass also decreases
295 soil C erosion during rainstorms (Mchunu and Chaplot, 2012). C increase may also be due to
296 a greater proportion of N fixing grass species that helps building soil organic matter as
297 indicated by Gao et al., (2009). CHG also resulted in higher soil compaction than all reference

298 treatments, which might be attributed to soil trampling by livestock hooves. The associated
299 decrease in soil porosity lessens oxygen and CO₂ exchanges between the atmosphere and
300 soils, thus limiting organic matter decomposition and C fluxes out of soils as pointed out by
301 previous studies such as these by Nsabimana et al., (2004), Pandey et al. (2014) and Chaplot
302 et al. (2015).

303 Using data from many sites across the world Dlamini et al (2016) showed that CHG perform
304 best in terms of SOC_c under dry and sandy soil conditions, which may be another reason for
305 discrepancies. In fact, under these conditions that C losses due to degradation were also
306 highest (-16% for dry climates vs -8% for wet; -10% for sandy vs -1% for clayey) due to a
307 combination of harsh climate and low aggregate stability, which subsequently result in poor
308 soil organic matter protection. In addition to SOC_c, CHG also enhanced SON_c and increased
309 C/N ratios.

310 Grazer exclusion (E) had an insignificant impact on soil carbon stocks under arid to semi-arid
311 and moist to humid climates (Table 7) as well under coarse textured soils, but was detrimental
312 to soil C under wet climate and clayey soils. A possible explanation for the latter is likely to
313 be the low number of data points for E under wet climates (n=4) and clayey soils (n=4).

314 Additionally, E tended to enhance soil nitrogen content but its impacts on the C/N ratio and
315 on soil bulk density were insignificant. Such results can be attributed to dominance by
316 nitrogen-fixing grass species that help building soil C or the presence of C₃ or C₄ grass
317 species of different C behavior (Ritchie and Raina, 2016; Mahaney et al., 2008).

318 In general, fire had detrimental effects on SOC, irrespective of climate and soil texture
319 (except for moist-humid climate with a 3.7% increase), but the differences with free grazing
320 were not significant. The loss of soil C can be explained by high C losses in gaseous forms
321 during combustion of organic matter (Oluwole et al., 2008) and through preferential erosion
322 of black carbon formed during burning of the grass (Rumpel et al., 2006). The loss of soil C

323 lessens the stability of soil structure (Heydari et al., 2017), which in turn increases soil C
324 losses to the atmosphere as gases from organic matter decomposition and/or via erosion by
325 water. Burning can also decrease soil basal cover and thus lessen C inputs to soils. The gains
326 could be explained by removal of low productivity grass species and/or trees of lower soil C
327 input capacity.

328

329 **4.2 Recommendations for appropriate grassland management**

330 The study pointed to controlled grazing at high intensity for short duration as the best
331 performing grassland management practice worldwide with, however, greater benefits in Arid
332 to humid climates than in the wet regions of the world. This practice was first suggested by
333 Voisin (1957) following his work under temperate climate and reworked as holistic grassland
334 management by Savory and Parsons (1980) in the dry southern Africa. Controlled high
335 density grazing (CHG) proved to be more efficient than free grazing or rotational grazing for
336 forage production (higher quality and quantity) and for rehabilitation of degraded grasslands
337 where quasi-bare soils are a common feature (Savory and Parsons, 1980). In his work Voisin
338 (1957) demonstrated that grass grazed at 3-leaf stage was not only of higher nutritional value
339 but also recovered faster as enough energy was accumulated in its stem and roots. Grazing at
340 the 3-leaf stage allows more grass to be produced as flowering is prevented, which helps
341 keeping their rooting systems and their associated water and nutrient supply functions during
342 droughts for long, thus potentially explaining the higher increase in C content and stocks in
343 high intensity short duration grazing. Voisin (1957) also demonstrated in Western Europe that
344 a grazing period of two to three days every 21 days in spring during when grass growth rate is
345 faster and every 35 days afterward was optimal. This is probably due to development of
346 higher rooting systems that allocate C to soils for a longer period as exudates. The fact that
347 more forage is harvested by cows under CHG practices and further transformed into dung

348 which has high potential to become soil organic matter is another potential process for
349 building soil C.

350 This study showed the benefits of CHG were higher for dry and sandy environments where
351 vegetative growth is low because of scarce rains and high temperatures. Improving grazing
352 management in these areas is not only important to mitigate against climate change but also to
353 ensure food security in already highly vulnerable regions. However, CHG implies a drastic
354 shift in livestock management from traditional free grazing as it requires subdivisions of
355 entire grassing areas into paddocks of similar size which need to be fenced and supplied with
356 water. Since CHG requires more labour and investment for paddock construction, accurate
357 estimation of grass stages and biomass over time for grazing planning, and moving the grazers
358 from paddock to paddock the practice has mostly been applied to highly degraded grasslands.
359 Rehabilitation of degraded grasslands using livestock requires extreme care to avoid
360 unintended consequences such as increased soil bulk density and reduced water infiltration,
361 especially nearby watering points.

362 The study results further suggests the lack of impact of Soil Order on soil C recovery, which
363 was surprising, probably owing to the that different soil orders may exhibit similar soil
364 properties. Finally, fire as a method for managing grasslands did not appear to be beneficial
365 for building soil C stocks, irrespective of the soil and environmental conditions. The fact that
366 burning is less detrimental under dry and sandy conditions than under clayey and wet ones is
367 probably due to the fact that the soils are already highly C depleted.

368

369 **Conclusions**

370 The main conclusions from the current study performed using 235 grassland experimental
371 sites from 18 countries across the world were that (i) grassland soil C and N was confirmed to
372 be significantly affected by grassland management practice but that large discrepancies in

373 such effect existed from site to site; (ii) sandy soils in grasslands of arid to semi-arid and
374 moist-humid climates have greater potential for C and N recovery following changes in
375 management practices as than clayey soils of the tropics; (iii) the best performing grassland
376 management practice was controlled rotational grazing with high density grazers and short
377 duration which gave rise to average soil C stock increase of 5.9% yr⁻¹ (iv) excluding grazers
378 or burning grasslands were not beneficial for soil C with exclosure being highly detrimental
379 under wet and clayey soils. The current widespread use of these practices needs to be
380 reassessed.

381 These results improve our knowledge on the impact of grassland management on soil C and
382 the links to climate change and soil texture. Adoption of rotational high-density grazing for
383 short duration in place of free grazing grasslands, especially on degraded grasslands, should
384 be seriously considered by policy and decision makers to mitigate against climate change
385 while fostering economic and social development. This study benefitted from the existence of
386 numerous publications that could assist decision and policy makers on determining proper
387 grassland management techniques for rehabilitating grasslands across the world. However,
388 most of the studies only reported on soil carbon content, little on N content and even less on C
389 and N stocks. In addition, only 11 studies assessed the impact of controlled high density
390 grazing thus pointing to the need to coordinate research on key soil variables and on this very
391 promising practice across different environmental conditions in order to ascertain its
392 underlying mechanisms and associated ecosystem functions of food and water security. The
393 research to come should thus consider critical variables and routinely use consistent standards
394 and inform on durations of experiments.

395 Finally, the fact that none of the study techniques was truly beneficial in terms of soil C and N
396 in the wet climate points to the need to test other alternatives.

397

398 **Acknowledgements**

399 The research received financial support from the Water Research Commission of the Republic
400 of South Africa (WRC K5/2266).

401

402 **References**

- 403 Abdalla K, Chivenge P, Everson C, Mathieu O, Thevenot M, Chaplot V. 2016. Long-term
404 annual burning of grassland increases CO₂ emissions from soils. *Geoderma* 282, 80-
405 86.
- 406 Aciego-Pietri JC, Brookes PC. 2008. Relationships between soil pH and microbial properties
407 in a UK arable soil. *Soil Biology and Biochemistry* 40, 1856-1861.
- 408 Al-Seekh SH, Mohammad AG, Amro YA. 2009. Effect of grazing on soil properties at
409 southern part of West Bank Rangeland. *Hebron University Research Journal* 1, 35-53.
- 410 Amiri F, Ariapour A, Fadai S. 2008. Effects of Livestock Grazing on Vegetation Composition
411 and Soil Moisture Properties in Grazed and Non-Grazed Range Site. *Journal of*
412 *Biological Sciences* 8 (8), 1289-1297.
- 413 Andersson S, Ingvar-Nilsson S. 2001. Influence of pH and temperature on microbial activity,
414 substrate availability of soil-solution bacteria and leaching of dissolved organic C in a
415 mor humus. *Soil Biology and Biochemistry* 33, 1181-1191.
- 416 Baldock, J., and Skjemstad, J. 1999. Soil organic carbon/soil organic matter.
- 417 Barger NN, Ojima DS, Belnap J, Shiping W, Yanfen W, Chens Z. 2004. Changes in plant
418 functional groups, litter quality, and soil Carbon and Nitrogen mineralization with
419 sheep grazing in an Inner Mongolian Grassland. *Journal of Range Management* 57 (6),
420 613-619.
- 421 Blair JM. 1997. Fire, N availability and plant response in grasslands: a test of the transient
422 maxima hypothesis. *Ecology* 78, 2359-2368.
- 423 Burke IC, Yonker CM, Parton WJ, Cole CV, Flach K, Schimel DS. 1989. Texture, climate,
424 and cultivation effects on soil organic matter in US grassland soils. *Soil Science*
425 *Society of American Journal* 53, 800-805.

- 426 Cao J, Wang X, Sun X, Zhang L, Tian Y. 2013. Effects of grazing intensity on soil labile
427 organic carbon fractions in a desert steppe area in Inner Mongolia. *SpringerPlus* 2 (1),
428 doi: 10.1186/2193-1801-2-S1-S1.
- 429 Carneis Filho AC, Crusciol CA, Guimarães TM, Calonego JC, Mooney SJ. (2016). Impact of
430 amendments on the physical properties of soil under tropical long-term no till
431 conditions. *PloS one* 11, e0167564.
- 432 Chaplot V, Dlamini P, Chivenge P. 2016. Potential of grassland rehabilitation through high
433 density-short duration grazing to sequester atmospheric carbon. *Geoderma* 271, 10-17.
- 434 Chartier MP, Rostagno CM, Pazos GE. 2011. Effects of soil degradation on infiltration rates
435 in grazed semiarid rangelands of northeastern Patagonia, Argentina. *Journal of Arid*
436 *Environment*. 75, 656-661.
- 437 Chen J, Tang H. 2016. Effect of grazing exclusion on vegetation characteristics and soil
438 organic carbon of *Leymus chinensis* grassland in northern China. *Sustainability* 8, doi:
439 10.3390/su8010056.
- 440 Chidhumayo EN. 1997. Annual and spatial variation in herbaceous biomass production in a
441 Zambian dry miombo woodland. *South African Journal of Botany* 63 (2), 74-81.
- 442 Coleman DC, Wall DH. 2015. Soil Fauna: Occurrence, Biodiversity, and Roles in Ecosystem
443 Function. In: Paul EA (Ed.), *Soil Microbiology, Ecology and Biochemistry*. 4th
444 Edition, ISBN: 978-0-12-415955-6, pp 111-149.
- 445 Conant RT, Cerri CEP, Osborne BB, Paustian K. 2017. Grassland management impacts on
446 soil carbon stocks: a new synthesis. *Ecological Applications*, 27 (2), 662–668
- 447 Conant RT, Paustian K, Elliot ET. 2001 Grassland management and conversion into
448 grasslands: Effects on soil carbon. *Ecological Applications* 11, 343-355.

449 Devi TI, Yadava PS, Garkoti SC. 2014. Cattle grazing influences soil microbial biomass in
450 sub-tropical grassland ecosystems at Nambol, Manipur, northeast India. *Tropical*
451 *Ecology* 55 (2), 195-206.

452 Dlamini P, Chivenge P, Manson A, Chaplot V. 2014. Land degradation impact on soil organic
453 carbon and nitrogen stocks. *Geoderma* 235, 372-381.

454 Donkor NT, Gedir JV, Hudson RJ, Bork EW, Chanasyk DS, Naeth MA. 2002. Impacts of
455 grazing systems on soil compaction and pasture production in Alberta. *Canadian*
456 *Journal of Soil Science* 82, 1-8.

457 du Toit GV, Snyman HA, Malan PJ. 2009. Physical impact of grazing by sheep on soil
458 parameters in the Nama Karoo subshrub/grass rangeland of South Africa. *Journal of*
459 *Arid Environments* 73, 804-810.

460 Evans CRW, Krzic M, Broersma K, Thompson DJ. 2012. Long-term grazing effects on
461 grassland soil properties in southern British Columbia. *Canadian Journal of Soil*
462 *Science* 92, 685-693.

463 Everson TM, Everson CS, Zuma KD. 2007. Community based research on the influence of
464 rehabilitation techniques on the management of degraded catchments. Report to the
465 Water Research Commission

466 Feller C, Beare MH. 1997. Physical control of soil organic matter dynamics in the Tropics.
467 *Geoderma* 79; 69-116.

468 Feng X, Fu B, Lu N, Zeng Y, Wu B. 2013. How ecological restoration alters ecosystem
469 services: an analysis of carbon sequestration in China's Loess Plateau. *Scientific*
470 *Report* 3, doi: 10.1038/srep02846.

471 Gasch C, Huzurbazar S, Stahl P. 2014. Measuring soil disturbance effects and assessing soil
472 restoration success by examining distributions of soil properties. *Applied Soil Ecology*
473 76, 102-111.

474 Guo LB, Gifford RM. 2002. Soil carbon stocks and land use change. *Global Change Biology*
475 8, 345-360.

476 Harris WN, Moretto AS, Distel RA, Bouttona TW, Boo RM. 2007. Fire and grazing in
477 grasslands of the Argentine Caldenal: Effects on plant and soil carbon and nitrogen.
478 *Acta Oecologica* 32, 207-214.

479 Hassink J. 1997. The capacity of soils to preserve organic C and N by their association with
480 clay and silt particles. *Plant Soil* 191; 77-87.

481 Heydari M, Rostamy A, Najafi F, Dey D. 2017. Effect of fire severity on physical and
482 biochemical soil properties in Zagros oak (*Quercus brantii* Lindl.) forests in Iran.
483 *Journal of Forestry Research* 28; 95-104.

484 Jambu M. 1991, *Exploratory and Multivariate Data Analysis*, Academic, Orlando, Fla.

485 Janssens IA, Dieleman W, Luysaert S, Subke JA, Reichstein M, Ceulemans R, Ciais P,
486 Dolman AJ, Grace J, Matteucci G, Papale D, Piao SL, Schulze ED, Tang J, Law BE.
487 2010. Reduction of forest respiration in response to nitrogen deposition. *Nature*
488 *Geoscience* 3, 315-322.

489 Jensen ES. 1994. Mineralization-immobilization of nitrogen in soil amended with low C:N
490 ratio plant residues with different particle sizes. *Soil Biology and Biochemistry* 26,
491 519-521.

492 Jiao T, Nie Z, Zhao G, Cao W. 2016. Changes in soil physical, chemical and biological
493 characteristics of a temperate desert steppe under different grazing regimes in
494 Northern China. *Communications in Soil Science and Plant Analysis* 47, 338-347.

495 Jobbágy EG, Jackson RB. 2000. The vertical distribution of soil organic carbon and its
496 relation to climate and vegetation. *Ecological Applications* 10, 423-436.

497 Kassahun A, Tegegne A, Aberra D. 2012. Impacts of rangeland degradation on soil physical,
498 chemical and seed bank properties along a gradient in three rangeland vegetation types

499 in Somali Region, Eastern Ethiopia. *Ethiopian Journal of Agricultural Science* 22, 84-
500 101.

501 Kelly RH, Burke IC, Lauenroth WK. 1996. Soil organic matter and nutrient availability
502 responses to reduced plant inputs in short grass steppe. *Ecology* 77, 2516-2527.

503 Kgosikoma OE, Mojeremane W, Harvie B. 2015. The impact of livestock grazing
504 management systems on soil and vegetation characteristics across savanna ecosystems
505 in Botswana. *African Journal of Range and Forage Science* 32 (4), 271-278.

506 Knapp AK, Fay PA, Blair JM, Collins SL, Smith MD, Carlisle JD, Harper CW, Danner BT,
507 Lett MS, McCarron JK. 2002. Rainfall variability, carbon cycling, and plant species
508 diversity in a mesic grassland. *Science* 298, 2202-2205.

509 Köppen W. 1936. Das geographische system der klimate. In: W Köppen, R Geiger (Eds),
510 Handbuch der Klimatologie. Verlag von Gebrüder Borntraeger, Berlin, Vol 1 Part C, 1
511 - 44 pp.

512 Kontopantelis and Reeves 2009. Metaeasy: a meta-analysis add-in for microsoft excel. *J Stat*
513 *Softw* 30, 1-25.

514 Koyama A, Kubo D, Yoshihara Y, Jamsran U, Okuro T. 2015. Response of degraded
515 vegetation to introduction of prescribed burning or mowing management in a
516 Mongolian steppe. *Grassland Science* 62, 37-44.

517 Krzic M, Lamagna SF, Newman RF, Bradfield G, Wallace BM. 2014. Long-term grazing
518 effects on rough fescue grassland soils in southern British Columbia. *Canadian Journal*
519 *of Soil Science* 94, 337-345.

520 Lado M, Ben-Hur M. 2004. Soil mineralogy effects on seal formation, run-off and soil loss.
521 *Applied Clay Science* 24, 209-224.

522 Lavado RS, Sierra JO, Hashimoto PN. 1996. Impact of grazing on soil nutrients in a Pampean
523 grassland. *Journal of Range Management* 49, 452-457.

524 Li SG, Harazono Y, Oikawa T, Zhao HL, He ZY, Chang XL. 2000. Grassland desertification
525 by grazing and the resulting micrometeorological changes in Inner Mongolia.
526 *Agricultural and Forest Meteorology* 102 (2-3), 125-137.

527 Li W, Huang H-Z, Zhang Z-N, Wu G-L. 2011. Effects of grazing on the soil properties and C
528 and N storage in relation to biomass allocation in an alpine meadow. *Journal of Soil*
529 *Science and Plant Nutrition* 11 (4), 27-39.

530 Liliana IP, Gabriela Q, Fernando OG, Pedro L. 2003. Biological and chemical response of a
531 grassland soil to burning. *Journal of Range Management* 56, 291-297.

532 Liu N, Zhang Y, Chang S, Kan H, Lin L. 2012. Impact of grazing on soil carbon and
533 microbial biomass in typical steppe and desert steppe of Inner Mongolia. *PLoS One* 7
534 (5), doi: 10.1371/journal.pone.0036434.

535 Mahaney WM, Smemo KA, Gross KL. 2008. Impacts of C 4 grass introductions on soil
536 carbon and nitrogen cycling in C 3-dominated successional systems. *Oecologia* 157;
537 295-305.

538 Manson AD, Jewitt D, Short AD. 2007. Effects of season and frequency of burning on soils
539 and landscape functioning in moist montane grassland. *African Journal of Range and*
540 *Forage Science* 24, 9-18.

541 Materechera SA, Mandringana OT, Mbokodi PM, Nyamapfene K. 1998. Organic matter, pH
542 and nutrient distribution in soil layer of a savannah thornveld subjected to different
543 burning frequencies at Alice, in the Eastern Cape. *South African Journal of Plant and*
544 *Soil* 15 (3), 109–115.

545 Mathew I, Shimelis H, Mutema M, Chaplot V. 2017. What crop type for atmospheric carbon
546 sequestration: Results from a global data analysis. *Agriculture, Ecosystems and*
547 *Environment* 243, 34-46.

548 McNaughton SJ, Banyikwa FF, McNaughton MM. 1997. Promotion of the cycling of diet-
549 enhancing nutrients by African grazers. *Science* 278, 1798-1800.

550 Miller RW, Donahue RL. 1990. *An Introduction to soils and plant growth*. 6th Edition,
551 Prentice-Hall International (UK) limited, London, pp. 110- 488.

552 Monleon VJ, Cromack Jr K, Landsberg JD. 1997. Short- and long-term effects of prescribed
553 under burning on nitrogen availability in ponderosa pine stands in central Oregon.
554 *Canadian Journal of Forest Research* 27, 369-378.

555 Mutema M, Chaplot V, Jewitt G, Chivenge P, Blöschl G. 2015. Annual water, sediment,
556 nutrient, and organic carbon fluxes in river basins: A global meta-analysis as a
557 function of scale, *Water Resources Research* 51, 8949-8972.

558 Mutema M, Mafongoya PL, Chikukura L, Nyagumbo I. 2013. Effects of crop residues and
559 reduced tillage on macrofauna abundance. *Journal of Organic Systems* ISSN 1177-
560 4258.

561 Nardoto GB, Bustamante, MNCB. 2003. Effects of fire on soil nitrogen dynamics and
562 microbial biomass in savannas of Central Brazil. *Pesquisa Agropecuária Brasileira* 38,
563 955-962.

564 Neff JC, Reynolds RL, Belnap J, Lamothe P. 2005. Multi-decadal impacts of grazing on soil
565 physical and biogeochemical properties in southeast Utah. *Ecological Applications* 15
566 (1), 87-95.

567 Niu D, Hall SJ, Fu H, Kang J, Qin Y, Else JJ. 2011. Grazing exclusion alters ecosystem
568 carbon pools in Alxa desert steppe. *New Zealand Journal of Agricultural Research* 54
569 (3), 127-142

570 Nüsse A, Linsler D, Kaiser M, Ebeling D, Tonn B, Isselstein J, Ludwig B. 2017. Effect of
571 grazing intensity and soil characteristics on soil organic carbon and nitrogen stocks in
572 a temperate long-term grassland. *Archives of Agronomy and Soil Science* 1476-3567.

573 Nyamadzawo G, Wuta M, Nyamangara J, Nyamugafata P, Tendayi T. 2014. Burning,
574 biomass removal and tillage effects on soil organic carbon and nutrients in seasonal
575 wetlands (Dambos) of Chiota smallholder farming area, Zimbabwe. *Archives of*
576 *Agronomy and Soil Science* 60 (10), 1411-1427.

577 O'Brien SL, Jastrow JD, Grimley DA, Gonzalez-Meler MA. 2010. Moisture and vegetation
578 controls on decadal-scale accrual of soil organic carbon and total nitrogen in restored
579 grasslands. *Global Change Biology*, 16 (9), 2573-2588.

580 Oğuz İ, Koçyiğit R, Erşahin S. 2015. The effect of range management on soil carbon content
581 in degraded soil. *Journal of Agricultural Faculty of Gaziosmanpasa* 32 (3), 133-137.

582 Ojima DS, Schimel DS, Parton WJ, Owensby CE. 1994. Long and short-term effects of fire
583 on nitrogen cycling in tall-grass prairie. *Biogeochemistry* 24, 67-84.

584 Oluwole FA, Sambo JM, Sikhalazo D. 2008. Long-term effects of different burning
585 frequencies on the dry savannah grassland in South Africa. *African Journal of*
586 *Agricultural Research* 3, 147-153.

587 Pribyl, D.W., 2010. A critical review of the conventional SOC to SOM conversion
588 factor. *Geoderma*, 156, 75-83.

589 Ritchie ME, Raina R. 2016. Effects of herbivores on nitrogen fixation by grass endophytes,
590 legume symbionts and free-living soil surface bacteria in the Serengeti. *Pedobiologia*
591 59, 233-241.

592 Rong Y, Yuan F, Ma L. 2014. Effectiveness of exclosures for restoring soils and vegetation
593 degraded by overgrazing in the Junggar Basin, China. *Grassland Science* 60, 118-124.

594 Sajedi T, Prescott CE, Seely B, Lavkulich LM. 2012. Relationships among soil moisture,
595 aeration and plant communities in natural and harvested coniferous forests in coastal
596 British Columbia, Canada. *Journal of ecology* 100; 605-618.

597 Sanjari G, Ghadiri H, Ciesiolka CAA, Yu B. 2008. Comparing the effects of continuous and
598 time-controlled grazing systems on soil characteristics in Southeast Queensland.
599 Australian Journal of Soil Research 46, 348-358.

600 Satrio AE, Gandaseca S, Ahmed OH, Majid NMA. 2009. The influence of chemical
601 properties on soil carbon storage of a tropical peat swamp forest. American Journal of
602 Applied Science 6, 1969-1972.

603 Savory, A. and Parsons, S.D. 1980. The Savory grazing method. Rangelands 2:234–237

604 Scheffer F, Schachtschabel P. 2002. Lehrbuch der Bodenkunde. Spektrum Akademischer
605 Verlag, Heidelberg, Germany. 593 pp.

606 Schimel DS, Braswell BH, Holland EA, McKeown R, Ojima DS, Painter TH, Parton WJ,
607 Townsend AR. 1994. Climatic, edaphic, and biotic controls over storage and turnover
608 of carbon in soils. Global Biogeochemical Cycles.8, 279-293.

609 Seidlheko EM, Allen BL, Wester DB. 1993. Effect of three grazing intensities on selected soil
610 properties in semi-arid west Texas. African Journal of Range and Forage Science 10
611 (2), 82-85.

612 Shimoda1 S, Takahashi Y. 2009. Differences in soil carbon storage due to mowing, burning
613 and uncontrolled management practices of grassland at the foot of Mount Sanbe,
614 Japan. Grassland Science 55, 175-180.

615 Six J, Conant RT, Paul EA, Paustian K. 2002. Stabilization mechanisms of soil organic
616 matter: implications for C-saturation. Plant Soil 241, 155-176.

617 Snyman HA. 2002. Fire and the dynamics of a semi-arid grassland influence on soil
618 characteristics. African Journal of Range and Forage Science 19, 137-145.

619 Steffens M, Kölbl A, Totsche KU, Kögel-Knabner I. 2008. Grazing effects on soil chemical
620 and physical properties in a semiarid steppe of Inner Mongolia (P.R. China).
621 Geoderma 143:63-72.

622 Su XK, Wu Y, Dong SK, Wen L, Li YY, Wang XX. 2015. Effects of grassland degradation
623 and re-vegetation on carbon and nitrogen storage in the soils of the Headwater Area
624 Nature Reserve on the Qinghai-Tibetan Plateau, China. *Journal of Mountain
625 Science* 12 (3), 582-591.

626 Su YZ, Li LY, Zhao HL. 2006. Soil properties and their spatial pattern in a degraded sandy
627 grassland under post-grazing restoration, Inner Mongolia, northern China.
628 *Biogeochemistry* 79, 297-314.

629 Su YZ, Li YL, Cui HY, Zhao WZ. 2005. Influences of continuous grazing and livestock
630 exclusion on soil properties in a degraded sandy grassland, Inner Mongolia, northern
631 China. *Catena* 59 (3), 267-278.

632 Su YZ, Zhao HL, Zhang TH, Zhao XY. 2004. Soil properties following cultivation and non-
633 grazing of a semi-arid sandy grassland in northern China. *Soil and Tillage Research* 75
634 (1), 27-36.

635 Suttie JM, Reynolds SG, Batello C. 2005. *Grasslands of the world*. Food and Agricultural
636 organization of the United Nations, Rome, Italy.

637 Teague WR, Dowhower SL, Baker SA, Ansley RJ, Kreuter UP, Conover DM, Waggoner JA.
638 2010. Soil and herbaceous plant responses to summer patch burns under continuous
639 and rotational grazing. *Agriculture, Ecosystems and Environment* 137, 113-123.

640 Thomas RJ, Logan KAB, Ironside AD, Bolton GR. 2006. The effects of grazing with and
641 without excretal returns on the accumulation of nitrogen by ryegrass in a continuously
642 grazed upland sward. *Grass Forage Science* 45, 65-75.

643 von Lützow M, Kögel-Knabner I, Ekschmitt K, Matzner E, Guggenberger G, Marschner B,
644 Flessa H. 2006. Stabilization of organic matter in temperate soils: mechanisms and
645 their relevance under different soil conditions - a review. *European Journal of Soil
646 Science* 57, 426-445.

647 Wan SQ, Hui DF, Luo YQ. 2001. Fire effects on nitrogen pools and dynamics in terrestrial
648 ecosystems-a meta-analysis. *Ecology Application* 11, 46-52

649 Wang Z, Johnson DA, Rong Y, Wang K. 2016. Grazing effects on soil characteristics and
650 vegetation of grassland in northern China. *Solid Earth* 7, 55-65.

651 Ward SE, Smart SM, Quirk H, Tallowin JRB, Mortimer SR, Shiel RS, Wilby A, Bardgett RD.
652 2016. Legacy effects of grassland management on soil carbon to depth. *Global Change*
653 *Biology* 22, 2929-2938.

654 Warren SD, Thurow TL, Blackburn WH, Garza NE. 1986. The influence of livestock
655 trampling under intensive rotation grazing on soil hydrological characteristics. *Journal*
656 *of Range Management* 39, 49-495.

657 Wegner T. 2000. *Applied business statistics: Methods and applications*. Juta and Co. Ltd Cape
658 Town, South Africa, ISBN 0702128732.

659 Weitz M, Wood MK. 1986. Short duration grazing in Central New Mexico: Effects on
660 infiltration rates. *Journal of Rangeland Management* 39, 365-368.

661 WRB. 2015. World reference base for soil resources 2014, update 2015. International soil
662 classification system for naming soils and creating legends for soil maps World Soil
663 Resources Reports No. 106, FAO, Rome.

664 Wu X, Li Z, Fu B, Lu F, Wang D, Liu H, Liu G. 2014. Effects of grazing exclusion on soil
665 carbon and nitrogen storage in semi-arid grassland in Inner Mongolia, China. *Chinese*
666 *Geographical Science* 24 (4), 479-487.

667 Xu M-Y, Wang K, Xie F. 2011. Effects of grassland management on soil organic carbon
668 density in agro-pastoral zone of northern China. *African Journal of Biotechnology* 10,
669 4844-4850.

670 Xu M-Y, Xie F, Wang K. 2014. Response of vegetation and soil carbon and nitrogen storage
671 to grazing Intensity in semi-arid grasslands in the agro-pastoral zone of Northern
672 China. *Plos One* 9 (5), doi: 10.1371/journal.pone.0096604.

673 Yalçın E, Doğan A, Akçin A. 2016. Effects of Grazing on Soil Parameters in Meadow
674 Ecosystems. *Hacettepe Journal of Biology Chemistry* 44 (2), 161-171.

675 Yan L, Chen S, Huang J, Lin G. 2010. Differential responses of auto- and heterotrophic soil
676 respiration to water and nitrogen addition in a semiarid temperate steppe. *Global*
677 *Change Biology* 16, 2345–2357.

678 Yang J, Kang Y, Sakurai K, Ohnishi K, Siriguleng. 2016. Change of soil microbial biomass,
679 C, N between long-time free grazing and enclosure pastures in semiarid grassland
680 ecosystem in Tongliao and Chifeng of Inner Mongolia. *Journal of Bioremediation*
681 *Biodegradation* 7 (3), doi: 10.4172/2155-6199.1000347.

682 Yusuf HM, Treydte AC, Sauerborn J. 2015. Managing semi-arid rangelands for Carbon
683 storage: Grazing and woody encroachment effects on Soil Carbon and Nitrogen. *PLoS*
684 *ONE* 10 (10), doi: 10.1371/journal.pone.0109063.

685 Zhao Y, Peth S, Kru¨mmelbeina J, Horn R, Wang Z, Steffens M, Hoffmann C, Peng X. 2007.
686 Spatial variability of soil properties affected by grazing intensity in Inner Mongolia
687 grassland. *Ecological Modelling* 205, 241-254.

688 Zhou G, Guan L, Wei X., Tang X, Liu S, Liu J, Zhang D, Yan J. 2008. Factors influencing
689 leaf litter decomposition: an intersite decomposition experiment across China. *Plant*
690 *and Soil*. 311, 1-2, p.61.

691 Zhou ZC, Gan ZT, Shangguan ZP, Dong ZB. 2010. Effects of grazing soil physical properties
692 and soil erodibility in semiarid grassland of the Northern Loess Plateau (China).
693 *Catena* 82, 87-91.




Figure 1 Global distribution of the sites where data used in the analysis




Figure 2. Soil response to selected grassland management practices as compared to free grazing at low density as control. A: Soil organic carbon stocks; B: Soil organic carbon content; C: soil bulk density. Error bars are effect size means \pm 95% bootstrap CIs. Where the CIs do not overlap the vertical $x=0$ line, the effect size for a practice is significant. n is the number of studies and the percentage corresponding to the bar indicates the mean percentage changes per year when changes are significant. Data for the surface soil layer are considered here.





Figure 3. Impact of soil orders (WRB, 2015) on the changes of soil organic carbon stocks following rehabilitation. The general statistics are median, 1st and 3rd quartile, non-outlier minimum and maximum.



-
-
- Figure 4. Impact of management duration (years) on the changes of soil organic carbon stocks following rehabilitation. The general statistics are median, 1st and 3rd quartile, non-outlier minimum and maximum.




Figure 3 Principal component analysis for selected grassland management practices (all treatments, A; grazing treatments, B; Fire treatments; C Exclosure treatments, D) and showing the correlation between the changes in selected soil properties and soil and environmental variables.

Table 1 Compilation of references (authors and years), country of reference, investigated practices (B, burn; G, Grazing; E, enclosure), mean treatment duration and selected environmental factors (MAP, mean annual precipitation; MAT, mean annual temperature), soil type as provided by the authors and equivalent WRB (2016) soil order.

n	Authors	Soil Type	Soil Order	Countries	Practices	Duration y ⁻¹	MAP mm y ⁻¹	MAT °C y ⁻¹
1	Abril and Bucher 1999	Aridic Haplustoll	Aridisols	Argentina	G	NA	550	23
2	Harris et al 2007	Calcistolls	Mollisols	Argentina	E, B	25,2	400	16
3	Lavado et al 1996	Typic Natraquoll	Mollisols	Argentina	G	38,8	950	15
4	Picone et al 2003	Typic Natraquoll,	Mollisols	Argentina	B	4,0	1029	15
5	Sanjari et al 2008	NA (clay to clay loam)	Alfisols	Australia	G, E	NA	645	15
6	Kgosikoma et al 2015	NA loam to sand	Entisols	Botswana	G, B	25,0	426	22
7	Conant et al 2017	NA	Oxisols	Brazil	G, E, B	16,0	1616	23
8	Nardoto and Bustamante 2003	Oxisols	Oxisols	Brazil	B	NA	1350	21
9	Donkor et al 2001	Luvissols	Alfisols	Canada	E	1,0	334	-8.5
10	Evans et al 2012	Chernozems	Mollisols	Canada	E	0,5	270	15
11	Krzic et al 2013	Chernozems	Mollisols	Canada	G	6,5	409	7.2
12	Cao et al 2013	NA	Mollisols	China	G, E	NA	280	3.4
13	Chen and Tang 2016	Calcic Chernozem,	Mollisols	China	G, E	1,3	279	2.4
14	Gao et al 2009	Mat Cry-gelic Cambisol	Mollisols	China	G	NA	750	1.1
15	Jiao et al 2016	"diluvial desert soil"	Aridisols	China	G	25,8	185	8.3
16	Liu et al 2012	NA	Mollisols	China	G, E	13,0	308	2.1
17	Pei et al 2007	Typic Calciorthid	Mollisols	China	E	13,5	134	9
18	Rong et al 2014	Calciorthid	Mollisols	China	G, E	25,0	100	3.5
19	Steffens et al 2008	Calcic Chernozems	Mollisols	China	G, E	17,0	343	0.7
20	Su et al. 2003	Sandy chestnut	Mollisols	China	G	5,0	366	6.5
21	Su et al 2004	Sandy chestnut	Mollisols	China	E	25,0	366	6.5
22	Su et al 2005	Sandy chestnut	Mollisols	China	E	10,0	366	6.5
23	Wang et al 2016	Orthic Aridisol	Aridisols	China	G, E	NA	400	1.5
24	Wei et al 2011	sub-alpine meadow	Entisols	China	G	2,0	620	1.2
25	Wu et al 2014	Chernozem/chestnut	Mollisols	China	G	25,0	339	-2.2
26	Xu et al 2011	Calcic-orthic	Aridisols	China	G, E	25,0	400	1.2
27	Xu et al 2014	Calcic-orthic	Aridisols	China	G, E	25,0	400	1.5
28	Yang et al 2016	Chernozem	Mollisols	China	G	12,0	378	7.3
29	Zhao et al 2007	Chernoziom	Mollisols	China	G, E	21,3	343	0.7
30	Zhou et al 2010	NA ("loess soils")	Mollisols	China	G, E	7,0	505	8.8
31	Kassahun et al 2012	Vertisols	Vertisols	Ethiopia	G	7,0	250	32
32	Yusuf et al 2015	Chromic Cambisol	Mollisols	Ethiopia	G, E	3,0	550	20
33	Nusse et al 2017	Vertisol	Vertisols	Germany	G	25,0	879	8.2
34	Devi et al 2014	NA (clayey loam)	Alfisols	India	G	30,6	1522	19
35	Zarekia et al 2012	NA (sandy loam)	Entisols	Iran	G, E	25,0	200	19
36	Koyama et al 2015	NA	Mollisols	Japan	B	30,0	232	0.2
37	Shimoda and Takahashi 2009	Gleyic	Ultisols	Japan	G	26,0	1724	16
38	Barger et al 2004	NA (coarse textured)	Mollisols	Mongolia	G	NA	399	0
39	Han et al 2008	Chernozem	Mollisols	Mongolia	G	NA	400	-0.2
40	Niu et al 2011	calcisols	Mollisols	Mongolia	G, E	21,8	210	8
41	Al-Seekh et al 2009	Rendzinas	Entisols	Palestine	E	6,5	325	32
42	Abdalla et al 2016	Westleigh	Alfisols	South Africa	B	42,5	694	16
43	Chaplot et al 2016	Acrisols	Alfisols	South Africa	G, E	4,0	684	13
44	Manson et al 2007	Hutton	Alfisols	South Africa	B	25,0	1380	25
45	Materechera et al 1998	Glenrosa	Alfisols	South Africa	B	13,5	500	21
46	Oluwole et al 2008	Glenrosa	Alfisols	South Africa	G, B	8,0	500	21
47	Snayman 2002	Bloemdal	Alfisols	South Africa	B	1,0	560	25
48	Oğuz, et al 2015	NA	Aridisols	Turkey	G, E	4,0	306	9.4
49	Yalçın et al 2016	Vertisol	Vertisols	Turkey	E	10,0	797	14
50	Ward et al 2016	NA	Alfisols	UK	G	25,0	885	9.7
51	Bauer et al 1987	Haploborolls	Mollisols	USA	E	9,0	412	6.3
52	Frank et al 1995	Haploborolls	Mollisols	USA	G, E	25,0	414	5.5
53	Gardner 1950	Whitehouse-	Entisols	USA	E	NA	431	11.6
54	Gill 2007	Cryoborolls	Entisols	USA	E	NA	932	1.1
55	Manley et al 1995	Ascalon/Altvan	Aridisols	USA	G	NA	405	7.6
56	Neff et al 2005	Haplocambids	Aridisols	USA	G, E	25	207	12
57	Seithheko et al 1993	Aridic	Aridisols	USA	G	3	400	15
58	Schuman et al 1999	Argiustolls	Mollisols	USA	G	8	387	8
59	Teague et al 2010	Tilvern and Wichita	Alfisols	USA	G, B	7,5	648	17
60	Weitz and Wood, 1986	Leeray clay	Vertisols	USA	G	25	361	9.9
61	Wood and Blackburn 1984	Leeray clay	Vertisols	USA	G	10	624	17.2
62	Nyamadzawo et al 2014	Haplic Lixisols	Alfisols	Zimbabwe	G, B	25	750	19

Table 2 Definitions of the environmental factors and soil properties used in the data-analysis

Environmental factors, soil properties and techniques	Symbols	Units	Definitions
Mean annual precipitation	MAP	mm year ⁻¹	Long-term (at least 30 year) mean annual precipitation for the study location from the papers
Mean annual temperature	MAT	°C	Long-term (at least 30 year) mean annual air temperature for the study location from the papers
Longitude	LONG	°	Longitude of the midpoint of study site as given in paper
Latitude	LAT	°	Latitude of the midpoint of study site as give in paper
Altitude	Z	m	Average elevation above sea level of the study site given in the papers
Clay content	Clay	%	Average clay content
Sand	Sand	%	Average sand content

Table 3 List of controlling factor classes describing the environmental conditions used in the analyses.

Environmental factors	Remarks	Class range	Name
Climate (MAP, mm year ⁻¹ ; MAT, °C year ⁻¹)	Warm and dry	MAT: 3.5-31.9 MAP 100-300	Arid
	Cool and dry	MAT: -2.2-25.0 MAP:300-550	Semi-arid
	Cool and moist	MAT: 1.2-18.6 MAP: 550-700	Moist
	Warm and wet	MAT: 8.2-25.0 MAP:700-1350	Humid
	Warm and wet	MAT:15.6-18.9 MAP: >1350	Wet
Clay (%)	Average clay	0-20	Low
		20-40	Medium
		>40	High
Sand (%)	Average sand content	0-20	Low
		20-40	Medium
		>40	High
Grazing intensity: number of animal units stocked per unit area per year (AU ha ⁻¹ yr ⁻¹)	Light grazing	0.4-2.5	LG
	High density	>2.5	HG
Livestock enclosure: period of time a plot has been excluded from grazing and/or burning (years)		≤10	E≤10
		10-20	E10-20
		20-30	E20-30
		≥30	E≥30
Burning frequencies: number of years from one incidence of burning grass to the other (years)		1-2	B1-2
		3-5	B3-5
		>5	B>5

Climate classes were adapted from Mathew et al. (2017); other factor classes were adapted from Mutema et al (2015).

Table 4 Overall descriptive statistics of the controlling factors selected for this study. Data from 235 experimental sites worldwide.

	Mean	Media	Min	Max	Q1	Q3	CV	SEM	Skew	Kurt
MAP	711	494	100	2326	343	879	77	36,1	1.4	1.4
MAT	14.0	15.0	-8.5	32.0	6.7	20.8	69.6	0.6	0.0	0.0
Z	956	1012	4	3500	418	1375	67	44.8	0.7	0.6
Clay	24.9	23.0	3.0	74.0	11.0	33.0	68.7	1.5	1.0	1.0

Mean; Median, Min=minimum value; Max=maximum value; Q1 and Q3=25th and 75th percentiles; CV=coefficient of variation; SEM=standard error of mean; Skew=skewness; Kurt=kurtosis; MAP=mean annual precipitation (mm yr⁻¹); MAT=mean annual air temperature (°C); Z=altitude (m); Clay= clay content (%).

Table 5 Overall descriptive statistics of the selected variables. Soil organic carbon content (SOC_C, %); soil organic nitrogen content (SON_C, %), Soil organic carbon stocks (SOC_s, kg C m³); soil bulk density (BD in g cm⁻³) ; variation in SOC_C, SON_C, SOC_s and BD (Δ SOC_C, Δ SON_C, Δ SOC_s and Δ BD in percentage difference per year from the reference (Ref) and the considered treatment (Treat), C/N ratio.

	n	Mean	median	min	max	Q1	Q3	CV	SEM	Skew	Kurt
SOC_C Ref	446	2.26	1.44	0.06	20.16	0.96	2.83	116	0.19	3.87	3.84
SOC_C Treat	446	2.34	1.42	0.12	26.04	0.95	2.53	135	0.22	4.75	4.72
ΔSOC_C	446	0.71	0.34	-38.7	42.9	-0.85	1.69	933	0.52	0.37	0.38
SON_C Ref	301	0.19	0.15	0.01	1.17	0.11	0.21	104	0.02	3.29	3.24
SON_C Treat	301	0.19	0.15	0.01	1.16	0.09	0.21	92	0.02	2.98	2.93
ΔSON_C	301	1.17	0.02	-25.9	28.9	-1.58	2.08	715	0.84	0.76	0.75
SOC_s Ref	367	3,02	1,93	0,08	12,70	1,34	4,27	86,96	0,17	1,72	1,71
SOC_s Treat	367	3,04	1,95	0,15	14,50	1,33	4,44	91,74	0,18	1,93	1,91
ΔSOC_s	367	0.24	0.089	-40.6	41.3	-1.05	0.98	748	0.54	0.08	0.08
C/N Ref	301	11.52	10.42	0.00	102	8.81	13.59	91	1.10	7.34	7.22
C/N Treat	301	11.72	10.10	0.00	103	8.26	14.37	92	1.14	6.78	6.67
ΔC/N	301	-0.43	-0.07	-56	87	-8.87	7.36	433	1.98	0.96	0.94
BD Ref	386	1.22	1.17	0.63	1.77	1.03	1.43	20	0.02	0.09	0.09
BD Treat	386	1.25	1.22	0.53	2.11	1.07	1.42	20	0.02	-0.04	-0.03
ΔBD	386	-0.40	-0,25	-10,71	13,91	-0,76	0,01	559	0,15	0,24	0,23

Table 6 Spearman rank correlation coefficients between environmental factors and the soil properties under different types of change in grassland management: grazing intensity, grazer exclusion (E), grass burning (B).

	-----Grazing-----				-----B-----				-----E-----			
	ΔSOC_c	ΔSON_c	$\Delta\text{C/N}$	ΔBD	ΔSOC_c	ΔSON_c	$\Delta\text{C/N}$	ΔBD	ΔSOC_c	ΔSON_c	$\Delta\text{C/N}$	ΔBD
MAP	0.27*	-0.39*	0.12	-0.49*	0.15	-0.14	0.53*	-0.33*	0.14	-0.07	0.23*	0.13
MAT	0.58*	0.54*	0.37*	-0.41*	0.02	-0.33*	0.02	-1.00*	-0.18	0.03	0.04	0.08
LAT	-0.20	0.31*	-0.04	0.27*	-0.17	-0.36*	0.21*	-0.33*	0.26*	0.18	0.04	-0.38*
Z	-0.06	-0.19	0.35*	0.04	0.02	-0.02	0.29*	-0.33*	-0.15	-0.12	0.03	0.31*
CLAY	0.67*	0.33*	0.47*	0.37*	-0.74*	-0.62*	0.24*	nd	0.10	0.43*	-0.45*	-0.32*
SOC_c ref	-0.05	-0.11	0.62*	0.23*	0.29*	-0.30*	0.41*	-1.00*	-0.26*	-0.18	0.08	-0.18
BD	0.52*	0.29*	-0.72*	-0.39*	0.74*	0.32*	0.32*	0.33*	-0.05	0.04	-0.05	0.20

* Significant correlations at $p < 0.05$

nd not determined due lack of adequate data

MAP=mean annual precipitation (mm yr^{-1}); MAT=mean annual air temperature ($^{\circ}\text{C}$); LAT=latitude ($^{\circ}$); Z=altitude above sea level (m); Clay=clay content (%); reference organic carbon content (SOC_c ref); soil bulk density (BD, g cm^{-3}).

Table 7 Climate and soil clay content impact on the change in soil organic carbon stocks (Δ SOCs).

<i>Climate</i>	CLG			CH			E		Burn			n
	g m⁻¹	%	n	g m⁻¹	%	n	g m⁻³	% yr⁻¹	n	g m⁻³	% yr⁻¹	
Arid/semi-arid	25.4	1.25	23	15.6*	1.9*	6	1.86	0.98	46	-819.2*	-15.3*	14
Moist/humid	65.3	1.98	26	233*	8.9*	3	38.7	3.8	20	-143.8*	-9.1*	13
Wet	47.9	3.12	245	6.2	0.5	1	-34.4*	-2.1*	4	63.73	0.24	31
<i>Clay content</i>												
0-20	43.1	2.77	148	211.7*	3.15*	6	53.35	4.81	13	-6.07	-1.51	13
20-40	166.3*	4.79*	61	nd	nd	4	-10.6	-2.44	12	-93.22	-6.35	17
>40	7.0	1.18	69	43.5	0.65	2	-51.71	-1.87	4	-17.71	-0.20	9

* Significant correlations at $p < 0.05$