



HAL
open science

Exposure to low to moderate doses of ionizing radiation induces a reduction of pro-inflammatory Ly6Chigh monocytes and a U-curved response of T cells in ApoE -/- mice

Nicolas Rey, Talin Ebrahimian, Celine Gloaguen, Dimitri Kerezeldze, Victor Magneron, Alice Bontemps, Christelle Demarquay, Gunilla Olsson, Siamak Haghdoost, Stephanie Lehoux, et al.

► To cite this version:

Nicolas Rey, Talin Ebrahimian, Celine Gloaguen, Dimitri Kerezeldze, Victor Magneron, et al.. Exposure to low to moderate doses of ionizing radiation induces a reduction of pro-inflammatory Ly6Chigh monocytes and a U-curved response of T cells in ApoE -/- mice. *Dose-Response*, 2021, 19 (2), pp.1-9. 10.1177/15593258211016237 . hal-03608570

HAL Id: hal-03608570

<https://hal.science/hal-03608570v1>

Submitted on 14 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Exposure to Low to Moderate Doses of Ionizing Radiation Induces A Reduction of Pro-Inflammatory Ly6^{Chigh} Monocytes and a U-Curved Response of T Cells in APOE ^{-/-} Mice

N. Rey¹, T. Ebrahimian², C. Gloaguen¹, D. Kereselidze¹, V. Magneron¹, C. A. Bontemps¹, C. Demarquay¹, G. Olsson³, S. Haghdoust^{3,4}, S. Lehoux², and Teni G. Ebrahimian¹ 

Dose-Response:
An International Journal
April-June 2021: 1-9
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15593258211016237
journals.sagepub.com/home/dos



Abstract

Low dose ionizing radiation (LDIR) is known to have a protective effect on atherosclerosis in rodent studies, but how it impacts different cells types involved in lesion formation remains incompletely understood. We investigated the immunomodulatory response of different doses and dose-rates of irradiation in ApoE^{-/-} mice. Mice were exposed to external γ rays at very low (1.4 mGy.h⁻¹) or low (50 mGy.h⁻¹) dose-rates, with cumulative doses spanning 50 to 1000 mGy. Flow cytometry of circulating cells revealed a significant decrease in pro-inflammatory Ly6^{Chigh} monocytes at all cumulative doses at low dose-rate, but more disparate effects at very low dose-rate with reductions in Ly6^{Chigh} cells at doses of 50, 100 and 750 mGy only. In contrast, Ly6^{Clow} monocytes were not affected by LDIR. Similarly, proportions of CD4⁺ T cell subsets in the spleen did not differ between irradiated mice and non-irradiated controls, whether assessing CD25⁺FoxP3⁺ regulatory or CD69⁺ activated lymphocytes. In the aorta, gene expression of cytokines such as IL-1 and TGF- β and adhesion molecules such as E-Selectin, ICAM-1, and VCAM-1 were reduced at the intermediate dose of 200 mGy. These results suggest that LDIR may reduce atherosclerotic plaque formation by selectively reducing blood pro-inflammatory monocytes and by impairing adhesion molecule expression and inflammatory processes in the vessel wall. In contrast, splenic T lymphocytes were not affected by LDIR. Furthermore, some responses to irradiation were nonlinear; reductions in aortic gene expression were significant at intermediate doses, but not at either highest or lowest doses. This work furthers our understanding of the impact of LDIR with different dose-rates on immune system response in the context of atherosclerosis.

Keywords

Low dose, dose-rate, immune system, atherosclerosis

Introduction

Despite multiple studies investigating the health impacts of ionizing radiation, questions remain regarding the effects of exposure on cardiovascular diseases. Both epidemiological¹⁻⁴ and experimental studies⁵⁻¹¹ revealed that exposure to high doses of ionizing radiation has detrimental cardiovascular outcomes. However, when it comes to low dose ionizing radiation (LDIR), epidemiological studies have not provided a clear message. For instance Schöllnberger et al¹² did not resolve whether or not there is a risk of heart disease for individuals exposed to doses below 2.6 Gy, and Azizova et al¹³ showed an

¹ IRSN, Institut de Radioprotection et de Sûreté Nucléaire, Laboratoire de Radiotoxicologie et Radiobiologie Experimentale, Fontenay-aux-Roses, France

² Lady Davis Institute, McGill University, Montreal, Canada

³ Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Stockholm, Sweden

⁴ University of Caen Normandie, Aria/Cimap, GANIL, Campus Jules Horowitz, Caen, France

Received 7 January 2021; received revised 8 April 2021; accepted 15 April 2021

Corresponding Author:

Teni G. Ebrahimian, Institut de Radioprotection et de Sûreté Nucléaire (IRSN), Laboratoire de Radiotoxicologie et Radiobiologie Experimentale, 92262, Fontenay-aux-Roses, France.

Email: teni.ebrahimian@irsn.fr



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

increased risk of developing ischemic heart disease for cumulative external doses above 1 Gy. There remains a lack of epidemiological data on the impact of radiation on cardiovascular diseases after exposure to low to moderate doses, especially under 500 mGy.¹⁴

Differing from the paucity of epidemiological information linking LDIR and cardiovascular diseases, several experimental studies have investigated this matter, revealing the importance of considering the dose-rate of radiation exposure. Such studies have been conducted mainly on mouse models of atherosclerosis, using genetically modified ApoE^{-/-} mice which become hyperlipidemic and develop atherosclerotic lesions similar to those found in humans when fed a chow diet.¹⁵ Atherosclerosis is a disease characterized by chronic and exacerbated inflammation in the wall of medium and large arteries, due to lipoprotein influx. Excessive growth or rupture of atherosclerotic plaques underlies most cardiovascular diseases. Mitchel et al¹⁶ demonstrated that exposure to LDIR, especially at low dose-rate, slowed plaque progression in mice. Mancuso et al¹⁷ indicated that acute irradiation at moderate doses (300 mGy) can have detrimental effects on atherosclerosis, but that chronic exposure to the same dose has less impact. Finally, both Le Gallic et al¹⁸ and Ebrahimian et al¹⁹ found that exposure to chronic internal LDIR enhances plaque stability in ApoE^{-/-} mice. These latter 2 studies even showed a decrease in inflammatory parameters after exposure to LDIR, including diminished plaque content of CD68⁺ foam cells and a shift in aortic mRNA expression favoring anti- rather than pro-inflammatory cytokines.^{18,19} These data corroborate the notion that ionizing radiation has an impact in the immune system, with high doses promoting inflammation²⁰ and proinflammatory macrophages,²¹ whereas lower doses lead to decreased inflammation.^{22,23}

Nevertheless, there is little information available regarding the impact of LDIR on monocytes and lymphocytes that contribute to atherosclerosis lesion formation in the first place. One study reported that *in vitro* irradiation of RAW264.7 monocytes/macrophages altered the binding of these cells to vascular cell adhesion molecule-1 (VCAM-1),²⁴ confirming a previous work demonstrating that LDIR reduced adhesion of monocytes to the endothelium.²⁵ It was also determined that LDIR affects cytokine gene expression by different T cells.²⁶ Only 1 study explored the effects of LDIR on white and red blood cell populations in the context of atherosclerosis,¹⁹ but specific monocyte and lymphocyte subgroups were not investigated. This represents a significant knowledge gap, since monocytes and T cells can polarize into distinct classes that are considered to be either pro-inflammatory or anti-inflammatory, with potentially opposing effects on atherosclerosis.^{27,28}

Hence, identifying how low doses of irradiation can modulate the inflammatory response in atherosclerosis is clearly essential. In order to investigate this process, ApoE^{-/-} mice were irradiated at 2 different dose-rates (50 mGy.h⁻¹ and 1.4 mGy.h⁻¹) and a wide range of cumulative doses of irradiation ranging from 50 mGy to 1000 mGy. Immunophenotyping of immune T cells in the spleen and monocyte subsets in

circulating blood were explored, and aortic mRNA expression of pro-inflammatory cytokines and adhesion molecules were evaluated.

Methods

Animals

All experiments and procedures were carried out in accordance with the Guide for the Care and Use of Laboratory Animals as published by the French regulations for animal experiments (Ministry of Agriculture Order No. B92-032-01, 2006) with European Directives (86/609/CEE), and approved by the local ethical committee of the Institute for Radiological Protection and Nuclear Safety (permit number P10-11, thematic number T29) and the Swedish Board of Agriculture (permit number N 134/16). Six to 8 week-old ApoE^{-/-} male mice on a C57BL/6 J background were obtained from Charles River Laboratory. Groups consisted of 6 to 8 mice. Animals were maintained in a specific-pathogen-free environment and monitored daily. Mice were fed a regular chow diet *ad libitum* and maintained in a 12 h light/dark cycle environment.

Irradiation

Mice were distributed into 6 groups according to total cumulative doses of γ rays (¹³⁷Cs): 50, 100, 200, 500, 750, or 1000 mGy (n = 6 each), and a control non-irradiated group (n = 8). Two dose rates were applied: very low dose rate (1.4 mGy.h⁻¹) for chronic exposure and low dose rate (50 mGy.h⁻¹) for acute exposure. Hence, it was possible to compare not only the impact of absolute doses to one another, but also compare equivalent doses administered at different dose-rates. Animals were irradiated at the Department of Molecular Biosciences, Experimental Animal Core Facility, Stockholm University, Sweden and then sent to IRSN. Application of correct dosimetry was assured by daily calibration using PTW UNIDOS E Universal Dosimeter equipped with a Farmer Ionization Chamber Type 30010 in the 9 different positions inside the cage thus limiting dose uncertainty to approximately +/- 7%. Mouse groups exposed to the same dose were sacrificed at the same age (Supplementary Figure 1).

Blood Sampling and Analyses

Mice were terminally anesthetized by intraperitoneal injection of ketamine/xylazine (Ketamine 500 Virbac, Rompun 2% Bayer). Blood was collected by cardiac puncture with a heparinized syringe. Blood was centrifuged for 10 minutes at 1000 g. Spleens were collected in FACS buffer (PBS1X-FBS 2%), whereas aortas were collected in Trizol (Sigma Aldrich LLC).

Flow Cytometry

Total blood and spleens were collected for flow cytometry experiments. Spleens were mashed, filtered and incubated with




Figure I. Exposure to low doses of irradiation decreases Ly6C^{Hi} pro-inflammatory circulating monocytes but does not affect anti-inflammatory Ly6C^{Lo} cells. ApoE^{-/-} mice were exposed to low doses of γ ionizing radiation at dose-rates of 1.4 and 50 mGy.h⁻¹. Mice were

sacrificed at day 12-50 post-irradiation and blood monocytes analyzed by flow cytometry. No effects of irradiation were noted in Ly6C^{Lo} cells. However, proportions of Ly6C^{Hi} monocytes (among CD11b⁺Gr1⁺ populations) were reduced at all doses of irradiation, most consistently in mice exposed to the dose-rate of 50 mGy.h⁻¹. Data are mean \pm SEM of n = 5-8. *P < 0.05, **P < 0.01 vs non-irradiated, age-matched control.

Real-time Polymerase Chain Reaction

Total RNA was extracted from aortas using Tri Reagent solution (Sigma Aldrich LLC). RNA quality (260/280 nm) was determined using a Nanodrop ND 1000 spectrophotometer. One microgram of total RNA was synthesized to 20 μ L complementary DNA (cDNA) using the high-capacity cDNA Reverse Transcription Kit from Applied Biosystems (Life Technologies) according to the manufacturer's protocol. Quantitative polymerase chain reaction analysis was performed with a QuantStudio 12 K Flex Real-Time PCR System (Life Technologies) using TaqMan 6 carboxyfluorescein-labeled probes and a standard thermal cycler protocol (50°C for 2 minutes before the first cycle, 95°C for 15 seconds, and 60°C for 1 minute repeated 45 times). Samples were run in duplicates and normalized with gapdh and hprt using geometric mean using the 2^{- $\Delta\Delta$ CT} method, control samples serving as a reference value of 1. We quantified the mRNA expression of intracellular adhesion molecule-1 (ICAM-1), vascular adhesion molecule-1 (VCAM-1), E-Selectin (E-Sel), tumor necrosis factor alpha (TNF- α), interleukin-1 (IL-1), and transforming growth factor beta (TGF β). The following assays were used: TGF β -Mm01178820_m1, ICAM-1-Mm00516023_m1, TNF α -Mm00443258_m1, E-Sel-Mm0441278_m1, IL-1rap-Mm00492638_m1, VCAM-1-Mm01320970-m1, GAPDH-Mm99999915_g1, and HPRT-Mm01545399_m1.

Figure I. (Continued). sacrificed at day 12-50 post-irradiation and blood monocytes analyzed by flow cytometry. No effects of irradiation were noted in Ly6C^{Lo} cells. However, proportions of Ly6C^{Hi} monocytes (among CD11b⁺Gr1⁺ populations) were reduced at all doses of irradiation, most consistently in mice exposed to the dose-rate of 50 mGy.h⁻¹. Data are mean \pm SEM of n = 5-8. *P < 0.05, **P < 0.01 vs non-irradiated, age-matched control.




Figure 2. Exposure to low doses of ionizing radiation has little impact on spleen T cells. ApoE^{-/-} mice were exposed to low doses of γ ionizing radiation at dose-rates of 1.4 and 50 mGy.h⁻¹. Mice were sacrificed at day 12-50 post-irradiation and spleen T cells analyzed by flow cytometry. (A) Proportions of regulatory T cells (CD25⁺Foxp3⁺ Tregs) quantified among CD4⁺CD8⁻ lymphocytes did not vary at any dose or dose-rate. (B) A reduced proportion of activated CD69⁺ T cells (among CD4⁺CD8⁻CD25⁻Foxp3⁻ lymphocytes) was observed at the single dose of 100 mGy administered at very low dose-rate. Data are mean \pm SEM of n = 5-8. **P < 0.01 vs non-irradiated, age-matched control.

Statistics

Results are presented as mean \pm SEM. Data were compared with 2-way analysis of variance (ANOVA) or 2-way ANOVA for repeated measures, with all ANOVA tests followed by a Student–Newman–Keuls posthoc test, or with an unpaired t-test, as appropriate. $P < 0.05$ was considered statistically significant.

Results

Low Doses of Ionizing Radiation Administered at Low or Very low Dose-Rates Decrease Blood pro-Inflammatory Ly6C^{Hi} Monocytes

Flow cytometry analysis of blood samples provided an overview of circulating monocyte response to irradiation. Ly6C^{Hi} monocytes are dominant in an inflammatory response and are often named “pro-inflammatory monocytes”. They accumulate preferentially in atherosclerotic plaques, compared with Ly6C^{Lo} monocytes.²⁹ We thus distinguished 2 circulating populations of monocytes according to expression of Ly6C, as depicted in Supplementary Figure 2. We observed that exposure to low doses of external gamma rays induced a significant reduction in the proportion of Ly6C^{Hi} pro-inflammatory monocytes in the blood (Figure 1, Supplementary Figure 3). This was especially evident when irradiation was administered at the 50 mGy.h⁻¹ dose-rate, which led to a 45%-78% decrease in circulating Ly6C^{Hi} cells at every dose tested ($P < 0.05$), compared with non-irradiated controls. At the 1.4 mGy.h⁻¹ dose-rate, significant reductions in Ly6C^{Hi} monocytes were also observed, although significance was only reached at doses of 50, 100 and 750 mGy (Figure 1, Supplementary Figure 3). In contrast, proportions of Ly6C^{Lo} “patrolling” monocytes in the blood were not affected by irradiation, whatever the dose or dose-rate applied. These results indicate that low doses of ionizing irradiation have a selective effect on circulating monocyte abundance, resulting in a reduced relative abundance of Ly6C^{Hi}, pro-inflammatory populations.

Low Doses of Ionizing Radiation Have Little Impact on T Cell Populations in the Spleen

The balance between regulatory (Treg) and effector (Teff) T cells can have a profound impact on atherosclerotic plaque formation and progression. Treg reduce the activation and proliferation of effector T cells and are important modulators of atherosclerotic lesion formation.³⁰ We compared the effects of different irradiation regimens on CD4⁺ T cells, either regulatory (CD25⁺FoxP3⁺) or effector (CD25⁺FoxP3⁻). The sorting strategy used to distinguish these populations are depicted in Supplementary Figure 4. Treg cells were not impacted by irradiation at all, whatever the dose or dose-rate tested (Figure 2, Supplementary Figures 4 and 5). However, we found that exposure to LDIR at very low dose and at low dose-rate induced a decrease in activated effector T cells in ApoE^{-/-} mice.




Figure 3. Exposure to low doses of ionizing radiation triggers a non-linear reduction in adhesion molecule and cytokine mRNA expression in the aorta. ApoE^{-/-} mice were exposed to low of γ ionizing radiation at dose-rates of 1.4 mGy.h⁻¹ (A) and 50 mGy.h⁻¹ (B). Mice were sacrificed at day 12-50 post-irradiation and aortic mRNA expression levels evaluated by Q-PCR. Samples were run in duplicates and normalized with gapdh and hprt using the 2^{- $\Delta\Delta$ CT} method. Results are expressed as fold change vs non-irradiated, age-matched control mice. Data are mean \pm SEM of n = 5-8 animals. **P* < 0.05 vs control.

Proportions of CD25⁺FoxP3⁺CD69⁺ cells were reduced in mice exposed to 100 mGy, to 3.4% \pm 0.6% of CD4⁺ cells, compared with 5.8% \pm 0.7% in control mice (*P* < 0.01). In mice irradiated at the lowest dose (50 mGy), reductions in CD25⁺FoxP3⁺CD69⁺ cells approached but did not reach significance. Hence, LDIR has little impact on splenic T cell populations, affecting only effector T cells at very low doses.

Low Doses of Ionizing Radiation Reduced the mRNA Expression of Cytokines and Adhesion Molecules in Aortas of ApoE^{-/-} Mice

Adhesion molecules, chemokines and cytokines have a direct impact in the ability of monocytes to be recruited to the vessel wall, to adhere and transmigrate through the endothelial layer, and to differentiate in inflammatory type macrophages. Their relative expression was investigated in aortas of ApoE^{-/-} mice. No effects were detected in animals exposed to the lowest and highest doses of LDIR, 50 and 1000 mGy (Figure 3). However, at the intermediate dose of 200 mGy, a significant impact on gene expression was noted. Among the adhesion molecules, VCAM-1 was significantly decreased at dose-rates of both 1.4 and 50 mGy.h⁻¹, by 51% and 54% respectively (*P* < 0.05). E-selectin was only reduced in aortas of mice exposed to at 50 mGy.h⁻¹. As for cytokine response, we observed a significant downregulation of TGFβ at both very low (51%) and low dose-rates (34%) compared with control. IL-1 expression was also reduced by half at the 50 mGy.h⁻¹ dose-rate

(*P* < 0.05). Hence, changes in the vasculature induced by low-dose irradiation point to a reduction in chemokine and adhesion molecule expression.

Discussion

In this study we show that very low and low doses of irradiation produce diverse responses in the different cell types we investigated, which are implicated in atherosclerosis plaque formation and progression. In general, we found that the low dose-rate (50 mGy.h⁻¹) had more impact than the very low dose-rate (1.4 mGy.h⁻¹) for a given total dose administered. Interestingly, LDIR mostly affected pro-inflammatory monocytes and aortic cells, resulting in reduced monocyte numbers and lowered expression of adhesion molecules and cytokines that would allow these cells to enter the plaque.

Cell components of both innate and adaptive immunity are involved in atherosclerosis. These cells originate in the bone marrow and are released in the circulation, from which they will migrate into tissues following chemotactic gradients, or transfer into lymphoid tissue such as the spleen for further maturation. Following entry and oxidation of low-density lipoproteins in the intimal layer of arteries, endothelial expression of adhesion molecules including ICAM-1, VCAM-1, and E-Selectin, as well as chemokine release, stimulate monocyte recruitment. As monocytes accumulate, they differentiate into macrophages that take up the oxidized lipids and produce pro-inflammatory cytokines such as TNF-α, IL-6, IL-1β, and interferon gamma (IFN-γ).³¹ This results in an alteration of the artery wall structure and enhanced recruitment of additional inflammatory cells including T lymphocytes.³²⁻³⁵

In mice, circulating blood monocytes can be separated in 2 subsets based on expression levels of Ly6C. ApoE^{-/-} mice have greater amounts of circulating monocytes than their wild-type counterparts,²⁹ and hypercholesterolemia enhances the proportion of the pro-inflammatory Ly6C^{Hi} monocytes in blood.^{29,36} Furthermore, Ly6C^{Hi} monocytes enter atherosclerotic plaques more readily than their Ly6C^{Lo} counterparts.²⁹ Our results establish that low-dose irradiation prompted a clear decrease in the levels of circulating Ly6C^{Hi} monocytes in the ApoE^{-/-} mice, at all applied doses. In contrast, no changes in Ly6C^{Lo} cells were observed. Our data support earlier reports that in vitro exposure of monocytes to LDIR favors an anti-inflammatory phenotype,³⁷⁻³⁹ whereas high doses of ionizing radiation have the opposite effect.⁴⁰ In addition, Sharma et al⁴¹ demonstrated that LDIR decreased the monocytes count in blood of rats. However, to the best of our knowledge, we are the first to report a specific suppressive effect of LDIR on the Ly6C^{Hi} monocyte subtype.

A second set of major actors in atherosclerosis is T lymphocytes, from the adaptive immune system. Specific CD4⁺ effector T cells can recognize oxidized LDL in the atherosclerotic intima and aggravate the inflammatory response by producing cytokines including IFN-γ.⁴² Inversely, regulatory T lymphocytes defined as CD4⁺CD25⁺Foxp3⁺ have the ability to modulate atherosclerosis³⁰ by producing the anti-inflammatory

cytokines IL-10 and transforming growth factor beta (TGF β).⁴³ Thus, the balance between inflammatory effector T cells and anti-inflammatory regulatory T cells will influence atherosclerosis progression or regression.⁴⁴ LDIR is broadly reported to alter spleen CD4⁺ T cell proliferation and numbers.⁴⁵⁻⁴⁹ Nevertheless, our data demonstrate that in ApoE^{-/-} mice, low dose irradiation had no impact in spleen Treg numbers, and only reduced activated CD69⁺ cells at a single low dose applied at very low dose-rate. In fact, irradiation appears to produce divergent effects in Tregs. In some studies, regulatory T cells were decreased after exposure of rodents to low dose of ionizing radiation, leading to a rather pro-inflammatory response.^{50,51} In contrast, Tregs were significantly increased following irradiation in mice affected with arthritis,⁵² asthma,⁵³ or prostate cancer.⁵⁴ Likewise, many,^{46,55} but not all⁵⁶ low dose radiation studies report enhanced CD69⁺ activated T cells. Regardless, our findings do not support a predominant role for regulatory or CD69⁺ T cells in the response to LDIR in the context of atherosclerosis.

Finally, we surveyed the chemokine and adhesion molecule profile of aortas in mice submitted to LDIR. Our experiments showed diminished expression levels of IL-1, TGF β , E-Sel, ICAM-1, and VCAM-1 at a cumulative dose of 200 mGy. Most effects were observed at the most elevated dose-rate, although reductions were also observed at 1.4 mGy.h⁻¹. Shin et al⁵⁷ found a variable serum cytokine profile in C57BL/6 mice irradiated at 200 mGy, noting that some interleukins were elevated but others reduced. Our study most closely resembles that of Mathias et al⁵⁸ who also irradiated ApoE^{-/-} mice at different doses and assessed cardiovascular outcomes after 3 and 6 months. They found that expression of some adhesion molecules was reduced in myocardial cross-sections of mice exposed to 25-500 mGy irradiation, but that VCAM-1 and E-Sel were enhanced after 2000 mGy. Variations in levels of circulating pro-inflammatory factors were also noted.⁵⁸ VCAM-1, ICAM-1, and E-selectin allow firm adhesion of monocytes to endothelial cells, which precedes their transmigration,⁵⁹ which contributes to atherosclerosis.^{60,61} Combined with reduced expression of adhesion molecules, the lowered IL-1 expression observed in our study would further favor atherosclerosis abatement, since it is pro-inflammatory and is associated with atherosclerosis progression.⁶²⁻⁶⁴ IL-1 is even involved in the upregulation of adhesion molecule expression on the endothelial cell surface,⁶³ suggesting that its decrease could explain the loss of VCAM-1, ICAM-1, and E-Sel we observed. However, the downregulation of TGF β , an anti-inflammatory and anti-atherogenic cytokine,⁶⁵ is a contradictory result that warrants further investigation.

Interestingly, the exposure of ApoE^{-/-} mice to low dose ionizing radiation resulted in nonlinear responses. At both dose rates, cytokine and adhesion molecule expression were reduced in mice exposed to an intermediate dose of 200 mGy, whereas neither the highest nor the lowest doses of irradiation affected mRNA expression. These nonlinear responses are in agreement with previous *in vitro* studies on gene expression in endothelial cells exposed to LDIR.⁶⁶ Also of interest, the dose-

rate at which LDIR was applied influenced monocyte subset abundance and aortic gene expression. Specifically, most significant effects were observed at 50 mGy.h⁻¹, suggesting that an accelerated dose-rate enhances the effectiveness of low dose irradiation. Previous *in vitro* results demonstrated that LDIR decreased inflammation when applied at an even higher dose rate of 1.15 Gy.min⁻¹.⁶⁷ To the best of our knowledge the dose-rate used in our study has been used in atherosclerosis models only in the works of Mancuso et al and Mitchel et al.^{16,67} However, those studies focused on phenotypical changes that occurred in plaque such as macrophage accumulation and lesion-associated macrophage lipids. They also occurred at a more prolonged time scale post-irradiation, spanning many months rather than days to weeks.

In conclusion, our observations are in line with the reported atheroprotective effect of low dose ionizing radiation observed in previous studies.^{16,18,19,67} This study provides new information pertaining to the impact on monocyte populations and aortic gene expression that further explain how LDIR may mitigate lesion formation. We used a wide range of doses and 2 dose-rates to demonstrate how diverse the responses to these stimuli can be. Low dose ionizing radiation clearly has modulating effects on different cell populations. Further testing of monocyte and lymphocyte subsets regulated by radiation could bring a better understanding of immune processes in the context of atherosclerosis.

Author Contributions

Lehoux and T.G. Ebrahimian are shared senior authors on this work.

Acknowledgments

The authors wish to thank K.Tack and D.Klokov for their support as a laboratory head and also thank D.Denis, M.Sebastien, R.Granger, S.Sache for experimental care.


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by grants from Electricité de France (EDF) to B.Leguen, from the Canadian Institutes of Health Research (CIHR) to S.Lehoux, and from the Swedish Radiation Protection Authority.

ORCID iD

T.G. Ebrahimian  <https://orcid.org/0000-0001-8413-4679>

Supplemental Material

Supplemental material for this article is available online.

References

1. Mulrooney DA, Armstrong GT, Huang S, et al. Cardiac outcomes in adult survivors of childhood cancer exposed to cardiotoxic therapy: a cross-sectional study. *Ann Int Med.* 2016;164(2): 93-101.

2. Boivin J-F, Hutchison GB, Lubin JH, Mauch P. Coronary artery disease mortality in patients treated for Hodgkin's disease. *Cancer*. 1992;69(5):1241-1247.
3. McGale P, Darby SC, Hall P, et al. Incidence of heart disease in 35,000 women treated with radiotherapy for breast cancer in Denmark and Sweden. *Radiother Oncol*. 2011;100(2):167-175.
4. Shimizu Y, Kodama K, Nishi N, et al. Radiation exposure and circulatory disease risk: Hiroshima and Nagasaki atomic bomb survivor data, 1950-2003. *BMJ*. 2010;340:b5349.
5. Yuan R, Sun Z, Cai J, et al. A novel anticancer therapeutic strategy to target autophagy accelerates radiation-associated atherosclerosis. *Int J Radiat Oncol Biol Phys*. 2021;109(2):540-552. doi:10.1016/j.ijrobp.2020.09.007
6. Lee C-L, Lee JW, Daniel AR, et al. Characterization of cardiovascular injury in mice following partial-heart irradiation with clinically relevant dose and fractionation. *Radiother Oncol*. 2021;157:155-162. doi:10.1016/j.radonc.2021.01.023
7. Hamada N, Kawano K-I, Yusoff FM, et al. Ionizing irradiation induces vascular damage in the aorta of wild-type mice. *Cancers (Basel)*. 2020;12(10):3030. doi:10.3390/cancers12103030
8. Stewart FA, Heeneman S, Te Poele J, et al. Ionizing radiation accelerates the development of atherosclerotic lesions in ApoE^{-/-} mice and predisposes to an inflammatory plaque phenotype prone to hemorrhage. *Am J Pathol*. 2006;168(2):649-658. doi:10.2353/ajpath.2006.050409
9. Newman AA, Baylis RA, Hess DL, et al. Irradiation abolishes smooth muscle investment into vascular lesions in specific vascular beds. *JCI Insight*. 2018;3(15):e121017. doi:10.1172/jci.insight.121017
10. Gabriels K, Hoving S, Seemann I, et al. Local heart irradiation of ApoE^(-/-) mice induces microvascular and endocardial damage and accelerates coronary atherosclerosis. *Radiother Oncol*. 2012;105(3):358-364. doi:10.1016/j.radonc.2012.08.002
11. Hoving S, Heeneman S, Gijbels MJJ, et al. Single-dose and fractionated irradiation promote initiation and progression of atherosclerosis and induce an inflammatory plaque phenotype in ApoE^(-/-) mice. *Int J Radiat Oncol Biol Phys*. 2008;71(3):848-857. doi:10.1016/j.ijrobp.2008.02.031
12. Schöllnberger H, Eidemüller M, Cullings HM, Simonetto C, Neff F, Kaiser JC. Dose-responses for mortality from cerebrovascular and heart diseases in atomic bomb survivors: 1950–2003. *Radiat Environ Biophys*. 2018;57(1):17-29. doi:10.1007/s00411-017-0722-5
13. Azizova TV, Muirhead CR, Druzhinina MB, et al. Cardiovascular diseases in the cohort of workers first employed at mayak PA in 1948–1958. *Radiat Res*. 2010;174(2):155-168. doi:10.1667/RR1789.1
14. Baselet B, Rombouts C, Benotmane AM, Baatout S, Aerts A. Cardiovascular diseases related to ionizing radiation: the risk of low-dose exposure (review). *Int J Molecul Med*. 2016;38(6):1623-1641. doi:10.3892/ijmm.2016.2777
15. Joven J, Rull A, Ferré N, et al. The results in rodent models of atherosclerosis are not interchangeable: the influence of diet and strain. *Atherosclerosis*. 2007;195(2):e85-e92.
16. Mitchel REJ, Hasu M, Bugden M, et al. Low-dose radiation exposure and atherosclerosis in ApoE^{-/-} mice. *Radiat Res*. 2011;175(5):665-676.
17. Mancuso M, Pasquali E, Braga-Tanaka I, et al. Acceleration of atherogenesis in ApoE^{-/-} mice exposed to acute or low-dose-rate ionizing radiation. *Oncotarget*. 2015;6(31):31263-31271. doi:10.18632/oncotarget.5075
18. Le Gallic C, Phalente Y, Manens L, et al. Chronic Internal Exposure to Low Dose ¹³⁷Cs Induces Positive Impact on the Stability of Atherosclerotic Plaques by Reducing Inflammation in ApoE^{-/-} Mice. *PLoS One*. 2015;10(6):e0128539. doi:10.1371/journal.pone.0128539
19. Ebrahimian TG, Beugnies L, Surette J, et al. Chronic exposure to external low-dose gamma radiation induces an increase in anti-inflammatory and anti-oxidative parameters resulting in atherosclerotic plaque size reduction in ApoE^{-/-} Mice. *Radiat Res*. 2018;189(2):187-196. doi:10.1667/RR14823.1
20. Williams J, Chen Y, Rubin P, Finkelstein J, Okunieff P. The biological basis of a comprehensive grading system for the adverse effects of cancer treatment. *Seminars in Radiat Oncol*. 2003;13(3):182-188. doi:10.1016/S1053-4296(03)00045-6
21. Gabriels K, Hoving S, Gijbels MJ, et al. Irradiation of existing atherosclerotic lesions increased inflammation by favoring pro-inflammatory macrophages. *Radiother Oncol*. 2014;110(3):455-460. doi:10.1016/j.radonc.2014.01.006
22. Seegenschmiedt MH, Makoski H-B, Trott K-R, Brady LW, eds. *Radiotherapy for Non-Malignant Disorders*. Springer; 2008. doi:10.1007/978-3-540-68943-0
23. Rodel F, Frey B, Gaipf U, et al. Modulation of inflammatory immune reactions by low-dose ionizing radiation: molecular mechanisms and clinical application. *CMC*. 2012;19(12):1741-1750. doi:10.2174/092986712800099866
24. Yuan Y, Lee SH, Wu S. The role of ROS in ionizing radiation-induced VLA-4 mediated adhesion of RAW264.7 cells to VCAM-1 under flow conditions. *Radiat Res*. 2013;179(1):62-68. doi:10.1667/RR3119.1
25. Kern PM, Keilholz L, Forster C, Hallmann R, Herrmann M, Seegenschmiedt MH. Low-dose radiotherapy selectively reduces adhesion of peripheral blood mononuclear cells to endothelium in vitro. *Radiother Oncol*. 2000;54(3):273-282. doi:10.1016/s0167-8140(00)00141-9
26. Cho S-J, Kang H, Hong E-H, Kim JY, Nam SY. Transcriptome analysis of low-dose ionizing radiation-impacted genes in CD4⁺ T-cells undergoing activation and regulation of their expression of select cytokines. *J Immunotoxicol*. 2018;15(1):137-146. doi:10.1080/1547691X.2018.1521484
27. Saigusa R, Winkels H, Ley K. T cell subsets and functions in atherosclerosis. *Nat Rev Cardiol*. 2020;17(7):387-401. doi:10.1038/s41569-020-0352-5
28. Narasimhan PB, Marcovecchio P, Hamers AAJ, Hedrick CC. Nonclassical Monocytes in Health and Disease. *Annu Rev Immunol*. 2019;37:439-456. doi:10.1146/annurev-immunol-042617-053119
29. Tacke F, Alvarez D, Kaplan TJ, et al. Monocyte subsets differentially employ CCR2, CCR5, and CX3CR1 to accumulate within

- atherosclerotic plaques. *J Clin Invest.* 2007;117(1):185-194. doi:10.1172/JCI28549
30. Klingenberg R, Gerdes N, Badeau RM, et al. Depletion of FOXP3+ regulatory T cells promotes hypercholesterolemia and atherosclerosis. *J Clin Invest.* 2013;123(3):1323-1334. doi:10.1172/JCI63891
 31. Rahman MS, Woollard K. Atherosclerosis. *Adv Exp Med Biol.* 2017;1003:121-144. doi:10.1007/978-3-319-57613-8_7
 32. Virmani R, Kolodgie FD, Burke AP, et al. Atherosclerotic plaque progression and vulnerability to rupture: angiogenesis as a source of intraplaque hemorrhage. *Arterioscler Thromb vasc Biol.* 2005;25(10):2054-2061.
 33. Moulton KS, Vakili K, Zurakowski D, et al. Inhibition of plaque neovascularization reduces macrophage accumulation and progression of advanced atherosclerosis. *Proc Natl Acad Sci.* 2003;100(8):4736-4741.
 34. Newby AC. Dual role of matrix metalloproteinases (matrixins) in intimal thickening and atherosclerotic plaque rupture. *Physiol Rev.* 2005;85(1):1-31.
 35. Fuster V, Moreno PR, Fayad ZA, Corti R, Badimon JJ. Atherothrombosis and high-risk plaque: part I: evolving concepts. *J Am Coll Cardiol.* 2005;46(6):937-954.
 36. Swirski FK, Libby P, Aikawa E, et al. Ly-6Chi monocytes dominate hypercholesterolemia-associated monocytosis and give rise to macrophages in atheromata. *J Clin Invest.* 2007;117(1):195-205. doi:10.1172/JCI29950
 37. El-Saghire H, Michaux A, Thierens H, Baatout S. Low doses of ionizing radiation induce immune-stimulatory responses in isolated human primary monocytes. *Int J Mol Med.* 2013;32(6):1407-1414. doi:10.3892/ijmm.2013.1514
 38. Frischholz B, Wunderlich R, Rühle P-F, et al. Reduced secretion of the inflammatory cytokine IL-1 β by stimulated peritoneal macrophages of radiosensitive Balb/c mice after exposure to 0.5 or 0.7 Gy of ionizing radiation. *Autoimmunity.* 2013;46(5):323-328. doi:10.3109/08916934.2012.747522
 39. Tsukimoto M, Homma T, Mutou Y, Kojima S. 0.5 Gy gamma radiation suppresses production of TNF-alpha through up-regulation of MKP-1 in mouse macrophage RAW264.7 cells. *Radiat Res.* 2009;171(2):219-224. doi:10.1667/RR1351.1
 40. Teresa Pinto A, Laranjeiro Pinto M, Patricia Cardoso A, et al. Ionizing radiation modulates human macrophages towards a pro-inflammatory phenotype preserving their pro-invasive and pro-angiogenic capacities. *Sci Rep.* 2016;6:18765. doi:10.1038/srep18765
 41. Sharma S, Singla N, Chadha VD, Dhawan D. A concept of radiation hormesis: stimulation of antioxidant machinery in rats by low dose ionizing radiation. *Hell J Nucl Med.* 2019;22(1):43-48.
 42. Zhou X, Nicoletti A, Elhage R, Hansson GK. Transfer of CD4(+) T cells aggravates atherosclerosis in immunodeficient apolipoprotein e knockout mice. *Circulation.* 2000;102(24):2919-2922. doi:10.1161/01.cir.102.24.2919
 43. Robertson A-KL, Rudling M, Zhou X, Gorelik L, Flavell RA, Hansson GK. Disruption of TGF- β signaling in T cells accelerates atherosclerosis. *J Clin Invest.* 2003;112(9):1342-1350. doi:10.1172/JCI18607
 44. Tabas I, Lichtman AH. Monocyte-Macrophages and T Cells in Atherosclerosis. *Immunity.* 2017;47(4):621-634. doi:10.1016/j.immuni.2017.09.008
 45. Liu S, Zhang Y, Zhao Y. Signal transduction in lymphocytes after low dose radiation. *Chin Med J.* 1994;107(6):431-436.
 46. Pandey R, Shankar BS, Sharma D, Sainis KB. Low dose radiation induced immunomodulation: effect on macrophages and CD8+ T cells. *Int J Radiat Biol.* 2005;81(11):801-812.
 47. Ishii K, Yamaoka K, Hosoi Y, Ono T, Sakamoto K. Enhanced mitogen-induced proliferation of rat splenocytes by low-dose whole-body X-irradiation. *Physiol Chem phys Med NMR.* 1995;27(1):17-23.
 48. Liu S, Han Z, Liu W. Changes in lymphocyte reactivity to modulatory factors following low dose ionizing radiation. *Biomed Environ Sci: BES.* 1994;7(2):130-135.
 49. Song K-H, Kim M-H, Kang S-M, et al. Analysis of immune cell populations and cytokine profiles in murine splenocytes exposed to whole-body low-dose irradiation. *Int J Radiat Biol.* 2015;91(10):795-803.
 50. Wang B, Li B, Dai Z, et al. Low-dose splenic radiation inhibits liver tumor development of rats through functional changes in CD4+ CD25+ Treg cells. *Int J Biochem Cell Biol.* 2014;55:98-108.
 51. Liu R, Xiong S, Zhang L, Chu Y. Enhancement of antitumor immunity by low-dose total body irradiation associated with selectively decreasing the proportion and number of T regulatory cells. *Cell Mol Immunol.* 2010;7(2):157-162.
 52. Nakatsukasa H, Tsukimoto M, Tokunaga A, Kojima S. Repeated gamma irradiation attenuates collagen-induced arthritis via up-regulation of regulatory T cells but not by damaging lymphocytes directly. *Radiat Res.* 2010;174(3):313-324.
 53. Park BS, Hong GU, Ro JY. Foxp3(+)-Treg cells enhanced by repeated low-dose gamma-irradiation attenuate ovalbumin-induced allergic asthma in mice. *Radiat Res.* 2013;179(5):570-583. doi:10.1667/RR3082.1
 54. Kachikwu EL, Iwamoto KS, Liao Y-P, et al. Radiation enhances regulatory T cell representation. *Int J Radiat Oncol Biol Phys.* 2011;81(4):1128-1135. doi:10.1016/j.ijrobp.2010.09.034
 55. Chen JC, Davis BH, Leon MA, Leong LC. Gamma radiation induces CD69 expression on lymphocytes. *Cytometry.* 1997;30(6):304-312.
 56. Song K-H, Jung S-Y, Park J-I, et al. Evaluation of anti-tumor effects of whole-body low-dose irradiation in metastatic mouse models. *Cancers (Basel).* 2020;12(5):1126. doi:10.3390/cancers12051126
 57. Shin SC, Lee K-M, Kang YM, et al. Alteration of cytokine profiles in mice exposed to chronic low-dose ionizing radiation. *Biochem Biophys Res Commun.* 2010;397(4):644-649.
 58. Mathias D, Mitchel REJ, Barclay M, et al. Low-dose irradiation affects expression of inflammatory markers in the heart of ApoE^{-/-} mice. *PLoS One.* 2015;10(3):e0119661. doi:10.1371/journal.pone.0119661
 59. Ley K, Laudanna C, Cybulsky MI, Nourshargh S. Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nat Rev Immunol.* 2007;7(9):678-689.

60. Ting HJ, Stice JP, Schaff UY, et al. Triglyceride-rich lipoproteins prime aortic endothelium for an enhanced inflammatory response to tumor necrosis factor- α . *Circ Res*. 2007;100(3):381-390.
61. Nageh MF, Sandberg ET, Marotti KR, et al. Deficiency of inflammatory cell adhesion molecules protects against atherosclerosis in mice. *Arterioscler Thromb Vasc Biol*. 1997;17(8):1517-1520.
62. Kamari Y, Shaish A, Shemesh S, et al. Reduced atherosclerosis and inflammatory cytokines in apolipoprotein-E-deficient mice lacking bone marrow-derived interleukin-1 α . *Biochem Biophys Res Commun*. 2011;405(2):197-203.
63. Kirii H, Niwa T, Yamada Y, et al. Lack of interleukin-1 β decreases the severity of atherosclerosis in ApoE-deficient mice. *Arterioscler Thromb Vasc Biol*. 2003;23(4):656-660.
64. Freigang S, Ampenberger F, Weiss A, et al. Fatty acid-induced mitochondrial uncoupling elicits inflammasome-independent IL-1 α and sterile vascular inflammation in atherosclerosis. *Nat Immun*. 2013;14(10):1045-1053.
65. Lutgens E, Daemen MJ. *Transforming Growth Factor- β : A Local or Systemic Mediator of Plaque Stability?* Am Heart Assoc; 2001.
66. Schröder S, Juerß D, Kriesen S, Manda K, Hildebrandt G. Immunomodulatory properties of low-dose ionizing radiation on human endothelial cells. *Int J Radiat Biol*. 2019;95(1):23-32.
67. Roedel F, Kley N, Beuscher HU, et al. Anti-inflammatory effect of low-dose X-irradiation and the involvement of a TGF-beta1-induced down-regulation of leukocyte/endothelial cell adhesion. *Int J Radiat Biol*. 2002;78(8):711-719. doi:10.1080/09553000210137671