



**HAL**  
open science

# Fast, slow convergence, and concentration in the house of cards replicator-mutator model

Bertrand Cloez, Pierre Gabriel

► **To cite this version:**

Bertrand Cloez, Pierre Gabriel. Fast, slow convergence, and concentration in the house of cards replicator-mutator model. 2022. hal-03608758v1

**HAL Id: hal-03608758**

**<https://hal.science/hal-03608758v1>**

Preprint submitted on 15 Mar 2022 (v1), last revised 18 Nov 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FAST, SLOW CONVERGENCE, AND CONCENTRATION IN THE HOUSE OF CARDS REPLICATOR-MUTATOR MODEL

BERTRAND CLOEZ AND PIERRE GABRIEL

ABSTRACT. We propose a fine analysis of the various possible long time behaviours of the solutions of the replicator-mutator equation with so-called Kingman’s house of cards mutations. In particular, we give what is to our knowledge the first concentration result for this model.

## CONTENTS

1. Introduction	1
2. A related conservative equation	6
2.1. Geometric convergence	9
2.2. Algebraic convergence	11
2.3. Weak convergence	14
3. The non-conservative linear equation	16
4. The nonlinear conservative equation	19
Acknowledgments	22
References	22

## 1. INTRODUCTION

We are interested in the long time behavior of the following non-linear integro-differential equation

$$\partial_t v_t(x) + a(x)v_t(x) = Q(x) \int_{\mathcal{X}} v_t(y)dy + v_t(x) \int_{\mathcal{X}} (a(y) - 1)v_t(y)dy \quad (1.1)$$

defined for  $t > 0$  and  $x \in \mathcal{X}$ , a measurable subset of  $\mathbb{R}^d$ , and complemented with an initial condition  $v_0(x)$ . The function  $Q(x)$  is measurable and strictly positive almost everywhere on  $\mathcal{X}$ , and the function  $a(x)$  is bounded below and continuous on  $\mathcal{X}$ .

This equation is a particular case of Kimura’s replicator-mutator model in evolutionary biology where  $\mathcal{X}$  stands for a set of phenotypic traits [42], and we refer to [25, 59] for a rigorous derivation from individual based models. The quantity  $-a(x)$  represents the fitness of the phenotype  $x$ , namely the difference between the birth and death rates. In the general model, the mutation term is  $\int_{\mathcal{X}} K(x, y)v(t, y)dy$ , where  $K(x, y)$  represents the creation rate of individuals with trait  $x$  from individual with trait  $y$ . The particular case  $K(x, y) = Q(x)$ , where the  $x$  distribution is the same whatever the original trait  $y$ , is known as the *house of cards* model of mutations after the work of Kingman [43], see [14, 13, 15, 18, 20, 58]. In [43], Kingman neglects the small mutations and only takes into account the mutations that have a significant effect on the population. The latter, often deleterious, destroy the biochemical ‘house of cards’ created by evolution. We are interested here in this model which, despite its apparent simplicity, captures the main features

---

2010 *Mathematics Subject Classification.* 45K05, 45M05, 92D15.

*Key words and phrases.* evolutionary genetics; selection-mutation; house of cards model; long time behavior; concentration phenomenon.

of the general case and in particular the possible concentration phenomenon, see below. The quadratic term in Equation (1.1) can be seen as a Lagrange multiplier ensuring that, if initially a probability distribution  $v_0(x)$ , then for all  $t > 0$  the solution  $v_t(x)$  is a probability distribution which represents the relative frequency of the traits in the population. A consequence of this conservativeness property is that the probability density solutions are insensitive to the addition of a constant to the fitness function  $a(x)$ . We will thus assume that this lower bounded function is actually nonnegative. This emphasizes that Equation (1.1) does not belong to the class of logistic type selection-mutation models which appear in evolutionary ecology. In these models, the competition for resources leads to negative quadratic terms, see for instance [10, 17, 19, 31, 35, 47, 48, 50].

Our aim is to give a precise description of the long time behavior of Equation (1.1) in the case when the fitness  $a(x)$  reaches its minimum – which can be assumed to be zero – at a unique point which, up to a translation of  $\mathcal{X}$ , can be assumed without loss of generality to be the origin. We thus make the following hypotheses on  $\mathcal{X}$ ,  $a$ , and  $Q$ :

- (HX) The trait space  $\mathcal{X}$  is a measurable subset of  $\mathbb{R}^d$  which contains a neighborhood of 0.
- (HQ) The mutation kernel  $Q : \mathcal{X} \rightarrow (0, \infty)$  is a positive measurable and integrable function.
- (Ha) The fitness  $a : \mathcal{X} \rightarrow [0, \infty)$  is a continuous function which satisfies

$$a(0) = 0, \quad a(x) > 0 \text{ for all } x \in \mathcal{X} \setminus \{0\}, \quad \text{and} \quad \int_{\mathcal{X} \cap \{|x| > 1\}} \frac{Q(x)}{a^2(x)} dx < \infty.$$

The last condition on  $a$  prevents the escape to infinity, a phenomenon that can occur in the replicator-mutator model [4, 5, 57].

Equation (1.1) is strongly related to the non-conservative linear equation

$$\partial_t u_t(x) = \mathcal{A}u_t(x) = -a(x)u_t(x) + Q(x) \int_{\mathcal{X}} u_t(y) dy. \quad (1.2)$$

For any nonnegative and non identically zero solution  $u(t, x)$  of Equation (1.2), the function

$$v_t(x) = \frac{u_t(x)}{\int_{\mathcal{X}} u_t(y) dy}$$

is solution to Equation (1.1). Reciprocally, if  $v(t, x)$  satisfies Equation (1.1), then the function

$$u_t(x) = v_t(x) \exp\left(\int_0^t \int_{\mathcal{X}} (1 - a(y)) v_s(y) dy ds\right)$$

verifies Equation (1.2). Also, finding a stationary probability distribution for Equation (1.1) is equivalent to find  $\lambda \in \mathbb{R}$  and a probability measure  $\gamma$  on  $\mathcal{X}$  such that  $\mathcal{A}\gamma = \lambda\gamma$ . This Perron-Frobenius eigenproblem can be easily solved and the explicit expression of the unique solution depends on whether the parameter

$$\rho = \int_{\mathcal{X}} \frac{Q(x)}{a(x)} dx,$$

which can be infinite, is larger or smaller than 1. More precisely, if  $\rho \geq 1$  then  $\gamma$  has a Lebesgue density given by

$$\gamma(dx) = \frac{Q(x)}{\lambda + a(x)} dx,$$

where  $\lambda \geq 0$  is the unique real number such that  $\int_{\mathcal{X}} \frac{Q}{\lambda + a} = 1$ . If  $\rho < 1$ , then  $\lambda = 0$  and  $\gamma$  has an atom at zero:

$$\gamma(dx) = (1 - \rho)\delta_0 + \frac{Q(x)}{a(x)} dx.$$

These computations are made in [15, 27, 28, 43] and the presence of a Dirac mass in the case  $\rho < 1$  suggests a concentration phenomenon, supported by numerical evidences [11], similarly as in [43]. However, to the best of our knowledge, there is no proof of convergence to a singular measure for the solutions of Equation (1.1) in the literature. Providing such a result is the main purpose of the present paper, but we also prove new convergence estimates in the non-singular case  $\rho \geq 1$ .

For characterizing the long time behavior of the solutions, the dual Perron eigenvalue problem is helpful. It consists in finding a non-negative and non-zero function  $h$  such that  $\mathcal{A}^*h = \lambda h$ , where the dual operator  $\mathcal{A}^*$  is given by

$$\mathcal{A}^*f(x) = -a(x)f(x) + \int_{\mathcal{X}} f(y)Q(y)dy. \quad (1.3)$$

Similarly as for the direct problem, the solutions can be computed explicitly. In the case  $\rho > 1$  they are positive and given by

$$h(x) = \frac{\alpha^{-1}}{\lambda + a(x)}$$

with  $\alpha$  any positive constant. The convenient choice we make is to take  $\alpha = \int_{\mathcal{X}} \frac{Q}{(\lambda+a)^2}$  so that  $h$  is the unique eigenfunction such that  $\langle \gamma, h \rangle = \int_{\mathcal{X}} h(x)\gamma(dx) = 1$ . In the case  $\rho < 1$  there is no strictly positive eigenfunction but only degenerated ones which are zero everywhere and positive at zero. Due to the singularity of  $\gamma$  in this case, we can nevertheless define a unique such eigenfunction  $h$  such that  $\langle \gamma, h \rangle = 1$  by setting

$$h(x) = \begin{cases} \frac{1}{1-\rho} & \text{if } x = 0, \\ 0 & \text{otherwise.} \end{cases}$$

In the critical case  $\rho = 1$ , both the functions  $\frac{1}{a}$  and  $\mathbf{1}_{\{0\}}$  are formally eigenfunctions of  $\mathcal{A}^*$ . Yet, the relevance of these eigenfunctions depends on the choice of the Banach space we consider. If studying Equation (1.2) in the  $L^1$  Lebesgue space associated to the measure  $\frac{dx}{a(x)}$ , then the eigenfunction  $\mathbf{1}_{\{0\}}$  corresponds to the zero linear form and the only non-trivial eigenfunction is  $\frac{1}{a}$ , which can be normalized to have  $\langle \gamma, h \rangle = 1$  provided that  $\int_{\mathcal{X}} \frac{Q}{a^2} < \infty$ . In contrast, if we work in a space of measures that contains the Dirac mass  $\delta_0$ , then  $\frac{1}{a}$  is not a bounded linear form on this space and  $\mathbf{1}_{\{0\}}$  is the relevant eigenfunction.

We work in various Banach spaces that we recall here. For a positive weight function  $\varphi$  on  $\mathcal{X}$  and  $p \in [1, \infty)$  we denote by  $L^p(\varphi) = L^p(\mathcal{X}, \varphi(x)dx)$  the standard Lebesgue space associated to the measure  $\varphi(x)dx$ . When  $\varphi = \mathbf{1}$ , the constant function equal to 1, we use the shorthand  $L^p$  for  $L^p(\mathbf{1})$  and  $L^\infty$  for  $L^\infty(\mathcal{X}, dx)$  endowed with the norm  $\|f\|_{L^\infty} = \text{ess sup}_{\mathcal{X}} |f|$ . For any  $p \in [1, \infty]$ , we use the standard notation  $p' \in [1, \infty]$  for the Hölder conjugate exponent, *i.e.* such that  $1/p + 1/p' = 1$ . We denote by  $\mathcal{M}(\mathcal{X})$  the space of finite signed measures on  $\mathcal{X}$ , endowed with the total variation norm  $\|\mu\|_{\text{TV}} = \mu_+(\mathcal{X}) + \mu_-(\mathcal{X}) = \sup_{\|f\|_\infty \leq 1} \langle \mu, f \rangle$  where the supremum is taken over the measurable or continuous functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  such that  $\|f\|_\infty = \sup_{\mathcal{X}} |f| \leq 1$  and the duality bracket is given by  $\langle \mu, f \rangle := \int_{\mathcal{X}} f d\mu$ . Due to the canonical injection  $L^1 \hookrightarrow \mathcal{M}(\mathcal{X})$ , we will often abuse notations and identify  $L^1$  functions to their associated Lebesgue density measure, and in particular for  $\mu \in L^1$ , we will use the notation  $d\mu$  for the measure  $\mu(dx) = \mu(x)dx$ . We denote by  $\mathcal{P}(\mathcal{X}) \subset \mathcal{M}(\mathcal{X})$  the subset of probability measures, and for a positive weight function  $\varphi$  we denote by  $\mathcal{M}(\varphi)$  the space of signed measures  $\mu$  such that  $\|\mu\|_{\mathcal{M}(\varphi)} = \langle \mu_+, \varphi \rangle + \langle \mu_-, \varphi \rangle = \sup_{\|f/\varphi\|_\infty \leq 1} \langle \mu, f \rangle < \infty$ . We also recall that vague convergence of measures means convergence for any test function in  $C_c(\mathcal{X})$  while narrow convergence means convergence for any test function in  $C_b(\mathcal{X})$ .

Our main results are summarized in the two following theorems. The first one is about the linear equation (1.2).

**Theorem 1.1.** *We suppose that (HX)-(HQ)-(Ha) are met. Then the following results hold:*

(1) **Exponential ergodicity.** *Assume that  $\rho = \int_X Q/a \in (1, +\infty]$ . Then  $\lambda > 0$  and*

(a) *For all  $u_0 \in \mathcal{M}(h)$  and all  $t \geq 0$ ,*

$$\|e^{-\lambda t} u_t - \langle u_0, h \rangle \gamma\|_{\mathcal{M}(h)} \leq e^{-\lambda t} \|u_0 - \langle u_0, h \rangle \gamma\|_{\mathcal{M}(h)}.$$

(b) *For all  $p \in [1, 2]$ , all  $u_0 \in L^p(\gamma^{1-p}h)$ , and all  $t \geq 0$ ,*

$$\|e^{-\lambda t} u_t - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)} \leq e^{-\lambda t} \|u_0 - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)}.$$

(c) *For all  $p > 2$ , all  $u_0 \in L^p(\gamma^{1-p}h)$ , and all  $t \geq 0$ ,*

$$\|e^{-\lambda t} u_t - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)} \leq 2 e^{-\lambda t} \|u_0 - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)}.$$

(d) *For all  $u_0 \in L^1(h)$  such that  $u_0/\gamma \in L^\infty$  and all  $t \geq 0$ ,*

$$\|e^{-\lambda t} u_t/\gamma - \langle u_0, h \rangle\|_{L^\infty} \leq 2 e^{-\lambda t} \|u_0/\gamma - \langle u_0, h \rangle\|_{L^\infty}.$$

(2) **Slow convergence and unboundedness.** *Assume that  $\rho = 1$  and  $1/a \in L^2(Q)$ . Then*

(a) *For all  $u_0 \in \mathcal{M}(h)$ ,*

$$u_t \xrightarrow[t \rightarrow +\infty]{} \langle u_0, h \rangle \gamma$$

*in the vague topology, and in the narrow topology if additionally  $a \in L^\infty$ .*

(a') *For  $u_0 = \delta_0$  we have for all  $f \in C_c(X)$ , and for all  $f \in C_b(X)$  if additionally  $a \in L^\infty$ ,*

$$\langle u_t, f \rangle \sim \|1/a\|_{L^2(Q)}^{-2} \langle \gamma, f \rangle t \quad \text{as } t \rightarrow +\infty.$$

(b) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$  and  $a \in L^\infty$ , then there exists  $C > 0$  such that for all  $u_0 \in \mathcal{M}((1+a^{-q})h)$  and all  $t \geq 0$ ,*

$$\|u_t - \langle u_0, h \rangle \gamma\|_{\mathcal{M}(h)} \leq C t^{1-q} \|u_0\|_{\mathcal{M}((1+a^{-q})h)}.$$

(c) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$ , then there exists  $C > 0$  such that for any  $u_0 \in L^1(h)$  such that  $u_0/\gamma \in L^\infty$ , any  $p \in [1, \infty)$ , and all  $t \geq 0$ ,*

$$\|u_t - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)} \leq C t^{-\frac{q-1}{p}} \|u_0/\gamma - \langle u_0, h \rangle\|_{L^\infty}.$$

(d) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$  and, for  $r > p$ , either  $p \in [1, 2]$  and  $a \in L^{2+r'}(Q)$ , or  $p \geq 2$  and  $a \in L^{2+p}(Q)$ , then there exists  $C > 0$  such that for all  $u_0 \in L^r(\gamma^{r-1}h)$  and all  $t \geq 0$ ,*

$$\|u_t - \langle u_0, h \rangle \gamma\|_{L^p(\gamma^{1-p}h)} \leq C t^{-(q-1)(1/p-1/r)} \|u_0 - \langle u_0, h \rangle\|_{L^r(\gamma^{r-1}h)}.$$

(3) **Degenerate convergence.**

(a) *If either  $\rho = 1$  and  $1/a \notin L^2(Q)$ , or if  $\rho < 1$ , then for all  $u_0 \in \mathcal{M}(1/a)$ , the convergence*

$$u_t \xrightarrow[t \rightarrow +\infty]{} 0$$

*holds in the vague topology, and in the narrow topology if additionally  $a \in L^\infty$ .*

(b) *If  $\rho < 1$ , then for all  $u_0 \in \mathcal{M}(X)$  we have in the narrow topology*

$$u_t \xrightarrow[t \rightarrow +\infty]{} \langle u_0, h \rangle \gamma = u_0(\{0\}) \left( \delta_0 + \frac{1}{1-\rho} \frac{Q}{a} \right).$$

It is worth precisising that for the definition of  $\mathcal{M}(1/a)$  in the above theorem, we have set  $(1/a)(0) = +\infty$ . In particular a measure in  $\mathcal{M}(1/a)$  has no atom at zero. The second main theorem is about the replicator-mutator equation (1.1).

**Theorem 1.2.** *Under Hypotheses (HX)-(HQ)-(Ha), the following results hold:*

(1) **Fast convergence.** *Assume that  $\rho = \int_{\mathcal{X}} Q/a \in (1, +\infty]$ .*

(a) *If  $a \in L^\infty$ , then for any  $v_0 \in \mathcal{P}(\mathcal{X}) \cap \mathcal{M}(h)$  there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t - \gamma\|_{\mathcal{M}(h)} \leq Ce^{-\lambda t}.$$

(b) *If  $p \in [1, 2)$  and  $a \in L^{p'-2}(Q)$ , or if  $p \geq 2$ , then for any  $v_0 \in \mathcal{P}(\mathcal{X}) \cap L^p(\gamma^{1-p}h)$  there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t - \gamma\|_{L^p(\gamma^{1-p}h)} \leq Ce^{-\lambda t}.$$

(c) *For all  $v_0 \in \mathcal{P}(\mathcal{X})$  such that  $v_0/\gamma \in L^\infty$  there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t/\gamma - 1\|_{L^\infty} \leq Ce^{-\lambda t}.$$

(2) **Slow convergence.** *Assume that  $\rho = 1$ .*

(a) *For any  $v_0 \in \mathcal{P}(\mathcal{X})$ , we have in the narrow topology*

$$v_t \xrightarrow[t \rightarrow +\infty]{} \gamma.$$

(b) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$  and  $a \in L^\infty$ , then for any  $v_0 \in \mathcal{P}(\mathcal{X}) \cap \mathcal{M}((1+a^{-q})h)$  there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t - \gamma\|_{\mathcal{M}(h)} \leq Ct^{1-q}.$$

(c) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$  and either  $p \in [1, 2)$  with  $a \in L^{p'-2}(Q)$ , or  $p \geq 2$ , then for any  $v_0 \in \mathcal{P}(\mathcal{X})$  such that  $v_0/\gamma \in L^\infty$ , there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t - \gamma\|_{L^p(\gamma^{1-p}h)} \leq Ct^{-\frac{q-1}{p}}.$$

(d) *If  $1/a \in L^{1+q}(Q)$  for some  $q > 1$  and, for  $r > p$ , either  $p \in [1, 2)$  and  $a \in L^{2+r'}(Q) \cap L^{p'-2}(Q)$ , or  $p \geq 2$  and  $a \in L^{2+p}(Q)$ , then for all  $v_0 \in \mathcal{P}(\mathcal{X}) \cap L^r(\gamma^{r-1}h)$  there exists  $C > 0$  such that for all  $t \geq 0$ ,*

$$\|v_t - \gamma\|_{L^p(\gamma^{1-p}h)} \leq Ct^{-(q-1)(1/p-1/r)}.$$

(3) **Concentration.** *If  $\rho < 1$  and  $\lim_{|x| \rightarrow \infty} a(x) = +\infty$ , then for all  $v_0 \in \mathcal{P}(\mathcal{X})$*

$$\frac{1}{t} \int_0^t v_s ds \xrightarrow[t \rightarrow +\infty]{} \gamma = (1 - \rho)\delta_0 + \frac{Q}{a}$$

*in the narrow topology.*

To our knowledge, these results are new in the literature. The concentration phenomenon for Equation (1.1) in the case  $\rho < 1$ , which is proved here to occur in Cesàro mean, was expected since the works of Kingman [43], Bürger and Bomze [15] and more recently Coville *et al.* [11, 28], but never rigorously established until now. Concentration phenomena are very relevant in evolutionary biology. It was proved to occur for logistic type nonlinearities but only in the pure selection case [1, 2, 3, 29, 40, 45, 52, 55] or in the vanishing mutation regime [10, 46, 53, 54]. For Kimura's replicator-mutator equation, a result similar to ours is proved in [36, 37] for the specific case  $\mathcal{X} = \mathbb{R}$  with  $a(x) = x$  and purely deleterious convolutive mutations, *i.e.*  $K(x, y) = J(x - y)$  with  $\text{supp } J \subset (-\infty, 0]$ , by means of an explicit formulation of the solutions through Laplace

transform. This approach cannot be adapted to house of cards mutations and our proof rather uses the explicit expression of the stationary distribution.

The convergence of the solutions of Equation (1.1) in the cases  $\rho > 1$  and  $\rho = 1$  with  $1/a \in L^2(Q)$  are consequences of the convergence results of the linear equation (1.2). This scheme of proof has for instance been used in [30, 51]. Exponential convergences are deduced from functional inequalities for suitable entropies, in the spirit of [7, 8, 9, 22, 24]. Polynomial convergences also rely on the use of entropies, but with weaker and somewhat more original functional inequalities; see however the closely related approaches [12, 16, 41, 56]. In the cases  $\rho = 1$  with  $1/a \notin L^2(Q)$  and  $\rho < 1$ , the results on the linear equation are not enough for deriving the convergence of the nonlinear equation. We then work directly on Equation (1.1) and prove fine upper and lower bounds of the solutions when times goes to infinity.

Our results extend those of Kingman [43] to the time continuous setting and to general trait spaces  $\mathcal{X}$  and coefficients  $a$  and  $Q$ . The convergence in the case  $\rho \geq 1$  corresponds to what Kingman calls democracy; the effect of the selection is simply to modify the shape of the distribution. The concentration in the case  $\rho < 1$  is named meritocracy by Kingman; the Dirac mass emerges from the growth of a new class of highly fitted individuals, and the smooth part consists of the descendants of these mutants. Yet another regime is considered by Kingman, which is the case where  $\text{supp } Q \subset \{x \in \mathcal{X}, a(x) \geq \epsilon\}$  for some  $\epsilon > 0$ , and  $\text{supp } \nu_0 \cap \{x \in \mathcal{X}, a(x) < \epsilon\} \neq \emptyset$ . In this situation, called aristocracy, some initial individuals are always better adapted than the mutants. In our study we do not consider this non-irreducible case, where the long time behavior strongly depends on  $\nu_0$ .

The paper is organized as follows. We first analyse in Section 2 a conservative linear equation which is closely related to Equation (1.2). Then we use the results of Section 2 to prove Theorem 1.1 in Section 3. Finally, in Section 4, we prove Theorem 1.2 by taking advantage of the results in Theorem 1.1.

## 2. A RELATED CONSERVATIVE EQUATION

In this section, we focus on the closely related and simpler conservative equation given by

$$\partial_t u_t(x) = \mathcal{L}u_t(x) = -\mathbf{a}(x)u_t(x) + \left( \int_{\mathcal{X}} \mathbf{a}(y)u_t(y)dy \right) \mathbf{Q}(x), \quad x \in \mathcal{X}, \quad (2.1)$$

where  $\mathbf{Q}(x)dx$  is a probability measure and  $\mathbf{a}$  is a continuous nonnegative function such that

$$\forall x \in \mathcal{X} \setminus \{0\}, \mathbf{a}(x) > \mathbf{a}(0) \geq 0. \quad (2.2)$$

It is worth noticing that, unlike Equation (1.2) for which adding a constant to  $a$  only translates the spectrum to the left or to the right, adding a constant to  $\mathbf{a}$  really modifies the equation. We consequently do not assume that  $\mathbf{a}(0) = 0$ . At infinity, we impose the integrability condition

$$\int_{\mathcal{X} \cap \{|x|>1\}} \frac{\mathbf{Q}(x)}{\mathbf{a}(x)} dx < \infty. \quad (2.3)$$

Equation (2.1) is a pure mutation equation, *i.e.* a pure jump process, without birth nor death. It is thus conservative in the sense that it leaves invariant the set  $\mathcal{P}(\mathcal{X})$  of probability measures. Another way to see it is to remark that the constants are solutions to the dual equation

$$\partial_t \varphi_t(x) = \mathcal{L}^* \varphi_t(x) = \mathbf{a}(x) \left( \int_{\mathcal{X}} \varphi_t(y) \mathbf{Q}(dy) - \varphi_t(x) \right) \quad (2.4)$$

since  $\mathcal{L}^*\mathbf{1} = 0$ . The operator  $\mathcal{L}^*$  thus defines a Markov semigroup  $(P_t)_{t \geq 0}$  in  $\mathcal{L}^\infty(X)$  or  $L^\infty(X, dx)$ , *i.e.* a positive semigroup with the property that  $P_t\mathbf{1} = \mathbf{1}$ , through the Duhamel formula

$$P_t f(x) = f(x)e^{-\mathbf{a}(x)t} + \int_0^t \mathbf{a}(x)e^{-\mathbf{a}(x)s} \langle \mathbf{Q}, P_{t-s} f \rangle ds. \quad (2.5)$$

This means that  $\varphi_t = P_t f$  is the solution to Equation (2.4), in the sense that

$$\partial_t P_t f = \mathcal{L}^* P_t f = P_t \mathcal{L}^* f,$$

with initial datum  $\varphi_0 = f$ . We then define by duality the left action of  $P_t$  on  $\mathcal{M}(X)$  through

$$\langle \mu P_t, f \rangle = \langle \mu, P_t f \rangle \quad (2.6)$$

for  $\mu \in \mathcal{M}(X)$  and any  $f \in \mathcal{L}^\infty(X)$ , which yields the solutions of the direct equation (2.1), *i.e.*  $u_t = \mu P_t$  is the solution with initial datum  $u_0 = \mu$ .

When  $\mathbf{Q}/\mathbf{a}$  is integrable, we assume without loss of generality, by rescaling time, that

$$\pi(dx) = \frac{\mathbf{Q}(x)}{\mathbf{a}(x)} dx$$

is a probability measure. It is easily seen that  $\mathcal{L}\pi = 0$  and  $\pi$  is thus an invariant measure of  $(P_t)_{t \geq 0}$ . Moreover we have  $\mathcal{L}(f\pi) = (\mathcal{L}^* f)\pi$  for all  $f \in L^\infty$ , and consequently

$$P_t f = \frac{(f\pi)P_t}{\pi}. \quad (2.7)$$

This relation, that will be useful in our study, allows extending the right action of  $P_t$  to  $L^1(\pi)$ . We now introduce the so-called  $\Phi$ -entropies, see for instance [24]. For a convex function  $\Phi : \mathcal{I} \rightarrow \mathbb{R}$ , where  $\mathcal{I} \subset \mathbb{R}$ , a function  $f : X \rightarrow \mathcal{I}$  and a probability measure  $\mu$ , the  $\Phi$ -entropy associated to  $\mu$  is defined by

$$\text{Ent}_\mu^\Phi(f) = \int \Phi(f) d\mu - \Phi\left(\int f d\mu\right) = \langle \mu, \Phi(f) \rangle - \Phi(\langle \mu, f \rangle)$$

and is nonnegative due to Jensen's inequality. Besides, Jensen's inequality also guarantees that for all  $t \geq s \geq 0$  and all  $x \in X$ ,

$$\Phi(P_t f)(x) = \Phi(\langle \delta_x P_{t-s}, P_s f \rangle) \leq \langle \delta_x P_{t-s}, \Phi(P_s f) \rangle = P_{t-s} \Phi(P_s f)(x),$$

and consequently, since  $\pi P_{t-s} = \pi$ ,

$$\text{Ent}_\pi^\Phi(P_t f) \leq \text{Ent}_\pi^\Phi(P_s f).$$

In other words, any  $\Phi$ -entropy associated to  $\pi$  decreases along the solutions of Equation (2.4). In particular, taking  $\Phi(x) = |x|^p$  with  $p \geq 1$ , we get that  $L^p(\pi^{1-p})$  is invariant under the right action of  $(P_t)_{t \geq 0}$ . If we consider  $\Phi(x) = x \log x$ , we get that for any positive functions  $\mu \in \mathcal{P}(X) \cap L^1$ , the Kullback-Leibler divergence  $D_{\text{KL}}(\mu P_t \| \pi)$  from  $\mu P_t$  to  $\pi$ , where  $D_{\text{KL}}(f \| g)$  is defined by

$$D_{\text{KL}}(f \| g) = \int_X f(x) \log \left( \frac{f(x)}{g(x)} \right) dx,$$

is non-increasing along time. Indeed, from (2.7) we have  $\mu P_t = P_t \left( \frac{\mu}{\pi} \right) \pi$  and then  $D_{\text{KL}}(\mu P_t \| \pi) = \text{Ent}_\pi^{x \mapsto x \log x} \left( P_t \left( \frac{\mu}{\pi} \right) \right)$ . Note that  $D_{\text{KL}}$  is not a distance since it is not symmetric and it does not satisfy the triangle inequality, but it controls the  $L^1$  distance due to Pinsker's inequality

$$\|f - g\|_{L^1} \leq \sqrt{2D_{\text{KL}}(f \| g)}.$$

When  $\mathbf{a}(0) > 0$ ,  $\pi$  is the unique invariant probability measure of  $(P_t)_{t \geq 0}$ . In contrast, when  $\mathbf{a}(0) = 0$ , the Dirac mass  $\delta_0$  is also invariant. In the case where  $\mathbf{Q}/\mathbf{a}$  is not integrable, we can no longer normalize  $\pi$  to be a probability measure, but due to Assumptions (2.2) and (2.3) and the

continuity of  $\mathbf{a}$ , we necessarily have  $\mathbf{a}(0) = 0$  and  $\delta_0$  is the unique invariant probability measure. We thus have three distinct situations:

- ◇ if  $\int_X \mathbf{Q}/\mathbf{a} = 1$  and  $\mathbf{a}(0) > 0$ , then  $\pi$  is the unique invariant probability measure;
- ◇ if  $\int_X \mathbf{Q}/\mathbf{a} = 1$  and  $\mathbf{a}(0) = 0$ , then  $\pi$  and  $\delta_0$  are the two unique invariant probability measures;
- ◇ if  $\int_X \mathbf{Q}/\mathbf{a} = \infty$ , then  $\delta_0$  is the unique invariant probability measure.

The following theorem gives estimates about the stability of these invariant measures. Most of them will be useful for investigating Equation (1.2).

**Theorem 2.1.** *Under Assumptions (2.2) and (2.3) on  $\mathbf{a}$  and  $\mathbf{Q}$ , we have the following results:*

(1) **Fast convergence.** *Assume that  $\int_X \mathbf{Q}/\mathbf{a} = 1$  and  $\mathbf{a}(0) = \inf(\mathbf{a}) > 0$ . Then  $\pi = \mathbf{Q}/\mathbf{a}$  is the unique invariant probability measure of  $(P_t)_{t \geq 0}$  and we have*

(a) *For all  $\mu \in \mathcal{P}(X)$  and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{\text{TV}} \leq e^{-\inf(\mathbf{a})t} \|\mu - \pi\|_{\text{TV}}.$$

(b) *For all  $p \in [1, 2]$ , all  $\mu \in \mathcal{P}(X) \cap L^p(\pi^{1-p})$ , and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{L^p(\pi^{1-p})} \leq e^{-\inf(\mathbf{a})t} \|\mu - \pi\|_{L^p(\pi^{1-p})}.$$

(c) *For all  $p > 2$ , all  $\mu \in \mathcal{P}(X) \cap L^p(\pi^{1-p})$ , and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{L^p(\pi^{1-p})} \leq 2 e^{-\inf(\mathbf{a})t} \|\mu - \pi\|_{L^p(\pi^{1-p})}.$$

(d) *For all  $\mu \in \mathcal{P}(X) \cap L^1$  such that  $\mu/\pi \in L^\infty$  and all  $t \geq 0$ ,*

$$\|\mu P_t/\pi - \mathbf{1}\|_{L^\infty} \leq 2 e^{-\inf(\mathbf{a})t} \|\mu/\pi - \mathbf{1}\|_{L^\infty}.$$

(e) *For all  $\mu \in \mathcal{P}(X) \cap L^1$  such that  $D_{KL}(\mu|\pi) < \infty$  and all  $t \geq 0$ ,*

$$D_{KL}(\mu P_t|\pi) \leq e^{-\inf(\mathbf{a})t} D_{KL}(\mu|\pi).$$

(2) **Slow convergence.** *Assume that  $\int_X \mathbf{Q}/\mathbf{a} = 1$  and  $\mathbf{a}(0) = \inf(\mathbf{a}) = 0$ . Then  $\pi$  and  $\delta_0$  are the two invariant probability measures of  $(P_t)_{t \geq 0}$ . Moreover*

(a) *If  $\mathbf{a} \in L^1(\mathbf{Q})$ , then for all  $\mu \in \mathcal{P}(X)$*

$$\mu P_t \xrightarrow[t \rightarrow +\infty]{} \mu(\{0\}) \delta_0 + (1 - \mu(\{0\})) \pi$$

*in the narrow topology.*

(b) *If  $1/\mathbf{a} \in L^q(\mathbf{Q})$  for some  $q > 1$  and  $\mathbf{a} \in L^\infty$ , then there exists  $C > 0$  such that for all  $\mu \in \mathcal{P}(X) \cap \mathcal{M}(\mathbf{a}^{-q})$  and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{\text{TV}} \leq C t^{1-q} \|\mu\|_{\mathcal{M}(\mathbf{a}^{-q})}.$$

(c) *If  $1/\mathbf{a} \in L^q(\mathbf{Q})$  for some  $q > 1$ , then there exists  $C > 0$  such that for any  $\mu \in \mathcal{P}(X) \cap L^1$  such that  $\mu/\pi \in L^\infty$ , any  $p \in [1, \infty)$ , and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{L^p(\pi^{1-p})} \leq C t^{-\frac{q-1}{p}} \|\mu/\pi - \mathbf{1}\|_{L^\infty}.$$

(d) *If  $1/\mathbf{a} \in L^q(\mathbf{Q})$  for some  $q > 1$  and, for  $r > p$ , either  $p \in [1, 2]$  and  $\mathbf{a} \in L^{r'}(\pi)$ , or  $p \geq 2$  and  $\mathbf{a} \in L^p(\pi)$ , then there exists  $C > 0$  such that for all  $\mu \in L^r(\pi^{r-1})$  and all  $t \geq 0$ ,*

$$\|\mu P_t - \pi\|_{L^p(\pi^{1-p})} \leq C t^{-(q-1)(1/p-1/r)} \|\mu - \pi\|_{L^r(\pi^{1-r})}.$$

(3) **Concentration.** If  $1/\mathbf{a} \notin L^1(\mathbf{Q})$ , then  $\delta_0$  is the unique invariant probability measure of  $(P_t)_{t \geq 0}$ . If furthermore  $\mathbf{a} \in L^1(\mathbf{Q})$ , then for all  $\mu \in \mathcal{P}(\mathcal{X})$  we have the narrow convergence

$$\mu P_t \xrightarrow[t \rightarrow +\infty]{} \delta_0.$$

Note that the constants  $C$  in (2c) and (2d) are explicitly calculable, see the proofs in Section 2.2.

**2.1. Geometric convergence.** We treat here the case of a function  $\mathbf{a}$  such that

$$\inf_{\mathcal{X}} \mathbf{a} = \mathbf{a}(0) > 0 \quad \text{and} \quad \int_{\mathcal{X}} \frac{\mathbf{Q}}{\mathbf{a}} = 1. \quad (2.8)$$

We recall that in this case  $\pi = \mathbf{Q}/\mathbf{a}$  is the unique invariant probability measure of  $(P_t)_{t \geq 0}$ . The first result of Theorem 2.1-(1) is proved through a coupling approach. We refer to [26, 33, 34, 49] for details on coupling techniques.

*Proof of Theorem 2.1-(1a).* Consider  $(\mathbb{P}_t)$  the semigroup on  $\mathcal{X} \times \mathcal{X}$  generated by

$$\begin{aligned} \mathbb{L}f(x, y) = & \min(\mathbf{a}(x), \mathbf{a}(y)) \left( \int f(z, z) \mathbf{Q}(dz) - f(x, y) \right) \\ & + (\mathbf{a}(x) - \mathbf{a}(y))_+ \left( \int f(z, y) \mathbf{Q}(dz) - f(x, y) \right) + (\mathbf{a}(y) - \mathbf{a}(x))_+ \left( \int f(x, z) \mathbf{Q}(dz) - f(x, y) \right). \end{aligned}$$

This means that  $(\mathbb{P}_t)$  is the solution to

$$\partial_t \mathbb{P}_t = \mathbb{P}_t \mathbb{L} = \mathbb{L} \mathbb{P}_t.$$

This semigroup is a coupling of  $(P_t)$  because if  $f(x, y) = g(x)$  (resp.  $g(y)$ ) then  $\mathbb{P}_t f(x, y) = P_t g(x)$  (resp.  $P_t g(y)$ ) for any function  $g : \mathcal{X} \rightarrow \mathbb{R}$ . Consequently,

$$\|\mu P_t - \pi\|_{\text{TV}} = \|\mu P_t - \pi P_t\|_{\text{TV}} = \sup_{\|g\|_{\infty} \leq 1} |\langle \mu P_t, g \rangle - \langle \pi P_t, g \rangle| = \sup_{\|g\|_{\infty} \leq 1} |\langle \eta P_t, F_g \rangle| \leq 2 \langle \eta P_t, \mathbf{1}_{x \neq y} \rangle,$$

where  $F_g(x, y) = g(x) - g(y)$ ,  $\eta$  is any coupling measure of  $\mu$  and  $\pi$  (i.e. a probability measure on  $\mathcal{X} \times \mathcal{X}$  with marginals  $\mu$  and  $\pi$ ), and  $\mathbf{1}_{x \neq y}$  is the function on  $\mathcal{X}^2$  which is 0 on the diagonal and 1 outside. Since  $\mathbb{L} \mathbf{1}_{x \neq y} \leq -\inf(\mathbf{a}) \mathbf{1}_{x \neq y}$ , Grönwall's lemma entails  $\mathbb{P}_t \mathbf{1}_{x \neq y} \leq e^{-\inf(\mathbf{a})t} \mathbf{1}_{x \neq y}$  and then

$$\|\mu P_t - \pi\|_{\text{TV}} \leq \langle \eta, \mathbb{P}_t \mathbf{1}_{x \neq y} \rangle \leq 2 e^{-\inf(\mathbf{a})t} \eta(\{x \neq y\}).$$

Recalling that  $2(\mu - \pi)_+(\mathcal{X}) = 2(\pi - \mu)_+(\mathcal{X}) = \|\mu - \pi\|_{\text{TV}}$ , we choose the coupling measure

$$\eta(dx, dy) = \frac{2(\mu - \pi)_+(dx)(\pi - \mu)_+(dy)}{\|\mu - \pi\|_{\text{TV}}} + \delta_x(dy)(\mu \wedge \pi)(dx),$$

where  $\mu \wedge \pi = \mu - (\mu - \pi)_+ = \pi - (\pi - \mu)_+$ , and get the result as  $\eta(\{x \neq y\}) = \|\mu - \pi\|_{\text{TV}}/2$ .  $\square$

The convergences in stronger norms of Theorem 2.1-(1) are proved through functional inequalities, of Poincaré or logarithmic Sobolev type, that are known to be a powerful tool for deriving the exponential decay of  $\Phi$ -entropies [7, 8, 9, 22, 24]. We have already seen that the  $\Phi$ -entropies associated to the invariant measure  $\pi$  decrease along the solutions of Equation (2.4). This can also be obtained by differentiating the entropy along the trajectories and defining the dissipation of entropy as the opposite of this derivative

$$\frac{d}{dt} \text{Ent}_{\pi}^{\Phi}(P_t f) = \int \Phi'(P_t f) \mathbb{L} P_t f d\pi = -D_{\pi}^{\Phi}(P_t f). \quad (2.9)$$

The convexity of  $\Phi$  and some calculations then ensure the non-negativity of the dissipation

$$\begin{aligned} D_\pi^\Phi(f) &= - \int \Phi'(f) Lf \, d\pi = \int [L(\Phi(f)) - \Phi'(f) Lf] \, d\pi \\ &= \int [\langle \mathbf{Q}, \Phi(f) \rangle - \Phi(f) - \Phi'(f)(\langle \mathbf{Q}, f \rangle - f)] \, d\mathbf{Q} \\ &= \iint [\Phi(f(x)) - \Phi(f(y)) - \Phi'(f(y))(f(x) - f(y))] \mathbf{Q}(x) \mathbf{Q}(y) \, dx dy \geq 0. \end{aligned}$$

When  $\Phi'$  is concave we have additionally, using Jensen's inequality,

$$\begin{aligned} D_\pi^\Phi(f) &= - \int \Phi'(f) Lf \, d\pi = \int \Phi'(f)(f - \mathbf{Q}(f)) \, d\mathbf{Q} \\ &= \langle \mathbf{Q}, f \Phi'(f) \rangle - \langle \mathbf{Q}, f \rangle \langle \mathbf{Q}, \Phi'(f) \rangle \\ &\geq \langle \mathbf{Q}, f \Phi'(f) \rangle - \langle \mathbf{Q}, f \rangle \Phi'(\langle \mathbf{Q}, f \rangle) = \text{Ent}_{\mathbf{Q}}^{x \mapsto x \Phi'(x)}(f). \end{aligned}$$

In the case  $\Phi(x) = |x|^p$  with  $p \geq 1$  we have  $x \Phi'(x) = p \Phi(x)$  and consequently

$$D_\pi^\Phi(f) \geq p \text{Ent}_{\mathbf{Q}}^\Phi(f). \quad (2.10)$$

In the specific case  $p = 2$  it is even an equality. For  $\Phi(x) = x \log x$  we have  $x \Phi'(x) = \Phi(x) + x$  and so for all  $f > 0$

$$D_\pi^\Phi(f) \geq \text{Ent}_{\mathbf{Q}}^\Phi(f) + \text{Ent}_{\mathbf{Q}}^{\text{Id}}(f) = \text{Ent}_{\mathbf{Q}}^\Phi(f). \quad (2.11)$$

When  $\mathbf{a}$  is constant,  $\text{Ent}_{\mathbf{Q}}^\Phi = \mathbf{a} \text{Ent}_\pi^\Phi$  and Inequality (2.11) ensures exponential decay of the entropy. This can be generalized for non-constant  $\mathbf{a}$  as shown in the following result.

**Proposition 2.2** ( $\Phi$ -entropy decay). *Under Assumption (2.8), if  $\Phi(x) = |x|^p$  with  $p \in [1, 2]$  or  $\Phi(x) = x \log x$ , then for all  $f \in L^1(\pi)$  such that  $\Phi(f) \in L^1(\pi)$  we have*

$$\text{Ent}_\pi^\Phi(f) \leq \frac{1}{p \inf \mathbf{a}} D_\pi^\Phi(f), \quad (2.12)$$

where  $p = 1$  when  $\Phi(x) = x \log x$ . Consequently, for all  $t \geq 0$ ,

$$\text{Ent}_\pi^\Phi(P_t f) \leq e^{-p(\inf \mathbf{a})t} \text{Ent}_\pi^\Phi(f). \quad (2.13)$$

Before proving this result, let us recall that Inequality (2.12) is a powerful inequality that yields further properties than exponential convergence. Let us cite for instance concentration of measures [44] or hypercontractivity [7, Théorème 2.8.2]: for every  $r > 1$

$$\|P_t f\|_{L^{r(t)}} \leq \|f\|_{L^r}$$

where  $r(t) = 1 + (r - 1)e^{4t \inf(\mathbf{a})}$ . Even if we prove (2.12) by a simple argument, a classical way for proving such inequality is the Bakry-Emery  $\Gamma_2$  criterion [8] which implies local Poincaré type inequality but seems not applying here.

*Proof of Proposition 2.2.* The exponential entropy decay (2.13) readily follows from the combination of (2.9) and (2.12). For proving (2.12), we use a similar argument to the classical perturbation result of Holley and Stroock for logarithmic Sobolev inequalities, see [39] or [24, Proposition 3.2]. Let  $f$  such that  $\text{Ent}_\pi^\Phi(f) < \infty$ . The functions  $\Phi$  under interest are all differentiable outside of zero, so we can define for all  $z \neq 0$

$$\begin{aligned} F(z) &= \int (\Phi(f) - \Phi(z) - \Phi'(z)(f - z)) \, d\pi \\ &= \langle \pi, \Phi(f) \rangle - \Phi(z) - \Phi'(z)(\langle \pi, f \rangle - z) \\ &= \text{Ent}_\pi^\Phi(f) + \Phi(\langle \pi, f \rangle) - \Phi(z) - \Phi'(z)(\langle \pi, f \rangle - z). \end{aligned}$$

The convexity of  $\Phi$  thus guarantees that  $\text{Ent}_\pi^\Phi(f) = \inf_{z \neq 0} F(z)$ . Consequently,

$$\begin{aligned} \text{Ent}_\pi^\Phi(f) &= \inf_{z \neq 0} \int (\Phi(f) - \Phi(z) - \Phi'(z)(f - z)) d\pi \\ &\leq \frac{1}{\inf \mathbf{a}} \inf_{z \neq 0} \int (\Phi(f) - \Phi(z) - \Phi'(z)(f - z)) d\mathbf{Q} = \frac{1}{\inf \mathbf{a}} \text{Ent}_{\mathbf{Q}}^\Phi(f) \end{aligned}$$

which, combined with (2.10) or (2.11), yields (2.12).  $\square$

*Proof of Theorem 2.1-(1b)-(1c)-(1d)-(1e).* For  $\Phi(x) = |x|^p$ ,  $1 \leq p \leq 2$ , the entropy decay (2.13) reads for  $f$  such that  $\langle \pi, f \rangle = 0$

$$\|P_t f\|_{L^p(\pi)} \leq e^{-\inf(\mathbf{a})t} \|f\|_{L^p(\pi)}. \quad (2.14)$$

Choosing  $f = \frac{\mu}{\pi} - \langle \mu, \mathbf{1} \rangle$  and using the duality relation  $\mu P_t - \pi = (P_t \frac{\mu}{\pi} - \mathbf{1})\pi$  yields (1b). For  $p > 2$ , we argue by the duality representation of the  $L^p$  norms. Using (2.14) for  $p' \in (1, 2)$ , we have for  $\mu \in \mathcal{P}(\mathcal{X}) \cap L^p(\pi^{1-p})$

$$\begin{aligned} \|\mu P_t - \pi\|_{L^p(\pi^{1-p})} &= \sup_{\|f\|_{L^{p'}(\pi)} \leq 1} \langle \mu P_t - \pi, f \rangle = \sup_{\|f\|_{L^{p'}(\pi)} \leq 1} \langle \mu - \pi, P_t f \rangle \\ &= \sup_{\|f\|_{L^{p'}(\pi)} \leq 1} \langle \mu - \pi, P_t f - \langle \pi, f \rangle \rangle \\ &\leq \|\mu - \pi\|_{L^p(\pi^{1-p})} \sup_{\|f\|_{L^{p'}(\pi)} \leq 1} \|P_t f - \langle \pi, f \rangle\|_{L^{p'}(\pi)} \\ &\leq \|\mu - \pi\|_{L^p(\pi^{1-p})} e^{-\inf(\mathbf{a})t} \sup_{\|f\|_{L^{p'}(\pi)} \leq 1} \|f - \langle \pi, f \rangle\|_{L^{p'}(\pi)} \\ &\leq 2 e^{-\inf(\mathbf{a})t} \|\mu - \pi\|_{L^p(\pi^{1-p})}. \end{aligned}$$

The same method yields (1d) since  $\|\mu/\pi\|_{L^\infty} = \sup_{\|f\|_{L^1(\pi)} \leq 1} \langle \mu, f \rangle$ . Finally, the Kullback-Leibler divergence decay also comes from (2.13) and the duality relation  $\mu P_t = (P_t \frac{\mu}{\pi})\pi$ , which yields

$$D_{\text{KL}}(\mu P_t \|\pi) = \text{Ent}_\pi^{x \mapsto x \log x} \left( P_t \frac{\mu}{\pi} \right).$$

$\square$

**2.2. Algebraic convergence.** In this subsection, we assume that

$$\inf(\mathbf{a}) = 0 \quad \text{and} \quad \int_{\mathcal{X}} \frac{\mathbf{Q}}{\mathbf{a}} = 1. \quad (2.15)$$

In particular, we have two invariant probability distributions  $\pi = \mathbf{Q}/\mathbf{a}$  and  $\delta_0$ . We start by proving (2b). To do so we use a subgeometric result from [38], which is inspired from [32] and was recently revisited in [23]. The result of (2b) is a direct consequence of the next proposition.

**Proposition 2.3.** *Suppose that (2.15) is verified, that  $\mathbf{a} \in L^\infty$ , and that  $1/\mathbf{a} \in L^q(\mathbf{Q})$  for some  $q > 1$ . Then there exists  $C > 0$  such that for any  $x \in \mathcal{X}$  and  $t \geq 0$*

$$\|\delta_x P_t - \pi\|_{\text{TV}} \leq \frac{C \mathbf{a}(x)^{-q}}{(t/q + 1)^q} + \frac{C}{(t/q + 1)^{q-1}} \leq \frac{C(1 + \mathbf{a}(x)^{-q})}{(t + 1)^{q-1}}.$$

*Proof.* It follows from [38, Theorem 4.1], provided that

$$\mathcal{L}^* V \leq K - \varphi(V)$$

is verified for some  $K > 0$  with  $V = \mathbf{a}^{-q}$  and  $\varphi : x \mapsto x^{\frac{q-1}{q}}$ , and that for every  $C > 0$ , there exists  $\alpha > 0$  and  $T > 0$  such that

$$\|\delta_x P_T - \delta_y P_T\|_{\text{TV}} \leq 2(1 - \alpha)$$

for every  $(x, y) \in \mathcal{X}^2$  such that  $V(x) + V(y) \leq C$ . Note that the precompactness assumption of [38, Theorem 4.1] is only needed to prove the existence of the invariant measure, which is known here. The first condition is clearly verified since for  $V = \mathbf{a}^{-q}$  we have

$$\mathcal{L}^*V(x) = \|1/\mathbf{a}\|_{L^q(\mathbf{Q})}^q \mathbf{a}(x) - \mathbf{a}^{1-q}(x)$$

and  $\mathbf{a}$  is supposed to be bounded. For the second condition, we first remark that for every  $C > 0$ , the bound  $V(x) + V(y) \leq C$  implies the existence of  $c > 0$  such that  $\mathbf{a}(x) \geq c$  and  $\mathbf{a}(y) \geq c$ . Then, for every  $(x, y)$  such that  $V(x) + V(y) \leq C$  and any  $z \in \{x, y\}$ , we have from Duhamel's formula (2.5) that for all  $f \geq 0$  and all  $t \geq 0$

$$P_t f(z) \geq c \int_0^t e^{-Cs} \langle \mathbf{Q}, P_{t-s} f \rangle ds = \alpha \langle \nu, f \rangle$$

where  $\nu(dx) = C(1 - e^{-Ct})^{-1} \int_0^t e^{-Cs} (\mathbf{Q} P_{t-s})(dx) ds$  is a probability measure and  $\alpha = \frac{c(1 - e^{-Ct})}{C} > 0$ . Hence, the measures  $\delta_x P_t - \alpha \nu$  and  $\delta_y P_t - \alpha \nu$  are positive and Doeblin's argument yields

$$\|\delta_x P_t - \delta_y P_t\|_{\text{TV}} \leq \|\delta_x P_t - \alpha \nu\|_{\text{TV}} + \|\delta_y P_t - \alpha \nu\|_{\text{TV}} = \langle \delta_x P_t - \alpha \nu, \mathbf{1} \rangle + \langle \delta_y P_t - \alpha \nu, \mathbf{1} \rangle = 2(1 - \alpha).$$

We can then apply [38, Theorem 4.1] which yields the result since

$$H_\varphi(u) = \int_1^u \frac{ds}{\varphi(s)} = q(u^{1/q} - 1) \quad \text{and} \quad H_\varphi^{-1}(t) = (t/q + 1)^q.$$

□

As for the fast convergence, we can give stronger convergence results depending on the tails of the initial condition. This sub-geometric bounds in entropy seems more original than the previous bound in total variation distance. Let us nevertheless cite [21, 56] which use others weakening of entropic inequalities to obtain algebraic convergence of some diffusion processes.

**Proposition 2.4.** *Suppose that (2.15) is verified and that  $1/\mathbf{a} \in L^q(\mathbf{Q})$  for some  $q > 1$ . Then for all  $p \in [1, 2]$ , all  $f \in L^1(\pi)$  such that  $f/\pi \in L^\infty$ , and all  $t \geq 0$ , we have*

$$\text{Ent}_\pi^{|\cdot|^p}(P_t f) \leq 2p(p(q-1))^{q-1} \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})}^q \|f/\pi\|_{L^\infty}^p t^{-(q-1)}. \quad (2.16)$$

If additionally  $\mathbf{a} \in L^{r'}(\pi)$  for some  $r > p$ , then for all  $f \in L^r(\pi)$  and all  $t \geq 0$  we have

$$\text{Ent}_\pi^{|\cdot|^p}(P_t f) \leq C_r \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})}^{q+p\frac{1-q}{r}} \max\left(1, \|\mathbf{a}\|_{L^{r'}(\pi)}^p\right) \|f\|_{L^r(\pi)}^p t^{-(q-1)(1-\frac{p}{r})} \quad (2.17)$$

where

$$C_r = 2p \left( p(q-1) \left( 1 - \frac{p}{r} \right) \right)^{(q-1)(1-\frac{p}{r})}.$$

*Proof.* We only prove (2.17), the proof of (2.16) being exactly the same by replacing  $r$  by  $\infty$ ,  $r'$  by 1, and  $\|f\|_{L^r(\pi)}$  by  $\|f/\pi\|_{L^\infty}$ .

Consider  $\Phi(x) = |x|^p$  with  $p \in [1, 2]$ , so that

$$D_\pi^\Phi(f) \geq p \text{Ent}_\pi^\Phi(f),$$

and start from

$$\begin{aligned} \text{Ent}_\pi^\Phi(f) &= \inf_{z \in \mathbb{R}} \int [\Phi(f) - \Phi(z) - \Phi'(z)(f-z)] d\pi \\ &\leq \int [\Phi(f) - \Phi(\langle \mathbf{Q}, f \rangle) - \Phi'(\langle \mathbf{Q}, f \rangle)(f - \langle \mathbf{Q}, f \rangle)] d\pi. \end{aligned}$$

Let  $q > 1$  such that  $1/\mathbf{a} \in L^q(\mathbf{Q})$  and denote

$$\Xi(f) = \Phi(f) - \Phi(\langle \mathbf{Q}, f \rangle) - \Phi'(\langle \mathbf{Q}, f \rangle)(f - \langle \mathbf{Q}, f \rangle).$$

For any  $\alpha \in (0, 1]$ , splitting  $\mathbf{a} = \mathbf{a}^\alpha \mathbf{a}^{1-\alpha}$  and using Hölder's inequality, we have

$$\text{Ent}_\pi^\Phi(f) \leq \int \Xi(f) d\pi \leq \left( \int (\Xi(f))^{\frac{q}{q-\alpha}} \mathbf{a}^{-\frac{(1-\alpha)q}{q-\alpha}} d\mathbf{Q} \right)^{1-\frac{\alpha}{q}} \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})}.$$

Using again Hölder's inequality we get

$$\text{Ent}_\pi^\Phi(f) \leq \left( \int \Xi(f) d\mathbf{Q} \right)^{\frac{\alpha}{q'}} \left( \int (\Xi(f))^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}} \mathbf{a}^{-1} d\mathbf{Q} \right)^{1-\alpha} \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})}$$

which also reads

$$\text{Ent}_\pi^\Phi(f) \leq \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})} \|\Xi(f)\|_{L^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}}(\pi)}^{1-\alpha+\frac{\alpha}{q}} (\text{Ent}_\mathbf{Q}^\Phi(f))^{\frac{\alpha}{q'}}.$$

Now we control  $\Xi(f)$  by

$$\begin{aligned} \Xi(f) &= \Phi(f) - \Phi(\langle \mathbf{Q}, f \rangle) - \Phi'(\langle \mathbf{Q}, f \rangle)(f - \langle \mathbf{Q}, f \rangle) \\ &= |f|^p - |\langle \mathbf{Q}, f \rangle|^p - p \text{sign}(\langle \mathbf{Q}, f \rangle) |\langle \mathbf{Q}, f \rangle|^{p-1} (f - \langle \mathbf{Q}, f \rangle) \\ &\leq |f|^p + (p-1) |\langle \mathbf{Q}, f \rangle|^p + p |\langle \mathbf{Q}, f \rangle|^{p-1} |f| \\ &= 2p \left( \frac{1}{2p} |f|^p + \frac{p-1}{2p} |\langle \mathbf{Q}, f \rangle|^p + \frac{1}{2} |\langle \mathbf{Q}, f \rangle|^{p-1} |f| \right) \end{aligned}$$

to get by convexity

$$\begin{aligned} &(\Xi(f))^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}} \\ &\leq (2p)^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}} \left( \frac{1}{2p} |f|^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})} + \frac{p-1}{2p} |\langle \mathbf{Q}, f \rangle|^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})} + \frac{1}{2} |\langle \mathbf{Q}, f \rangle|^{(p-1)(1+\frac{\alpha}{1-\alpha}\frac{1}{q})} |f|^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}} \right), \end{aligned}$$

so that  $\|\Xi(f)\|_{L^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}}(\pi)}^{1-\alpha+\frac{\alpha}{q}}$  is bounded by

$$(2p)^{1-\alpha+\frac{\alpha}{q}} \left( \frac{1}{2p} \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})} + \frac{p-1}{2p} |\langle \mathbf{Q}, f \rangle|^{p(1-\alpha+\frac{\alpha}{q})} + \frac{1}{2} |\langle \mathbf{Q}, f \rangle|^{(p-1)(1-\alpha+\frac{\alpha}{q})} \|f\|_{L^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}}(\pi)}^{1-\alpha+\frac{\alpha}{q}} \right).$$

Then we estimate  $\langle \mathbf{Q}, f \rangle$  by

$$|\langle \mathbf{Q}, f \rangle| \leq \int |f| \mathbf{a} d\pi \leq \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)} \|\mathbf{a}\|_{L^{[p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})]'}(\pi)}$$

to obtain, using that  $\|f\|_{L^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}}(\pi)} \leq \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}$  since  $p \geq 1$  and  $\pi$  is probability measure,

$$\|\Xi(f)\|_{L^{1+\frac{\alpha}{1-\alpha}\frac{1}{q}}(\pi)}^{1-\alpha+\frac{\alpha}{q}} \leq C \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})}$$

with

$$C = (2p)^{1-\alpha+\frac{\alpha}{q}} \max \left( 1, \|\mathbf{a}\|_{L^{[p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})]'}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})} \right).$$

Since  $t \mapsto \|P_t f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})}$  is nonincreasing by entropy property, we finally get for all  $t \geq 0$

$$\text{Ent}_\pi^\Phi(P_t f) \leq C \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})} \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})} (\text{Ent}_\mathbf{Q}^\Phi(P_t f))^{\frac{\alpha}{q'}}.$$

This yields the differential inequality

$$\frac{d}{dt} \text{Ent}_\pi^\Phi(P_t f) = -D_\pi^\Phi(P_t f) \leq -\frac{1}{p} \text{Ent}_\mathbf{Q}^\Phi(P_t f) \leq -C' (\text{Ent}_\pi^\Phi(P_t f))^{\frac{\alpha'}{\alpha}},$$

where

$$C' = \frac{1}{p} \left( C \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})} \|f\|_{L^{p(1+\frac{\alpha}{1-\alpha}\frac{1}{q})}(\pi)}^{p(1-\alpha+\frac{\alpha}{q})} \right)^{-\frac{\alpha'}{\alpha}},$$

which ensures that

$$\text{Ent}_\pi^\Phi(P_t f) \leq \left( (\text{Ent}_\pi^\Phi(f))^{1-\frac{q'}{\alpha}} + \left(\frac{q'}{\alpha} - 1\right) C' t \right)^{\frac{1}{1-\frac{q'}{\alpha}}}.$$

Choosing  $\alpha$  such that  $r = p(1 + \frac{\alpha}{1-\alpha} \frac{1}{q}) > p$  we have

$$C' = \frac{1}{p} \left( C \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})} \|f\|_{L^r(\pi)}^{\frac{pr}{p+q(r-p)}} \right)^{-\frac{q+r-p}{q-1}} \quad \text{with} \quad C = (2p)^{\frac{r}{p+q(r-p)}} \max \left( 1, \|\mathbf{a}\|_{L^r(\pi)}^{\frac{pr}{p+q(r-p)}} \right),$$

and the speed of convergence is given by

$$-\frac{1}{1-\frac{q'}{\alpha}} = \frac{(q-1)(r-p)}{r} = (q-1) \left( 1 - \frac{p}{r} \right).$$

We thus finally have

$$\text{Ent}_\pi^\Phi(P_t f) \leq C'' \left\| \frac{1}{\mathbf{a}} \right\|_{L^q(\mathbf{Q})}^{q+p\frac{1-q}{r}} \max \left( 1, \|\mathbf{a}\|_{L^r(\pi)}^p \right) \|f\|_{L^r(\pi)}^p t^{-(q-1)(1-\frac{p}{r})}$$

with

$$C'' = 2p \left( p(q-1) \left( 1 - \frac{p}{r} \right) \right)^{(q-1)(1-\frac{p}{r})}.$$

□

*Proof of Theorem 2.1 - (2c) and (2d).* We only give the proof of (2d), since (2c) can be seen as the limit case “ $r = \infty$ ” and proved in the exact same way, using (2.16) instead of (2.17).

Suppose that (2.15) is verified, let  $p \in [1, 2]$ ,  $q > 1$ ,  $r > p$ , and assume that  $\mathbf{a} \in L^r(\pi)$  and  $1/\mathbf{a} \in L^q(\mathbf{Q})$ . Then (2.17) in Proposition 2.4 ensures the existence of  $C > 0$  such that for all  $f \in L^r(\pi)$  with  $\langle \pi, f \rangle = 0$  and all  $t \geq 0$

$$\|P_t f\|_{L^p(\pi)} \leq C \max(1, \|\mathbf{a}\|_{L^r(\pi)}) \|f\|_{L^r(\pi)} t^{-(q-1)(\frac{1}{p}-\frac{1}{r})}.$$

This inequality applied to  $f = \frac{\mu}{\pi} - \langle \mu, \mathbf{1} \rangle$  yields (2d) for  $p \in [1, 2]$ , since  $\mu P_t - \pi = (P_t \frac{\mu}{\pi} - \mathbf{1})\pi$ . For the case  $p > 1$  in (2d) we argue by duality, similarly as in the proof of Theorem 2.1-(1c)

$$\begin{aligned} \|\mu P_t - \pi\|_{L^{p'}(\pi^{1-p'})} &= \sup_{\|f\|_{L^r(\pi)} \leq 1} \langle \mu P_t - \pi, f \rangle = \sup_{\|f\|_{L^r(\pi)} \leq 1} \langle \mu - \pi, P_t f - \langle \pi, f \rangle \rangle \\ &\leq \|\mu - \pi\|_{L^{p'}(\pi^{1-p'})} \sup_{\|f\|_{L^r(\pi)} \leq 1} \|P_t f - \langle \pi, f \rangle\|_{L^p(\pi)} \\ &\leq \|\mu - \pi\|_{L^{p'}(\pi^{1-p'})} C \max(1, \|\mathbf{a}\|_{L^r(\pi)}) t^{-(q-1)(1/p-1/r)} \sup_{\|f\|_{L^r(\pi)} \leq 1} \|f - \langle \pi, f \rangle\|_{L^r(\pi)} \\ &\leq 2C \max(1, \|\mathbf{a}\|_{L^r(\pi)}) t^{-(q-1)(1/p-1/r)} \|\mu - \pi\|_{L^{p'}(\pi^{1-p'})}. \end{aligned}$$

This gives the conclusion by replacing  $r'$  by  $p$ , and consequently  $p'$  by  $r$ , since  $\frac{1}{p} - \frac{1}{r} = \frac{1}{r'} - \frac{1}{p'}$ . □

**2.3. Weak convergence.** In this subsection, we consider that  $\mathbf{a}(0) = 0$  and  $\mathbf{a} \in L^1(\mathbf{Q})$ , or equivalently  $\mathbf{a} \in L^2(\mathbf{Q}/\mathbf{a})$ , and we prove the weak convergence results, *i.e.* for the narrow topology, of Theorem 2.1. To do so, we work in the space  $L^2(\mathbf{Q}/\mathbf{a})$  and use the quadratic entropy functional

$$H_2[f] = \|f\|_{L^2(\mathbf{Q}/\mathbf{a})}^2 = \int_{\mathcal{X}} \frac{f^2(x)}{\mathbf{a}(x)} \mathbf{Q}(dx)$$

which is, up to the addition of the mean, the variance under  $\pi = \mathbf{Q}/\mathbf{a}$  (which is not necessarily a finite measure here). As for the  $\Phi$ -entropies, we have the dissipation property

$$\frac{d}{dt} H_2[P_t f] = -D_2[P_t f], \quad \text{with} \quad D_2[f] = \iint (f(x) - f(y))^2 \mathbf{Q}(dx) \mathbf{Q}(dy) \geq 0.$$

In particular, the semigroup  $(P_t)_{t \geq 0}$  is well defined on  $L^2(\mathbf{Q}/\mathbf{a})$ . We start by proving a useful result which is a consequence of the above entropy property. It is useful to define the domain of  $\mathcal{L}^*$  in  $L^2(\mathbf{Q}/\mathbf{a})$  by

$$\mathcal{D}(\mathcal{L}^*) = \{f \in L^2(\mathbf{Q}/\mathbf{a}), \mathcal{L}^* f \in L^2(\mathbf{Q}/\mathbf{a})\}.$$

**Lemma 2.5.** *Assume that  $\mathbf{a} \in L^1(\mathbf{Q})$  and let  $f \geq 0$  in  $\mathcal{D}(\mathcal{L}^*)$ . For any sequence  $(t_n)_{n \geq 0}$  of positive real numbers which is increasing and tends to  $+\infty$ , there exists a sub-sequence  $(t'_n)_{n \geq 0}$  and a continuous function  $\rho : \mathbb{R} \rightarrow [0, \infty)$  such that for any  $\varphi \in L^2(\mathbf{a}\mathbf{Q})$  the convergence*

$$\int_{\mathcal{X}} P_{t+t'_n} f(x) \varphi(x) \mathbf{Q}(dx) \rightarrow \rho(t) \langle \mathbf{Q}, \varphi \rangle \quad (2.18)$$

holds locally uniformly in time.

*Proof.* Let  $f \geq 0$  in  $\mathcal{D}(\mathcal{L}^*)$ . By virtue of the entropy property, we have for all  $t \geq 0$

$$\|P_t f\|_{L^2(\mathbf{Q}/\mathbf{a})} \leq \|f\|_{L^2(\mathbf{Q}/\mathbf{a})} \quad \text{and} \quad \|\partial_t P_t f\|_{L^2(\mathbf{Q}/\mathbf{a})} = \|P_t \mathcal{L}^* f\|_{L^2(\mathbf{Q}/\mathbf{a})} \leq \|\mathcal{L}^* f\|_{L^2(\mathbf{Q}/\mathbf{a})}.$$

Consequently, due to Arzelà-Ascoli and Banach-Alaoglu theorems, we can extract, from the sequence  $(f_n)_{n \geq 0}$ , defined by  $f_n(t, x) = P_{t+t_n} f(x)$ , a sub-sequence, still denoted  $(f_n)$ , such that for all  $\varphi \in L^2(\mathbf{a}\mathbf{Q})$  the convergence

$$\int f_n(t, x) \varphi(x) \mathbf{Q}(dx) \rightarrow \int f_\infty(t, x) \varphi(x) \mathbf{Q}(dx),$$

holds locally uniformly in time, for some  $f_\infty : \mathbb{R} \rightarrow L^2(\mathbf{Q}/\mathbf{a})$  non-negative and weakly continuous. Now we check that  $f_\infty$  is constant in  $x$ . Let  $(g_n)_{n \geq 0}$  be the sequence of three variables functions defined by  $g_n(t, x, y) = f_n(t, x) - f_n(t, y)$ . For all  $\psi \in L^2(\mathbf{Q}(dx) \otimes \mathbf{Q}(dy))$  and  $T > 0$ , we have

$$\left| \int_0^T \iint g_n(t, x, y) \psi(x, y) \mathbf{Q}(dx) \mathbf{Q}(dy) dt \right| \leq \int_0^T \sqrt{D_2[f_n(t, \cdot)]} dt \sqrt{\iint \psi^2(x, y) \mathbf{Q}(dx) \mathbf{Q}(dy)} \xrightarrow{n \rightarrow \infty} 0,$$

because

$$\int_0^T \sqrt{D_2[f_n(t, \cdot)]} dt = \int_n^{T+n} \sqrt{D_2[f(t, \cdot)]} dt \leq T \int_n^{T+n} D_2[f(t, \cdot)] dt$$

and

$$\int_0^\infty D_2[f(t, \cdot)] dt \leq H_2[f] < \infty.$$

Choosing now  $\psi(x, y) = \xi(x)$  with  $\xi \in L^1(\mathbf{Q})$  such that  $\langle \mathbf{Q}, \xi \rangle = 0$ , we obtain

$$\int_0^T \int f_\infty(t, x) \xi(x) \mathbf{Q}(dx) dt = \lim_{n \rightarrow \infty} \int_0^T \int f_n(t, x) \xi(x) \mathbf{Q}(dx) dt = 0,$$

This ensures that  $f_\infty(t, x) = \rho(t)$ , for every  $t \geq 0$  and almost all  $x \in \mathcal{X}$ , for some non-negative continuous function  $\rho$ .  $\square$

We are now in position to prove the weak-\* convergences results of Theorem 2.1.

*Proof of Theorem 2.1-(2a).* We assume here that  $\int \mathbf{Q}/\mathbf{a} = 1$ ,  $\mathbf{a}(0) = 0$ , and  $\mathbf{a} \in L^1(\mathbf{Q})$ . In this case  $\pi = \mathbf{Q}/\mathbf{a}$  is an invariant probability measures and  $\langle \pi, P_t f \rangle = \langle \pi, f \rangle$  for all  $f \in \mathcal{D}(\mathcal{L}^*)$ . This enforces, by taking  $\varphi = 1/\mathbf{a}$  in (2.18), that  $\rho(t) = \langle \pi, f \rangle$  for all  $t$ , and we can infer that for any  $\varphi \in L^2(\mathbf{a}\mathbf{Q})$  and  $f \in \mathcal{D}(\mathcal{L}^*)$ ,

$$\int_{\mathcal{X}} P_t f(x) \varphi(x) \mathbf{Q}(dx) \xrightarrow{t \rightarrow +\infty} \langle \pi, f \rangle \langle \mathbf{Q}, \varphi \rangle.$$

In particular by choosing  $\varphi = \mathbf{1}$ , which is admissible since we supposed  $\mathbf{a} \in L^1(\mathbf{Q})$ , this ensures that  $\langle \mathbf{Q}, P_t f \rangle \rightarrow \langle \pi, f \rangle$  as  $t \rightarrow \infty$ . We deduce from Duhamel's formula (2.5), by using a Cesàro type argument, that for any continuous function  $f \in \mathcal{D}(\mathcal{L}^*)$  and for all  $x \in \mathcal{X} \setminus \{0\}$

$$P_t f(x) = f(x)e^{-\mathbf{a}(x)t} + \int_0^t \mathbf{a}(x)e^{-\mathbf{a}(x)s} \langle \mathbf{Q}, P_{t-s} f \rangle ds \xrightarrow[t \rightarrow +\infty]{} \langle \pi, f \rangle.$$

On the other hand, since  $\mathbf{a}(0) = 0$ , we have  $P_t f(0) = f(0)$  for all  $t \geq 0$ . Finally, as  $C_b(\mathcal{X}) \subset \mathcal{D}(\mathcal{L}^*)$  and  $(P_t)_{t \geq 0}$  is a Markov semigroup, we obtain by dominated convergence that for any  $\mu \in \mathcal{P}(\mathcal{X})$  and any  $f \in C_b(\mathcal{X})$

$$\langle \mu P_t, f \rangle \xrightarrow[t \rightarrow +\infty]{} \mu(\{0\}) + (1 - \mu(\{0\})) \langle \pi, f \rangle$$

which is the desired result.  $\square$

*Proof of Theorem 2.1-(3).* Here we assume that  $\mathbf{a} \in L^1(\mathbf{Q})$  and  $\int \mathbf{Q}/\mathbf{a} = +\infty$ . Then  $\mathbf{Q}/\mathbf{a}$  is not a finite measure, and a continuous function  $f$  that belongs to  $L^2(\mathbf{Q}/\mathbf{a})$  necessarily vanishes at 0. Since we know from the proof of Lemma 2.5 that for all  $t$  the constant function  $x \mapsto f_\infty(t, x) = \rho(t)$  belongs to  $L^2(\mathbf{Q}/\mathbf{a})$ , we must have  $\rho(t) = 0$  and we deduce that for any  $\varphi \in L^2(\mathbf{a}\mathbf{Q})$  and  $f \in \mathcal{D}(\mathcal{L}^*)$ ,

$$\int_{\mathcal{X}} P_t f(x) \varphi(x) \mathbf{Q}(dx) \xrightarrow[t \rightarrow +\infty]{} 0.$$

In particular by choosing  $\varphi = \mathbf{1}$ , which is admissible since we supposed  $\mathbf{a} \in L^1(\mathbf{Q})$ , this ensures that  $\langle \mathbf{Q}, P_t f \rangle \rightarrow 0$  as  $t \rightarrow \infty$ . We deduce from Duhamel's formula (2.5), by using a Cesàro type argument, that for any continuous function  $f \in \mathcal{D}(\mathcal{L}^*)$  and for all  $x \in \mathcal{X} \setminus \{0\}$

$$P_t f(x) = f(x)e^{-\mathbf{a}(x)t} + \int_0^t \mathbf{a}(x)e^{-\mathbf{a}(x)s} \langle \mathbf{Q}, P_{t-s} f \rangle ds \xrightarrow[t \rightarrow +\infty]{} 0. \quad (2.19)$$

Since a continuous function  $f$  which belongs to  $\mathcal{D}(\mathcal{L}^*)$  necessarily verifies  $f(0) = 0$ , and since  $\mathbf{a}(0) = 0$ , this convergence actually holds for all  $x \in \mathcal{X}$ .

Now, let  $\mu \in \mathcal{P}(\mathcal{X})$ . Due to (2.3), we can find in  $\mathcal{D}(\mathcal{L}^*)$  a continuous function  $f \geq 0$  such that  $f(x) = 1$  for all  $|x| \geq 1$ . We deduce from (2.19) and by dominated convergence, since  $(P_t)_{t \geq 0}$  is a Markov semigroup, that  $\langle \mu P_t, f \rangle = \langle \mu, P_t f \rangle \rightarrow 0$  and this prevents the mass of  $\mu P_t$  to go to infinity. As a consequence, we can extract from  $(\mu P_t)_{t \geq 0}$  a sub-sequence which converges narrowly to some  $\mu_\infty \in \mathcal{P}(\mathcal{X})$ . Since  $\langle \mu P_t, f \rangle \rightarrow 0$  for all bounded continuous functions  $f$  with support that does not contain zero, this limit  $\mu_\infty$  must be  $\delta_0$  and finally the whole trajectory  $(\mu P_t)_{t \geq 0}$  converges to  $\delta_0$  in the narrow topology.  $\square$

### 3. THE NON-CONSERVATIVE LINEAR EQUATION

We consider now the non-conservative linear equation (1.2) and we prove Theorem 1.1. Denoting by  $(M_t)_{t \geq 0}$  the semigroup generated by  $\mathcal{A}^*$  defined in (1.3), we have similarly as for the conservative equation (2.1) that the unique solution to (1.2) with initial data  $u_0$  is given by  $u_t = u_0 M_t$ , in the suitable Banach spaces. For proving Theorem 1.1, we perform a so-called  $h$ -transform of  $(M_t)_{t \geq 0}$  and use the results of Theorem 2.1.

*Proof of Theorem 1.1-(1).* Here we consider the case  $\rho = \int_{\mathcal{X}} \frac{\mathbf{Q}}{\mathbf{a}} \in (1, +\infty]$ , so that  $\lambda$ , defined by the relation  $\int_{\mathcal{X}} \frac{\mathbf{Q}}{\lambda + \mathbf{a}} = 1$ , is strictly positive. We have already seen in the introduction that the function

$$h(x) = \frac{\alpha^{-1}}{\lambda + \mathbf{a}(x)}$$

verifies  $\mathcal{A}^*h = \lambda h$  and that choosing  $\alpha = \int_X \frac{Q}{(\lambda+a)^2}$  we have  $\langle \gamma, h \rangle = 1$  where

$$\gamma(dx) = \frac{Q(dx)}{\lambda + a(x)}$$

verifies  $\mathcal{A}\gamma = \lambda\gamma$ . Performing a  $h$ -transform of  $(M_t)_{t \geq 0}$  consists in defining

$$P_t f = \frac{M_{\alpha t}(fh)}{M_{\alpha t}h} = e^{-\lambda \alpha t} \frac{M_{\alpha t}(fh)}{h}.$$

The new family  $(P_t)_{t \geq 0}$  is a Markov semigroup with infinitesimal generator given by

$$\alpha \frac{\mathcal{A}^*(fh) - \lambda fh}{h} = \mathbf{a} \left( \int f d\mathbf{Q} - f \right) = \mathcal{L}^* f,$$

where

$$\mathbf{a}(x) = \alpha \frac{\langle Q, h \rangle}{h(x)} = \alpha(\lambda + a(x)) \quad \text{and} \quad \mathbf{Q}(dx) = \frac{h(x)Q(dx)}{\langle Q, h \rangle} = \gamma(dx).$$

The time scaling with parameter  $\alpha$  in the  $h$ -transform ensures that the invariant measure of  $\mathcal{L}^*$

$$\pi(dx) = \frac{\mathbf{Q}(dx)}{\mathbf{a}(x)}$$

is a probability measure. The hypotheses of Theorem 2.1-(1) are then satisfied, and it yields the results of Theorem 1.1-(1) by using the relation

$$e^{-\lambda t} u_t = e^{-\lambda t} u_0 M_t = \frac{(u_0 h) P_{t/\alpha}}{h}.$$

□

*Proof of Theorem 1.1-(2).* We consider now that  $\rho = \int_X \frac{Q}{a} = 1$ , so that  $\lambda = 0$ , together with the assumption that  $1/a \in L^2(Q)$ . We can then still define  $h$  and  $\gamma$  such that  $\langle \gamma, h \rangle = 1$  by setting

$$h(x) = \frac{\alpha^{-1}}{a(x)} \quad \text{and} \quad \gamma(dx) = \frac{Q(dx)}{a(x)}$$

with  $\alpha = \|1/a\|_{L^2(Q)}^2$ . Similarly as for the proof of Theorem 1.1-(1), the estimates in Theorem 1.1-(2b)-(2c)-(2d) are then direct consequences of Theorem 2.1-(2b)-(2c)-(2d).

The vague and narrow convergences in Theorem 1.1-(2a) readily follow from Theorem 2.1-(2a) since if  $u_0 \in \mathcal{M}(h)$  then necessarily  $u_0(\{0\}) = 0$ , and if  $a \in L^\infty$  then any  $f \in C_b(X)$  satisfies  $f/h \in C_b(X)$ . For Theorem 1.1-(2a'), we use the Duhamel formula

$$M_t f(x) = f(x) e^{-a(x)t} + \int_0^t e^{-a(x)s} \langle Q, M_{t-s} f \rangle ds.$$

The vague convergence in Theorem 1.1-(2a) ensures that  $\langle Q, M_t f \rangle \rightarrow \langle Q, h \rangle \langle \gamma, f \rangle = \alpha^{-1} \langle \gamma, f \rangle$  as  $t \rightarrow +\infty$  for all  $f \in C_c(X)$  and consequently, since  $a(0) = 0$ ,

$$M_t f(0) = f(0) + \int_0^t \langle Q, M_{t-s} f \rangle ds \sim \alpha^{-1} \langle \gamma, f \rangle t \quad \text{as } t \rightarrow +\infty.$$

When  $a \in L^\infty$ , we can replace  $C_c(X)$  by  $C_b(X)$  since the convergence in Theorem 1.1-(2a) holds narrowly, and the proof is complete. □

*Proof of Theorem 1.1-(3).* We consider that either  $\rho = 1$  and  $1/a \notin L^2(Q)$ , or  $\rho < 1$ . If  $\rho = 1$  and  $1/a \notin L^2(Q)$ , the function  $h(x) = 1/a(x)$  is an eigenfunction of  $\mathcal{A}^*$  associated to the eigenvalue  $\lambda = 0$ , but it cannot be normalized in such a way that  $\langle \gamma, h \rangle = 1$ . If  $\rho < 1$  the function  $1/a$  is not an eigenfunction of  $\mathcal{A}^*$ . Nevertheless, in both cases, we can perform the  $\frac{1}{a}$ -transform

$$P_t f = \frac{M_t(f/a)}{M_t(1/a)}.$$

The family  $(P_t)_{t \geq 0}$  is a Markov semigroup with infinitesimal generator given by

$$\mathcal{L}^* f = \mathbf{a} \left( \int f d\mathbf{Q} - f \right),$$

where

$$\mathbf{a}(x) = \rho a(x) \quad \text{and} \quad \mathbf{Q}(dx) = \frac{Q(dx)}{\rho a(x)}.$$

We are then in a situation where  $1/\mathbf{a} \notin L^1(\mathbf{Q})$ , so Theorem 2.1-(3) ensures that for all  $\mu \in \mathcal{M}(\mathcal{X})$  and all  $f \in C_c(\mathcal{X})$ , or for all  $f \in C_b(\mathcal{X})$  if  $a$  is bounded,  $\langle \mu, P_t(af) \rangle \rightarrow \mu(\{0\})a(0)f(0) = 0$  as  $t \rightarrow +\infty$ . Besides, we have

$$\partial_t M_t(1/a) = M_t \mathcal{L}^*(1/a) = (\rho - 1)M_t \mathbf{1} \leq 0,$$

from which we get that  $M_t(1/a) \leq 1/a$  for all  $t \geq 0$ . Finally, we deduce that for any  $u_0 \in \mathcal{M}(1/a)$  and all  $f \in C_c(\mathcal{X})$ , or all  $f \in C_b(\mathcal{X})$  if  $a \in L^\infty$ ,

$$|\langle u_t, f \rangle| = |\langle u_0, M_t f \rangle| = |\langle M_t(1/a)u_0, P_t(af) \rangle| \leq \langle |u_0|/a, P_t(a|f) \rangle \xrightarrow{t \rightarrow +\infty} 0,$$

which concludes the proof of Theorem 1.1-(3a). For (3b), we have on the one hand from Duhamel's formula that

$$M_t \mathbf{1}(x) = e^{-a(x)t} + \int_0^t e^{-a(x)s} \langle Q, M_{t-s} \mathbf{1} \rangle ds \leq 1 + \frac{1}{a(x)} \sup_{s \in [0, t]} \langle Q, M_s \mathbf{1} \rangle,$$

which gives by integration against  $Q$

$$\langle Q, M_t \mathbf{1} \rangle \leq 1 + \rho \sup_{s \in [0, t]} \langle Q, M_s \mathbf{1} \rangle,$$

from which we infer that

$$\sup_{s \geq 0} \langle Q, M_s \mathbf{1} \rangle \leq \frac{1}{1 - \rho}. \quad (3.1)$$

On the other hand, coming back to  $\partial_t M_t(1/a) = (1 - \rho)M_t \mathbf{1}$ , we deduce that for all  $t \geq 0$

$$\int_0^t M_s \mathbf{1} ds = \frac{1}{1 - \rho} (1/a - M_t(1/a)) \leq \frac{1}{(1 - \rho)a},$$

which yields

$$\int_0^{+\infty} \langle Q, M_s \mathbf{1} \rangle ds \leq \frac{\rho}{(1 - \rho)}, \quad (3.2)$$

thus guaranteeing that the positive measure defined by

$$\mu(dx) = \int_0^\infty (QM_s)(dx) ds$$

is finite. Using (3.1) in Duhamel's formula, we get that for all  $x \neq 0$

$$\begin{aligned} M_t \mathbf{1}(x) &= e^{-a(x)t} + \int_0^{t/2} e^{-a(x)s} \langle Q, M_{t-s} \mathbf{1} \rangle ds + \int_{t/2}^t e^{-a(x)s} \langle Q, M_{t-s} \mathbf{1} \rangle ds \\ &\leq e^{-a(x)t} + \int_{t/2}^t \langle Q, M_s \mathbf{1} \rangle ds + \frac{1}{1 - \rho} \int_{t/2}^t e^{-a(x)s} ds, \end{aligned}$$

and the last three terms tend to zero as  $t$  goes to infinity, by using (3.2) for the second one. Consequently, for all  $f \in C_b(\mathcal{X})$  and all  $x \neq 0$  we have  $|M_t f(x)| \leq \|f\|_\infty M_t \mathbf{1}(x) \rightarrow 0$  as  $t \rightarrow +\infty$ . Since for  $x = 0$  we have

$$M_t f(0) = f(0) + \int_0^t \langle Q, M_s f \rangle ds \xrightarrow{t \rightarrow +\infty} f(0) + \langle \mu, f \rangle.$$

We can apply this convergence result to  $M_s f$ , for any  $s \geq 0$ , instead of  $f$ , because  $\|M_s \mathbf{1}\|_\infty < +\infty$ , to obtain that

$$\langle \delta_0 + \mu, f \rangle = \langle \delta_0 + \mu, M_s f \rangle.$$

Consequently  $\delta_0 + \mu$  is an eigenvector of  $\mathcal{A}$  and, by uniqueness, is then equal to  $\gamma$  up to the multiplicative constant  $(1 - \rho)$ . The proof of Theorem 1.1-(3b) is complete.  $\square$

**Remark 3.1** (An example). *Duhamel's formula gives that  $M_t f(x) \geq e^{-ta(x)} f(x)$  for any non-negative function  $f$  and  $t \geq 0$ . When  $X = [0, 1]$ ,  $Q(x) = 1$ ,  $a(x) = x^p$  with  $p \in (0, 1)$  and  $f(x) = x^q$  with  $q > -1$ , this gives*

$$\int_0^1 M_t f(x) dx \geq \int_0^1 e^{-tx^p} x^q dx = \frac{t^{-\frac{1+q}{p}}}{p} \int_0^t e^{-y} y^{\frac{q+1}{p}-1} dy.$$

We proved that the left-hand side tends to 0, since  $\rho = 1/(p+1) < 1$ , and we see that the rate of convergence in total variation distance is slower than any polynomial rate.

#### 4. THE NONLINEAR CONSERVATIVE EQUATION

We turn now to the nonlinear replicator-mutator equation (1.1). For  $v_0 \in \mathcal{P}(X)$ , we say that  $(v_t)_{t \geq 0}$  is solution to Equation (1.1) if it belongs to  $C([0, T], \mathcal{P}(X))$ ,  $t \mapsto av_t$  belongs to  $L^1([0, T], \mathcal{M}(X))$ , and for all  $t \geq 0$

$$v_t = e^{-(a+1)t} v_0 + \int_0^t e^{-(a+1)(t-s)} (Q + \langle v_s, a \rangle v_s) ds. \quad (4.1)$$

We can prove, as in [6] where the replicator-mutator equation with convolutive mutations is studied, that for any  $v_0 \in \mathcal{P}(X)$  there is a unique solution to Equation (1.1) which is given by

$$v_t = \frac{u_t}{\langle u_t, \mathbf{1} \rangle},$$

where  $u_t = v_0 M_t$  is the unique solution to Equation (1.2) with initial datum  $v_0$ . Now we give the proof of Theorem 1.2.

*Proof of Theorem 1.2-(1)-(2b)-(2c)-(2d).* These results are consequences of Theorem 1.1-(1)-(2b)-(2c)-(2d) by writing

$$\begin{aligned} \|v_t - \gamma\| &= \left\| \frac{e^{-\lambda t} u_t}{\langle e^{-\lambda t} u_t, \mathbf{1} \rangle} - \frac{\langle v_0, h \rangle \gamma}{\langle v_0, h \rangle} \right\| \\ &= \left\| \frac{\langle \langle v_0, h \rangle \gamma - e^{-\lambda t} u_t, \mathbf{1} \rangle e^{-\lambda t} u_t + \langle e^{-\lambda t} u_t, \mathbf{1} \rangle (e^{-\lambda t} u_t - \langle v_0, h \rangle \gamma)}{\langle e^{-\lambda t} u_t, \mathbf{1} \rangle \langle v_0, h \rangle} \right\| \\ &\leq \frac{\|e^{-\lambda t} u_t\|}{\langle e^{-\lambda t} u_t, \mathbf{1} \rangle} \frac{|\langle e^{-\lambda t} u_t - \langle v_0, h \rangle \gamma, \mathbf{1} \rangle|}{\langle v_0, h \rangle} + \frac{\|e^{-\lambda t} u_t - \langle v_0, h \rangle \gamma\|}{\langle v_0, h \rangle}, \end{aligned}$$

if the convergence  $\|e^{-\lambda t} u_t - \langle v_0, h \rangle \gamma\| \rightarrow 0$  yields the convergence  $|\langle e^{-\lambda t} u_t - \langle v_0, h \rangle \gamma, \mathbf{1} \rangle| \rightarrow 0$  with the same speed. It is true for the norm of  $\mathcal{M}(h)$  if  $a \in L^\infty$  due to the inequality

$$\|u\|_{\text{TV}} \leq \|1/h\|_\infty \|u\|_{\mathcal{M}(h)} = \alpha \|\lambda + a\|_\infty \|u\|_{\mathcal{M}(h)}.$$

In  $L^p(\gamma^{1-p}h)$  with  $p \in [1, 2)$ , it is true if  $a \in L^{p'-2}(Q)$  by virtue of Hölder's inequality

$$\|u\|_{L^1} \leq \|1/h\|_{L^{p'}} \|u/\gamma\|_{L^p(\gamma h)} = \alpha \left( \int_X (\lambda + a)^{p'-2} Q \right)^{1/p'} \|u\|_{L^p(\gamma^{1-p}h)}.$$

For  $p \in [2, \infty]$  we have by Jensen's inequality

$$\int_X (\lambda + a)^{p'-2} Q \leq \left( \int_X \frac{Q}{\lambda + a} \right)^{2-p'} = 1,$$

so that  $\|\cdot\|_{L^1} \leq \alpha \|\cdot\|_{L^p(\gamma^{1-p}h)}$  without needing further conditions than **(Ha)** and **(HQ)**.  $\square$

It remains to prove **(2a)** and **(3)** of Theorem 1.2. Note that when  $v_0(\{0\}) > 0$ , these results easily follow from Theorem 1.1-**(3b)**. We even get a better result than **(3)** in this case, namely a convergence without Cesàro mean. However the situation is trickier when  $v_0(\{0\}) = 0$  and we need the two following lemmas.

**Lemma 4.1.** *Assume that  $\rho \leq 1$ . Then, for all  $v_0 \in \mathcal{P}(X)$  and all  $f \in C_b(X)$  non-negative,*

$$\liminf_{t \rightarrow +\infty} \langle v_t, f \rangle \geq \langle Q/a, f \rangle.$$

*Proof.* Since  $v_t \geq 0$ , we deduce from (4.1) that for all  $t \geq 0$

$$v_t \geq Q \int_0^t e^{-(a+1)(t-s)} ds = \frac{Q}{a+1} (1 - e^{-(a+1)t}) \geq (1 - e^{-t}) \frac{Q}{a+1}.$$

Consequently, for any  $\epsilon \in (0, 1)$ , we have for all  $t \geq t_1 = -\log \epsilon$ ,

$$v_t \geq (1 - \epsilon) \frac{Q}{a+1}.$$

Using this estimate to bound from below the quantity  $\langle v_s, a \rangle$  in (1.1), we get that for all  $t \geq t_1$

$$v_t \geq \frac{Q}{a+1} (1 - e^{-(a+1)(t-t_1)}) + \alpha_1 \int_{t_1}^t e^{-(a+1)(t-s)} v(s) ds$$

where

$$\alpha_1 = (1 - \epsilon) \int_X \frac{aQ}{a+1} < 1 - \epsilon.$$

Grönwall's lemma then yields that for all  $t \geq t_1$ ,

$$v(t) \geq \frac{Q}{a+1-\alpha_1} (1 - e^{-(a+1-\alpha_1)(t-t_1)}).$$

Setting  $t_2 = t_1 + \frac{\log \epsilon}{\alpha_1 - 1}$  we get that for all  $t \geq t_2$

$$v_t \geq (1 - \epsilon) \frac{Q}{a+1-\alpha_1}.$$

Defining the sequences  $(t_n)$  and  $(\alpha_n)$  by  $\alpha_0 = t_0 = 0$  and

$$\alpha_{n+1} = (1 - \epsilon) \int_X \frac{aQ}{a+1-\alpha_n} < 1 - \epsilon \quad \text{and} \quad t_{n+1} = t_n + \frac{\log \epsilon}{\alpha_n - 1}$$

we have by induction that for all  $t \geq t_n$ ,

$$v_t \geq (1 - \epsilon) \frac{Q}{a+1-\alpha_n}$$

and consequently, for all  $f \geq 0$  in  $C_b(X)$ , all  $\epsilon \in (0, 1)$ , and all  $n \in \mathbb{N}$ ,

$$\liminf_{t \rightarrow +\infty} \langle v_t, f \rangle \geq \langle Q/(a+1-\alpha_n), f \rangle.$$

We now study the sequence  $(\alpha_n)$ . We deduce by a simple induction from

$$\alpha_{n+1} - \alpha_n = (\alpha_n - \alpha_{n-1}) (1 - \epsilon) \int_X \frac{aQ}{(a+1-\alpha_n)(a+1-\alpha_{n-1})},$$

that  $\alpha_{n+1} - \alpha_n > 0$ . The sequence  $(\alpha_n)_{n \geq 0}$  is increasing and bounded; it then converges to a limit  $\ell_\epsilon \in (0, 1 - \epsilon]$  which satisfies

$$\ell_\epsilon = (1 - \epsilon) \int \frac{aQ}{a+1-\ell_\epsilon} = (1 - \epsilon) \left( 1 - (1 - \ell_\epsilon) \int_X \frac{Q}{a+1-\ell_\epsilon} \right).$$

We now study the function  $\epsilon \mapsto \ell_\epsilon$ . For  $0 < \epsilon_1 < \epsilon_2 < 1$  we have

$$\begin{aligned} \ell_{\epsilon_1} - \ell_{\epsilon_2} &= (1 - \epsilon_1)(\ell_{\epsilon_1} - \ell_{\epsilon_2}) \int_{\mathcal{X}} \frac{aQ}{(a+1-\ell_{\epsilon_1})(a+1-\ell_{\epsilon_2})} + (\epsilon_2 - \epsilon_1) \int_{\mathcal{X}} \frac{aQ}{a+1-\ell_{\epsilon_2}} \\ &= A(\ell_{\epsilon_1} - \ell_{\epsilon_2}) + B \end{aligned}$$

with

$$A = (1 - \epsilon_1) \int_{\mathcal{X}} \frac{aQ}{(a+1-\ell_{\epsilon_1})(a+1-\ell_{\epsilon_2})} < \rho \leq 1 \quad \text{and} \quad B > 0,$$

which yields  $\ell_{\epsilon_1} - \ell_{\epsilon_2} \geq \frac{B}{1-A} > 0$ . The function  $\epsilon \mapsto \ell_\epsilon$  is then decreasing and it converges, when  $\epsilon \rightarrow 0$ , to a limit  $\ell_0 \in (0, 1]$  which satisfies

$$1 - \ell_0 = (1 - \ell_0) \int_{\mathcal{X}} \frac{Q}{a+1-\ell_0}.$$

Since  $\int_{\mathcal{X}} \frac{Q}{a+1-\ell_0} < \rho \leq 1$  if  $\ell_0 < 1$ , we necessarily have  $\ell_0 = 1$ , and finally for any  $f \geq 0$  in  $C_b(\mathcal{X})$

$$\liminf_{t \rightarrow +\infty} \langle v_t, f \rangle \geq \sup_{\epsilon \in (0,1)} \sup_{n \geq 0} \langle Q/(a+1-\alpha_n), f \rangle = \sup_{\epsilon \in (0,1)} \langle Q/(a+1-\ell_\epsilon), f \rangle = \langle Q/a, f \rangle.$$

□

**Lemma 4.2.** *For all  $v_0 \in \mathcal{P}(\mathcal{X})$  we have*

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t \langle v_s, a \rangle ds \leq 1.$$

*Proof.* Let  $t_0 > 0$ . From (4.1) we readily see that  $v_{t_0} \geq e^{-t_0(a+1)}Q$  and so, by virtue of (HX)-(HQ)-(Ha), for any  $\epsilon > 0$  there exists a set  $A_\epsilon \subset \mathcal{X}$  such that

$$\int_{A_\epsilon} v_{t_0} > 0 \quad \text{and} \quad \sup_{A_\epsilon} a \leq \epsilon.$$

Defining  $w_t = v_t - Q/a$ , Equation (1.1) also reads

$$\partial_t w_t = \partial_t v_t = -aw_t + \langle w_t, a \rangle v_t,$$

and so for all  $t \geq t_0$

$$v_t = v_{t_0} + \int_{t_0}^t (\langle w_s, a \rangle - a) v_s ds.$$

Integrating over  $A_\epsilon$  we get by Grönwall's lemma

$$\left( \int_{A_\epsilon} v_{t_0} \right) e^{\int_{t_0}^t \langle w_s, a \rangle ds - \epsilon t} \leq \int_{A_\epsilon} v_t \leq 1$$

and consequently

$$\frac{1}{t} \int_{t_0}^t \langle w_s, a \rangle ds \leq \epsilon - \frac{\log \left( \int_{A_\epsilon} v_0 \right)}{t},$$

Taking first the lim sup as  $t \rightarrow +\infty$  and letting then  $\epsilon$  go to zero, we find that

$$\limsup_t \frac{1}{t} \int_0^t \langle w_s, a \rangle ds \leq 0,$$

which is the desired result by definition of  $w_t$ . □

We are now in position to finish the proof of Theorem 1.2.

*Proof of Theorem 1.2 (2a).* Assume that  $\rho = \int_{\mathcal{X}} \frac{Q}{a} = 1$ . Since  $(v_t)_{t \geq 0}$  is a family of probability measures, there exists by weak-\* compactness a sub-sequence which converges in the vague topology to a positive measure  $\mu$  with mass  $\langle \mu, \mathbf{1} \rangle \leq 1$ . Due to Lemma 4.1, this limit must verify  $\mu \geq Q/a$ , and since the mass of  $Q/a$  is 1, the measure  $\mu$  is necessarily equal to  $Q/a$ . The uniqueness of the limit guarantees that the whole family  $(v_t)$  converges to  $Q/a$  for the vague topology and, since in  $\mathcal{P}(\mathcal{X})$  vague convergence is equivalent to narrow convergence, the proof is complete.  $\square$

*Proof of Theorem 1.2 (3).* Assume that  $\rho = \int_{\mathcal{X}} \frac{Q}{a} < 1$  and  $\lim_{|x| \rightarrow \infty} a(x) = +\infty$ , and define for all  $t \geq 0$  the probability measure

$$\bar{v}_t = \frac{1}{t} \int_0^t v_s ds.$$

Lemma 4.2 and the fact that  $a$  tends to  $+\infty$  at infinity guarantee that the family  $(\bar{v}_t)_{t \geq 0}$  is tight. Prokhorov's theorem then ensures the existence of a sub-sequence which converges in the narrow topology to a probability measure  $\mu$ , which must satisfy  $\mu \geq Q/a$  by virtue of Lemma 4.1. Since Lemma 4.2 also ensures that  $\langle \mu, a \rangle \leq \langle Q/a, a \rangle$ , and  $a(x) > 0$  for all  $x \neq 0$ , we deduce that  $\mu - Q/a$  must be supported by  $\{0\}$ . This means that  $\mu = \rho \delta_0 + Q/a$ , and the proof is complete since the whole family  $(\bar{v}_t)$  must converge to this unique limit.  $\square$

#### ACKNOWLEDGMENTS

The authors are grateful to the Jérôme Coville and Tristan Roget for discussion on the subject. The authors have been supported by ANR projects, funded by the French Ministry of Research ANR NOLO (ANR-20-CE40-0015). B.C. also received the support of the Chair ‘‘Modélisation Mathématique et Biodiversité’’ of VEOLIA-Ecole Polytechnique-MnHn-FX.

#### REFERENCES

- [1] A. S. Ackleh, J. Cleveland, and H. R. Thieme. Population dynamics under selection and mutation: long-time behavior for differential equations in measure spaces. *J. Differential Equations*, 261(2):1472–1505, 2016.
- [2] A. S. Ackleh, B. G. Fitzpatrick, and H. R. Thieme. Rate distributions and survival of the fittest: a formulation on the space of measures. *Discrete Contin. Dyn. Syst. Ser. B*, 5(4):917–928, 2005.
- [3] A. S. Ackleh, D. F. Marshall, H. E. Heatherly, and B. G. Fitzpatrick. Survival of the fittest in a generalized logistic model. *Math. Models Methods Appl. Sci.*, 9(9):1379–1391, 1999.
- [4] M. Alfaro and R. Carles. Explicit solutions for replicator-mutator equations: extinction versus acceleration. *SIAM Journal on Applied Mathematics*, 74(6):1919–1934, 2014.
- [5] M. Alfaro and R. Carles. Replicator-mutator equations with quadratic fitness. *Proceedings of the American Mathematical Society*, 145(12):5315–5327, 2017.
- [6] M. Alfaro, P. Gabriel, and O. Kavian. Confining integro-differential equations originating from evolutionary biology: ground states and long time dynamics. Oct. 2021. arXiv:2110.07214.
- [7] C. Ané, D. Bakry, and M. Ledoux. *Sur les inégalités de Sobolev logarithmiques*, volume 10. Société mathématique de France Paris, 2000.
- [8] D. Bakry. L'hypercontractivité et son utilisation en théorie des semigroupes. In *Lectures on probability theory*, pages 1–114. Springer, 1994.
- [9] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348. Springer Science & Business Media, 2013.
- [10] G. Barles, S. Mirrahimi, and B. Perthame. Concentration in Lotka-Volterra parabolic or integral equations: a general convergence result. *Methods Appl. Anal.*, 16(3):321–340, 2009.
- [11] O. Bonnefon, J. Coville, and G. Legendre. Concentration phenomenon in some non-local equation. *Discrete Contin. Dyn. Syst. Ser. B*, 22(3):763–781, 2017.
- [12] E. Bouin, J. Dolbeault, L. Lafleche, and C. Schmeiser. Hypocoercivity and sub-exponential local equilibria. *Monatshefte für Mathematik*, 194(1):41–65, 2021.
- [13] R. Bürger. Perturbations of positive semigroups and applications to population genetics. *Mathematische Zeitschrift*, 197(2):259–272, 1988.
- [14] R. Bürger. *The mathematical theory of selection, recombination, and mutation*. Wiley Series in Mathematical and Computational Biology. John Wiley & Sons, Ltd., Chichester, 2000.

- [15] R. Bürger and I. M. Bomze. Stationary distributions under mutation-selection balance: structure and properties. *Advances in applied probability*, 28(1):227–251, 1996.
- [16] J. A. Cañizo and A. Molino. Improved energy methods for nonlocal diffusion problems. *Discrete Contin. Dyn. Syst.*, 38(3):1405–1425, 2018.
- [17] A. Calsina and S. Cuadrado. Small mutation rate and evolutionarily stable strategies in infinite dimensional adaptive dynamics. *J. Math. Biol.*, 48(2):135–159, 2004.
- [18] A. Calsina and S. Cuadrado. Asymptotic stability of equilibria of selection-mutation equations. *J. Math. Biol.*, 54(4):489–511, 2007.
- [19] A. Calsina, S. Cuadrado, L. Desvillettes, and G. Raoul. Asymptotics of steady states of a selection-mutation equation for small mutation rate. *Proc. Roy. Soc. Edinburgh Sect. A*, 143(6):1123–1146, 2013.
- [20] A. Calsina, S. Cuadrado, L. Desvillettes, and G. Raoul. Asymptotic profile in selection-mutation equations: Gauss versus Cauchy distributions. *J. Math. Anal. Appl.*, 444(2):1515–1541, 2016.
- [21] P. Cattiaux, I. Gentil, and A. Guillin. Weak logarithmic sobolev inequalities and entropic convergence. *Probability theory and related fields*, 139(3):563–603, 2007.
- [22] P. Cattiaux, A. Guillin, and C. Roberto. Poincaré inequality and the  $L^p$  convergence of semi-groups. *Electron. Commun. Probab.*, 15:270–280, 2010.
- [23] J. A. Cañizo and S. Mischler. Harris-type results on geometric and subgeometric convergence to equilibrium for stochastic semigroups. Oct. 2021. arXiv:2110.09650.
- [24] D. Chafaï. Entropies, convexity, and functional inequalities, on  $\phi$ -entropies and  $\phi$ -sobolev inequalities. *Journal of Mathematics of Kyoto University*, 44(2):325–363, 2004.
- [25] N. Champagnat, R. Ferrière, and S. Méléard. From individual stochastic processes to macroscopic models in adaptive evolution. *Stochastic Models*, 24(sup1):2–44, 2008.
- [26] M. Chen. *From Markov chains to non-equilibrium particle systems*. World Scientific, 2004.
- [27] J. Coville. On a simple criterion for the existence of a principal eigenfunction of some nonlocal operators. *Journal of Differential Equations*, 249(11):2921–2953, 2010.
- [28] J. Coville. Singular measure as principal eigenfunction of some nonlocal operators. *Applied Mathematics Letters*, 26(8):831–835, 2013.
- [29] L. Desvillettes, P.-E. Jabin, S. Mischler, and G. Raoul. On selection dynamics for continuous structured populations. *Commun. Math. Sci.*, 6(3):729–747, 2008.
- [30] P. Diaconis and L. Miclo. On quantitative convergence to quasi-stationarity. *Ann. Fac. Sci. Toulouse Math. (6)*, 24(4):973–1016, 2015.
- [31] O. Diekmann, P.-E. Jabin, S. Mischler, and B. Perthame. The dynamics of adaptation: an illuminating example and a hamilton–jacobi approach. *Theoretical population biology*, 67(4):257–271, 2005.
- [32] R. Douc, G. Fort, and A. Guillin. Subgeometric rates of convergence of  $f$ -ergodic strong Markov processes. *Stochastic Process. Appl.*, 119(3):897–923, 2009.
- [33] N. Fournier and B. Perthame. Transport distances for PDEs: the coupling method. *EMS Surv. Math. Sci.*, 7(1):1–31, 2020.
- [34] N. Fournier and B. Perthame. A nonexpanding transport distance for some structured equations. *SIAM J. Math. Anal.*, 53(6):6847–6872, 2021.
- [35] S. Genieys, V. Volpert, and P. Auger. Pattern and waves for a model in population dynamics with nonlocal consumption of resources. *Math. Model. Nat. Phenom.*, 1(1):65–82, 2006.
- [36] M.-E. Gil, F. Hamel, G. Martin, and L. Roques. Mathematical properties of a class of integro-differential models from population genetics. *SIAM Journal on Applied Mathematics*, 77(4):1536–1561, 2017.
- [37] M.-E. Gil, F. Hamel, G. Martin, and L. Roques. Dynamics of fitness distributions in the presence of a phenotypic optimum: an integro-differential approach. *Nonlinearity*, 32(10):3485, 2019.
- [38] M. Hairer. P@w course on the convergence of markov processes. <http://www.hairer.org/Teaching.html>, 2010.
- [39] R. Holley and D. W. Stroock. Logarithmic sobolev inequalities and stochastic ising models. 1986.
- [40] P.-E. Jabin and G. Raoul. On selection dynamics for competitive interactions. *J. Math. Biol.*, 63(3):493–517, 2011.
- [41] O. Kavian, S. Mischler, and M. Ndao. The Fokker-Planck equation with subcritical confinement force. *J. Math. Pures Appl. (9)*, 151:171–211, 2021.
- [42] M. Kimura. A stochastic model concerning the maintenance of genetic variability in quantitative characters. *Proceedings of the National Academy of Sciences of the United States of America*, 54(3):731, 1965.
- [43] J. F. Kingman. A simple model for the balance between selection and mutation. *Journal of Applied Probability*, 15(1):1–12, 1978.
- [44] M. Ledoux. Concentration of measure and logarithmic sobolev inequalities. In *Seminaire de probabilités XXXIII*, pages 120–216. Springer, 1999.

- [45] T. Lorenzi and C. Pouchol. Asymptotic analysis of selection-mutation models in the presence of multiple fitness peaks. *Nonlinearity*, 33(11):5791–5816, 2020.
- [46] A. Lorz, S. Mirrahimi, and B. Perthame. Dirac mass dynamics in multidimensional nonlocal parabolic equations. *Communications in Partial Differential Equations*, 36(6):1071–1098, 2011.
- [47] P. Magal. Mutation and recombination in a model of phenotype evolution. *J. Evol. Equ.*, 2(1):21–39, 2002.
- [48] P. Magal and G. F. Webb. Mutation, selection, and recombination in a model of phenotype evolution. *Discrete Contin. Dynam. Systems*, 6(1):221–236, 2000.
- [49] F. Malrieu. Some simple but challenging Markov processes. *Ann. Fac. Sci. Toulouse Math. (6)*, 24(4):857–883, 2015.
- [50] S. Mirrahimi, B. Perthame, and J. Y. Wakano. Direct competition results from strong competition for limited resource. *J. Math. Biol.*, 68(4):931–949, 2014.
- [51] W. Oçafrain. Convergence to quasi-stationarity through poincaré inequalities and bakry-emery criteria. *Electronic Journal of Probability*, 26:1–30, 2021.
- [52] B. Perthame. *Transport equations in biology*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [53] B. Perthame and G. Barles. Dirac concentrations in lotka-volterra parabolic pdes. *Indiana University Mathematics Journal*, pages 3275–3301, 2008.
- [54] G. Raoul. Long time evolution of populations under selection and vanishing mutations. *Acta Appl. Math.*, 114(1-2):1–14, 2011.
- [55] G. Raoul. Local stability of evolutionary attractors for continuous structured populations. *Monatsh. Math.*, 165(1):117–144, 2012.
- [56] M. Röckner and F.-Y. Wang. Weak poincare inequalities and l2-convergence rates of markov semigroups. *Journal of Functional Analysis*, 185(2):564–603, 2001.
- [57] L. S. Tsimring, H. Levine, and D. A. Kessler. Rna virus evolution via a fitness-space model. *Physical review letters*, 76(23):4440, 1996.
- [58] M. Turelli. Heritable genetic variation via mutation-selection balance: Lerch’s zeta meets the abdominal bristle. *Theoretical Population Biology*, 25(2):138–193, 1984.
- [59] J. Y. Wakano, T. Funaki, and S. Yokoyama. Derivation of replicator–mutator equations from a model in population genetics. *Japan Journal of Industrial and Applied Mathematics*, 34(2):473–488, 2017.

(B. CLOEZ) MISTEA, INRAE, INTITUT AGRO, UNIV. MONTPELLIER, 2 PLACE PIERRE VIALA, 34060 MONTPELLIER, FRANCE.

*Email address:* `bertrand.cloez@inrae.fr`

(P. GABRIEL) LABORATOIRE DE MATHÉMATIQUES DE VERSAILLES, UVSQ, CNRS, UNIVERSITÉ PARIS-SACLAY, 45 AVENUE DES ÉTATS-UNIS, 78035 VERSAILLES CEDEX, FRANCE.

*Email address:* `pierre.gabriel@uvsq.fr`