



Chemical Composition and Antimicrobial and Antioxidant Activities of Mentha (longifolia L. and viridis) Essential Oils.

Mounira Mkaddem, Jalloul Bouajila, Monia Ennajar, Ahmed Lebrihi, Florence Mathieu, Mehrez Romdhane

► To cite this version:

Mounira Mkaddem, Jalloul Bouajila, Monia Ennajar, Ahmed Lebrihi, Florence Mathieu, et al.. Chemical Composition and Antimicrobial and Antioxidant Activities of Mentha (longifolia L. and viridis) Essential Oils.. Journal of Food Science, 2009, 74 (7), pp.M358-M363. <10.1111/J.1750-3841.2009.01272.X>. <hal-03608871>

HAL Id: hal-03608871

<https://hal.science/hal-03608871v1>

Submitted on 15 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 5905

To link to this article: DOI:10.1111/J.1750-3841.2009.01272.X
URL: <http://dx.doi.org/10.1111/J.1750-3841.2009.01272.X>

To cite this version: Mkaddem, Mounira and Bouajila, Jalloul and Ennajar, Monia and Lebrihi, Ahmed and Mathieu, Florence and Romdhane, Mehrez (2009) Chemical Composition and Antimicrobial and Antioxidant Activities of Mentha (longifolia L. and viridis) Essential Oils. *Journal of Food Science*, vol. 74 (n°7). pp. M358-M363. ISSN 0022-1147

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes.diff.inp-toulouse.fr

Chemical Composition and Antimicrobial and Antioxidant Activities of *Mentha* (*longifolia* L. and *viridis*) Essential Oils

MOUNIRA MKADDEM, JALLOUL BOUAJILA, MONIA ENNAJAR, AHMED LEBRIHI, FLORENCE MATHIEU, AND MEHREZ ROMDHANE

ABSTRACT: The study was aimed to investigate essential oil chemical composition (gas chromatography/flame ionization detection [GC-FID] and gas chromatography/mass spectrometry [GC-MS]) and antioxidant (1,1-diphenyl-2-picrylhydrazyl free radical (DPPH) and 2,2'-azinobis-3-ethylbenzothiazoline-6-sulphonate [ABTS] assays) and antimicrobial (Gram-positive and Gram-negative bacteria, fungi, and yeast) activities of essential oils extracted from leaves of *Mentha longifolia* L. and *Mentha viridis*. GC-MS analysis revealed that *M. longifolia* was constituted by pulegone (54.41%) as a major component followed by isomenthone (12.02%), 1,8-cineole (7.41%), borneol (6.85%), and piperitenone oxide (3.19%). *M. viridis* was rich in carvone (50.47%), 1,8-cineole (9.14%), and limonene (4.87%). The antioxidant activity by ABTS assay showed IC_{50} values of 476.3 ± 11.7 and 195.1 ± 4.2 mg/L for *M. longifolia* and *M. viridis*, respectively, the DPPH assays have resulted in a moderate IC_{50} (>8000 mg/L and 3476.3 ± 133 mg/L for *M. longifolia* and *M. viridis*, respectively). Antimicrobial activity showed that *Listeria monocytogenes* and *Klebsiella pneumoniae* bacteria were more inhibited by the 2 essential oils tested. *Escherichia coli* was least susceptible. A strong activity was also observed on fungi and yeasts. Carvone, thymol, and piperitenone oxide have not been detected in Tunisian *M. longifolia*. Camphor is reported for the 1st time for *M. viridis*. Antioxidant and antibacterial activities were correlated to chemical composition.

Keywords: antioxidant activity, antimicrobial activity, essential oil, *Mentha longifolia* L., *Mentha viridis* L.

Introduction

The genus *Mentha* L. (Lamiaceae), is widely distributed in all continents (except in South America and Antarctica) (Chambers 1992). The centers of variety of this genus are Europe, Australia, Central Asia, and North Africa (Harley and Brighton 1977; Chambers 1992). It groups spontaneous and cultivated forms. It is represented by 18 species and about 100 varieties and cultivars divided into 5 sections: Audibertia, Eriodontes, Mentha, Preslia, and Pulegium. The systematic of the genus is not very elucidated because of the strong morphologic variations, levels of ploidy ($2n = 2x = 24$ to $2n = 6x = 96$), and hybridizations intra- and interspecific (or between spontaneous and cultivated forms).

Several studies have reported the research of phylogenetic relations between different species on the basis of morphologic markers (Malinvaud 1880; Hüsnü 2002), chemical (Lawrence 1978; Chalchat and others 1997; Ræis Vasco and others 1999), caryologic (Harley and Brighton 1977; Singh and Sharma 1986), or molecular. Others researchers have aimed also the identification of specific composites of essential oils for its economic value and its use in pharmaceutical, cosmetic, food, confectionary, and beverage industries (Hendriks 1998; Kanatt and others 2007).

In Tunisia, the genus *Mentha* L. is represented by the species *M. rotundifolia* L., *M. longifolia* L. Huds., *M. spicata* (M. *viridis*) L., *M. aquatica* L., and *M. pulegium* L. (Maire 1938; Bonnier and De-Layens 1953; Pottier-Alapetite 1981). Many naturalized species (*M. piperita*, *M. spicata*, *M. longifolia*, and so on) are cultivated for family usages or for small commerce.

M. longifolia or wild mint is a fast-growing, perennial herb, growing wild in the northeastern part of Tunisia and cultivated in all regions. The species possesses antimicrobial and antioxidant properties (Economou and others 1991; Kaur and Kapoor 2002; Daferera and others 2003; Gulluce and others 2007). However, there has been no attempt to study antimicrobial and antioxidant activities of essential oils and extracts of *M. longifolia* from Tunisia.

M. viridis (green mint) is cultivated everywhere in Tunisia (humid to arid and semi-arid climate). It is considered as a simple variety obtained by culture of the wild mint. Green mint essential oil contains monoterpenoids like carvone, limonene, menthone, menthol, pulegone, and dihydrocarveol. Some of them were found to possess high antioxidant activity (Elmasta and others 2006). This composition can vary considerably according to the origin of the plant.

The green mint acquires a very powerful action on the nervous system (Bruneton 1993; Adersen and others 2006). The boiled leaves extract has an anti-infectious antifatulence effect and anti-inflammatory action (notably of the digestive system), it was counseled in the viral hepatitis and colitis, gastric acidities, aerophagia, to stimulate the digestion; furthermore, it presents some invigorating and stimulating qualities (Saleem and others 2000; Kouhila and others 2001; Kumar and Chattopadhyay 2007; Arumugam and others 2008).

Authors Mkaddem, Ennajar, and Romdhane are with Laboratoire de Modélisation Analyse et Commande des Systèmes. Ecole Natl. d'Ingénieurs de Gabès. Rue Omar elkhatab-ZRIG-6029 Gabès Tunisie. Author Bouajila is with Laboratoire de Synthèse et de Physicochimie de Molécules d'Intérêt Biologique, UMR CNRS 5068, Univ. Paul-Sabatier, 118 route de Narbonne, F-31062 Toulouse, France. Authors Lebrihi and Mathieu are with Univ. de Toulouse, LGC UMR 5503 (CNRS/INPT/UPS), ENSAT/ INPT, 1, Ave. de l'Agrobiopole, BP 32 607, Auzeville-Tolosane, 31 326 Castanet-Tolosan, France. Direct inquiries to author Bouajila (E-mail: bouajila@cict.fr).

Apart from being a stimulant and carminative, the mint plant is also known for its insecticidal (Papachristos and Stamopoulos 2002), antimicrobial, antispasmodic, and antiplatelet properties (Samarth and Kumar 2003; Ozgen and others 2006; Tognolini and others 2006).

Actually, essential oils and their components are gaining increasing interest because of their relatively safe status, their wide acceptance by consumers, and their exploitation for potential multi-purpose functional use (Ormancey and others 2001; Yadegarinia and others 2006). Many researchers have reported the antimicrobial, antifungal, and antioxidant properties of essential oils (Baratta and others 1998; Vardar-Unlü and others 2003; Mimica-Dukic and others 2004; Singh and others 2005; Sacchetti and others 2005). In fact, essential oils from mint species have strong antioxidant activity and good antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans*, and *Salmonella enteritidis* (Yadegarinia and others 2006; Gulluce and others 2007; Mahboubi and Haghi 2008). However, activity against fungi *Aspergillus ochraceus* and yeast species as *Saccharomyces cerevisiae* has not been reported previously only for *Menta arvensis* (Basilico and Basilico 1999). The *Mucor ramannianus* has not been tested until now with essential oils of *M. longifolia* L. and *M. viridis* L. The search for antioxidant and antimicrobial activities of Tunisian *Mentha* species essential oils is also not explored.

Therefore, the objectives of this study were to analyze chemical composition of hydrodistilled essential oils of *M. longifolia* and *M. viridis* by a GC-MS system to determine the essential oils chemotypes and investigate their antimicrobial and antioxidant activities.

Materials and Methods

Chemicals used

All chemicals used were of analytical reagent grade. All reagents were purchased from Sigma-Aldrich and Fluka (Saint-Quentin, France).

Essential oil isolation

Fresh leaves of *M. longifolia* L. and *M. viridis* L. cultivated species were collected (May 2007) from the region of Gabes Tunisia.

About 100 g of fresh leaves from each sample of *Mentha* were placed in 400 mL distilled water and submitted to hydrodistillation for 3 h using a Clevenger-type 5 apparatus (British Pharmacopoeia 1980). The condensation of each essential oil gives 2 phases: an organic phase was formed practically by essential oils (Hpo) to which we added anhydrous sodium sulfate to eliminate water traces and an aqueous phase that contains a negligible quantity of essential oil. Essential oils were stored at 4 °C until tested and analyzed.

Gas chromatography and gas chromatography-mass spectrometry

Quantitative and qualitative analysis of essential oils was carried out by gas chromatography-flame ionization detection (GC-FID) and gas chromatography-mass spectrometry (GC-MS). Gas chromatography analyses were carried out on a Varian Star 3400 (Les Ulis, France) Cx chromatograph fitted with a fused silica capillary DB-5MS (5% phenylmethylpolysiloxane, 30 m × 0.25 mm, film thickness 0.25 µm). Chromatographic conditions were a 60 to 260 °C temperature rise with a gradient of 5 °C/min and 15 min isotherm at 260 °C. A 2nd gradient was applied to 340 °C at 40 °C/min. Total analysis time was 57 min. For analysis, essential oils were dissolved in petroleum ether. One microliter of sample was injected in the split mode ratio of 1 : 10. Helium (purity

99.999%) was used as carrier gas at 1 mL/min. The injector was operated at 200 °C. The mass spectrometer (Varian Saturn GC-MS/MS 4D) was adjusted for an emission current of 10 µA and electron multiplier voltage between 1400 and 1500 V. Trap temperature was 150 °C and that of the transfer line was 170 °C. Mass scanning was from 40 to 650 amu.

Compounds were identified by comparison of their KI (Kovats indices) relative to C5-C24 *n*-alkanes obtained on a nonpolar DB-5MS column, with those provided in the literature, by comparison of their mass spectra with those recorded in NIST 08 (Natl. Inst. of Standards and Technology) and reported in published articles and by co-injection of available reference compounds. The samples were analyzed in duplicate. The percentage composition of essential oils was computed by the normalization method from the GC peak areas, assuming identical mass response factor for all compounds. Results were calculated as mean values of 2 injections from essential oil, without using correction factors.

Free radical scavenging activity: DPPH test

Antioxidant scavenging activity was studied using 1,1-diphenyl-2-picrylhydrazyl free radical (DPPH) as described by Blois (1958) with some modifications. Briefly, 1.5 mL of various dilutions of the test materials (2 mg/mL vitamin C were diluted 1/200, 1/400, 1/800, 1/1000, 1/1200; 16 mg/mL of essential oil *M. longifolia* were diluted 1/2, 1/3, 1/4, 1/5, 1/10; 16 mg/mL of essential oil *M. viridis* were diluted 1/2, 1/3, 1/4, 1/5, 1/10) were mixed with 1.5 mL of 0.2 mM methanolic DPPH solution. After an incubation period of 30 min at 25 °C, the absorbance at 520 nm was measured. The wavelength of maximum absorbance of DPPH was recorded as $A_{(\text{sample})}$, using a spectrophotometer (Helios, Unicam, Cambridge, U.K.). A blank experiment was also carried out applying the same procedure to a solution without the test material and the absorbance was recorded as $A_{(\text{blank})}$. The free radical scavenging activity of each solution was then calculated as percent inhibition according to the following equation:

$$\% \text{ inhibition} = [(A_{(\text{blank})} - A_{(\text{sample})}) / A_{(\text{blank})}] \times 100$$

Antioxidant activity of standard or essential oils was expressed as IC_{50} , defined as the concentration of the test material required to cause a 50% decrease in initial DPPH concentration. Ascorbic acid was used as a standard. All measurements were performed in triplicate.

ABTS radical scavenging assay

The radical scavenging capacity of the 2,2-AzinoBis-3-ethylbenzoThiazoline-6-Sulphonate radical cation (ABTS) was determined as described by Re and others (1999). ABTS was generated by mixing a 7 mM of ABTS at pH 7.4 (5 mM NaH_2PO_4 , 5 mM Na_2HPO_4 , and 154 mM NaCl) with 2.5 mM potassium persulfate (final concentration) followed by storage in the dark at room temperature for 16 h before use. The mixture was diluted with ethanol to give an absorbance of 0.70 ± 0.02 units at 734 nm using a spectrophotometer. For each sample, diluted methanol solution (2 mg/mL of vitamin C were diluted 1/200, 1/400, 1/800, 1/1000, 1/1200; 6 mg/mL of essential oil *M. longifolia* were diluted 1/4, 1/5, 1/10, 1/20, 1/50; 2 mg/mL of essential oil *M. viridis* were diluted 1/4, 1/5, 1/10, 1/20, 1/50) of the essential oil (100 µL) was allowed to react with fresh ABTS solution (900 µL), and then the absorbance was measured 6 min after initial mixing. Ascorbic acid was used as a standard. The capacity of free radical scavenging was expressed by IC_{50} (mg/L) values, which represents the

concentration required to scavenge 50% of ABTS radicals. The capacity of free radical scavenging IC_{50} was determined using the same equation previously used for the DPPH method. All measurements were performed in triplicate.

Microbial strains

All strains were obtained from the Laboratory of Chemical Engineering, Bioprocess Systems Microbiens Dept. of High Natl. School of Agronomy of Toulouse. The different essential oils were

Table 1 – Chemical composition (percent) of essential oils extracted from *Mentha longifolia* and *Mentha viridis* leaves using hydrodistillation methods.

Nr	KI ^a	Compounds	<i>M. longifolia</i> (%)	<i>M. viridis</i> (%)
1	936	α -pinene	0.52	0.50
2	976	sabinene	0.70	0.20
3	985	β -myrcene	0.04	0.19
4	990	3-octanol	—	0.54
5	1011	δ -3-carene	0.18	0.34
6	1025	p-cymene	0.04	0.33
7	1028	limonene	0.12	4.87
8	1030	1,8-cineole	7.41	9.14
9	1057	γ -terpinene	0.63	0.78
10	1072	linalool oxide	1.17	—
11	1086	α -terpinolene	—	0.05
12	1094	linalool	0.04	0.17
13	1105	p-menth-2-en-1-ol	—	1.20
14	1141	trans-pinocarveol	0.09	—
15	1143	camphor	1.67	3.68
16	1148	isopulegol	0.04	0.28
17	1155	menthone	0.18	—
18	1159	β -terpineol	—	0.30
19	1164	isomenthone	12.02	1.13
20	1166	borneol	6.85	0.74
21	1178	terpinen-4-ol	0.06	2.97
22	1185	α -terpineol	1.52	1.97
23	1190	dihydrocarveol ^b	—	0.45
24	1193	neodihydrocarveol ^b	—	0.69
25	1202	carveol	—	0.83
26	1205	verbenone	0.19	—
27	1238	pulegone	54.41	1.11
28	1241	carvone	—	50.47
29	1285	bornyl acetate	0.37	0.64
30	1330	1,2,4-trimethoxybenzene ^b	0.65	—
31	1350	eugenol	—	0.27
32	1352	terpinyl acetate	0.61	0.43
33	1366	dihydroeugenol	0.18	0.18
34	1369	piperitenone oxide	3.19	0.48
35	1389	β -elemene	—	1.00
36	1390	β -cubebene	—	0.12
37	1394	cis-jasmone	0.16	1.16
38	1406	methyl eugenol	—	0.36
39	1415	β -caryophyllene	2.04	3.00
40	1432	β -gurjunene	—	0.51
41	1438	geranyl acetone	0.04	—
42	1439	α -guaiene	—	1.01
43	1450	α -himachalene	0.17	0.17
44	1454	α -humulene	2.97	0.62
45	1482	β -selinene	0.33	1.86
46	1486	α -selinene	—	0.51
47	1499	α -muurolene	0.34	—
48	1512	calamenene ^b	—	0.22
49	1513	γ -cadinene	0.31	1.12
50	1524	δ -cadinene	—	0.26
51	1534	α -cadinene	—	0.20
52	1580	β -caryophyllene oxide	0.24	0.86
53	1636	T-muurolol	0.26	0.51
54	1678	cis- α -santalol	—	0.42
55	ND	grandisol ^b	—	0.27
56	ND	dimethylfuran lactone ^b	—	0.11
Total			99.72	99.18
Monoterpene hydrocarbons			2.19	6.93
Monoterpenes oxygenated			89.18	77.22
Sesquiterpenes hydrocarbons			6.16	10.38
Sesquiterpenes Oxygenated			0.5	1.79
Other			1.71	2.90
Phenolics			0.18	0.45

^aDB-5MS nonpolar column relative to C8-C24 n-alkanes.

^bTentatively identified supported by good match of MS.

ND = not determined.

individually tested against a panel of microorganisms including 2 Gram-positive (*S. aureus* CIP7625, *Listeria monocytogenes* Scott A 724), 2 Gram-negative (*E. coli* ATCC10536, *Klebsiella pneumoniae* CIP8291), 2 yeasts (*S. cerevisiae* ATCC 4226 A and *C. albicans* IPA 200), and 2 fungi (*M. ramannianus* ATCC 9314 and *A. ochraceus* NRRL 3174). The bacterial strains were cultured on nutrient broth for 48 h at 37 °C, while fungi and yeasts were propagated in CYB at 37 °C for 48 h to 3 d before use. All microorganisms were stocked at –20 °C in appropriate conditions and regenerated twice before use in the manipulations.

Well-diffusion assay

Agar well-diffusion method was employed for the determination of antimicrobial activity of essential oils (Feyza and others 2009). Briefly, 15 mL of nutrient agar and CYA, respectively, for bacterial and fungal strains were inoculated with 100 µL of microbial suspension containing 10⁸ CFU/mL of the indicator strain and then poured onto a petri dish and allowed to solidify. The dish was left to cool down and to solidify at room temperature for 30 min. Wells of 10 mm dia were aseptically bored into the culture medium. A sterile cox borer was used to punch wells and 15 µL of each essential oil were added to the wells. Each plate contained only 1 essential oil. Ampicillin and nalidixic acid (30 µg/well) were used as positive reference standards to determine the sensitivity of Gram-positive and Gram-negative bacterial species, respectively. Nystatin (30 µg/well) was used as positive reference standard to determine the sensitivity of fungi and yeast species. These plates, after staying at 4 °C for 2 h to allow dispersal, were incubated for 24 h at 37 °C for bacteria and 48 h at 30 °C for yeasts and fungi. The antimicrobial activity was visually appraised as inhibition zones surrounding the wells. The diameters of the inhibition zones were measured in millimeters. Tests were carried out in triplicate.

Statistical analysis

All data were expressed as means ± standard deviations (SD) of triplicate measurements. The confidence limits were set at $P < 0.05$. Standard deviations did not exceed 5% for the majority of the values obtained.

Results and Discussion

The extraction yield of hydrodistilled *M. longifolia* essential oil was 1.3% whereas the yield of *M. viridis* oil was 0.8%. In the literature (Kokkini and others 1995; Elmasta and others 2006), yield values were between 1.2% and 3.2% for the 2 species essential oils. The yield variability in the total essential oil content within the 2 species of mint could have resulted from the variation of climatic factors (Lawrence 1989; Kokkini and others 1995).

Chemical composition

The components identified in the 2 species of mint are listed in Table 1. Thirty-five components representing 99.72% of the total volatiles were identified for *M. longifolia* essential oil. The main components were oxygenated monoterpenes (89.18%). Pulegone (54.41% of the total essential oil) was the major compound followed by isomenthone (12.02%), 1,8-cineole (7.41%), borneol (6.85%), and piperitenone oxide (3.19%). *M. longifolia* including a high percentage of pulegone is reported also by Asekun and others (2007). Several compounds such as thymol (Gullus and others 2007), carvone (Kokkini and others 1995), or piperitone oxide (Ghoulami and others 2001) previously found in *M. longifolia* essential oil, have not been found in our samples.

Forty-nine components representing 99.18% of the total *M. viridis* essential oil were identified and characterized mainly by car-

vone (50.47%), followed by 1,8-cineole (9.14%), limonene (4.87%), camphor (3.68%), and β-caryophyllene (3%). Oxygenated monoterpenes corresponded to the main fraction (77.22%) of the total essential oil. *M. viridis* was rich in carvone as reported by other studies (Akhila and others 1980; Kokkini and others 1995). Camphor content for *M. viridis* is reported for the 1st time.

The 2 species showed an important variation in essential oil composition due to geographical and bioclimatic factors of the region. Different essential oil chemotypes have been described for *M. viridis* (Kokkini and others 1995) and *M. longifolia* (Gulluce and others 2007).

Antioxidant activity

Free radical scavenging activity of essential oils from 2 species of mint was investigated by ABTS assay. The ABTS has not been formerly tested for *Mentha* species. A moderate antioxidant activity was obtained to scavenge the ABTS radical cation (Table 2). *M. longifolia* essential oil (476.3 mg/L) was less active than *M. viridis* (195.1 mg/L).

The DPPH radical scavenging activity of essential oils confirms the result obtained with ABTS assay. Therefore, the DPPH activity of *M. viridis* essential oil (IC_{50} of 3476.3 mg/L) was more important than for *M. longifolia* (>8000 mg/L, Table 2). IC_{50} values obtained for Tunisian *M. longifolia* essential oil (>8000 mg/L) are comparable with Turkish *M. longifolia* essential oil (10700 ± 5.01 µg/mL) (Gulluce and others 2007). A previous study showed that methanol extract of *M. longifolia* was more able to reduce the stable free radical DPPH than that of essential oil (Gulluce and others 2007).

The essential oil of *M. viridis* contains more antioxidants than *M. longifolia* essential oil. Activity might be related to its phenolic content as reported in previous studies (Mimica-Dukic and others 1999; Gulluce and others 2007). Our study confirmed this result; in effect, the percentage of phenolics in *M. viridis* was 0.45% (dihydroeugenol and eugenol) relatively superior to that in *M. longifolia* essential oil that attained a percentage of 0.18% (dihydroeugenol).

Antimicrobial activity

Antimicrobial activity of essential oils was evaluated against 2 Gram-positive bacteria, 2 Gram-negative bacteria, 2 fungi, and 2 yeast species by zone diameter (Table 3). These microorganisms are morphologically and physiologically different. Essential oils exhibited a great potential for antimicrobial activity against 2 fungi (*M. ramannianus* and *A. ochraceus*). In fact, inhibition zone diameters for *M. ramannianus* (35 to 40 mm) and *A. ochraceus* (32 to 43 mm) were higher (Table 3).

This is the 1st study of antimicrobial activity of *M. longifolia* and *M. viridis* essential oils against yeast species (*C. albicans* and *S. cerevisiae*). Inhibition zones were obtained by essential oils of 2 species against yeast *C. albicans* (19 to 21 mm) and *S. cerevisiae* (25 to 28 mm). These activities may be related to major compounds of the *M. longifolia* and *M. viridis* essential oils (pulegone and carvone, respectively) but there is no previous study that reported antimicrobial activity of pulegone and carvone components against yeast.

Table 2 – Antioxidant activity of *Mentha longifolia* and *Mentha viridis* using ABTS and DPPH radical scavenging assay.

Samples	ABTS IC_{50} (mg/L)	DPPH IC_{50} (mg/L)
<i>Mentha longifolia</i>	476.3 ± 11.7	>8000
<i>Mentha viridis</i>	195.1 ± 4.2	3476.3 ± 133
Vitamin C	1.9 ± 0.1	4.4 ± 0.2

Table 3 – Antimicrobial activity (diameter of the inhibition zones in mm) of *Mentha longifolia* and *Mentha viridis* essential oils.

Antimicrobials	Microorganisms	Essential oils		References standard		
		<i>Mentha longifolia</i>	<i>Mentha viridis</i>	Ampicillin	Nalidixic acid	Nystatin
Fungi	<i>Mucor ramannianus</i> ATCC 9314	35	40	—	—	31
	<i>Aspergillus ochraceus</i> NRRL 3174	32	43	—	—	24
Yeast	<i>Candida albicans</i> IPA 200	19	21	—	—	25
	<i>Saccharomyces cerevisiae</i> ATCC 4226 A	28	25	—	—	29
Gram-positive bacteria	<i>Klebsiella pneumoniae</i> CIP8291	20	25	—	28	—
	<i>Escherichia coli</i> ATCC10536	—	—	—	30	—
Gram-negative bacteria	<i>Staphylococcus aureus</i> CIP7625	—	—	20	—	—
	<i>Listeria monocytogenes</i> Scott A 724	24	29	35	—	—

— = absence of inhibition zone detected; essential oils concentration: 15 μ L/well; reference standards concentration: 30 μ g/well.

The maximal inhibition zones for bacterial strains were 25 to 29 mm for *M. viridis* essential oil and 20 to 24 mm for *M. longifolia* essential oil against, respectively, *K. pneumoniae* and *L. monocytogenes*. However, no antimicrobial activity of essential oils tested was noted against *E. coli* and *S. aureus*. Results obtained in this study show that mint essential oils do not have selective antimicrobial activity on the basis of the cell-wall differences of bacterial microorganisms (Gram-positive or Gram-negative bacteria) as reported previously (Cosentino and others 1999; Karaman and others 2003). Referring to the literature, the antimicrobial activity of the tested essential oils may be due to monoterpene hydrocarbons (Carlo and others 2005). *M. viridis* essential oil contains more terpene hydrocarbons (monoterpenes and sesquiterpenes) than *M. longifolia* (Table 3). This result is in agreement with the greatest diameter of inhibition obtained for about 7/8 of microorganisms tested (except *S. cerevisiae*) for *M. viridis* essential oil (Table 3). Despite the feeble percentage of the monoterpene and sesquiterpene hydrocarbons for the 2 essential oils (17.31% for *M. viridis* and 8.35% for *M. longifolia*) the samples presented an interesting antimicrobial activity. Although these compounds are not abundant in essential oil, activity was important. It is necessary to signal that other compounds can contribute to improve this activity.

Conclusions

Carvone, thymol, and piperitone oxide have not been detected in Tunisian *M. longifolia*. Camphor is reported for the 1st time for *M. viridis*. Essential oils of *M. viridis* and *M. longifolia* have a medium antioxidant activity against ABTS and DPPH free radical scavenging. This activity is related to low phenolic content. Essential oils of Tunisian *M. longifolia* and *M. viridis* cultivated species exhibited a high antimicrobial activity against fungi and yeast species.

Essential oil represents a complex mixture of different chemical components. The antimicrobial effect of total essential oil cannot be reduced to major components. The results of this study showed that *M. longifolia* and *M. viridis* possess essential oils with important antimicrobial properties. Antibacterial activity could be influenced by terpene hydrocarbons (monoterpenes and sesquiterpenes) but other components can contribute to this activity. *In vivo* assay is also necessary to confirm antimicrobial activity of *M. longifolia* and *M. viridis*. These studies can be used in pharmaceuticals and natural therapies of infectious diseases for humans, and management of plant diseases.

Separation of terpene hydrocarbons, responsible for the antimicrobial activity, would be an interesting study to identify the molecule generating the good efficacy.

References

- Adersen A, Gauguin B, Gudiksen L, Jager AK. 2006. Screening of plants used in Danish folk medicine to treat memory dysfunction for acetylcholinesterase inhibitory activity. *J Ethnopharmacol* 104:418–22.
- Akhila A, Banthorpe DV, Rowan MG. 1980. Biosynthesis of Carvone in *Mentha*-*Spicata*. *Phytochemistry* 19:1433–8.
- Arumugam P, Gayatri Priya N, Subathra M, Ramesh A. 2008. Anti-inflammatory activity of four solvent fractions of ethanol extract of *Mentha spicata* L. investigated on acute and chronic inflammation induced rats. *Environmen Toxicol Pharmacol* 26:92–5.
- Asekun OT, Grierson DS, Afolayan AJ. 2007. Effects of drying methods on the quality and quantity of the essential oil of *Mentha longifolia* L. subsp. *Capensis*. *Food Chem* 101:995–8.
- Baratta MT, Dorman HJD, Deans SG, Figueiredo AC, Barroso JG, Ruberto G. 1998. Antimicrobial and antioxidant properties of some commercial essential oils. *Flav Fragr J* 13:235–44.
- Basilico MZ, Basilico JC. 1999. Inhibitory effects of some spice essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxin A production. *Lett Appl Microbiol* 29:238–41.
- Blois MS. 1958. Antioxidant determinations by the use of stable free radical. *Nature* 181:1199–200.
- Bonnier G, De-Layens G. 1953. Nouvelle flore pour la détermination facile des plantes de la région parisienne. Paris: Librairie Générale de l'Enseignement. 285 p.

- British Pharmacopoeia. 1980. Adsorbed tetanus vaccine. London: HMSO. 880 p.
- Bruneton J. 1993. Pharmacognosie-Phytochimie des Plantes Médicinales. London, Paris, and New York: Lavoisier Tech Doc. 915 p.
- Carlo IGT, Adam K, Valentina C, Maria TR, Sandro D, Paolo C. 2005. Chemical composition and antioxidant, antimicrobial, and antifungal activities of the essential oil of *Achillea ligustica* All. J Agric Food Chem 53:10148–53.
- Chalchat JC, Muhayimana A, Habimana JL, Chabard JL. 1997. Aromatic plants of Rwanda. II. Chemical composition of essential oils of ten *Eucalyptus* species growing in Ruhande arboretum, Butare, Rwanda. J Essen Oil Res 9:159–65.
- Chambers H. 1992. *Mentha*: genetic resources and the collection at USDA-ARSNCGR-Corvallis. Lam Newsl 1:3–4.
- Cosentino S, Tuberose CIG, Pisano B, Satta M, Mascia V, Arzedi E. 1999. *In vitro* antimicrobial activity and chemical composition of Sardinian *Thymus* essential oils. Lett Appl Microbiol 29:130–5.
- Daferera DJ, Ziogas BN, Polissiou MG. 2003. The effectiveness of plant essential oils on the growth of *Botrytis cinerea*, *Fusarium* sp. and *Clavibacter michiganensis* ssp. *Michiganensis*. Crop Prot 22:39–44.
- Economou KD, Oreopolou O, Thomopoulos CD. 1991. Antioxidant activity of some plant extracts of the family *Labiatae*. J Am Oil Chem Soc 68:109–13.
- Elmasta M, Dermirtas I, Isildak O, Aboul-Enein, HY. 2006. Antioxidant activity of *S-carvone* isolated from spearmint (*Mentha spicata* L. Fam. *Lamiaceae*). J Liq Chromatogr Relat Technol 29:1465–75.
- Feyza O, Belma A, Sahlan O, Senol A. 2009. Essential oil composition, antimicrobial and antioxidant activities of *Satureja cuneifolia* Ten. Food Chem 112:874–9.
- Ghoulami S, Il Idrissi A, Fkih-Tetouani S. 2001. Phytochemical study of *Mentha longifolia* of Morocco. Fitoterap 72(5):596–8.
- Gulluce M, Sahin F, Sokmen M, Ozer H, Daferera D, Sokmen A, Polissiou M, Adiguzel A, Ozkan H. 2007. Antimicrobial and antioxidant properties of the essential oils and methanol extract from *Mentha longifolia* L. ssp. *Longifolia*. Food Chem 103:1449–56.
- Harley RM, Brighton CA. 1977. Chromosome number in the genus *Mentha* L. Bot J Lin Soc 74:71–96.
- Hendriks H. 1998. Pharmaceutical aspects of some *Mentha* herbs and their essential oils. Perfum Flavor 23:15–23.
- Hüsnü CB. 2002. Aromatic biodiversity among the flowering plant taxa of Turkey. Pure Appl Chem 74:527–45.
- Kanatt SR, Chander R, Sharma A. 2007. Antioxidant potential of mint (*Mentha spicata* L.) in radiation-processed lamb meat. Food Chem 100:451–8.
- Karaman I, Sahin F, Gulluce M, Ogutcu H, Sengul M, Adiguzel A. 2003. Antimicrobial activity of aqueous and methanol extracts of *Juniperus oxycedrus* L. J Ethnopharmacol 85:231–5.
- Kaur C, Kapoor HC. 2002. Anti-oxidant activity and total phenolic content of some Asian vegetables. Int J Food Sc Tech 37:153–61.
- Kokkini S, Vokou D. 1989. *Mentha spicata* (Lamiaceae) chemotypes growing wild in Greece. Econom Bot 43:192–202.
- Kokkini S, Karousou R, Lanaras T. 1995. Essential oils of spearmint (Carvone-rich) plants from the island of Crete (Greece). Biochem Syst Ecol 23:425–30.
- Kouhila M, Belghit A, Daguén M, Boutaleb BC. 2001. Experimental determination of the sorption isotherms of mint (*Mentha viridis*), sage (*Salvia officinalis*) and verben (*Lippia citriodora*). J Food Engr 47:281–7.
- Kumar A, Chattopadhyay S. 2007. DNA damage protecting activity and antioxidant potential of pudina extract. Food Chem 100:1377–84.
- Lawrence BM. 1978. A study of the monoterpene interrelationships in the genus *Mentha* with special reference to the origin of pulegone and menthofuran [PhD thesis]. Groningen, The Netherlands: State Univ.
- Lawrence BM. 1989. Labiatae oils-mother nature's chemical factory. Paper XIth, 11th Int Congress of Essential Oils, Fragrance and Flavour, New Delhi. 71 p.
- Mahboubi M, Haghi G. 2008. Antimicrobial activity and chemical composition of *Mentha pulegium* L. essential oil. J Ethnopharmacol 119:325–7.
- Maire R. 1938. Sur quelques plantes du Sahara occidentale L. Bull Soc Hist Nat Afr N XXVIII (8–9):1–480.
- Malinvaud E. 1880. Simple aperçu des hybrides dans le genre *Mentha*. Bull Soc Botan France 27:332–47.
- Mimica-Dukic N, Popovic M, Jakovljevic V, Szabo A, Gasic O. 1999. Pharmacological studies of *Mentha longifolia* phenolic extracts II. Hepatoprotective activity. Pharma Biol 37(3):221–4.
- Mimica-Dukic N, Bozin B, Sokovic M, Simin N. 2004. Antimicrobial and antioxidant activities of *Melissa officinalis* L. (Lamiaceae) essential oil. J Agric Food Chem 52:2485–9.
- Ormancey X, Sisalli S, Coutiere P. 2001. Formulation of essential oils in functional perfumery. Parf Cosm Actu 157:30–40.
- Ozgen U, Mavi A, Terzi Z, Yildirim A, Coskum M, Houghton PJ. 2006. Antioxidant properties of some medicinal Lamiaceae (Labiatae) species. Pharma Biol 44:107–12.
- Papachristos DP, Stamopoulos DC. 2002. Repellent, toxic and reproduction inhibitory effects of essential oil vapours on *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). J Stored Prod Res 38:117–28.
- Pottier-Alapetite G. 1981. Flore de la Tunisie, Angiospermes dicotylédones. Gamopétales. Pub Sci Tun 1:1–654.
- Raeis vasco EMC, Coelho JAP, Palavra AMF. 1999. Comparison of pennyroyal oils obtained by supercritical CO₂ extraction and hydrodistillation. Flav Frag J 14:156–60.
- Re R, Pellegrini N, Protaggente A, Pannala A, Yang M, Rice-Evans C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med 26:1231–7.
- Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R. 2005. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. Food Chem 91:621–32.
- Saleem M, Alam A, Sultana S. 2000. Attenuation of benzoyl peroxide-mediated cutaneous oxidative stress and hyperproliferative response by the prophylactic treatment of mice with spearmint (*Mentha spicata*). Food Chem Toxicol 38:939–48.
- Samarth RM, Kumar A. 2003. *Mentha piperita* (Linn) leaf extract provides protection against radiation induced chromosomal damage in bone marrow of mice. Indian J Exp Biol 41:229–37.
- Singh G, Marimuthu P, De Heluani Carola S, Catalan C. 2005. Antimicrobial and antioxidant potentials of essential oil and acetone extract of *Myristica fragrans* Houtt. (Ail Part). J Food Sci 70:M141–8.
- Singh TP, Sharma AK. 1986. *Mentha*-taxonomic status as interpreted through cytology, genetics and phytochemistry. Ind J Genet 46:198–208.
- Tognolini M, Barocelli E, Ballabeni V, Bruni R, Bianchi A, Chiavarini M, Impicciatore M. 2006. Comparative screening of plant essential oils: phenylpropanoid moiety as basic core for antiplatelet activity. Life Sci 78:1419–32.
- Yadegarinia D, Gachkar L, Rezaei MB, Taghizadeh M, Aastaneh SA, Rasooli I. 2006. Biochemical activities of Iranian *Mentha piperita* L. and *Myrtus communis* L. essential oils. Phytochemistry 67:1249–55.
- Vardar-Unlü G, Candan F, Sökmén A, Daferera D, Polissiou M, Sökmén M, Dönmez E, Tepe B. 2003. Antimicrobial and antioxidant activity of the essential oil and methanol extracts of *Thymus pectinatus* Fisch. et Mey. Var. *pectinatus* (Lamiaceae). J Agric Food Chem 51:63–7.