



HAL
open science

Moduli space of irregular rank two parabolic bundles over the Riemann sphere and its compactification

Arata Komyo, Frank Loray, Masa-Hiko Saito

► **To cite this version:**

Arata Komyo, Frank Loray, Masa-Hiko Saito. Moduli space of irregular rank two parabolic bundles over the Riemann sphere and its compactification. 2022. hal-03614445v1

HAL Id: hal-03614445

<https://hal.science/hal-03614445v1>

Preprint submitted on 20 Mar 2022 (v1), last revised 12 Oct 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MODULI SPACE OF IRREGULAR RANK TWO PARABOLIC BUNDLES OVER THE RIEMANN SPHERE AND ITS COMPACTIFICATION

ARATA KOMYO, FRANK LORAY, AND MASA-HIKO SAITO

ABSTRACT. In this paper, we study rank 2 (quasi) parabolic bundles over the Riemann sphere with an effective divisor and these moduli spaces. First we consider a criterium when a parabolic bundle admits a unramified irregular singular parabolic connection. Second, to give a good compactification of the moduli space of semistable parabolic bundles, we introduce a generalization of parabolic bundles, which is called refined parabolic bundles. Third, we discuss a stability condition of refined parabolic bundles and define elementary transformations of the refined parabolic bundles. Finally, we describe the moduli spaces of refined parabolic bundles when the dimensions of the moduli spaces are two. These are related to geometry of some weak del Pezzo surfaces.

1. INTRODUCTION

In this paper, we investigate the moduli space of rank 2 (quasi) parabolic bundles over the Riemann sphere with an effective divisor D and extend some results obtained in [21] when the effective divisor D is reduced. Here, let ν be a positive integer, fix points t_1, t_2, \dots, t_ν on \mathbb{P}^1 , and set $D = n_1[t_1] + \dots + n_\nu[t_\nu]$. A parabolic bundle (E, \mathbf{l}) on (\mathbb{P}^1, D) is a pair where E is a rank 2 vector bundle over \mathbb{P}^1 , $\mathbf{l} = \{l_i\}_{1 \leq i \leq \nu}$, and $l_i \subset E|_{n_i[t_i]}$ is a free $\mathcal{O}_{n_i[t_i]}$ -submodule of length n_i for each i .

Parabolic (vector) bundles over an irreducible smooth complex projective curve with a reduced effective divisor were introduced in Mehta–Seshadri [23]. To construct a good moduli space, a stability condition for the parabolic weights α was introduced. The moduli spaces of α -semistable parabolic bundles were constructed in [23], [6], and [9]. The effect on the moduli space of varying the parabolic weights was studied in [11], [5], [25], and [14]. This effect is interesting from the point of view of the birational geometry (in the sense of Mori’s program). A generalization of parabolic bundles was studied in [7] and [8]. Here the (generalized) parabolic bundles are defined over an irreducible smooth complex projective curve with an effective divisor, which is not necessary reduced. A stability condition of such parabolic bundles was introduced and the moduli spaces of α -semistable parabolic bundles were constructed in [8]. Finally a more general notion of parabolic bundles was studied by Yokogawa ([27] and [28]).

Usually, a parabolic bundle consists of a vector bundle E , filtrations $\mathbf{l} = \{l_i\}_{1 \leq i \leq \nu}$, and parabolic weights α . But in this paper, we call a pair (E, \mathbf{l}) parabolic bundle and call the filtration \mathbf{l} a parabolic structure. Parabolic structures are important for study of geometry of isomonodromic

Date: March 20, 2022.

2010 Mathematics Subject Classification. Primary 14H60, Secondary 14D20 14Q10.

Key words and phrases. Parabolic bundle, refined parabolic bundle, projective line, elementary transformation, moduli space, weak del Pezzo surface.

The first author is supported by JSPS KAKENHI: Grant Numbers JP17H06127 and JP19K14506. The second author is supported by CNRS and ANR-16-CE40-0008 project “Foliage”. The third author is supported by JSPS KAKENHI: Grant Number JP17H06127.

deformations (or Painlevé type equations). For example, parabolic connections were studied in [16], [17], and [15]. A parabolic connection is a pair of a logarithmic connection and a parabolic structure, which is compatible with the connection. Remark that parabolic connections are defined over an irreducible smooth complex projective curve with a reduced effective divisor. A stability condition for the parabolic weights α was introduced in [16] and [15]. Now we fix generic α so that α -stable = α -semistable. The moduli spaces of α -stable parabolic connections were constructed in [16] and [15]. The cases where the underlying effective divisors are not necessary reduced were studied in [18]. That is, unramified irregular singular parabolic connections were studied. A stability condition for the parabolic weights α was also introduced and the moduli spaces of α -stable unramified irregular singular parabolic connections were constructed in [18]. These moduli spaces play important roles to prove the geometric Painlevé property of the isomonodromic deformations of these connections ([16], [15], and [18]). To study geometry of the moduli spaces of parabolic connections (or unramified irregular singular parabolic connections), the moduli spaces of parabolic bundles are useful. Indeed, we have a correspondence from (unramified irregular singular) parabolic connections to parabolic bundles by forgetting the connections. Remark that the corresponding parabolic bundles are not necessary α -stable if (unramified irregular singular) parabolic connections are α -stable. But, this correspondence induces maps from Zariski open subsets on the moduli spaces of connections to the moduli spaces of α -stable parabolic bundles. We may expect that these maps have nice properties (for example, these are affine bundles and Lagrangian fibrations).

In this paper, we impose that the ranks of underlying vector bundles are 2 and base curves are the projective line \mathbb{P}^1 . So we will consider only special situations. But isomonodromic deformations in this situation are interesting. Indeed, these isomonodromic deformations correspond to

- Painlevé equation P_{VI} when $D = [t_1] + [t_2] + [t_3] + [t_4]$;
- Garnier systems when $\deg(D) \geq 5$ and D is reduced;
- Painlevé equations $P_V, P_{IV}, P_{III(D_6^{(1)})}$ and P_{II} when $D = [t_1] + [t_2] + 2[t_3]$, $D = [t_1] + 3[t_2]$, $D = 2[t_1] + 2[t_2]$, and $D = 4[t_1]$, respectively; and
- some irregular Garnier systems when $\deg(D) \geq 5$ and D is not reduced.

On the other hand, the special situation was studied in the point of view of the geometric Langlands problem [3], [4], and [13].

1.1. When the effective divisor is reduced. Our purpose is to extend some result on parabolic bundles in the cases where D is reduced. Now we recall some results on parabolic bundles with a reduced effective divisor $D = [t_1] + \cdots + [t_\nu]$, which are concerned with the contents of the present paper. Let $\Lambda = (\lambda_i^+, \lambda_i^-)_{1 \leq i \leq \nu}$ be a tuple of complex numbers such that $\sum_{i=1}^{\nu} (\lambda_i^+ + \lambda_i^-) + d = 0$. Here we impose $\sum_{i=1}^{\nu} \lambda_i^{\epsilon_i} \notin \bar{\mathbb{Z}}$ for any choice $(\epsilon_i)_{1 \leq i \leq \nu} \in \{+, -\}$. We say Λ is generic if Λ satisfies this condition. If two residual eigenvalues of a parabolic connection are λ_i^+ and λ_i^- for each t_i , then we call a Λ -parabolic connection the parabolic connection. We say a parabolic bundle is Λ -flat if it admits a Λ -parabolic connection. We have a criterium of Λ -flatness given by [3, Proposition 3]:

$$(1.1) \quad \text{End}(E, \mathbf{1}) = \mathbb{C} \iff (E, \mathbf{1}) \text{ is } \Lambda\text{-flat} \iff (E, \mathbf{1}) \text{ is undecomposable.}$$

Let $\mathfrak{Bun}(D, d)$ be the coarse moduli space of undecomposable parabolic bundles of degree d over (\mathbb{P}^1, D) . By the flatness criterium, the image of the moduli space of Λ -parabolic connections under the forgetful map is just $\mathfrak{Bun}(D, d)$. However, as observed in [3], [2] and [21], the coarse moduli space $\mathfrak{Bun}(D, d)$ is a non Hausdorff topological space, or a nonseparated scheme. We introduce a stability condition with respect to weights $\mathbf{w} \in (w_i) \in [0, 1]^\nu$ (as in [21, Definition 2.2]). The moduli space $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ of \mathbf{w} -semistable parabolic bundles of degree d is a normal irreducible projective

variety. The open subset of \mathbf{w} -stable bundles is smooth. For generic weights \mathbf{w} , \mathbf{w} -semistable = \mathbf{w} -stable and we get a smooth projective variety. If $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is not empty, the dimension of $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is $n - 3$. On the other hand, we have a following fact given by [21, Proposition 3.4]:

$$(1.2) \quad (E, \mathbf{1}) \text{ is indecomposable} \iff \mathbf{w}\text{-stable for a convenient choice of weights } \mathbf{w}.$$

As a consequence, the coarse moduli space $\mathfrak{Bun}(D, d)$ of indecomposable parabolic bundles of degree d is covered by those smooth projective varieties $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ when \mathbf{w} runs. When $\deg(D) = 4$ or $\deg(D) = 5$, the dimension of $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is 1 or 2, respectively. In these cases, we can give explicit description of $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ for some weights \mathbf{w} . We fix a real number $w \in [0, 1]$. We impose that weights \mathbf{w} satisfy the condition $w_1 = w_2 = \dots = w_\nu = w$. We call weights which satisfy this condition *democratic weights*. Now, we assume that $\deg(D) = 4$ and $d = -1$. Then we have

- $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$ is empty when $0 < w < \frac{1}{4}$, $\frac{3}{4} < w < 1$;
- $\mathfrak{Bun}_{\mathbf{w}}(D, -1) = \mathbb{P}^1$ when $\frac{1}{4} < w < \frac{1}{2}$;
- $\mathfrak{Bun}_{\mathbf{w}}(D, -1) = \mathbb{P}^1$ when $\frac{1}{2} < w < \frac{3}{4}$.

The coarse moduli space $\mathfrak{Bun}(D, d)$ is constructed by gluing $\mathfrak{Bun}_{\mathbf{w}}(D, -1) \cong \mathbb{P}^1$ ($\frac{1}{4} < w < \frac{1}{2}$) and $\mathfrak{Bun}_{\mathbf{w}}(D, -1) \cong \mathbb{P}^1$ ($\frac{1}{2} < w < \frac{3}{4}$) outside t_1, \dots, t_4 (for instance, see [21, Section 3.7]). Next, we assume that $\deg(D) = 5$ and $d = -1$. Then we have the following description of the moduli spaces $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$:

- $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$ is empty when $0 < w < \frac{1}{5}$;
- $\mathfrak{Bun}_{\mathbf{w}}(D, -1) = \mathbb{P}^2$ when $\frac{1}{5} < w < \frac{1}{3}$;
- $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$ is a del Pezzo surface of degree 4 when $\frac{1}{3} < w < \frac{3}{5}$;
- $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$ is a del Pezzo surface of degree 5 when $\frac{3}{5} < w < 1$.

(See [21, Section 6.1] and [13, Proposition 3.11]). When d is even, descriptions of the moduli spaces are given by [13, Theorem 3.2]. To cover the coarse moduli space $\mathfrak{Bun}(D, -1)$, thinking only democratic weights is not enough. By applying elementary transformations on $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$ with democratic weights \mathbf{w} , we can cover the coarse moduli space $\mathfrak{Bun}(D, -1)$ by the moduli spaces $\mathfrak{Bun}_{\mathbf{w}}(D, -1)$. Remark that the stability of parabolic bundles changes by applying the elementary transformations. If $w_1 = w_2 = \dots = w_\nu = \frac{1}{2}$, the stability of parabolic bundles is preserved by applying the elementary transformations. So some elementary transformations induce automorphisms on the moduli space $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ with $w = \frac{1}{2}$. In [1], a modular interpretation for the automorphism group of the moduli space $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ with any $\deg(D)$ and with any $\deg(E)$ is given by this point of view.

1.2. Results. Now we describe results of the present paper. The effective divisor D is not necessary reduced. First we will discuss a counterpart of (1.1). Let $\Lambda = (\lambda^+, \lambda^-)$ where $\lambda^\pm \in \Omega^1(D)/\Omega^1$. As in the reduced divisor cases, we impose that Λ satisfies some conditions. If the diagonalization of the principal parts of an unramified irregular singular parabolic connection at D are a diagonal matrix whose diagonal entries are λ^+ and λ^- , then we call a Λ -unramified irregular singular parabolic connection this unramified irregular singular parabolic connection. We say a parabolic bundle is Λ -flat if it admits a Λ -unramified irregular singular parabolic connection. If $\deg(D) = 4$ and $\deg(E)$ is odd, then

$$\text{End}(E, \mathbf{1}) = \mathbb{C} \iff (E, \mathbf{1}) \text{ is } \Lambda\text{-flat}$$

which is proved in [4, Proposition 4.8]. We can define indecomposability for parabolic bundles when the effective divisor D is not necessary reduced. If $(E, \mathbf{1})$ is Λ -flat, then $(E, \mathbf{1})$ is indecomposable. But the converse proposition is not true even when $\deg(D) = 4$ and $\deg(E)$ is odd. So in addition

to undecomposability, we assume the *admissibility* for parabolic bundles, which is defined in Definition 7 below. Using some material of [4, Section 4] and ideas of [3, Proposition 3], we can show that

$$(1.3) \quad (E, \mathbf{l}) \text{ is simple} \implies (E, \mathbf{l}) \text{ is } \Lambda\text{-flat} \implies (E, \mathbf{l}) \text{ is undecomposable and admissible.}$$

This is a counterpart of (1.1). Here we say (E, \mathbf{l}) is *simple* if $\text{End}(E, \mathbf{l}) = \mathbb{C}$. Counterexamples of the converse of the first implication of (1.3) are described in Proposition 11 and Proposition 13 below. Counterexamples of the converse of the second implication of (1.3) also are described in Proposition 11 and Proposition 13 (see also Example 31). If $\deg(D) = 4$ and $\deg(E)$ is odd, then we have

$$(1.4) \quad (E, \mathbf{l}) \text{ is undecomposable and admissible} \implies \text{End}(E, \mathbf{l}) = \mathbb{C}.$$

The implications (1.3) and (1.4) are described in Proposition 8 below. These are the first results.

Second we will discuss a counterpart of (1.2). We can introduce a stability condition with respect to weights $\mathbf{w} = (w_i) \in [0, 1]^\nu$, which is defined in Section 4 below. Let $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ be the moduli space of \mathbf{w} -semistable parabolic bundles. However $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is not projective in general, since we impose that each l_i is free in the definition of parabolic bundles. So, to construct a good compactification, we will define *refined parabolic bundles*, which are generalization of parabolic bundles. Let $l_{i, \bullet} = \{l_{i,k}\}_{1 \leq k \leq n_i}$ be a filtration $E|_{n_i[t_i]} \supset l_{i,n_i} \supset l_{i,n_i-1} \supset \cdots \supset l_{i,1} \supset 0$ of $\mathcal{O}_{n_i[t_i]}$ -modules such that the length of $l_{i,k}$ is k . We call $l_{i, \bullet} = \{l_{i,k}\}_{1 \leq k \leq n_i}$ a refined parabolic structure at t_i . If l_{i,n_i} is free, then the notion of refined parabolic structures coincides with the notion of parabolic structures. Refined parabolic bundles are vector bundles with refined parabolic structures. For weights

$$\mathbf{w} = (\mathbf{w}_1, \dots, \mathbf{w}_\nu), \quad \mathbf{w}_i = (w_{i,n_i}, \dots, w_{i,1}) \in [0, 1]^{n_i} \quad (1 \leq i \leq \nu),$$

where $0 \leq w_{i,n_i} \leq \cdots \leq w_{i,1} \leq 1$, we can define \mathbf{w} -stability and \mathbf{w} -semistability of a refined parabolic bundle (Definition 28 below).

Remark 1. *A compactification of the moduli space of α -stable Λ -parabolic connections is constructed in [16, Proposition 5.6]. This compactification is the moduli space of α -stable Λ - ϕ -parabolic connections. When D is reduced, $\deg(D) = 4$ and $\text{rank}(E) = 2$, the compactification with the boundary divisor is isomorphic to the Okamoto–Painlevé pair of P_{VI} ([17]). Here the notion of Okamoto–Painlevé pairs was introduced in [26]. In [24], a compactification of the moduli space of α -stable Λ -unramified irregular singular parabolic connections was studied when D is not reduced, $\deg(D) = 4$ and $\text{rank}(E) = 2$. However, the moduli space of α -stable Λ - ϕ -unramified irregular singular parabolic connections is projective but is not smooth. So, in [24], Miyazaki introduced refined parabolic structures, which are same as our refined parabolic structures. By considering refined parabolic structures, we have smooth compactifications, which are isomorphic to the Okamoto–Painlevé pairs of P_V , P_{IV} , $P_{III(D_6^{(1)})}$, and P_{II} .*

Remark 2. *The notion of the refined parabolic bundles is contained in the notion of the parabolic bundles in Yokogawa’s papers [27] and [28].*

We can define undecomposable refined parabolic bundles and admissible refined parabolic bundles, naturally.

Remark 3. *We have defined the Λ -flatness for (ordinary) parabolic bundles. On the other hand, we do not define the Λ -flatness for refined parabolic bundles, since unramified irregular singular parabolic connections have (ordinary) parabolic structures.*

To give a counterpart of (1.2), the conditions undecomposableness and admissibility are not sufficient. That is, there exist undecomposable and admissible refined parabolic bundles which is not \mathbf{w} -stable for any weights \mathbf{w} (Example 31 below).

Remark 4. *Moreover, there exist examples of Λ -flat parabolic bundles, which are \mathbf{w} -unstable for any weights \mathbf{w} .*

So we will define *tame* refined parabolic bundles in Definition 32. We will check that

$$(E, \mathbf{1}) \text{ is undecomposable and tame } \implies (E, \mathbf{1}) \text{ is admissible}$$

(described in Proposition 33 below). Then we have a counterpart of (1.2) as follows:

Theorem A (Theorem 1 and Corollary 2). *Let $(E, \mathbf{1})$ be a refined parabolic bundle of degree d over (\mathbb{P}^1, D) . We have the following equivalence relation:*

$$(E, \mathbf{1}) \text{ is undecomposable and tame } \iff \mathbf{w}\text{-stable for a convenient choice of weights } \mathbf{w}.$$

Let $(E, \mathbf{1})$ be a parabolic bundle of degree d over (\mathbb{P}^1, D) . Then

$$(E, \mathbf{1}) \text{ is simple } \iff \mathbf{w}\text{-stable for a convenient choice of weights } \mathbf{w}.$$

Third, we will discuss elementary transformations of refined parabolic bundles. We will define elementary transformations of refined parabolic bundles in Section 5.3 and we will show some properties of the elementary transformations in Section 5.4. For example, we will discuss a relation between the elementary transformations and the stability condition (Proposition 42).

Fourth, we will describe the moduli space of \mathbf{w} -stable refined parabolic bundles when $\deg(D) = 5$, $\deg(E) = 1$, and \mathbf{w} is generic democratic weights. Let $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, d)$ be the moduli space of \mathbf{w} -stable refined parabolic bundles of degree d . Here \mathbf{w} -semistable = \mathbf{w} -stable, since \mathbf{w} is generic. In this case, the dimension of $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, d)$ is 2. Since $\deg(D) = 5$, the effective divisor D is one of the following effective divisors:

$$\begin{aligned} D_{2111} &:= 2[t_1] + [t_2] + [t_3] + [t_4], & D_{221} &:= 2[t_1] + 2[t_2] + [t_3], & D_{311} &:= 3[t_1] + [t_2] + [t_3], \\ D_{32} &:= 3[t_1] + 2[t_2], & D_{41} &:= 4[t_1] + [t_2], & D_5 &:= 5[t_1]. \end{aligned}$$

For these cases, by Theorem A, we can make a list of refined parabolic bundles which are \mathbf{w} -stable for a convenient choice of weights \mathbf{w} . This list is described in the appendix. We will define types of the refined parabolic bundles by some numerical data. This type is related to behavior of the stability index when weights are varied. By Proposition 46 below, we have that $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) = \mathbb{P}^2$ when $\frac{1}{5} < w < \frac{1}{3}$. By this fact and the relation between the types and behavior of the stability index, we have the following theorem.

Theorem B (Section 6.1). *Assume that $\deg(D) = 5$, $\deg(E) = 1$, and weight $\mathbf{w} = (w_i)_i$ satisfies $w_{i,1} = w_{i,2} = \dots = w_{i,n_i} = w$ for any i ($0 < w < 1$), that is, \mathbf{w} are democratic weights.*

(i) *We have the following explicit descriptions of $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$:*

- $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ is empty when $0 < w < \frac{1}{5}$;
- $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) = \mathbb{P}^2$ when $\frac{1}{5} < w < \frac{1}{3}$;
- $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ is a weak del Pezzo surface of degree 4 when $\frac{1}{3} < w < \frac{3}{5}$;
- $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ is a weak del Pezzo surface of degree 5 when $\frac{3}{5} < w < 1$.

Let $\Pi_{-2}(D)$ and $\Pi'_{-2}(D)$ be the effective divisors consisting of all (-2) -curves on $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ when $\frac{1}{3} < w < \frac{3}{5}$ and $\frac{3}{5} < w < 1$, respectively. Then the configurations of $\Pi_{-2}(D)$ and $\Pi'_{-2}(D)$ are same, and the configuration is $A_1, 2A_1, A_2, A_2 + A_1, A_3$ and A_4 when $D = D_{2111}, D_{221}, D_{311}, D_{32}, D_{41},$ and D_5 , respectively.

- (ii) If $(E, \{l_{i,\bullet}\}_i)$ be a refined parabolic bundle corresponding to a point on $\Pi_{-2}(D)$ or $\Pi'_{-2}(D)$, then there exists i ($1 \leq i \leq \nu$) such that l_{i,n_i} is not free, that is, $l_{i,\bullet}$ is not a parabolic structure for this i .
- (iii) If $(E, \{l_{i,\bullet}\}_i)$ be a refined parabolic bundle corresponding to a point on $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ ($\frac{1}{5} < w < \frac{1}{3}$), $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \setminus \Pi_{-2}(D)$ ($\frac{1}{3} < w < \frac{3}{5}$), or $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \setminus \Pi'_{-2}(D)$ ($\frac{3}{5} < w < 1$), then l_{i,n_i} is free for any i , and $(E, \{l_{i,n_i}\}_i)$ is a simple parabolic bundle. In particular, $(E, \{l_{i,n_i}\}_i)$ is a Λ -flat for generic Λ .

These weak del Pezzo surfaces are listed in [12, Section 8.6.3]. Descriptions of the corresponding isomonodromic deformations are given by Kimura [20] (see also [19]). The corresponding isomonodromic deformations are the following:

the divisor	Π_{-2}	the isomonodromic deformations
D_{2111}	A_1	$H(1, 1, 1, 2)$
D_{221}	$2A_1$	$H(1, 2, 2)$
D_{311}	A_2	$H(1, 1, 3)$
D_{32}	$A_2 + A_1$	$H(2, 3)$
D_{41}	A_3	$H(1, 4)$
D_5	A_4	$H(5)$

Here the notations in the last column are in [20, pp.39–40].

Fifth, we will recover the geometry of the weak del Pezzo surfaces of degree 4 from the modular point of view. When weights \mathbf{w} are the democratic weights with $w = \frac{1}{2}$, the moduli spaces of refined parabolic bundles are weak del Pezzo surfaces of degree 4. We may reinterpret the negative curves on these surfaces as the loci of special refined parabolic bundles. Since weights \mathbf{w} are the democratic weights with $w = \frac{1}{2}$, we may show that the stability of refined parabolic bundles is preserved by applying the elementary transformations (Section 6.2). So some elementary transformations induce automorphisms on the moduli space $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, d)$ with $w = \frac{1}{2}$. We will reconstruct automorphisms of the weak del Pezzo surfaces of degree 4 via these elementary transformations of refined parabolic bundles.

The organization of this paper is as follows. In Section 2, we will define parabolic bundles over (\mathbb{P}^1, D) , where D is not necessary reduced. Next, in Section 2.1, we will show the implications (1.3) and (1.4), and will give a flatness criterium. In Section 2.2, we will give examples of the flatness criterium. In Section 3, we will define refined parabolic bundles and will discuss combinatorial structure of refined parabolic structures at a multiple point. This structure is related to Young tableaux. In Section 4, we will define a stability condition of refined parabolic bundles, and we will show Theorem A. In Section 5.3, we will define elementary transformations of refined parabolic bundles. In Section 5.4, we will show some properties of the elementary transformations of refined parabolic bundles. In Section 6.1, we will describe the moduli spaces when $\deg(D) = 5$, $\deg(E) = 1$, and weights \mathbf{w} are democratic weights. In Section 6.3, we will recover the geometry of the weak del Pezzo surfaces of degree 4 from the modular point of view. In the appendix, we will give a list of special refined parabolic bundles, which appear when $\deg(D) = 5$ and $\deg(E) = 1$.

Acknowledgments. The authors would like to warmly thank Professor Roman Fedorov for illuminating discussion helping us to understand the flatness criterium in Proposition 8.

2. PARABOLIC BUNDLES

Let D be an effective divisor on \mathbb{P}^1 , defined by

$$D = \sum_{i=1}^{\nu} n_i [t_i], \quad \deg(D) = \sum_{i=1}^{\nu} n_i =: n.$$

Take a generator f_i of the maximal ideal of $\mathcal{O}_{\mathbb{P}^1, t_i}$. Set $\mathcal{O}_{n_i[t_i]} := \mathcal{O}_{\mathbb{P}^1, t_i} / (f_i^{n_i})$. For a vector bundle E on \mathbb{P}^1 , we denote by $E|_{n_i[t_i]}$ the tensor product $E \otimes \mathcal{O}_{n_i[t_i]}$. Let $I = \{1, 2, \dots, \nu\}$ be the set of indexes of D . Take $\Lambda = (\lambda^+, \lambda^-)$, with $\lambda^\pm \in \Omega^1(D)/\Omega^1$. Set $\lambda_i^\pm := \lambda^\pm|_{n_i[t_i]} \in \Omega^1(D)/\Omega^1 \otimes \mathcal{O}_{n_i[t_i]}$ for each $i \in I$. We assume (as in [4, Section 2.1]):

- (a) the order of pole of $\lambda_i^+ - \lambda_i^-$ is precisely n_i ,
- (b) $\sum_i \text{Res}_{t_i} \lambda_i^+ + \lambda_i^- = d \in \mathbb{Z}$,
- (c) $\sum_i \text{Res}_{t_i} \lambda_i^\pm \notin \mathbb{Z}$ whatever are the signs \pm ,
- (d) if $n_i = 1$ then $\lambda_i^+ - \lambda_i^- \notin \mathbb{Z}$.

The relation in (b) is called Fuchs' relation and the condition (d) is called non resonant condition. Let E be a rank 2 vector bundle on \mathbb{P}^1 and $\nabla: E \rightarrow E \otimes \Omega^1(D)$ be a meromorphic connection with polar divisor D . If we say (E, ∇) is Λ -connection if its principal part at each pole $n_i[t_i]$ is given by

$$d + \begin{pmatrix} \lambda_i^+ & 0 \\ 0 & \lambda_i^- \end{pmatrix} : E|_{n_i[t_i]} \longrightarrow E|_{n_i[t_i]} \otimes \Omega^1(D)/\Omega^1$$

up to holomorphic gauge transformation. We call Λ the formal (or spectral) data of the meromorphic connection. A Λ -connection (E, ∇) is non-resonant by (a) and (d). The degree of E is d by (b). The following is [4, Lemma 4.6]:

Proposition 5. *Let Λ satisfying (a),(b),(c),(d) as above. Then, any Λ -connection (E, ∇) is irreducible.*

Proof. If $L \subset E$ is a ∇ -invariant line bundle, then $\nabla|_L : L \rightarrow L \otimes \Omega^1(D)$ is a connection with principal parts λ_i^\pm at each t_i . Therefore, Fuchs' relation for $\nabla|_L$ yields a contradiction with generic assumption (c) on residual eigenvalues. \square

Let (E, ∇) be a Λ -connection. By the condition of Λ , there is a unique free $\mathcal{O}_{n_i[t_i]}$ -submodule $l_i \subset E|_{n_i[t_i]}$ of length n_i for each i by the compatible with ∇ in the following sense: $\nabla(s) - \lambda_i^\pm s$ is regular at t_i for any i and every section s of E in a neighborhood of t_i such that $s|_{n_i[t_i]} \in l_i$. Let $\mathbf{l} = \{l_i\}_{i \in I}$ be the set of the free $\mathcal{O}_{n_i[t_i]}$ -submodule $l_i \subset E|_{n_i[t_i]}$ of length n_i given by a Λ -connection for each i . We call the triple (E, ∇, \mathbf{l}) *unramified irregular singular Λ -parabolic connection* (see [18, Definition 2.1]). After fixing convenient weights α , the moduli space of α -stable unramified irregular singular Λ -parabolic connections of rank 2 on (\mathbb{P}^1, D) forms a smooth quasi-projective scheme of dimension $2(n-3)$ (see [18, Section 2]). In our case, by Proposition 5, unramified irregular singular Λ -parabolic connections induced by Λ -connections are irreducible. So these unramified irregular singular Λ -parabolic connections satisfy the stability condition in [18, Definition 2.2]. So we identify α -stable unramified irregular singular Λ -parabolic connections with Λ -connections. Let $\mathfrak{Con}_\Lambda(D)$ be the moduli space of Λ -connections. In this paper, we are concerned with forgetting the connections from unramified irregular singular Λ -parabolic connections.

Definition 6. *We say (E, \mathbf{l}) (where $\mathbf{l} = \{l_i\}_{i \in I}$) is a parabolic bundle of rank 2 and of degree d over (\mathbb{P}^1, D) if E is a rank 2 vector bundle on \mathbb{P}^1 with $\deg(E) = d$ and $l_i \subset E|_{n_i[t_i]}$ is a free $\mathcal{O}_{n_i[t_i]}$ -submodule of length n_i for each i . We call l_i a parabolic structure at t_i .*

2.1. Flatness criterium. We say a parabolic bundle (E, \mathbf{l}) is Λ -flat if there exists a connection on E compatible with parabolic structure \mathbf{l} and formal data Λ . We say that a parabolic bundle $(E, \{l_i\}_{i \in I})$ is *decomposable* if there exists a decomposition $E = L_1 \oplus L_2$ such that $l_i = l_i^{(1)}$ or $l_i = l_i^{(2)}$ for any $i \in I$, where we set $l_i^{(1)} := l_i \cap (L_1|_{n_i[t_i]})$ and $l_i^{(2)} := l_i \cap (L_2|_{n_i[t_i]})$. We say that $(E, \{l_i\}_{i \in I})$ is *undecomposable* if $(E, \{l_i\}_{i \in I})$ is not decomposable. If D is a reduced divisor, then

$$\text{End}(E, \mathbf{l}) = \mathbb{C} \iff (E, \mathbf{l}) \text{ is } \Lambda\text{-flat} \iff (E, \mathbf{l}) \text{ is undecomposable}$$

(see [3, Proposition 3] and [21, Proposition 3.1]). The identification of these three conditions is not necessarily true for any curves. So for reduced divisor cases, the target of the forgetting map from $\mathbf{Con}_\Lambda(D)$ is the moduli space of undecomposable parabolic bundles. For $n = 4$ (that is, $D = t_1 + \dots + t_4$), the moduli space of rank 2 undecomposable parabolic bundles with degree 0 (or 1) can be obtained by glueing two copies of \mathbb{P}^1 outside t_1, \dots, t_4 (see [4, Section 2.3]). For $n = 5$ (that is, $D = t_1 + \dots + t_5$), Loray–Saito have provided a precise description of the moduli space of rank 2 undecomposable parabolic bundles with degree -1 . This moduli space is closely related to the geometry of degree 4 del Pezzo surfaces.

Now we consider cases where D is not necessary reduced. It is proved in [4, Proposition 4.8] that, if $n = 4$ (Painlevé case) and $\deg(E)$ is odd, then

$$\text{End}(E, \mathbf{l}) = \mathbb{C} \iff (E, \mathbf{l}) \text{ is } \Lambda\text{-flat}.$$

We will check the following claim: If (E, \mathbf{l}) is Λ -flat, then (E, \mathbf{l}) is undecomposable. But the converse proposition is not true even when $\deg(D) = 4$ and $\deg(E)$ is odd. So in addition to undecomposability, we assume the *admissibility* defined below.

Remind that, on \mathbb{P}^1 , any vector bundle is decomposable by Birkhoff:

$$E \simeq \mathcal{O}(d_1) \oplus \mathcal{O}(d_2) \quad \text{with} \quad d_1 \leq d_2 \text{ unique.}$$

The subbundle $\mathcal{O}(d_1) \hookrightarrow E$ is not unique, but if $d_1 < d_2$, then $\mathcal{O}(d_2) \hookrightarrow E$ is unique. When $d_1 = d_2$, the two subbundles of the decomposition can be arbitrarily chosen in a 1-parameter family. In any case, the (possible) factor $L = \mathcal{O}(d_2)$ of the decomposition is characterized by $\deg(E) \leq 2 \deg(L)$.

Definition 7. We will say that (E, \mathbf{l}) is *admissible* if the following condition is satisfied:

$$(2.1) \quad \forall L \subset E, \quad \deg(E) \leq 2 \deg(L) \implies \sum_{i \in I} \text{length}(l_i \cap L|_{n_i[t_i]}) \leq n + \deg(E) - 2 \deg(L) - 2.$$

Using some material of [4, Section 4] and ideas of [3, Proposition 3], we can prove:

Proposition 8. Let Λ satisfying (a),(b),(c),(d) as above, and (E, \mathbf{l}) a parabolic bundle of degree d . Then, we have implications (i) \Rightarrow (ii) \Rightarrow (iii) where:

- (i) $\text{End}(E, \mathbf{l}) = \mathbb{C}$,
- (ii) (E, \mathbf{l}) is Λ -flat,
- (iii) (E, \mathbf{l}) is undecomposable and admissible.

Moreover, for $n = 4$ and $\deg(E)$ odd, these are equivalent.

Proof. (ii) \Rightarrow (iii). Let ∇ be a connection on E compatible with the parabolic structure \mathbf{l} and formal data Λ . Assume, by contradiction, that $(E, \mathbf{l}) = (L_1, D_1) \oplus (L_2, D_2)$ where $D = D_1 + D_2$ with D_1, D_2 having disjoint support. Then, with respect to that decomposition, the connection ∇ writes

$$\begin{pmatrix} \nabla_1 & \theta_{1,2} \\ \theta_{2,1} & \nabla_2 \end{pmatrix}$$

where $\theta_{i,j} \in \text{Hom}(L_j, L_i \otimes \Omega^1(D))$ and $\nabla_i : L_i \rightarrow L_i \otimes \Omega^1(D)$ a connection. Then, the principal part at t_i writes:

$$\text{either } \begin{pmatrix} \lambda_i^+ & * \\ 0 & \lambda_i^- \end{pmatrix}, \quad \text{or } \begin{pmatrix} \lambda_i^- & 0 \\ * & \lambda_i^+ \end{pmatrix}.$$

Therefore, Fuchs' relation for L_1 (or L_2) yields a contradiction with generic assumption (c) on residual eigenvalues.

On the other hand, let $L \subset E$ and consider the composition map

$$L \hookrightarrow E \xrightarrow{\nabla} E \otimes \Omega^1(D) \rightarrow (E/L) \otimes \Omega^1(D).$$

This map φ_L is \mathcal{O} -linear and it is not the zero map, otherwise L would be ∇ -invariant, therefore contradicting Proposition 5. We deduce that

$$m := \deg(E) - 2 \deg(L) + n - 2 \geq 0.$$

Moreover, m is the number of zeroes of φ_L (counted with multiplicities). If $l_i \cap L|_{n_i[t_i]} \neq 0$, then $l_i \cap L|_{n_i[t_i]}$ contributes for zeroes of φ_L , and we therefore deduce

$$\sum_{i \in I} \text{length}(l_i \cap L|_{n_i[t_i]}) \leq \deg(E) - 2 \deg(L) + n - 2$$

which implies, when applied to the case $\deg(E) \leq 2 \deg(L)$, that $(E, \mathbf{1})$ is admissible.

(i) \Rightarrow (ii). Let $(E, \mathbf{1})$ be as in (i). We can locally trivialize $E|_{U_\alpha} \cong \mathbb{C}^2$ with parabolic structure normalized as $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, so that we can locally define a Λ -connection compatible with the parabolic structure on U_α by

$$\nabla_\alpha = d + \begin{pmatrix} \lambda_\alpha^+ & 0 \\ 0 & \lambda_\alpha^- \end{pmatrix}$$

where λ_α^\pm is a 1-form on U_α with principal part $\lambda^\pm|_{U_\alpha}$. On intersections $U_\alpha \cap U_\beta$, the difference $\nabla_\alpha - \nabla_\beta = \theta_{\alpha\beta}$ is a section of the sheaf of parabolic Higgs fields $\mathcal{Higgs}(E, \mathbf{1})$ which is the subsheaf of $\mathcal{E}nd(E, \mathbf{1}) \otimes \Omega^1(D)$ whose principal parts are nilpotent. In above trivialization, we have

$$\theta_{\alpha\beta} = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix} + \text{holomorphic}.$$

This defines an element of $\{\theta_{\alpha\beta}\} \in H^1(\mathbb{P}^1, \mathcal{Higgs}(E, \mathbf{1}))$ and the existence of a global Λ -connection ∇ compatible with the parabolic structure is equivalent to the vanishing of this cocycle: $\nabla_{U_\alpha} = \nabla_\alpha - \theta_\alpha$ with $\theta_\alpha - \theta_\beta = \theta_{\alpha\beta}$.

In order to prove that $\{\theta_{\alpha\beta}\} \equiv 0$, we are going to use Serre duality, i.e. there is a perfect pairing

$$\begin{aligned} H^1(\mathbb{P}^1, \mathcal{Higgs}(E, \mathbf{1})) \times H^0(\mathbb{P}^1, \mathcal{E}nd(E, \mathbf{1})) &\rightarrow H^1(\mathbb{P}^1, \Omega^1) \xrightarrow{\sim} \mathbb{C} \\ (\{\theta_{\alpha\beta}\}, A) &\mapsto \{\omega_{\alpha\beta} := \text{tr}(\theta_{\alpha\beta} \cdot A)\} \end{aligned}$$

where $\mathcal{E}nd(E, \mathbf{1})$ is the sheaf of parabolic endomorphisms. If $\{\omega_{\alpha\beta}\} \equiv 0$ for all A , then we can conclude that $\{\theta_{\alpha\beta}\} \equiv 0$ and therefore that there exists a global unramified irregular singular Λ -parabolic connection on $(E, \mathbf{1})$. By assumption (i), each A takes the form $c \cdot \text{Id}_E$, and by linearity, it is therefore enough to test the vanishing of $\{\omega_{\alpha\beta}\}$ for $A = \text{Id}_E$.

Fix any global meromorphic connection ∇^0 on E ; for instance, one can pull-back the trivial connection d from any birational trivialization $E \dashrightarrow \mathbb{C}^2$. It obviously satisfies Fuchs' relation $\sum \text{Res } \nabla^0 = -\deg(E)$. Now consider the following meromorphic resolution of the cocycle

$$\theta_{\alpha\beta} = (\nabla_\alpha - \nabla^0) - (\nabla_\beta - \nabla^0);$$

therefore, $\{\omega_{\alpha\beta}\}$ can also be viewed as the meromorphic Čech coboundary

$$\{\omega_{\alpha\beta}\} = \partial \{\omega_\alpha := \text{tr}((\nabla_\alpha - \nabla^0) \cdot A)\}.$$

Then we have:

$$\{\omega_{\alpha\beta}\} \sim 0 \quad \iff \quad \sum \text{Res}(\omega_\alpha) = 0$$

(note that residue does not depend on the choice of α as residue of $\nabla_\alpha - \nabla_\beta$ is nilpotent). Indeed, the vanishing of residue is equivalent to the existence of a global meromorphic 1-form ω such that $\omega - \omega_\alpha$ is holomorphic, i.e. such that $\{\omega_{\alpha\beta}\}$ is a holomorphic Čech coboundary:

$$\{\omega_{\alpha\beta}\} = \partial\{\omega_\alpha - \omega\}$$

meaning that $\{\omega_{\alpha\beta}\} \sim 0$ in $H^1(\mathbb{P}^1, \Omega^1)$. If $A = \text{Id}_E$, then we have by Fuchs' relation for ∇^0 and assumption (b) for Λ , we have

$$\sum \text{Res}(\omega_\alpha) = \sum \text{Res}(\text{tr}(\nabla_\alpha)) - \sum \text{Res}(\text{tr}(\nabla^0)) = 0.$$

(iii) \Rightarrow (i) in the case $n = 4$ and $\deg(E)$ odd. Let A be an endomorphism of the parabolic bundle $(E, \mathbf{1})$. Since eigenvalues of $A|_{E_x}$ are holomorphic functions of $x \in X$, they are constant $c_1, c_2 \in \mathbb{C}$. If $c_1 \neq c_2$, then the corresponding eigenspaces define a decomposition $E = L_1 \oplus L_2$ and parabolic structure, since it is A -invariant, must be contained in $L_1 \cup L_2$; this implies that $(E, \mathbf{1})$ is decomposable, contradiction. We therefore have $c_1 = c_2 =: c$, and any endomorphism of the parabolic bundle takes the form $A = c \cdot \text{Id}_E + N$ with N nilpotent. Clearly, $c \cdot \text{Id}_E \in \text{End}(E, \mathbf{1})$, and (i) is equivalent to say that there is no non-zero nilpotent endomorphism in $\text{End}(E, \mathbf{1})$.

Assume, by contradiction, that N is a non-zero nilpotent element of $\text{End}(E, \mathbf{1})$. Let $L_0 = \ker(N) = \text{image}(N)$. Since N is not trivial, it induces a non trivial morphism $E/L_0 \rightarrow L_0$. This implies that $\deg(E) - 2 \deg(L_0) \leq 0$. In particular, we have a decomposition $E = L \oplus L_0$ with $\deg(L) \leq \deg(L_0)$ and, through that decomposition, the nilpotent endomorphism N writes:

$$N = \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix}$$

where $f = f(x)$ is a polynomial of degree $2 \deg(L_0) - \deg(E)$. Locally at $n_i t_i$, the parabolic structure is given, through the splitting and local coordinate $z = x - t_i$, by

$$\begin{pmatrix} z^{m_i} g(z) \\ 1 \end{pmatrix}$$

where $m_i = \sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i t_i})$, and $g(0) \neq 0$. The action of N on the parabolic writes

$$\begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \begin{pmatrix} z^{m_i} g \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ z^{m_i} f g \end{pmatrix}$$

so that N preserves the parabolic structure if, and only if, f (and therefore N) vanish at t_i at order $n_i - 2m_i$. Indeed, we have

$$(2.2) \quad \det \begin{pmatrix} z^{m_i} g & 0 \\ 1 & z^{m_i} f g \end{pmatrix} = z^{2m_i} f g^2 = 0 \quad \text{mod } z^{n_i} \quad \text{iff } f = z^{n_i - 2m_i} \tilde{f}$$

and the induced action of N is by multiplication

$$\begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \begin{pmatrix} z^{m_i} g \\ 1 \end{pmatrix} = z^{m_i} f g \cdot \begin{pmatrix} z^{m_i} g \\ 1 \end{pmatrix} \quad \text{mod } z^{n_i}.$$

We promptly deduce that

$$(2.3) \quad n - 2 \sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i[t_i]}) \leq \deg(f) \leq 2 \deg(L_0) - \deg(E).$$

On the other hand, from (iii), $(E, \mathbf{1})$ is admissible and we have that

$$\sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i[t_i]}) \leq n + \deg(E) - 2 \deg(L_0) - 2.$$

We deduce

$$\frac{n + \deg(E) - 2 \deg(L_0)}{2} \leq \sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i[t_i]}) \leq n + \deg(E) - 2 \deg(L_0) - 2.$$

This implies, by comparing left and right hand-sides, that

$$n \geq \underbrace{2 \deg(L_0) - \deg(E)}_{\geq 0} + 4$$

which is impossible when $n = 4$ and $\deg(E)$ odd. Therefore, $N = 0$ and $\text{End}(E, \mathbf{1}) = \mathbb{C}$. \square

The undecomposability and admissibility are independent of Λ . So, when D is reduced or when $n = 4$ and $\deg(E)$ is odd, the Λ -flatness is independent of Λ . But, in general, the Λ -flatness depends on Λ . Indeed, we have the following proposition.

Proposition 9 ([4, Corollary 4.4]). *Let Λ satisfying (a),(b),(c),(d) as above, and $(E, \mathbf{1})$ a parabolic bundle of degree d . Assume $(E, \mathbf{1})$ is undecomposable. Then $(E, \mathbf{1})$ is Λ -flat if, and only if, for any nilpotent endomorphism N of $(E, \mathbf{1})$, we have*

$$(2.4) \quad \text{Res}(N|\mathbf{1} \cdot (\lambda^+ - \lambda^-)) = 0$$

where $N|\mathbf{1}$ denotes the action induced by N on the parabolic structure, and Res is the sum of residues.

Proof. We rephrase and continue the part [(iii) \Rightarrow (i)] of the proof of Proposition 8. The obstruction to the existence of a connection is defined by an element $\{\nabla_\alpha - \nabla_\beta\} \in H^1(\mathbb{P}^1, \mathcal{Higgs}(E, \mathbf{1}))$. By Serre duality, it is zero if, and only if $\text{Res}(\text{tr}(\nabla_\alpha - \nabla^0) \cdot A) = 0$ for all endomorphism A of $(E, \mathbf{1})$; here ∇^0 is a global meromorphic connection that we have fixed *a priori*. Since $(E, \mathbf{1})$ is undecomposable, any endomorphism takes the form $A = c \cdot \text{Id}_E + N$ with N nilpotent. We already know that the condition is satisfied for $A = \text{Id}_E$ because of Fuchs' relation and assumption (b) for Λ ; by linearity, it is enough to consider $A = N$ nilpotent. A non zero nilpotent endomorphism takes the form

$$N = \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix}$$

in a decomposition $E = L \oplus L_0$ with $\deg(L) \leq \deg(L_0)$. In fact, we can choose ∇^0 such that it has one single pole with residue

$$\begin{pmatrix} k_1 & 0 \\ 0 & k_2 \end{pmatrix} \frac{dz}{z}$$

so that the contribution of ∇^0 in $\text{Res}(\text{tr}(\nabla_\alpha - \nabla^0) \cdot A)$ is zero. Let us compute the contribution of ∇_α . Let the parabolic structure be defined by

$$\begin{pmatrix} g \\ 1 \end{pmatrix}$$

(e.g. in local coordinate near a pole). Since it is N -invariant, we have $fg^2 \sim 0$ on D (or $n_i t_i$) and

$$\begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix} \cdot \begin{pmatrix} g \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ fg \end{pmatrix} \sim fg \begin{pmatrix} g \\ 1 \end{pmatrix}$$

so that the induced action by (nilpotent) multiplication $N|_D$ is the multiplication by fg . For instance, if g vanishes at t_i with multiplicity m_i , then f vanishes with multiplicity $n_i - 2m_i$, and $N|_D = fg$ vanishes with multiplicity $n_i - m_i$. By gauge transformation, we can normalize

$$\begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} \rightsquigarrow \mathbf{1} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad N = \begin{pmatrix} -fg & 0 \\ f & fg \end{pmatrix}, \quad \nabla_\alpha = \begin{pmatrix} \lambda^- & 0 \\ \star & \lambda^+ \end{pmatrix}$$

so that we finally get

$$(\nabla_\alpha - \nabla^0) \cdot N = \begin{pmatrix} -fg\lambda^- & 0 \\ \star & fg\lambda^+ \end{pmatrix}$$

so that

$$\text{Res}(\text{tr}(\nabla_\alpha - \nabla^0) \cdot N) = \text{Res}(fg(\lambda^+ - \lambda^-)) = \text{Res}(N|\mathbf{1} \cdot (\lambda^+ - \lambda^-))$$

must be zero for any nilpotent endomorphism N of $(E, \mathbf{1})$. \square

2.2. Examples. We will give counterexamples of the implication [(ii) \Rightarrow (i)] and the implication [(iii) \Rightarrow (ii)], which are converses of the implications in Proposition 8. To give counterexamples, we will use the criterium of Proposition 9. First, we consider an example of this criterium.

Example 10. Assume $n = 5$ with divisor $2[0] + 2[1] + [\infty]$, $E = \mathcal{O} \oplus \mathcal{O}(1)$ and the parabolic structure $\mathbf{1}$ intersect the destabilizing line bundle $\mathcal{O}(1)$ only at $[0] + [1]$ (without multiplicity). In other words, the parabolic structure is given by

$$\begin{pmatrix} c_0 x \\ 1 \end{pmatrix} \pmod{x^2}, \quad \begin{pmatrix} c_1(x-1) \\ 1 \end{pmatrix} \pmod{(x-1)^2}, \quad \text{and} \quad \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ at } \infty.$$

Then $(E, \mathbf{1})$ is indecomposable provided that $(c_0, c_1) \neq (0, 0)$. On the other hand, there is (up to homothecy) one non trivial nilpotent endomorphism

$$N = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

that preserves the parabolic structure. It induces the multiplication by $c_0 x$ and $c_1(x-1)$ on parabolic directions over $x = 0$ and $x = 1$ respectively. Up to bundle isomorphism, we see that only the projective variable $[c_0 : c_1] \in \mathbb{P}^1$ makes sense. Criterium of Proposition 9 writes

$$\text{Res}(c_0 x(\lambda_0^+ - \lambda_0^-) + c_1(x-1)(\lambda_1^+ - \lambda_1^-)) = 0.$$

Recall that dominant terms of $\lambda^+ - \lambda^-$ are non zero at $x = 0$ and $x = 1$, given say by $a_0 \frac{dx}{x^2}$ and $a_1 \frac{dx}{(x-1)^2}$ respectively. Flatness condition writes

$$c_0 a_0 + c_1 a_1 = 0$$

which determines the parabolic bundle uniquely.

Now we consider the case where $n = \deg(D) = 4$ and $\deg(E)$ is even. We give counterexamples of the implication [(ii) \Rightarrow (i)] and the implication [(iii) \Rightarrow (ii)].

Proposition 11. Let $n = \deg(D) = 4$, Λ as above, $(E, \mathbf{1})$ a parabolic bundle, $\deg(E)$ any even number.

- Assume that $(E, \mathbf{1})$ is indecomposable and admissible. If $(E, \mathbf{1})$ is not simple, then $E = L \otimes (\mathcal{O} \oplus \mathcal{O})$ and

(i) $D = 2[t_1] + 2[t_2]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x - t_1) \\ 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} c_2(x - t_2) \\ 1 \end{pmatrix}$$

where $[c_1 : c_2] \in \mathbb{P}^1$;

(ii) $D = 4[t_1]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x - t)^2 + c_2(x - t)^3 \\ 1 \end{pmatrix}$$

where $[c_1 : c_2] \in \mathbb{P}^1$.

- The only Λ -flat parabolic bundles that are not simple are the following ones: $E = L \otimes (\mathcal{O} \oplus \mathcal{O})$ and
 - $D = 2[t_1] + 2[t_2]$ and parabolic structure writes as in (i) where $[c_1 : c_2] \in \mathbb{P}^1$ is a point determined by Λ ;
 - $D = 4[t_1]$ and parabolic structure writes as in (ii) where $[c_1 : c_2] \in \mathbb{P}^1$ is a point determined by Λ .

Proof. Setting $k = \max_L(2 \deg(L) - \deg(E))$, L_0 be maximizing and $m = \sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i[t_i]})$. Then, end of proof of Proposition 8 yields

$$\frac{n - k}{2} \leq m \leq n - k - 2 \quad \text{so that in particular} \quad k + 4 \leq n.$$

It follows, when $n = 4$ that $k = 0$ and $m = 2$. Since $k = 0$, then $E = L \otimes (\mathcal{O} \oplus \mathcal{O})$. On the other hand, assuming $(E, \mathbf{1})$ is not simple, we know that there is a non trivial nilpotent endomorphism which writes

$$N = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

and L_0 is the invariant subspace. We deduce that all parabolics intersect L_0 , and at each pole t_i , we must have $n_i - 2m_i \leq 0$; otherwise, parabolic structure would not be invariant. Together with above inequalities, we get $n_i = 2m_i$ at each t_i , whence the only two possibilities of the statement. In each case, writing down the criterium of Proposition 9, as in Example 10, we get the restriction on $(c_1 : c_2)$. \square

Now we consider the case where $n = \deg(D) = 5$ and $\deg(E)$ is odd.

Lemma 12. *Let $n = \deg(D) = 5$ and $(E, \mathbf{1})$ a parabolic bundle with odd degree. Assume that $(E, \mathbf{1})$ is indecomposable and admissible. If $(E, \mathbf{1})$ is not simple, then $E = L \otimes (\mathcal{O} \oplus \mathcal{O}(1))$ and*

(i) $D = 2[t_1] + 2[t_2] + [t_3]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x - t_1) \\ 1 \end{pmatrix}, \quad \begin{pmatrix} c_2(x - t_2) \\ 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

where $[c_1 : c_2] \in \mathbb{P}^1$ satisfies $c_1 c_2 \neq 0$;

(ii) $D = 3[t_1] + 2[t_2]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x - t_1) + c_2(x - t_1)^2 \\ 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} c_3(x - t_2) \\ 1 \end{pmatrix}$$

where $[c_1 : c_2 : c_3] \in \mathbb{P}^2$ satisfies $c_1 c_3 \neq 0$;

(iii) $D = 4[t_1] + [t_2]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x-t_1)^2 + c_2(x-t_1)^3 \\ 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

where $[c_1 : c_2] \in \mathbb{P}^1$ satisfies $c_1 \neq 0$;

(iv) $D = 5[t_1]$ and parabolic structure writes

$$\begin{pmatrix} c_1(x-t_1)^2 + c_2(x-t_1)^3 + c_3(x-t_1)^4 \\ 1 \end{pmatrix}$$

where $[c_1 : c_2 : c_3] \in \mathbb{P}^2$ satisfies $c_1 \neq 0$.

In particular, when $D = 2t_1 + t_2 + t_3 + t_4$ or $D = 3t_1 + t_2 + t_3$, we have that

$$\text{End}(E, \mathbf{1}) = \mathbb{C} \iff (E, \mathbf{1}) \text{ is } \Lambda\text{-flat} \iff (E, \mathbf{1}) \text{ is undecomposable and admissible.}$$

Proof. Setting $k = \max_L(2 \deg(L) - \deg(E))$, L_0 be maximizing and $m = \sum_{i \in I} \text{length}(l_i \cap L_0|_{n_i[t_i]})$. Since $(E, \mathbf{1})$ is not simple, we have $k + 4 \leq n$ as in the proof of the previous proposition. By this inequality, we have $k = 0$ or $k = 1$. Since E has odd degree, $k \neq 0$. So we have $k = 1$. This means that $E = L \otimes (\mathcal{O} \oplus \mathcal{O}(1))$. Since $(E, \mathbf{1})$ is not simple, we know that there is a non trivial nilpotent endomorphism which writes

$$N = \begin{pmatrix} 0 & 0 \\ f(x) & 0 \end{pmatrix}$$

and L_0 is the invariant subspace. Here $f(x)$ is a polynomial in x with $\deg(f) \leq 1$. For one of indexes in I , m_i satisfies a condition $n_i - 2m_i \leq 1$, and, for other indexes, m_i satisfies a condition $n_i - 2m_i \leq 0$. In particular $\sum m_i \geq 2$. Since $(E, \mathbf{1})$ is admissible, we have $\sum m_i \leq 2$. Then for one of indexes in I , $n_i - 2m_i = 1$, and, for other indexes, $n_i - 2m_i = 0$. There exist the only four possibilities of the statement. \square

Now we give counterexamples when $n = \deg(D) = 5$ and $\deg(E)$ is odd.

Proposition 13. *Let $n = \deg(D) = 5$, Λ as above, and $(E, \mathbf{1})$ a parabolic bundle with odd degree. Then, the only Λ -flat parabolic bundles that are not simple are the following ones: $E = L \otimes (\mathcal{O} \oplus \mathcal{O}(1))$ and*

- $D = 2[t_1] + 2[t_2] + [t_3]$ and parabolic structure writes as in (i) of Lemma 12 where $[c_1 : c_2]$ is a point determined by Λ ;
- $D = 3[t_1] + 2[t_2]$ and parabolic structure writes as in (ii) of Lemma 12 where $[c_1 : c_2 : c_3]$ lies along some line determined by Λ ;
- $D = 4[t_1] + [t_2]$ and parabolic structure writes as in (iii) of Lemma 12 where $[c_1 : c_2]$ is a point determined by Λ ;
- $D = 5[t_1]$ and parabolic structure writes as in (iv) of Lemma 12 where $[c_1 : c_2 : c_3]$ lies along some line determined by Λ .

Proof. By Proposition 8, we have that $(E, \mathbf{1})$ is undecomposable and admissible. Then, the parabolic structure write as in Lemma 12. In each case, writing down the criterium of Proposition 9, as in Example 10. \square

3. REFINED PARABOLIC BUNDLES

In the previous section, we have defined parabolic bundles of rank 2 and of degree d over (\mathbb{P}^1, D) . In this section, we will consider a generalization of parabolic bundles, which will be called *refined parabolic bundles*.

3.1. Cases where D is a reduced effective divisor. Here we assume that D is a reduced effective divisor. In [21], the moduli space of indecomposable parabolic bundles of rank 2 and of degree d was studied. Since now we assume that D is reduced, this moduli space coincides with the moduli space of Λ -flat parabolic bundles of rank 2 and of degree d . We denote by $\mathfrak{Bun}_\Lambda(D, d)$ the moduli space of Λ -flat parabolic bundles with degree d . This moduli space is the image of the forgetting map:

$$\begin{aligned} \mathfrak{Con}_\Lambda(D) &\longrightarrow \mathfrak{Bun}_\Lambda(D, d) \\ (E, \nabla) &\longmapsto (E, \mathbf{1}). \end{aligned}$$

Here $\mathfrak{Con}_\Lambda(D)$ is the moduli space of Λ -connections. This forgetting map is important for understanding the moduli space $\mathfrak{Con}_\Lambda(D)$. But the moduli space $\mathfrak{Bun}_\Lambda(D, d)$ is a non-separated scheme. To get a good moduli space, we have to impose a stability condition with respect to $\mathbf{w} = (w_i) \in [0, 1]^n$. We denote by $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ the moduli space of \mathbf{w} -semistable parabolic bundles of rank 2 and of degree d . The moduli space $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is a normal irreducible projective variety. The open subset of \mathbf{w} -stable parabolic bundles is smooth. For generic weight, we have that \mathbf{w} -semistable = \mathbf{w} -stable. So $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is a smooth projective variety. An important fact is that $(E, \{l_i\}_{i \in I})$ is stable for a convenient choice of weights \mathbf{w} if, and only if, it is indecomposable (see [21, Proposition 3.4]). So the moduli space $\mathfrak{Bun}_\Lambda(D, d)$ is covered by $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ for some \mathbf{w} .

3.2. Definition of refined parabolic bundles. We try to extend the story in Section 3.1 for cases where D are not necessary reduced. When D is not a reduced divisor, there are some problems: (1) indecomposable parabolic bundles are not necessary Λ -flat and are not necessary \mathbf{w} -stable for some weight \mathbf{w} ; (2) $\mathfrak{Bun}_{\mathbf{w}}(D, d)$ is not necessary projective for generic weight \mathbf{w} . (Remark that we impose that l_i is free). Here the stability condition for parabolic bundles when D is not necessary reduced is defined in Section 4 below.

For the problem (1), in Section 4.1 and Section 4.2 below, we will discuss on a necessary and sufficient condition for the condition that there exist weights \mathbf{w} such that a parabolic bundle is \mathbf{w} -stable. The present section is concerned with the problem (2), that is, constructing a smooth compactification of $\mathfrak{Bun}_{\mathbf{w}}(D, d)$. To take a good compactification, we consider a generalization of parabolic bundles as follows.

Definition 14. We say $(E, \{l_{i, \bullet}\}_{i \in I})$ is a refined parabolic bundle of rank 2 and of degree d if E is a rank 2 vector bundle on \mathbb{P}^1 with $\deg(E) = d$ and $l_{i, \bullet} = \{l_{i, k}\}_{1 \leq k \leq n_i}$ is a filtration

$$E|_{n_i[t_i]} \supset l_{i, n_i} \supset l_{i, n_i-1} \supset \cdots \supset l_{i, 1} \supset 0$$

of $\mathcal{O}_{n_i[t_i]}$ -modules where the length of $l_{i, k}$ is k . We call this filtration a refined parabolic structure.

Let $(E, \{l_i\}_{i \in I})$ be a parabolic bundle of rank 2 and of degree d . For $(E, \{l_i\}_{i \in I})$, we can define a refined parabolic bundle of rank 2 and of degree d as follows: We define a filtration $l_{i, \bullet}$ as

$$l_{i, \bullet}: E|_{n_i[t_i]} \supset l_i \supset f_i \cdot l_i \supset f_i^2 \cdot l_i \supset \cdots \supset f_i^{n_i-1} \cdot l_i \supset 0$$

for each i . Here f_i is a generator of the maximal ideal of $\mathcal{O}_{\mathbb{P}^1, t_i}$. Then we have a refined parabolic bundle $(E, \{l_{i, \bullet}\}_{i \in I})$. Conversely if a refined parabolic structure $E|_{n_i[t_i]} \supset l_{i, n_i} \supset l_{i, n_i-1} \supset \cdots \supset l_{i, 1} \supset 0$ where l_{i, n_i} is free, then the refined parabolic structure coincides with the refined parabolic structure induced by the parabolic structure l_{i, n_i} . So refined parabolic bundles are generalization of parabolic bundles. In Section 4 below, we will define a stability condition for refined parabolic bundles. First of all, in present section, we discuss refined parabolic structures at a multiple point. We will describe a combinatorial structure on a refined parabolic structure at a multiple point.

3.3. Refined parabolic structure at a multiple point. Let E be a rank 2 vector bundle on \mathbb{P}^1 . We take a point t on \mathbb{P}^1 and a positive integer n such that $n > 1$. First, we will define the *type* of an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$. We consider a filtration of sheaves

$$E \supset E(-[t]) \supset E(-2[t]) \supset \cdots \supset E(-(n-1)[t]) \supset E(-n[t]).$$

Note that $E|_{n[t]} = E/E(-n[t])$. Set $V_j := E(-(n-j+1)[t])/E(-n[t])$ for $j = 1, 2, \dots, n+1$. Then we have a filtration

$$E|_{n[t]} = V_{n+1} \supset V_n \supset V_{n-1} \supset \cdots \supset V_2 \supset V_1 = 0$$

of $\mathcal{O}_{n[t]}$ -modules.

Definition 15. We fix a positive integer k where $1 \leq k \leq n$. Let l be an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ with $\text{length}(l) = k$. We define a tuple of integers $\lambda := (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$ by

$$\lambda_j := \text{length}((V_{j+1} \cap l)/(V_j \cap l))$$

for $j = 1, 2, \dots, n$. We call the tuple λ the type of the $\mathcal{O}_{n[t]}$ -submodule l with $\text{length}(l) = k$.

We have the equality $\sum_{j=1}^n \lambda_j = k$. The inequalities $\lambda_j \geq 0$ and

$$\lambda_j \leq \text{length}(V_{j+1}/V_j) = 2$$

follow for $j = 1, 2, \dots, n$. Let f be a generator of the maximal ideal of $\mathcal{O}_{\mathbb{P}^1, t}$. The map from V_{j+1} to V_j defined by multiply by f induces an injective map

$$(V_{j+1} \cap l)/(V_j \cap l) \xrightarrow{\text{multiply by } f} (V_j \cap l)/(V_{j-1} \cap l)$$

for $j = 2, 3, \dots, n$. Then we have the following inequality

$$\lambda_j = \text{length}((V_{j+1} \cap l)/(V_j \cap l)) \leq \text{length}((V_j \cap l)/(V_{j-1} \cap l)) = \lambda_{j-1}$$

for $j = 2, 3, \dots, n$. For a tuple of integers $\lambda = (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$, we set

$$(3.1) \quad \begin{aligned} a_1(\lambda) &:= \#\{j \mid j \in \{1, 2, \dots, n\}, \lambda_j = 1\} \quad \text{and} \\ a_2(\lambda) &:= \#\{j \mid j \in \{1, 2, \dots, n\}, \lambda_j = 2\}. \end{aligned}$$

Proposition 16. If a tuple of integers $\lambda = (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$ satisfies the condition

$$(3.2) \quad 0 \leq \lambda_n \leq \lambda_{n-1} \leq \cdots \leq \lambda_1 \leq 2 \quad \text{and} \quad \sum_{j=1}^n \lambda_j = k,$$

then there exists an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ with $\text{length}(l) = k$ whose type is λ .

Proof. We define \mathbf{v} and \mathbf{v}' as

$$\mathbf{v} = f^{n-a_1(\lambda)-a_2(\lambda)} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \in \mathcal{O}_{n[t]} \oplus \mathcal{O}_{n[t]} \quad \mathbf{v}' = f^{n-a_2(\lambda)} \begin{pmatrix} v'_1 \\ v'_2 \end{pmatrix} \in \mathcal{O}_{n[t]} \oplus \mathcal{O}_{n[t]}$$

so that $\mathbf{v} \in ((f^{n-a_1(\lambda)-a_2(\lambda)})/((f^{n-a_1(\lambda)-a_2(\lambda)+1}))^{\oplus 2}) \cong \mathbb{C}^2$ and $\mathbf{v}' \in ((f^{n-a_2(\lambda)})/((f^{n-a_2(\lambda)+1}))^{\oplus 2}) \cong \mathbb{C}^2$ are linearly independent. Let l be the $\mathcal{O}_{n[t]}$ -submodule of $\mathcal{O}_{n[t]} \oplus \mathcal{O}_{n[t]}$ generated by \mathbf{v} and \mathbf{v}' . The elements $f^{a_1(\lambda)}\mathbf{v}$ and \mathbf{v}' are linearly independent in $((f^{n-a_2(\lambda)})^{\oplus 2} \cap l)/((f^{n-a_2(\lambda)+1})^{\oplus 2} \cap l)$. In other words,

$$\text{length}(((f^{n-a_2(\lambda)})^{\oplus 2} \cap l)/((f^{n-a_2(\lambda)+1})^{\oplus 2} \cap l)) = 2.$$

If we take an isomorphism $E|_{n[t]} \cong \mathcal{O}_{n[t]} \oplus \mathcal{O}_{n[t]}$, then l induces a refined parabolic structure on $E|_{n[t]}$ via this isomorphism. The type of this refined parabolic structure is λ . \square

Let $\lambda = (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$ be the type of an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$. The type λ is a partition of the positive integer k :

$$k = \overbrace{0 + \dots + 0}^{a_1(\lambda)} + \overbrace{1 + \dots + 1}^{a_2(\lambda)} + \overbrace{2 + \dots + 2}^{a_2(\lambda)}.$$

So there exists the corresponding Young diagram to the type λ . The number of columns of the Young diagram is 1 or 2.

Lemma 17. *Let l be an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ with $\text{length}(l) = k$. Let $\lambda = (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$ be the type of l . If the number of columns of the corresponding Young diagram is 1, then the $\mathcal{O}_{n[t]}$ -submodule l is generated by one element of $E|_{n[t]}$. If the number of columns of the corresponding Young diagram is 2, then the $\mathcal{O}_{n[t]}$ -submodule l is generated by two elements of $E|_{n[t]}$.*

Proof. Let $a_1(\lambda)$ and $a_2(\lambda)$ be the integers defined in (3.1). We take \mathbf{v} such that the image of \mathbf{v} under the map $V_{a_1(\lambda)+a_2(\lambda)+1} \rightarrow V_{a_1(\lambda)+a_2(\lambda)+1}/V_{a_1(\lambda)+a_2(\lambda)}$ is nonzero.

Now we assume that the number of columns of the corresponding Young diagram is 1. That is, $a_2(\lambda) = 0$. We consider an element $\mathbf{v}' \in l$. There exists an integer $n_{\mathbf{v}'}$ such that $\mathbf{v}' \in V_{n_{\mathbf{v}'}+1}$ and the image of \mathbf{v}' under the map $V_{n_{\mathbf{v}'}+1} \rightarrow V_{n_{\mathbf{v}'}+1}/V_{n_{\mathbf{v}'}}$ is nonzero. We have $a_1(\lambda) \geq n_{\mathbf{v}'}$. The element $f^{a_1(\lambda)-n_{\mathbf{v}'}}\mathbf{v}$ is in $V_{n_{\mathbf{v}'}+1}$. Since $\lambda_{n_{\mathbf{v}'}} = 1$, the images of $f^{a_1(\lambda)-n_{\mathbf{v}'}}\mathbf{v}$ and \mathbf{v}' under the map $V_{n_{\mathbf{v}'}+1} \rightarrow V_{n_{\mathbf{v}'}+1}/V_{n_{\mathbf{v}'}}$ are linearly dependent. Then there exist complex numbers α_1 and α_2 such that $\alpha_1 f^{a_1(\lambda)-n_{\mathbf{v}'}}\mathbf{v} + \alpha_2 \mathbf{v}' \in V_{n_{\mathbf{v}'}}$. Moreover, the images of $f^{a_1(\lambda)-n_{\mathbf{v}'}+1}\mathbf{v}$ and $\alpha_1 f^{a_1(\lambda)-n_{\mathbf{v}'}}\mathbf{v} + \alpha_2 \mathbf{v}'$ under the map $V_{n_{\mathbf{v}'}} \rightarrow V_{n_{\mathbf{v}'}}/V_{n_{\mathbf{v}'}-1}$ are linearly dependent, since $\lambda_{n_{\mathbf{v}'}-1} = 1$. By repeating this argument, we may write the element $\mathbf{v}' \in l$ as a linear combination of $\mathbf{v}, f\mathbf{v}, f^2\mathbf{v}$, and so on. Then l is generated by \mathbf{v} .

Next we assume that the number of columns of the corresponding Young diagram is 2. That is, $a_2(\lambda) > 0$. There exists an element \mathbf{w} such that the image of \mathbf{w} under the map $V_{a_2(\lambda)+1} \rightarrow V_{a_2(\lambda)+1}/V_{a_2(\lambda)}$ is nonzero and the images of $f^{a_1(\lambda)}\mathbf{v}$ and \mathbf{w} under the map $V_{a_2(\lambda)+1} \rightarrow V_{a_2(\lambda)+1}/V_{a_2(\lambda)}$ are linearly independent. We may show that l is generated by \mathbf{v} and \mathbf{w} by using the argument as in the case $a_2(\lambda) = 0$. \square

Let \mathbf{v} be an element of $E|_{n[t]}$. We denote by $\langle \mathbf{v} \rangle_{\mathcal{O}_{n[t]}}$ the $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ generated by \mathbf{v} . If we take another generator \mathbf{v}' of $\langle \mathbf{v} \rangle_{\mathcal{O}_{n[t]}}$, then there is an element $F \in \mathcal{O}_{n[t]}$ such that $\mathbf{v} = F \cdot \mathbf{v}'$. We assume that l is generated by elements $\mathbf{v}_1, \mathbf{v}_2$ of $E|_{n[t]}$, where $\text{length}(\langle \mathbf{v}_1 \rangle_{\mathcal{O}_{n[t]}}) \geq \text{length}(\langle \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}})$. If we take another generator \mathbf{v}'_1 and \mathbf{v}'_2 of l , where $\text{length}(\langle \mathbf{v}'_1 \rangle_{\mathcal{O}_{n[t]}}) \geq \text{length}(\langle \mathbf{v}'_2 \rangle_{\mathcal{O}_{n[t]}})$, then there is an element $F_1, F_2, G_1, G_2 \in \mathcal{O}_{n[t]}$ such that $\mathbf{v}_1 = F_1 \cdot \mathbf{v}'_1 + F_2 \cdot \mathbf{v}'_2$ and $\mathbf{v}_2 = G_1 \cdot f^d \cdot \mathbf{v}'_1 + G_2 \cdot \mathbf{v}'_2$, where $d = \text{length}(\langle \mathbf{v}'_1 \rangle) - \text{length}(\langle \mathbf{v}'_2 \rangle)$.

Let $l_\bullet = \{l_k\}_{1 \leq k \leq n}$ be a filtration

$$E|_{n[t]} \supset l_n \supset l_{n-1} \supset \dots \supset l_1 \supset 0$$

of $\mathcal{O}_{n[t]}$ -submodules of $E|_{n[t]}$ where the length of l_k is k . Let $\lambda^{(k)} = (\lambda_n^{(k)}, \lambda_{n-1}^{(k)}, \dots, \lambda_1^{(k)})$ be the type of the $\mathcal{O}_{n[t]}$ -module l_k for $k = 1, 2, \dots, n$. Then we have a sequence of Young diagrams

$$\lambda^{(n)} \supset \lambda^{(n-1)} \supset \dots \supset \lambda^{(1)},$$

where the skew diagram $\lambda^{(k)}/\lambda^{(k-1)}$ consists of one box.

Definition 18. *Let $l_\bullet = \{l_k\}_{1 \leq k \leq n}$ be a filtration*

$$E|_{n[t]} \supset l_n \supset l_{n-1} \supset \dots \supset l_1 \supset 0 \quad (\text{length}(l_k) = k \text{ for } k = 1, 2, \dots, n)$$

of $\mathcal{O}_{n[t]}$ -modules. We denote by T_{l_\bullet} the corresponding standard Young tableau to the sequence of Young diagrams as above. We call T_{l_\bullet} the standard tableau of l_\bullet .

3.4. Parameter space of refined parabolic structures at a multiple point. Next, we will consider families of refined parabolic structures at a multiple point and their parameter spaces. Here, when we consider families of refined parabolic structures, we fix the first submodule l_n of refined parabolic structures. We will construct parameter spaces of families of refined parabolic structures. We will see that there exists a correspondence between their irreducible components and standard Young tableaux whose shapes are the type of l_n .

Let $\lambda = (\lambda_n, \lambda_{n-1}, \dots, \lambda_1)$ be a Young diagram which satisfies the condition (3.2). Let $a_1(\lambda)$ and $a_2(\lambda)$ be the integers defined in (3.1). We may describe the Young diagram λ as $(1^{a_1(\lambda)}, 2^{a_2(\lambda)})$. Let l be an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ with type λ . First we consider a parameter space of $\mathcal{O}_{n[t]}$ -submodules l' of l with $\text{length}(l/l') = 1$. We consider the case $a_2(\lambda) = 0$. In this case, l is generated by an element $\mathbf{v} \in E|_{n[t]}$ (see Lemma 17). An $\mathcal{O}_{n[t]}$ -submodule l' of l with $\text{length}(l/l') = 1$ is unique: $l' = f \cdot l$. Here f is a generator of the maximal ideal of $\mathcal{O}_{\mathbb{P}^1, t}$. For the case $a_2(\lambda) \neq 0$, we have the following lemma.

Lemma 19. *Assume that $a_2(\lambda) \neq 0$. By taking generators of l , we may construct a family of $\mathcal{O}_{n[t]}$ -submodules l' of l with $\text{length}(l/l') = 1$ parametrized by \mathbb{P}^1 such that,*

- when $a_1(\lambda) \neq 0$:
 - the type of the submodule l' corresponding to $[0 : 1]$ is $(1^{a_1(\lambda)-1}, 2^{a_2(\lambda)})$, and
 - the type of the submodule l' corresponding to $[\alpha_1 : \alpha_2]$, where $\alpha_1 \neq 0$, is $(1^{a_1(\lambda)+1}, 2^{a_2(\lambda)-1})$;
- when $a_1(\lambda) = 0$:
 - the type of the submodule l' corresponding to any $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$ is $(1^{a_1(\lambda)+1}, 2^{a_2(\lambda)-1})$.

This family gives a bijection

$$\mathbb{P}^1 \longrightarrow \{l' \mid l' \text{ is an } \mathcal{O}_{n[t]}\text{-submodule of } l \text{ such that } \text{length}(l/l') = 1\}.$$

Proof. We take generators \mathbf{v}_1 and \mathbf{v}_2 of l such that \mathbf{v}_1 is nonzero in $V_{a_1(\lambda)+a_2(\lambda)+1}/V_{a_1(\lambda)+a_2(\lambda)}$. Let $[\alpha_1 : \alpha_2]$ be a point of \mathbb{P}^1 . We set

$$\mathbf{v}'_0 := \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2, \quad \mathbf{v}'_1 = f \mathbf{v}_1, \quad \text{and} \quad \mathbf{v}'_2 := f \mathbf{v}_2.$$

Let l' be an $\mathcal{O}_{n[t]}$ -submodule of l generated by $\mathbf{v}'_0, \mathbf{v}'_1$, and \mathbf{v}'_2 . We may check that $\text{length}(l/l') = 1$. Then we have an $\mathcal{O}_{n[t]}$ -submodule of l corresponding to the point $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$. When $a_1(\lambda) \neq 0$, the type of the submodule l' corresponding to $[0 : 1]$ is $(1^{a_1(\lambda)-1}, 2^{a_2(\lambda)})$, and the type of the submodule l' corresponding to $[1 : \alpha_2/\alpha_1]$ is $(1^{a_1(\lambda)+1}, 2^{a_2(\lambda)-1})$. When $a_1(\lambda) = 0$, the type of the submodule l' corresponding to $[\alpha_1 : \alpha_2]$ is $(1^{a_1(\lambda)+1}, 2^{a_2(\lambda)-1})$. So we have a map from \mathbb{P}^1 to the set of $\mathcal{O}_{n[t]}$ -submodules l' of l with $\text{length}(l/l') = 1$.

We will construct the inverse map as follows. Let l' be an $\mathcal{O}_{n[t]}$ -submodule of l and $\text{length}(l/l') = 1$. Let λ' be the type of l' . First we assume that $a_2(\lambda) > a_2(\lambda')$. We have $a_1(\lambda') = a_1(\lambda) + 1$ and $a_2(\lambda') = a_2(\lambda) - 1$. In this case, we may take an element \mathbf{w} of l' such that \mathbf{w} is nonzero in $V_{a_1(\lambda)+a_2(\lambda)+1}/V_{a_1(\lambda)+a_2(\lambda)}$. Since elements of l' are also elements of l , there exist complex numbers α_1 and α_2 such that $\mathbf{w} - \alpha_1 \mathbf{v}_1 - \alpha_2 \mathbf{v}_2 \in \langle f \mathbf{v}_1, f \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}$. Since $(\alpha_1, \alpha_2) \neq (0, 0)$, we have the ratio $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$ by \mathbf{w} . But we may check that the ratio $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$ is unique for the choice of such an element \mathbf{w} . Then the ratio $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$ corresponds to l' . Second we assume that $a_2(\lambda) = a_2(\lambda')$. We have $a_1(\lambda') = a_1(\lambda) - 1$. In particular, $a_1(\lambda) \neq 0$. In this case, we may take an element \mathbf{w} of l' such that \mathbf{w} is nonzero in $V_{a_1(\lambda)+a_2(\lambda)}/V_{a_1(\lambda)+a_2(\lambda)-1}$. We may check that $\mathbf{w} \in \langle f \mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}$. Since $\text{length}(l/l') = 1$, there exists a nonzero complex number α such that

$\mathbf{w} - \alpha \mathbf{v}_2 \in \langle f\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}$. So we have the ratio $[0 : \alpha] \in \mathbb{P}^1$ by \mathbf{w} . But we may check that the ratio $[0 : \alpha] \in \mathbb{P}^1$ is unique for the choice of such an element \mathbf{w} . Then the ratio $[0 : \alpha] \in \mathbb{P}^1$ corresponds to l' . By these arguments, we have the inverse map. \square

Let $\lambda^{(n)} = (1^{a_1}, 2^{a_2})$ be a Young diagram and l_n be an $\mathcal{O}_{n[t]}$ -submodule of $E|_{n[t]}$ with type $\lambda^{(n)}$. Before we consider families and parameter spaces of refined parabolic structures, we discuss families and parameter spaces of filtrations $l_n \supset l_{n-1} \supset \cdots \supset l_{n-m+1}$ ($2 \leq m \leq n$) with $\text{length}(l_{n-k+1}/l_{n-k}) = 1$ ($k = 1, 2, \dots, m-1$). If $m = n$, then this filtration is a refined parabolic structure. We consider examples with $m = 3$. We will consider the following four sequences of Young diagrams:

$$\begin{aligned} T: \lambda^{(n)} &= (1^{a_1}, 2^{a_2}) \supset (1^{a_1-1}, 2^{a_2}) \supset (1^{a_1-2}, 2^{a_2}) \\ T': \lambda^{(n)} &= (1^{a_1}, 2^{a_2}) \supset (1^{a_1+1}, 2^{a_2-1}) \supset (1^{a_1}, 2^{a_2-1}) \\ T'': \lambda^{(n)} &= (1^{a_1}, 2^{a_2}) \supset (1^{a_1-1}, 2^{a_2}) \supset (1^{a_1}, 2^{a_2-1}) \\ T''': \lambda^{(n)} &= (1^{a_1}, 2^{a_2}) \supset (1^{a_1+1}, 2^{a_2-1}) \supset (1^{a_1+2}, 2^{a_2-2}). \end{aligned}$$

Here we assume that

- $a_1 \geq 2$ when we consider the sequence T ,
- $a_2 \geq 1$ when we consider the sequences T' ,
- $a_1 \geq 1$ and $a_2 \geq 1$ when we consider the sequences T'' , and
- $a_2 \geq 2$ when we consider the sequence T''' .

We will construct families of filtrations with fixed l_n corresponding to these sequences of Young diagrams. Here for a sequence of Young diagrams, the ‘‘corresponding’’ means that filtrations parametrized by a Zariski open subset of the parameter space have this sequence of Young diagrams. We take generators \mathbf{v}_1 and \mathbf{v}_2 of l_n such that $\text{length}(\langle \mathbf{v}_1 \rangle_{\mathcal{O}_{n[t]}}) = a_1 + a_2$ and $\text{length}(\langle \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}) = a_2$.

Example 20. *First we will construct a family corresponding to T . This is the easiest case. Filtrations with T are unique. This filtration is*

$$l_n = \langle \mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} \supset \langle f\mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} \supset \langle f^2\mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}.$$

So the parameter space corresponding to T is a point.

Example 21. *Second we will construct a family corresponding to T' as follows. We define a family of filtrations with fixed l_n :*

$$\begin{aligned} l_{n-1} &= \langle \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2, f\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} \\ l_{n-2} &= \langle \alpha_1 f\mathbf{v}_1 + \alpha_2 f\mathbf{v}_2, f\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} = \langle f\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}, \end{aligned}$$

which is parametrized by $[\alpha_1 : \alpha_2] \in \mathbb{P}^1$. We denote by $C_{l_n, T'}$ this parameter space \mathbb{P}^1 . If $\alpha_1 \neq 0$, the sequences of types of the parametrized filtrations are T' . So filtrations parametrized by the affine line \mathbb{A}^1 have the sequence T' . If $\alpha_1 = 0$, the sequence of types of the parametrized filtration is T'' . So we have the parameter space $C_{l_n, T'}$ corresponding to T' .

Example 22. *Third we will construct a family corresponding to T'' as follows. We define a family of filtrations with fixed l_n :*

$$l'_{n-1} = \langle f\mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} \quad \text{and} \quad l'_{n-2} = \langle \alpha'_1 f\mathbf{v}_1 + \alpha'_2 \mathbf{v}_2, f^2\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}$$

which is parametrized by $[\alpha'_1 : \alpha'_2] \in \mathbb{P}^1$. So we have the parameter space $C_{l_n, T''}$ corresponding to T'' . This parameter space $C_{l_n, T''}$ is \mathbb{P}^1 .

We may check that $[0 : 1] \in C_{l_n, T'}$ and $[1 : 0] \in C_{l_n, T''}$ parametrize the same filtration:

$$l'_{n-1} = \langle f\mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}} \quad \text{and} \quad l'_{n-2} = \langle f\mathbf{v}_1, f\mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}.$$

So the parameter spaces $C_{l_n, T'}$ and $C_{l_n, T''}$ are the projective lines and the intersection $C_{l_n, T'} \cap C_{l_n, T''}$ is a point.

Proposition 23. *Let $\mathbb{F}_2 \rightarrow \mathbb{P}^1$ be the Hirzebruch surface of degree two. We denote by s_0 and p_∞ the 0-section and the fiber of $\mathbb{F}_2 \rightarrow \mathbb{P}^1$ at ∞ , respectively. We can construct a family of filtrations with fixed l_n corresponding to T''' . More precisely, by this family, s_0 is identified with $C_{l_n, T'}$, p_∞ is identified with $C_{l_n, T''}$, and $\mathbb{F}_2 \setminus (s_0 \cup p_\infty)$ is identified with*

$$\left\{ l_n \supset l_{n-1} \supset l_{n-2} \left| \begin{array}{l} l_{n-k} \text{ is an } \mathcal{O}_{n[t]}\text{-submodule of } l_{n-k+1} \\ \text{such that } \text{length}(l_{n-k+1}/l_{n-k}) = 1 \ (k = 1, 2) \\ \text{and the sequence of Young diagram is } T''' \end{array} \right. \right\}.$$

Proof. Set $U_0 := \{[1 : x_1] \mid x_1 \in \mathbb{C}\} \subset \mathbb{P}^1$ and $U_\infty := \{[y_1 : 1] \mid y_1 \in \mathbb{C}\} \subset \mathbb{P}^1$. We consider \mathbb{F}_2 as the gluing of $U_0 \times \mathbb{P}^1$ and $U_\infty \times \mathbb{P}^1$ as follows:

$$\begin{aligned} (U_0 \times \mathbb{P}^1)|_{U_0 \cap U_\infty} &\longrightarrow (U_\infty \times \mathbb{P}^1)|_{U_0 \cap U_\infty} \\ ([1 : x_1], [\alpha_{1,3} : \alpha_{2,3}]) &\longmapsto ([y_1 : 1], [\beta_{1,3} : \beta_{2,3}]) \end{aligned}$$

where $y_1 = 1/x_1$, $\beta_{1,3} = \alpha_{1,3}$, and $\beta_{2,3} = -x_1^{-2}\alpha_{2,3}$. We will construct a family of filtrations $l_n \supset l_{n-1} \supset l_{n-2}$ with $\text{length}(l_{n-k+1}/l_{n-k}) = 1$ ($k = 1, 2$) parametrized by \mathbb{F}_2 as follows. We take generators \mathbf{v}_1 and \mathbf{v}_2 of l_n such that $\text{length}(\langle \mathbf{v}_1 \rangle_{\mathcal{O}_{n[t]}}) = a_1 + a_2$ and $\text{length}(\langle \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}) = a_2$. For $([1 : x_1], [\alpha_{1,3} : \alpha_{2,3}]) \in U_0 \times \mathbb{P}^1$, we define l_{n-1} and l_{n-2} as

$$\begin{aligned} l_{n-1} &= \langle \mathbf{v}_1 + x_1\mathbf{v}_2, f\mathbf{v}_2 \rangle \\ l_{n-2} &= \langle \alpha_{1,3}(\mathbf{v}_1 + x_1\mathbf{v}_2) + \alpha_{2,3}f\mathbf{v}_2, f\mathbf{v}_1 + x_1f\mathbf{v}_2, f^2\mathbf{v}_2 \rangle. \end{aligned}$$

On the other hand, for $([y_1 : 1], [\beta_{1,3} : \beta_{2,3}]) \in U_\infty \times \mathbb{P}^1$ we define l'_{n-1} and l'_{n-2} as

$$\begin{aligned} l'_{n-1} &= \langle y_1\mathbf{v}_1 + \mathbf{v}_2, f\mathbf{v}_1 \rangle \\ l'_{n-2} &= \langle \beta_{1,3}(y_1\mathbf{v}_1 + \mathbf{v}_2) + \beta_{2,3}f\mathbf{v}_1, y_1f\mathbf{v}_1 + f\mathbf{v}_2, f^2\mathbf{v}_1 \rangle. \end{aligned}$$

On $U_0 \cap U_\infty$, we put $y_1 = 1/x_1$, $\beta_{1,3} = \alpha_{1,3}$, and $\beta_{2,3} = -x_1^{-2}\alpha_{2,3}$. Then we have that $l_{n-1}|_{U_0 \cap U_\infty} = l'_{n-1}|_{U_0 \cap U_\infty}$ and $l_{n-2}|_{U_0 \cap U_\infty} = l'_{n-2}|_{U_0 \cap U_\infty}$. So we obtain a family of filtrations parametrized by \mathbb{F}_2 . Points on the Zariski open subset $\alpha_{1,3} \neq 0$ in $U_0 \times \mathbb{P}^1$ parametrize filtrations whose sequences of types are T''' . So we say that the parameter space \mathbb{F}_2 corresponds to T''' . The curve $C_{l_n, T'}$ defined above is embedded into \mathbb{F}_2 as the 0-section. Here the 0-section is defined by $\alpha_{1,3} = 0$ over U_0 and $\beta_{1,3} = 0$ over U_∞ . The curve $C_{l_n, T''}$ defined above is embedded into \mathbb{F}_2 as the fiber of the point $y_1 = 0$ under $\mathbb{F}_2 \rightarrow \mathbb{P}^1$. \square

We observe that the dimension of the parameter space corresponding to a sequence of Young diagrams $\lambda^{(n)} \supset \lambda^{(n-1)} \supset \lambda^{(n-2)}$ is

$$\# \left\{ k \mid n-1 \leq k \leq n, a_2(\lambda^{(k-1)}) = a_2(\lambda^{(k)}) - 1 \right\}.$$

Next we consider an example with $m = 4$. That is, we will construct a family of filtrations $l_n \supset l_{n-1} \supset l_{n-2} \supset l_{n-3}$ with $\text{length}(l_{n-k+1}/l_{n-k}) = 1$ ($k = 1, 2, 3$).

Proposition 24. *We can construct a \mathbb{P}^1 -bundle over \mathbb{F}_2 such that this bundle has a 0-section and we can construct a family corresponding to*

$$T^4: \lambda^{(n)} = (1^{a_1}, 2^{a_2}) \supset (1^{a_1+1}, 2^{a_2-1}) \supset (1^{a_1+2}, 2^{a_2-2}) \supset (1^{a_1+3}, 2^{a_2-3})$$

over this \mathbb{P}^1 -bundle. Here we assume that $a_2 \geq 3$.

Proof. Let $l_n \supset \tilde{l}_{n-1} \supset \tilde{l}_{n-2}$ be the family of filtrations parametrized by \mathbb{F}_2 defined in the case corresponding to T''' . We set $U^1 := U_0 \times U_0$, $U^2 := U_0 \times U_\infty$, $U^3 := U_\infty \times U_0$, $U^4 := U_\infty \times U_\infty$. These products give an affine open covering of \mathbb{F}_2 . We consider the restriction $\tilde{l}_{n-2}|_{U^i}$ for $1 \leq i \leq 4$. Since we assume that $a_2 \geq 3$, we may take two elements \tilde{v}_1^i and \tilde{v}_2^i of $\tilde{l}_{n-2}|_{U^i}$ such that $\tilde{l}_{n-2}|_{U^i} = \langle \tilde{v}_1^i, \tilde{v}_2^i \rangle_{\mathcal{O}_{n[t]}}$. For each U^i , we define a family of filtrations $l_n \supset \tilde{l}_{n-1}^i \supset \tilde{l}_{n-2}^i \supset \tilde{l}_{n-3}^i$ (so $m = 4$) parametrized by $U^i \times \mathbb{P}^1$ as follows. We set $\tilde{l}_{n-1}^i := \tilde{l}_{n-1}|_{U^i}$, $\tilde{l}_{n-2}^i := \tilde{l}_{n-2}|_{U^i}$, and

$$(3.3) \quad \tilde{l}_{n-3}^i := \langle \alpha_{1,4}^i \tilde{v}_1^i + \alpha_{2,4}^i \tilde{v}_2^i, f \tilde{v}_1^i, f \tilde{v}_2^i \rangle_{\mathcal{O}_{n[t]}}.$$

where $[\alpha_{1,4}^i : \alpha_{2,4}^i] \in \mathbb{P}^1$. Then we glue $\{U^i \times \mathbb{P}^1\}_{1 \leq i \leq 4}$ so that these families patch together. Then we have a \mathbb{P}^1 -bundle over \mathbb{F}_2 . We denote by $\mathbb{F}_2^{(4)} \rightarrow \mathbb{F}_2$ this \mathbb{P}^1 -bundle. By the gluing the families, we have a family of filtrations $l_n \supset \tilde{l}_{n-1} \supset \tilde{l}_{n-2} \supset \tilde{l}_{n-3}$ parametrized by $\mathbb{F}_2^{(4)}$. Points on a Zariski open subset of $\mathbb{F}_2^{(4)}$ parametrize filtrations whose sequences of types are T^4 .

We may check that $\mathbb{F}_2^{(4)} \rightarrow \mathbb{F}_2$ has the 0-section. We may assume that $\text{length}(\langle \tilde{v}_1^i \rangle_{\mathcal{O}_{n[t]}}) \geq \text{length}(\langle \tilde{v}_2^i \rangle_{\mathcal{O}_{n[t]}})$ for each i . Then this 0-section is defined by $\alpha_{1,4}^i = 0$ in (3.3) for each i ($1 \leq i \leq 4$). That is, by the patching of the families, the local sections $\alpha_{1,4}^i = 0$ over U^i patch together. \square

To describe parameter spaces of refined parabolic structures $l_n \supset l_{n-1} \supset \dots \supset l_1 \supset 0$, we consider sequences of maps, which are \mathbb{P}^1 -bundles or identity maps as follows. For T^4 , we define a sequence of maps:

$$\mathbb{F}_{l_n, T^4}^{(4)} \xrightarrow{p_3} \mathbb{F}_{l_n, T^4}^{(3)} \xrightarrow{p_2} \mathbb{F}_{l_n, T^4}^{(2)} \xrightarrow{p_1} \mathbb{F}_{l_n, T^4}^{(1)} = \{\text{pt}\}$$

so that $\mathbb{F}_{l_n, T^4}^{(2)}$ is the projective line, p_2 is the Hirzebruch surface of degree two, and p_3 is the \mathbb{P}^1 -bundle in Proposition 24. We identify $\mathbb{F}_{l_n, T^4}^{(4)}$ with the parameter space of this family corresponding to T^4 . Let

$$T^5: \lambda = (1^{a_1}, 2^{a_2}) \supset (1^{a_1+1}, 2^{a_2-1}) \supset (1^{a_1+2}, 2^{a_2-2}) \supset (1^{a_1+1}, 2^{a_2-2}).$$

For T^5 , we define

$$\mathbb{F}_{l_n, T^5}^{(4)} \xrightarrow{p_3} \mathbb{F}_{l_n, T^5}^{(3)} \xrightarrow{p_2} \mathbb{F}_{l_n, T^5}^{(2)} \xrightarrow{p_1} \mathbb{F}_{l_n, T^5}^{(1)} = \{\text{pt}\}$$

so that $\mathbb{F}_{l_n, T^5}^{(2)}$ is the projective line, p_2 is the Hirzebruch surface of degree two, $\mathbb{F}_{l_n, T^4}^{(3)} = \mathbb{F}_{l_n, T^4}^{(4)}$, and p_3 is the identity map. We can construct a family corresponding to T^5 over the 0-section of the \mathbb{P}^1 -bundle in Proposition 24. So we may identify $\mathbb{F}_{l_n, T^5}^{(4)}$ with the parameter space of this family corresponding to T^5 .

Now we consider families and parameter spaces of refined parabolic structures.

Proposition 25. *We set $\lambda^{(n)} = (1^{a_1}, 2^{a_2})$. We fix an $\mathcal{O}_{n[t]}$ -submodule l_n of $E|_{n[t]}$ with $\text{length}(l_n) = n$ whose type is $\lambda^{(n)}$. Let \mathcal{T}_n be the set of standard Young tableaux whose shapes are the Young diagram $\lambda^{(n)}$. For $T = (\lambda^{(n)}, \lambda^{(n-1)}, \dots, \lambda^{(1)}) \in \mathcal{T}_n$, we can construct a sequence of maps:*

$$\mathbb{F}_{l_n, T}^{(n)} \xrightarrow{p_{n-1}} \mathbb{F}_{l_n, T}^{(n-1)} \xrightarrow{p_{n-2}} \dots \xrightarrow{p_2} \mathbb{F}_{l_n, T}^{(2)} \xrightarrow{p_1} \mathbb{F}_{l_n, T}^{(1)} = \{\text{pt}\}$$

such that

- $\mathbb{F}_{l_n, T}^{(k+1)} = \mathbb{F}_{l_n, T}^{(k)}$ and p_k is the identity when $a_2(\lambda^{(n-k+1)}) = a_2(\lambda^{(n-k)})$
- p_k is a \mathbb{P}^1 -bundle when $a_2(\lambda^{(n-k+1)}) = a_2(\lambda^{(n-k)}) + 1$
- there exists a family of refined parabolic structures $\tilde{l}_{\bullet, T}: E|_{n[t]} \supset l_n \supset \tilde{l}_{n-1, T} \supset \cdots \supset \tilde{l}_{1, T} \supset 0$ with the fixed l_n parametrized by $\mathbb{F}_{l_n, T}^{(n)}$ such that on a Zariski open subset $\mathbb{C}^{a_2} \subset \mathbb{F}_{l_n, T}^{(n)}$ the restriction $\tilde{l}_{\bullet, T}|_{\mathbb{C}^{a_2}}$ gives a bijection

$$\mathbb{C}^{a_2} \longrightarrow \left\{ l_{\bullet}: l_n \supset l_{n-1} \supset \cdots \supset l_1 \supset 0 \mid \begin{array}{l} l_{\bullet} \text{ is a refined parabolic structure on } n[t] \\ \text{whose standard tableau is } T \end{array} \right\}.$$

Proof. We will construct a sequence of maps inductively. We assume that we have a sequence of maps

$$\mathbb{F}_{l_n, T}^{(k)} \xrightarrow{p_{k-1}} \mathbb{F}_{l_n, T}^{(k-1)} \xrightarrow{p_{k-2}} \cdots \xrightarrow{p_2} \mathbb{F}_{l_n, T}^{(2)} \xrightarrow{p_1} \mathbb{F}_{l_n, T}^{(1)} = \{\text{pt}\}$$

and such a family $l_n \supset \tilde{l}_{n-1, T} \supset \cdots \supset \tilde{l}_{n-k+1, T}$ of filtrations parametrized by $\mathbb{F}_{l_n, T}^{(k)}$.

First we consider the case $a_2(\lambda^{(n-k+1)}) \neq 0$. We take an affine open covering $\{U^i\}_i$ of $\mathbb{F}_{l_n, T}^{(k)}$ so that we may take two elements $\tilde{\mathbf{v}}_1^i$ and $\tilde{\mathbf{v}}_2^i$ of $\tilde{l}_{n-k+1, T}|_{U^i}$ such that $\tilde{l}_{n-k+1, T}|_{U^i} = \langle \tilde{\mathbf{v}}_1^i, \tilde{\mathbf{v}}_2^i \rangle_{\mathcal{O}_{n[t]}}$ with $\text{length}\langle \tilde{\mathbf{v}}_1^i \rangle_{\mathcal{O}_{n[t]}} \geq \text{length}\langle \tilde{\mathbf{v}}_2^i \rangle_{\mathcal{O}_{n[t]}}$. For each U^i , we set

$$\tilde{l}_{n-k}^i := \langle \alpha_{1, k+1}^i \tilde{\mathbf{v}}_1^i + \alpha_{2, k+1}^i \tilde{\mathbf{v}}_2^i, f \tilde{\mathbf{v}}_1^i, f \tilde{\mathbf{v}}_2^i \rangle_{\mathcal{O}_{n[t]}}.$$

where $[\alpha_{1, k+1}^i : \alpha_{2, k+1}^i] \in \mathbb{P}^1$. We glue $\{U^i \times \mathbb{P}^1\}_i$ so that these families patch together. Then we have a \mathbb{P}^1 -bundle over $\mathbb{F}_{l_n, T}^{(k)}$. If $a_2(\lambda^{(n-k+1)}) = a_2(\lambda^{(n-k)})$, then we define $\mathbb{F}_{l_n, T}^{(k+1)}$ by the 0-section of this \mathbb{P}^1 -bundle and define $\tilde{l}_{n-k, T}$ by the gluing of $\langle \tilde{\mathbf{v}}_1^i, f \tilde{\mathbf{v}}_1^i \rangle_{\mathcal{O}_{n[t]}}$. If $a_2(\lambda^{(n-k+1)}) = a_2(\lambda^{(n-k)}) + 1$, then we define $\mathbb{F}_{l_n, T}^{(k+1)}$ by this \mathbb{P}^1 -bundle and define $\tilde{l}_{n-k, T}$ by the gluing of \tilde{l}_{n-k}^i .

Second we consider the case $a_2(\lambda^{(n-k+1)}) = 0$. In particular $a_2(\lambda^{(n-k+1)}) = a_2(\lambda^{(n-k)})$. We take an affine open covering $\{U^i\}_i$ of $\mathbb{F}_{l_n, T}^{(k)}$ so that we may take an element $\tilde{\mathbf{v}}^i$ such that $\tilde{l}_{n-k+1, T}|_{U^i} = \langle \tilde{\mathbf{v}}^i \rangle_{\mathcal{O}_{n[t]}}$. For each U^i , we set $\tilde{l}_{n-k}^i := \langle f \tilde{\mathbf{v}}^i \rangle_{\mathcal{O}_{n[t]}}$. We define $\mathbb{F}_{l_n, T}^{(k+1)}$ by $\mathbb{F}_{l_n, T}^{(k+1)} = \mathbb{F}_{l_n, T}^{(k)}$ and define $\tilde{l}_{n-k, T}$ by the gluing of \tilde{l}_{n-k}^i . \square

So the parameter space of refined parabolic structures with fixed l_n is $\bigcup_{T \in \mathcal{T}_{l_n}} \mathbb{F}_{l_n, T}^{(n)}$. Remark that $\mathbb{F}_{l_n, T}^{(n)}$ is irreducible and the dimension of $\mathbb{F}_{l_n, T}^{(n)}$ is a_2 for each T . If $a_2 = 0$ or $a_2 = 1$, we can describe $\bigcup_{T \in \mathcal{T}_{l_n}} \mathbb{F}_{l_n, T}^{(n)}$ easily. First if $a_2 = 0$, that is, $\lambda^{(n)} = (1^n, 2^0)$, then $\#\mathcal{T}_{l_n} = 1$ and $\mathbb{F}_{l_n, T}^{(n)}$ is a point. So $\bigcup_{T \in \mathcal{T}_{l_n}} \mathbb{F}_{l_n, T}^{(n)}$ is a point. This point corresponds to the filtration

$$l_n \supset f \cdot l_n \supset f^2 \cdot l_n \supset \cdots \supset f^{n-1} \cdot l_n \supset 0.$$

Next we consider the case $a_2 = 1$, that is, $\lambda^{(n)} := (0, 1, \dots, 1, 2) = (1^{n-2}, 2^1)$. We define a standard tableau T_k so that the corresponding sequence of Young diagrams is

$$(3.4) \quad \begin{aligned} \lambda^{(n)} &= (1^{n-2}, 2^1) \supset (1^{n-3}, 2^1) \supset \cdots \supset (1^{n-k-1}, 2^1) \\ &\supset (1^{n-k}, 2^0) \supset (1^{n-k-1}, 2^0) \supset \cdots \supset (1^1, 2^0) \end{aligned}$$

for $k = 1, 2, \dots, n-1$. By Proposition 25 and the argument in Example 22, we obtain the following statement:

Corollary 26. *The irreducible component $\mathbb{F}_{l_n, T_k}^{(n)}$ corresponding to T_k is the projective line \mathbb{P}^1 . The parameter space of refined parabolic structures with fixed l_n is*

$$\mathbb{F}_{l_n, T_1}^{(n)} \cup \mathbb{F}_{l_n, T_2}^{(n)} \cup \cdots \cup \mathbb{F}_{l_n, T_{n-1}}^{(n)},$$

which is a chain of $(n-1)$ projective lines. The intersection $\mathbb{F}_{l_n, T_{k_1}}^{(n)} \cap \mathbb{F}_{l_n, T_{k_2}}^{(n)}$ is a point if $|k_1 - k_2| = 1$. When $|k_1 - k_2| \neq 1$, this intersection is empty.

4. STABILITY OF REFINED PARABOLIC BUNDLES

In the previous section, we have defined refined parabolic bundles. We would like to consider a moduli space of refined parabolic bundles. In order to obtain a good moduli space, we have to introduce a stability condition of refined parabolic bundles. For this purpose, first, we define the stability index as in [21, Section 2]. This stability condition is the same as the stability condition as in [27]. In particular, the notion of parabolic bundles is contained in the notion of refined parabolic bundles. So by the definition of the stability condition for refined parabolic bundles, we have a stability condition for parabolic bundles.

Let $(E, \{l_{i, \bullet}\}_{i \in I})$ be a refined parabolic bundle. We fix weights

$$\mathbf{w} = (\mathbf{w}_1, \dots, \mathbf{w}_\nu), \quad \mathbf{w}_i = (w_{i, n_i}, \dots, w_{i, 1}) \in [0, 1]^{n_i}$$

such that

$$(4.1) \quad 0 \leq w_{i, n_i} \leq \cdots \leq w_{i, 1} \leq 1$$

for any $i \in I$.

Definition 27. *Let L be a line subbundle of E . We define the \mathbf{w} -stability index of L to the real number*

$$\text{Stab}_{\mathbf{w}}(L) := \deg(E) - 2 \deg(L) + \sum_{i \in I} \sum_{k=1}^{n_i} w_{i, k} (1 - 2 \text{length}((l_{i, k} \cap L|_{n_i[t_i]}) / (l_{i, k-1} \cap L|_{n_i[t_i]})))$$

where $l_{i, 0} = 0$ for any i .

If we set $\tilde{I} := \{(i, k) \mid i \in I, k = 1, 2, \dots, n_i\}$ and

$$\epsilon_{i, k}(L) := \begin{cases} -1 & \text{when } \text{length}((l_{i, k} \cap L|_{n_i[t_i]}) / (l_{i, k-1} \cap L|_{n_i[t_i]})) \neq 0 \\ 1 & \text{when } \text{length}((l_{i, k} \cap L|_{n_i[t_i]}) / (l_{i, k-1} \cap L|_{n_i[t_i]})) = 0 \end{cases},$$

then we have the following equality:

$$\text{Stab}_{\mathbf{w}}(L) = \deg(E) - 2 \deg(L) + \sum_{(i, k) \in \tilde{I}} \epsilon_{i, k}(L) \cdot w_{i, k}.$$

Definition 28. *A refined parabolic bundle (E, \mathbf{l}) is \mathbf{w} -stable (resp. \mathbf{w} -semistable) if for any subbundle $L \subset E$, the following inequality holds:*

$$\text{Stab}_{\mathbf{w}}(L) > 0 \quad (\text{resp. } \geq 0).$$

Now we discuss on the stability condition by using the parabolic degree. First, we recall a definition of the parabolic degree. We fix parabolic weights $\boldsymbol{\alpha} = (\boldsymbol{\alpha}_1, \dots, \boldsymbol{\alpha}_\nu)$, where each $\boldsymbol{\alpha}_i$ is a

tuple of real numbers $\alpha_i = (\alpha_{i,n_i+1}, \alpha_{i,n_i}, \dots, \alpha_{i,1})$ with the condition $0 < \alpha_{i,n_i+1} < \alpha_{i,n_i} < \dots < \alpha_{i,1} < 1$. We define an α -parabolic degree of a line subbundle L of E as

$$\text{para-deg}(L) = \deg(L) + \sum_{i \in I} \sum_{k=1}^{n_i+1} \alpha_{i,k} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})),$$

where $l_{i,n_i+1} := E|_{n_i[t_i]}$. We can define the stability of refined parabolic bundles by using the parabolic degree: A refined parabolic bundle (E, L) is stable if for any subbundle $L \subset E$, the following inequality holds:

$$\frac{\text{para-deg}(E)}{\text{rank}(E)} > \frac{\text{para-deg}(L)}{\text{rank}(L)}.$$

Now we will discuss a relation between the stability defined by the parabolic degree and the stability defined by the stability index. We compute $\text{para-deg}(E)/\text{rank}(E) - \text{para-deg}(L)/\text{rank}(L)$ as follows.

$$\begin{aligned} \frac{\text{para-deg}(E)}{\text{rank}(E)} - \frac{\text{para-deg}(L)}{\text{rank}(L)} &= \frac{1}{2} \left(\deg(E) - 2 \deg(L) + \sum_{i \in I} \sum_{k=1}^{n_i+1} \alpha_{i,k} \text{length}((l_{i,k}) / (l_{i,k-1})) \right. \\ &\quad \left. - 2 \sum_{i \in I} \sum_{k=1}^{n_i+1} \alpha_{i,k} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \\ &= \frac{1}{2} \left(\deg(E) - 2 \deg(L) + \sum_{i \in I} \left(\alpha_{i,n_i+1} n_i + \sum_{k=1}^{n_i} \alpha_{i,k} \right) \right. \\ &\quad \left. - 2 \sum_{i \in I} \left(\alpha_{i,n_i+1} \left(n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \right. \right. \\ &\quad \left. \left. + \sum_{k=1}^{n_i} \alpha_{i,k} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \right) \\ &= \frac{1}{2} \left(\deg(E) - 2 \deg(L) + \sum_{i \in I} \left(\sum_{k=1}^{n_i} (\alpha_{i,k} - \alpha_{i,n_i+1}) \right) \right. \\ &\quad \left. - 2 \sum_{i \in I} \left(\sum_{k=1}^{n_i} (\alpha_{i,k} - \alpha_{i,n_i+1}) \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \right). \end{aligned}$$

So if we set $w_{i,k} := \alpha_{i,k} - \alpha_{i,n_i+1}$, then we have the following equality

$$\frac{\text{para-deg}(E)}{\text{rank}(E)} - \frac{\text{para-deg}(L)}{\text{rank}(L)} = \frac{\text{Stab}_{\mathbf{w}}(L)}{2}.$$

Then the stability defined by the stability index is the same as the stability defined by the parabolic degree.

4.1. Tame and undecomposable refined parabolic bundles. In Section 2, we have defined undecomposable parabolic bundles and admissible parabolic bundles. Here we define undecomposable refined parabolic bundles and admissible refined parabolic bundles.

Definition 29. Let $(E, \{l_{i,\bullet}\}_{i \in I})$ be a refined parabolic bundle of rank 2 and of degree d . We say that $(E, \{l_{i,\bullet}\}_{i \in I})$ is decomposable if there exists a decomposition $E = L_1 \oplus L_2$ and $l_{i,k} = l_{i,k}^{(1)} \oplus l_{i,k}^{(2)}$

for any $i = 1, \dots, \nu$ and $k = 1, \dots, n_i$, where we set $l_{i,k}^{(1)} := l_{i,k} \cap L_1|_{n_i[t_i]}$ and $l_{i,k}^{(2)} := l_{i,k} \cap L_2|_{n_i[t_i]}$. We say that $(E, \{l_{i,\bullet}\}_{i \in I})$ is undecomposable if $(E, \{l_{i,\bullet}\}_{i \in I})$ is not decomposable.

Definition 30. Let $(E, \{l_{i,\bullet}\}_{i \in I})$ be a refined parabolic bundle of rank 2 and of degree d . We say $(E, \{l_{i,\bullet}\}_{i \in I})$ is admissible if this refined parabolic bundle satisfies the following condition:

$$\sum_{i \in I} \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \leq n + \deg(E) - 2 \deg(L) - 2$$

for any line subbundles L such that $\deg(E) \leq 2 \deg(L)$.

If D is a reduced effective divisor, then we have the fact that $(E, \{l_{i,\bullet}\}_{i \in I})$ is stable for a convenient choice of weights \mathbf{w} if, and only if, it is undecomposable (see [21, Proposition 3.4]). On the other hand, when D is not necessarily reduced, there exist examples of (refined) parabolic bundles $(E, \{l_{i,\bullet}\}_{i \in I})$ such that it is undecomposable and it is not stable for any weights \mathbf{w} .

Example 31. Let D and $(E, \mathbf{1})$ be one of the four listed parabolic bundles in Lemma 12. Then $(E, \mathbf{1})$ is undecomposable and admissible. But, by the inequalities (4.1), the parabolic bundle $(E, \mathbf{1})$ is not stable for any weights \mathbf{w} for any cases in Lemma 12. In particular, there exist parabolic bundles $(E, \{l_{i,\bullet}\}_{i \in I})$ such that it is Λ -flat and that it is not stable for any weights \mathbf{w} . (Here Λ is defined as in Proposition 13).

Now we will find a necessary and sufficient condition of a refined parabolic bundle $(E, \{l_{i,\bullet}\}_{i \in I})$ for the condition that there exist weights \mathbf{w} such that $(E, \{l_{i,\bullet}\}_{i \in I})$ is \mathbf{w} -stable. For this purpose, we will define tame refined parabolic bundles. For a line subbundle $L \subset E$ and $i \in I$, we define an integer $N_i(L)$ as

$$(4.2) \quad N_i(L) = \max_{k' \in \{1, 2, \dots, n_i\}} \left\{ \sum_{k=1}^{k'} \epsilon_{i,k}(L) \right\}.$$

We set $I_L^+ = \{i \in I \mid N_i(L) > 0\}$. Moreover, we set

$$k_{i,\max}(L) := \max \left\{ k' \in \{1, 2, \dots, n_i\} \mid N_i(L) = \sum_{k=1}^{k'} \epsilon_{i,k}(L) \right\},$$

$$K_{i,\max}(L) := \{1, 2, \dots, k_{i,\max}(L)\}, \quad \text{and} \quad K_{i,\max}^c(L) := \{1, 2, \dots, n_i\} \setminus K_{i,\max}(L).$$

Definition 32. Let $(E, \{l_{i,\bullet}\}_{i \in I})$ be a refined parabolic bundle of rank 2 and of degree d . We say $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame if this refined parabolic bundle satisfies the following condition:

- I_L^+ is not empty, and
- $-\deg(E) + 2 \deg(L) + 1 \leq \sum_{i \in I_L^+} N_i(L)$

for any line subbundles L such that $\deg(E) \leq 2 \deg(L)$.

If E has a decomposition $E \cong \mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ where $d_1 < d_2$, then there exists a unique line subbundle L such that $\deg(E) \leq 2 \deg(L)$. This line subbundle is the destabilizing bundle $\mathcal{O}(d_2)$. When $d_1 = d_2$, there also exists a line subbundle L such that $\deg(E) \leq 2 \deg(L)$. Remark that this line bundle is not unique in this case. Now, we will show the implication (1.2), that is,

$$(E, \mathbf{1}) \text{ is undecomposable and tame} \implies (E, \mathbf{1}) \text{ is admissible.}$$

Moreover we will show that, when $(E, \mathbf{1})$ is a parabolic bundle,

$$(E, \mathbf{1}) \text{ is simple} \iff (E, \mathbf{1}) \text{ is undecomposable and tame.}$$

Proposition 33. *If a refined parabolic bundle $(E, \{l_{i, \bullet}\}_{i \in I})$ is tame and undecomposable, then it is admissible.*

Proof. For each $i \in I_L^+$, we have an inequality

$$N_i(L) \leq \sum_{k \in \{k | \epsilon_{i,k}(L)=1\}} \epsilon_{i,k}(L) = n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})).$$

Clearly $n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \geq 0$. Then we have that for any line subbundles L such that $\deg(E) \leq 2 \deg(L)$,

$$\begin{aligned} -\deg(E) + 2 \deg(L) + 1 &\leq \sum_{i \in I_L^+} N_i(L) \\ &\leq \sum_{i \in I_L^+} \left(n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \\ &\leq \sum_{i \in I} \left(n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) \right) \end{aligned}$$

We set $E = \mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ (with $d_1 \leq d_2$) and $L = \mathcal{O}(d_2)$. We assume that

$$(4.3) \quad -\deg(E) + 2 \deg(L) + 1 = \sum_{i \in I_L^+} N_i(L),$$

$$(4.4) \quad N_i(L) = \sum_{k \in \{k | \epsilon_{i,k}(L)=1\}} \epsilon_{i,k}(L) \quad (\text{for any } i \in I_L^+),$$

and

$$(4.5) \quad n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})) = 0 \quad (\text{for any } i \notin I_L^+).$$

By the equalities (4.4) and (4.5), we have

$$(4.6) \quad \epsilon_{i,k}(L) = \begin{cases} 1 & \text{for any } i \in I_L^+ \text{ and any } k \in K_{i,\max}(L) \\ -1 & \text{for any } i \in I_L^+ \text{ and any } k \in K_{i,\max}^c(L) \\ -1 & \text{for any } i \notin I_L^+ \text{ and any } k \in \{1, 2, \dots, n_i\} \end{cases}.$$

By the equality (4.3), we have the following equality:

$$(4.7) \quad d_2 - d_1 + 1 = \sum_{i \in I_L^+} \#K_{i,\max}(L) = \sum_{i \in I_L^+} k_{i,\max}(L).$$

If $d_1 = d_2$, then $\sum_{i \in I_L^+} k_{i,\max}(L) = 1$. So $\#I_L^+ = 1$ and $K_{i_0,\max}(L) = \{1\}$ for the $i_0 \in I_L^+$. We have that $\epsilon_{i,k}(L) = 1$ when $(i, k) = (i_0, 1)$ (with $i_0 \in I_L^+$) and $\epsilon_{i,k}(L) = -1$ otherwise. This implies

that $(E, \{l_{i,\bullet}\}_{i \in I})$ is decomposable. It is a contradiction. So $d_1 \neq d_2$. Then the vector bundle $\mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ has the following automorphism:

$$\begin{pmatrix} 1 & 0 \\ G & c \end{pmatrix},$$

where $c \in \mathbb{C}^*$ and $G = G(x)$ be a polynomial of degree less than or equal to $d_2 - d_1$. By (4.6) and (4.7), there exists an automorphism of $\mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ such that $l_{i,k} = l_{i,k}^{(1)} \oplus l_{i,k}^{(2)}$ for any $i = 1, \dots, \nu$ and $k = 1, \dots, n_i$. Here we set $l_{i,k}^{(1)} := l_{i,k} \cap \mathcal{O}(d_1)|_{n_i[t_i]}$ and $l_{i,k}^{(2)} := l_{i,k} \cap \mathcal{O}(d_2)|_{n_i[t_i]}$. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable, it is a contradiction. So we have

$$-\deg(E) + 2\deg(L) + 1 < \sum_{i \in I} \left(n_i - \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]}) \right).$$

This inequality implies that $(E, \{l_{i,\bullet}\}_{i \in I})$ is admissible. \square

Proposition 34. *Let $(E, \{l_i\}_{i \in I})$ be a parabolic bundle (Definition 6) and $(E, \{l_{i,\bullet}\}_{i \in I})$ be the corresponding refined parabolic bundle. The parabolic bundle $(E, \{l_i\}_{i \in I})$ is simple if, and only if, $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable and tame.*

Proof. First we will show that $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable and tame when $(E, \{l_i\}_{i \in I})$ is simple. If $(E, \{l_i\}_{i \in I})$ is simple, $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable by Proposition 8. So we will show that $(E, \{l_i\}_{i \in I})$ is not simple when $(E, \{l_{i,\bullet}\}_{i \in I})$ is not tame. We will show that there is a non-zero nilpotent element of $\text{End}(E, \{l_i\}_{i \in I})$. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is not tame, there exists a line subbundle L_0 with $\deg(E) \leq \deg(L_0)$ such that $I_{L_0}^+$ is empty or $-\deg(E) + 2\deg(L_0) \geq \sum_{i \in I_{L_0}^+} N_i(L_0)$. We decompose $E = L \oplus L_0$ with $\deg(L) \leq \deg(L_0)$. We take a nilpotent element $N \in \text{End}(L \oplus L_0)$ as follows:

$$N = \begin{pmatrix} 0 & 0 \\ f & 0 \end{pmatrix}$$

where $f = f(x)$ is a polynomial where $\deg(f) \leq 2\deg(L_0) - \deg(E)$. We set $m_i = \text{length}(l_i \cap \mathcal{O}(L_0)|_{n_i[t_i]})$ and $I^+ = \{i \in I \mid n_i - 2m_i \geq 0\}$. Since l_i is free, we may check that

$$\epsilon_{i,k}(L_0) = \begin{cases} 1 & (m_i < k \leq n_i) \\ -1 & (1 \leq k \leq m_i) \end{cases}$$

for any $i \in I$. So we have that $N_i(L_0) = n_i - 2m_i$ and $I^+ = I_{L_0}^+$. First we assume that $I_{L_0}^+$ is empty. Then, by using the equalities (2.2), we may check that there exists a non-zero polynomial f such that N preserves the parabolic structure $\{l_i\}_{i \in I}$. So if $I_{L_0}^+$ is empty, then $(E, \{l_i\}_{i \in I})$ is not simple. Second we assume that $I^+ \neq \emptyset$ and $-\deg(E) + 2\deg(L_0) \geq \sum_{i \in I_{L_0}^+} N_i(L_0)$. By this inequality, we have that

$$(4.8) \quad \sum_{i \in I^+} (n_i - 2m_i) \leq 2\deg(L_0) - \deg(E).$$

Then, by using the equalities (2.2), we may check that there exists a non-zero polynomial f such that N preserves the parabolic structure $\{l_i\}_{i \in I}$. Finally we obtain that $(E, \{l_i\}_{i \in I})$ is not simple when $(E, \{l_{i,\bullet}\}_{i \in I})$ is not tame.

Next we will show that $(E, \{l_i\}_{i \in I})$ is simple when $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable and tame. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is indecomposable, any endomorphism A of the $(E, \{l_i\}_{i \in I})$ takes the form $A = c \cdot \text{Id}_E + N$ with $c \in \mathbb{C}$ and N nilpotent. We assume that there exists a non-zero nilpotent

element N of $\text{End}(E, \{l_i\}_{i \in I})$. By using the argument as in the verification of the inequality (2.3), we may check that there exists a line subbundle L_0 with $\deg(E) \leq 2 \deg(L_0)$ such that $\sum_{i \in I^+} (n_i - 2m_i) \leq 2 \deg(L_0) - \deg(E)$. Here, m_i and I^+ are defined as above for this L_0 . By the argument as above, we have that $N_i(L_0) = n_i - 2m_i$ and $I_{L_0}^+ = I^+$. So we obtain that $I_{L_0}^+$ is empty or the inequality $-\deg(E) + 2 \deg(L_0) \geq \sum_{i \in I_{L_0}^+} N_i(L_0)$ holds. Since $(E, \{l_{i, \bullet}\}_{i \in I})$ is tame, it is a contradiction. So $(E, \{l_i\}_{i \in I})$ is simple. \square

4.2. Proof of Theorem A. Now, we start to show Theorem A (Theorem 1 and Corollary 2 below).

Proposition 35. *If a refined parabolic bundle $(E, \{l_{i, \bullet}\}_{i \in I})$ is stable for a convenient choice of weights \mathbf{w} , then it is tame and undecomposable.*

Proof. Assume that $(E, \{l_{i, \bullet}\}_{i \in I})$ is decomposable. There exists a decomposition $E \cong L_1 \oplus L_2$ such that $\text{Stab}_{\mathbf{w}}(L_1) + \text{Stab}_{\mathbf{w}}(L_2) = 0$ for any weight \mathbf{w} . Then $(E, \{l_{i, \bullet}\}_{i \in I})$ is not stable for any weights \mathbf{w} . Assume that $(E, \{l_{i, \bullet}\}_{i \in I})$ is not tame. There exists a line subbundle $L \subset E$ with $\deg(E) \leq 2 \deg(L)$ such that

- I_L^+ is empty, or
- I_L^+ is not empty and $-\deg(E) + 2 \deg(L) \geq \sum_{i \in I_L^+} N_i(L)$.

If I_L^+ is empty, then we have the inequality $\sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w_{i,k} \leq 0$ for any $i \in I$, since we have the inequalities (4.1). So we have

$$\begin{aligned} \text{Stab}_{\mathbf{w}}(L) &= \deg(E) - 2 \deg(L) + \sum_{i \in I} \sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w_{i,k} \\ &\leq \deg(E) - 2 \deg(L) \leq 0. \end{aligned}$$

Then $(E, \{l_{i, \bullet}\}_{i \in I})$ is not stable for any weights \mathbf{w} . Now we assume that I_L^+ is not empty. By the inequalities (4.1) and the definitions of $K_{i, \max}(L)$ and $K_{i, \max}^c(L)$, we have the following inequalities for each $i \in I_L^+$:

$$\sum_{k \in K_{i, \max}^c(L)} \epsilon_{i,k}(L) \cdot w_{i,k} \leq 0 \quad \text{and} \quad \sum_{k \in K_{i, \max}(L)} \epsilon_{i,k}(L) \cdot w_{i,k} \leq N_i(L).$$

If $i \notin I_L^+$, then $\sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w_{i,k} \leq 0$. By these inequalities, we have

$$\begin{aligned} \text{Stab}_{\mathbf{w}}(L) &= \deg(E) - 2 \deg(L) + \sum_{i \in I \setminus I_L^+} \sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w_{i,k} \\ &\quad + \sum_{i \in I_L^+} \sum_{k \in K_{i, \max}^c(L)} \epsilon_{i,k}(L) \cdot w_{i,k} + \sum_{i \in I_L^+} \sum_{k \in K_{i, \max}(L)} \epsilon_{i,k}(L) \cdot w_{i,k} \\ &\leq \deg(E) - 2 \deg(L) + \sum_{i \in I_L^+} N_i(L) \leq 0. \end{aligned}$$

Then $(E, \{l_{i, \bullet}\}_{i \in I})$ is not stable for any weights \mathbf{w} . \square

Proposition 36. *Assume that $E = \mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ with $d_1 < d_2$. If a refined parabolic bundle $(E, \{l_{i, \bullet}\}_{i \in I})$ is tame and undecomposable, then it is stable for a convenient choice of weights \mathbf{w} .*

Proof. We assume that $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame and undecomposable. Since $d_1 < d_2$, there exists a unique morphism $\varphi_{d_2}: \mathcal{O}(d_2) \subset E$. We set

$$N_i^c(\mathcal{O}(d_2)) := \begin{cases} \sum_{k \in K_{i,\max}^c(\mathcal{O}(d_2))} \epsilon_{i,k}(\mathcal{O}(d_2)) & \text{for } i \in I_{\mathcal{O}(d_2)}^+ \\ \sum_{k=1}^{n_i} \epsilon_{i,k}(\mathcal{O}(d_2)) & \text{for } i \notin I_{\mathcal{O}(d_2)}^+ \end{cases}.$$

For the unique morphism $\varphi_{d_2}: \mathcal{O}(d_2) \subset E$, we define weights $\mathbf{w} = (w_1, \dots, w_n)$ as follows:

$$w_{i,k} = \begin{cases} w & \text{for } i \in I_{\mathcal{O}(d_2)}^+ \text{ and } k \in K_{i,\max}(\mathcal{O}(d_2)) \\ w' & \text{for } i \in I_{\mathcal{O}(d_2)}^+ \text{ and } k \in K_{i,\max}^c(\mathcal{O}(d_2)) \\ w' & \text{for } i \notin I_{\mathcal{O}(d_2)}^+ \text{ and } k \in \{1, 2, \dots, n_i\} \end{cases}.$$

Here w and w' are real numbers such that

$$(4.9) \quad \begin{cases} w > -\frac{\sum_{i \in I} N_i^c(\mathcal{O}(d_2))}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} \cdot w' + \frac{d_2 - d_1}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} \\ w < -\frac{\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) - 2}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} \cdot w' + \frac{d_2 - d_1}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} \end{cases} \quad \text{and} \quad \begin{cases} 0 < w' < w < 1 \\ w + w' < 1 \end{cases}.$$

Remark that, since $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame and $d_1 < d_2$, we have the following inequalities:

$$0 < \frac{d_2 - d_1}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} < 1 \quad \text{and} \quad -\frac{\sum_{i \in I} N_i^c(\mathcal{O}(d_2))}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))} < -\frac{\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) - 2}{\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))}.$$

So there exist such real numbers w and w' .

We will check that $(E, \{l_{i,\bullet}\}_{i \in I})$ is \mathbf{w} -stable for such a weights \mathbf{w} . For the subbundle $\mathcal{O}(d_2) \subset E$, we have the following inequality:

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(d_2)) = d_1 - d_2 + \left(\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) \right) w' + \left(\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2)) \right) w > 0$$

by the inequalities (4.9). Next we calculate the stability index $\text{Stab}_{\mathbf{w}}(\mathcal{O}(d_1))$ for any subbundles $\mathcal{O}(d_1) \subset E$:

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(d_1)) \geq d_2 - d_1 - \left(\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) \right) w' - \left(\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2)) \right) w + 2w' > 0$$

by the inequalities (4.9). Here in the first inequality, we used the assumption that $(E, \{l_{i,\bullet}\}_{i \in I})$ is undecomposable and $w' < w$. Let L be a subbundle $L \subset E$ such that $\deg(L) \leq d_1 - 1$. We will calculate the stability index $\text{Stab}_{\mathbf{w}}(L)$. For the subbundle L , we set

$$K_{\delta_1, \delta_2}^{(i, d_2)}(L) := \{k \mid k \in \{1, 2, \dots, n_i\}, \text{ and } \epsilon_{i,k}(L) = \delta_1, \epsilon_{i,k}(\mathcal{O}(d_2)) = \delta_2\},$$

where $i \in I$ and $\delta_1, \delta_2 \in \{+, -\}$. We set $k_{\delta_1, \delta_2}^{(i, d_2)}(L) = \#K_{\delta_1, \delta_2}^{(i, d_2)}(L)$ for any $i \in I$, and

$$k_{\delta_1, \delta_2}^{(i, d_2)}(L)' = \#(K_{\delta_1, \delta_2}^{(i, d_2)}(L) \cap K_{i,\max}(\mathcal{O}(d_2))) \quad \text{and} \quad k_{\delta_1, \delta_2}^{(i, d_2)}(L)'' = \#(K_{\delta_1, \delta_2}^{(i, d_2)}(L) \cap K_{i,\max}^c(\mathcal{O}(d_2)))$$

for any $i \in I_L^+$. We have the following inequality:

$$\begin{aligned}
& \sum_{i \in I_{\mathcal{O}(d_2)}^+} \sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w_{i,k} + \sum_{i \in I \setminus I_{\mathcal{O}(d_2)}^+} \sum_{k=1}^{n_i} \epsilon_{i,k}(L) \cdot w' \\
&= \sum_{i \in I_{\mathcal{O}(d_2)}^+} (k_{+,+}^{(i,d_2)}(L)' - k_{-,+}^{(i,d_2)}(L)' + k_{+,-}^{(i,d_2)}(L)' - k_{-,-}^{(i,d_2)}(L)') \cdot w \\
&\quad + \sum_{i \in I_{\mathcal{O}(d_2)}^+} (k_{+,+}^{(i,d_2)}(L)'' - k_{-,+}^{(i,d_2)}(L)'' + k_{+,-}^{(i,d_2)}(L)'' - k_{-,-}^{(i,d_2)}(L)'') \cdot w' \\
&\quad + \sum_{i \in I \setminus I_{\mathcal{O}(d_2)}^+} (k_{+,+}^{(i,d_2)}(L) - k_{-,+}^{(i,d_2)}(L) + k_{+,-}^{(i,d_2)}(L) - k_{-,-}^{(i,d_2)}(L)) \cdot w' \\
&\geq \sum_{i \in I_{\mathcal{O}(d_2)}^+} (-N_i(\mathcal{O}(d_2)) - 2k_{-,-}^{(i,d_2)}(L)') \cdot w + \sum_{i \in I_{\mathcal{O}(d_2)}^+} (-N_i^c(\mathcal{O}(d_2)) - 2k_{-,-}^{(i,d_2)}(L)'') \cdot w' \\
&\quad + \sum_{i \in I \setminus I_{\mathcal{O}(d_2)}^+} (-N_i^c(\mathcal{O}(d_2)) - 2k_{-,-}^{(i,d_2)}(L)) \cdot w'
\end{aligned}$$

By this inequality and $w' < w$, we have the following inequality:

$$\begin{aligned}
& \text{Stab}_{\mathbf{w}}(L) \geq \deg(E) - 2 \deg(L) \\
(4.10) \quad & - \sum_{i \in I} N_i^c(\mathcal{O}(d_2))w' - \sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))w - 2 \left(\sum_{i \in I} k_{-,-}^{(i,d_2)}(L) \right) w
\end{aligned}$$

We describe the subbundle $L \hookrightarrow \mathcal{O}(d_1) \oplus \mathcal{O}(d_2)$ by a vector $(f_1^L(x), f_2^L(x))$, where $f_1^L(x)$ and $f_2^L(x)$ are polynomials in x such that $\deg(f_1^L(x)) \leq d_1 - \deg(L)$ and $\deg(f_2^L(x)) \leq d_2 - \deg(L)$. We have

$$[\text{the zero order of } f_1^L(x) \text{ at } t_i] \geq \#\{k \in \{1, 2, \dots, n_i\} \mid \epsilon_{i,k}(L) = \epsilon_{i,k}(\mathcal{O}(d_2)) = -1\}$$

for each $i \in I$. By this inequality, we have $d_1 - \deg(L) \geq \sum_{i \in I} k_{-,-}^{(i,d_2)}(L)$. When $\sum_{i \in I} k_{-,-}^{(i,d_2)}(L) \neq 0$ for the subbundle $L \subset E$, we have the following inequality:

$$\begin{aligned}
\text{Stab}_{\mathbf{w}}(L) &\geq d_2 - d_1 - \sum_{i \in I} N_i^c(\mathcal{O}(d_2))w' - \sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2))w + \left(\sum_{i \in I} k_{-,-}^{(i,d_2)}(L) \right) \cdot 2(1-w) \\
&\geq d_2 - d_1 - \left(\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) \right) w' - \left(\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2)) \right) w + 2(1-w) \\
&> 2(1-w-w') > 0.
\end{aligned}$$

Here in the first inequality, we used the inequality $d_1 - \deg(L) \geq \sum_{i \in I} k_{-,-}^{(i,d_2)}(L)$ and the inequality (4.10). In the second inequality, we used the inequalities $\sum_{i \in I} k_{-,-}^{(i,d_2)}(L) \geq 1$ and $w < 1$. In the third inequality and in the last inequality, we used the inequalities in (4.9). When $\sum_{i \in I} k_{-,-}^{(i,d_2)}(L) = 0$,

for the subbundle $L \subset E$, we have the following inequality:

$$\begin{aligned} \text{Stab}_{\mathbf{w}}(L) &\geq d_1 + d_2 - 2 \deg(L) - \left(\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) \right) w' - \left(\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2)) \right) w \\ &\geq d_2 - d_1 - \left(\sum_{i \in I} N_i^c(\mathcal{O}(d_2)) \right) w' - \left(\sum_{i \in I_{\mathcal{O}(d_2)}^+} N_i(\mathcal{O}(d_2)) \right) w + 1 \\ &> 1 - 2w' > 0. \end{aligned}$$

Here we used the inequality $\deg(L) \leq d_1 - 1$ in the second inequality. Then we obtain that $(E, \{l_{i,\bullet}\}_{i \in I})$ is \mathbf{w} -stable for the weights \mathbf{w} . \square

Proposition 37. *Assume that $E = \mathcal{O}(d) \oplus \mathcal{O}(d)$. If a refined parabolic bundle $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame and undecomposable, then it is stable for a convenient choice of weights \mathbf{w} .*

Proof. We take an element $i_1 \in I$. Since $E = \mathcal{O}(d) \oplus \mathcal{O}(d)$ and $\text{length}(l_{i_1,1}) = 1$, we can take a line subbundle $L_1 \subset E$ with $\deg(L_1) = d$ such that $\epsilon_{i_1,1}(L_1) = -1$. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame, there exists an element $i_2 \in I$ such that

$$N_{i_2}(L_1) = \sum_{k \in K_{i_2, \max}(L_1)} \epsilon_{i_2,k} \geq 1.$$

We divide into the following two cases:

- (A) the set $\{k \in K_{i_2, \max}(L_1) \mid \epsilon_{i_2,k}(L_1) = 1, \epsilon_{i_2,k}(L) = -1\}$ is empty for any line subbundles $L \subset E$ with $\deg(L) = d$,
- (B) the set $\{k \in K_{i_2, \max}(L_1) \mid \epsilon_{i_2,k}(L_1) = 1, \epsilon_{i_2,k}(L) = -1\}$ is not empty for some line subbundle $L \subset E$ with $\deg(L) = d$.

For the case (A), we determine weights \mathbf{w} as follows:

(A-i) When $i_1 \neq i_2$,

$$w_{i,k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_2 \text{ and } k \in K_{i_2, \max}(L_1) \\ 0 & \text{otherwise} \end{cases}$$

(A-ii) When $i_1 = i_2$,

$$w_{i,k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_1 \text{ and } k \in K_{i_1, \max}(L_1) \setminus \{1\} \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_2 < w_1 < 1$.

For the case (B), we determine weights \mathbf{w} as follows. We take a line subbundle L_2 of E with $\deg(L_2) = d$ so that

$$(4.11) \quad \#\{k \in K_{i_2, \max}(L_1) \mid \epsilon_{i_2,k}(L_1) = 1, \epsilon_{i_2,k}(L_2) = -1\}$$

is maximized. In particular $L_1 \neq L_2$ in $E = \mathcal{O}(d) \oplus \mathcal{O}(d)$. So it is impossible that $\epsilon_{i,k}(L_1) = \epsilon_{i,k}(L_2) = -1$ for any $(i, k) \in \tilde{I}$. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is undecomposable, the set

$$(4.12) \quad \{(i, k) \in \tilde{I} \mid \epsilon_{i,k}(L_1) = \epsilon_{i,k}(L_2) = 1\}$$

is not empty. We take an element (i_3, k_3) of the set (4.12) such that the pair satisfies the equality $k_3 = \min\{k \in \{1, 2, \dots, n_{i_3}\} \mid \epsilon_{i_3, k}(L_1) = \epsilon_{i_3, k}(L_2) = 1\}$. If the subset

$$(4.13) \quad \{k \in \{1, 2, \dots, n_i\} \mid \epsilon_{i_2, k}(L_1) = \epsilon_{i_2, k}(L_2) = 1\} \cap K_{i_2, \max}(L_1)$$

is not empty, we take the element (i_3, k_3) from this subset (4.13). So, if $i_2 \neq i_3$, we have that

$$(4.14) \quad \epsilon_{i_2, k}(L_1) + \epsilon_{i_2, k}(L_2) = 0 \quad (k \in K_{i_2, \max}(L_1)).$$

So, when $i_2 \neq i_3$, or when $i_2 = i_3$ and $k_3 > k_{i_2, \max}(L_1)$, we have that

$$(4.15) \quad \epsilon_{i_2, k}(L) = 1 \quad (k \in K_{i_2, \max}(L_1))$$

for any line subbundles L such that $\deg(L) = d$, $L \neq L_1$, and $L \neq L_2$.

(B-i) When i_1, i_2, i_3 are distinct from each other, we determine weights \mathbf{w} as follows:

$$w_{i, k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_2 \text{ and } k \in K_{i_2, \max}(L_1) \\ w_3 & \text{when } i = i_3 \text{ and } k = 1, 2, \dots, k_3 - 1 \\ w_4 & \text{when } i = i_3 \text{ and } k = k_3 \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_1, w_2 < 1$ and $0 < w_4 < w_3 < 1$.

(B-ii) When $i_1 \neq i_2 = i_3$ and $k_{i_2, \max}(L_1) \geq k_3$, we determine weights \mathbf{w} as follows:

$$w_{i, k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_2 \text{ and } k \in K_{i_2, \max}(L_1) \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_1, w_2 < 1$.

(B-iii) When $i_1 \neq i_2 = i_3$ and $k_3 > k_{i_1, \max}(L_1)$, we determine weights \mathbf{w} as follows:

$$w_{i, k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_2 \text{ and } k \in K_{i_2, \max}(L_1) \\ w_3 & \text{when } i = i_2 \text{ and } k = k_{i_2, \max}(L_1) + 1, \dots, k_3 - 1 \\ w_4 & \text{when } i = i_2 \text{ and } k = k_3 \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_4 < w_3 < w_2 < 1$ and $0 < w_1 < 1$.

(B-iv) When $i_1 = i_3 \neq i_2$, we determine weights \mathbf{w} as follows:

$$w_{i, k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_2 \text{ and } k \in K_{i_2, \max}(L_1) \\ w_3 & \text{when } i = i_1 \text{ and } k = 2, 3, \dots, k_3 - 1 \\ w_4 & \text{when } i = i_1 \text{ and } k = k_3 \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_2 < 1$ and $0 < w_4 < w_3 < w_1 < 1$.

(B-v) When $i_1 = i_2 \neq i_3$, we determine weights \mathbf{w} as follows:

$$w_{i,k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_1 \text{ and } k \in K_{i_1, \max}(L_1) \setminus \{1\} \\ w_3 & \text{when } i = i_3 \text{ and } k = 1, \dots, k_3 - 1 \\ w_4 & \text{when } i = i_3 \text{ and } k = k_3 \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_2 < w_1 < 1$ and $0 < w_4 < w_3 < 1$.

(B-vi) When $i_1 = i_2 = i_3$ and $k_{i_1, \max}(L_1) \geq k_3$, we determine weights \mathbf{w} as follows:

$$w_{i,k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_1 \text{ and } k \in K_{i_1, \max}(L_1) \setminus \{1\} \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_2 < w_1 < 1$.

(B-vii) When $i_1 = i_2 = i_3$ and $k_3 > k_{i_1, \max}(L_1)$, we determine weights \mathbf{w} as follows:

$$w_{i,k} := \begin{cases} w_1 & \text{when } i = i_1 \text{ and } k = 1 \\ w_2 & \text{when } i = i_1 \text{ and } k \in K_{i_1, \max}(L_1) \setminus \{1\} \\ w_3 & \text{when } i = i_1 \text{ and } k = k_{i_1, \max}(L_1) + 1, \dots, k_3 - 1 \\ w_4 & \text{when } i = i_1 \text{ and } k = k_3 \\ 0 & \text{otherwise} \end{cases}$$

with $0 < w_4 < w_3 < w_2 < w_1 < 1$.

If we take a convenient choice of w_1, w_2, w_3, w_4 (or w_1, w_2), we may check that $(E, \{l_{i, \bullet}\}_{i \in I})$ is \mathbf{w} -stable for the weights \mathbf{w} . Now we check this claim for the 3 cases: (A-i); (B-i); and (B-vi). For other cases, we will check $(E, \{l_{i, \bullet}\}_{i \in I})$ is \mathbf{w} -stable for some weights \mathbf{w} by the same argument as in the three cases.

We consider the case (A-i). We assume that

$$-w_1 + N_{i_2}(L_1) \cdot w_2 > 0.$$

Since $N_{i_2}(L_1) \geq 1$, there exists a pair w_1, w_2 satisfies this inequality and $0 < w_2, w_1 < 1$. We have the following inequality for $\text{Stab}_{\mathbf{w}}(L_1)$:

$$\text{Stab}_{\mathbf{w}}(L_1) = -w_1 + N_{i_2}(L_1) \cdot w_2 > 0.$$

For any line subbundles $L \subset E$ with $L_1 \neq L$, we have the following inequality:

$$\begin{aligned} \text{Stab}_{\mathbf{w}}(L) &= \deg(E) - 2 \deg(L) + w_1 + k_{i_2, \max}(L_1) \cdot w_2 \\ &\geq w_1 + k_{i_2, \max}(L_1) \cdot w_2 > 0. \end{aligned}$$

Then we obtain that $(E, \{l_{i, \bullet}\}_{i \in I})$ is \mathbf{w} -stable for the weights \mathbf{w} .

We consider the case (B-i). We assume that

$$\begin{cases} -w_1 + N_{i_2}(L_1)w_2 + \left(\sum_{k'=1}^{k_3-1} \epsilon_{i_3, k'}(L_1) \right) w_3 + w_4 > 0 \\ -w_1 + N_{i_2}(L_1)w_2 + \left(\sum_{k'=1}^{k_3-1} \epsilon_{i_3, k'}(L_1) \right) w_3 - w_4 < 0 \\ w_1 + k_{i_2, \max}(L_1) \cdot w_2 + (k_3 - 1)w_3 - w_4 > 0 \\ w_1 + k_{i_2, \max}(L_1) \cdot w_2 + (k_3 - 1)w_3 + w_4 < 2 \end{cases}.$$

We may check that there exist such real numbers w_1, w_2, w_3, w_4 . We have the following inequalities for $\text{Stab}_{\mathbf{w}}(L_1)$ and $\text{Stab}_{\mathbf{w}}(L_2)$:

$$\begin{aligned}\text{Stab}_{\mathbf{w}}(L_1) &= -w_1 + N_{i_2}(L_1)w_2 + \left(\sum_{k'=1}^{k_3-1} \epsilon_{i_3, k'}(L_1) \right) w_3 + w_4 > 0 \\ \text{Stab}_{\mathbf{w}}(L_2) &= w_1 - N_{i_2}(L_1)w_2 - \left(\sum_{k'=1}^{k_3-1} \epsilon_{i_3, k'}(L_1) \right) w_3 + w_4 > 0.\end{aligned}$$

Here for the equality in the computation of $\text{Stab}_{\mathbf{w}}(L_2)$, we used the equality (4.14) and the equality $\epsilon_{i_3, k}(L_1) + \epsilon_{i_3, k}(L_2) = 0$ for $k = 1, 2, \dots, k_3 - 1$. This equality is given by the equality $k_3 = \min\{k \in \{1, 2, \dots, n_{i_3}\} \mid \epsilon_{i_3, k}(L_1) = \epsilon_{i_3, k}(L_2) = 1\}$. For any line subbundles L such that $\deg(L) = d$, $L \neq L_1$, and $L \neq L_2$, we have the following inequalities:

$$\text{Stab}_{\mathbf{w}}(L) \geq w_1 + k_{i_2, \max}(L_1) \cdot w_2 + (k_3 - 1)w_3 - w_4 > 0.$$

by the equality (4.15). For any line subbundles L with $\deg(L) \leq d - 1$, we have the following inequalities

$$\begin{aligned}\text{Stab}_{\mathbf{w}}(L) &\geq \deg(E) - 2 \deg(L) - w_1 - k_{i_2, \max}(L_1) \cdot w_2 - (k_3 - 1)w_3 - w_4 \\ &\geq 2 - w_1 - k_{i_2, \max}(L_1) \cdot w_2 - (k_3 - 1)w_3 - w_4 > 0.\end{aligned}$$

Then we obtain that $(E, \{l_{i, \bullet}\}_{i \in I})$ is \mathbf{w} -stable for the weights \mathbf{w} .

We consider the case (B-vi). We assume that

$$\begin{cases} -w_1 + (N_{i_2}(L_1) + 1) \cdot w_2 > 0 \\ w_1 - (N_{i_2}(L_1) - 1) \cdot w_2 > 0 \\ w_1 + (k_{i_2, \max}(L_1) - 1) \cdot w_2 < 2 \end{cases}.$$

Since $N_{i_2}(L_1) \geq 1$, there exists a pair w_1, w_2 satisfies this inequality and $0 < w_2, w_1 < 1$. We have the following inequality for $\text{Stab}_{\mathbf{w}}(L_1)$:

$$\text{Stab}_{\mathbf{w}}(L_1) = -w_1 + (N_{i_2}(L_1) + 1) \cdot w_2 > 0,$$

since $\epsilon_{i_1, 1}(L_1) = -1$ and $N_{i_2}(L_1) = \sum_{k \in K_{i_2, \max}(L_1)} \epsilon_{i_2, k}(L_1)$. For any line subbundles $L \subset E$ with $\deg(L) = d$ and $L \neq L_1$, we have the following inequality:

$$\begin{aligned}\text{Stab}_{\mathbf{w}}(L) &\geq \text{Stab}_{\mathbf{w}}(L_2) \geq \deg(E) - 2 \deg(L_2) + w_1 + (-N_{i_2}(L_1) + 1) \cdot w_2 \\ &= w_1 - (N_{i_2}(L_1) - 1) \cdot w_2 > 0.\end{aligned}$$

Here for the first inequality, we used the fact that L_2 maximizes $\#\{k \in K_{i_2, \max}(L_1) \mid \epsilon_{i_2, k}(L_1) = 1, \epsilon_{i_2, k}(L_2) = -1\}$ and for the second inequality we used $k_{i_1, \max}(L_1) \geq k_3$. For any line subbundles $L \subset E$ with $\deg(L) \leq d - 1$, we have the following inequality:

$$\begin{aligned}\text{Stab}_{\mathbf{w}}(L) &\geq \deg(E) - 2 \deg(L) - w_1 - (k_{i_2, \max}(L_1) - 1) \cdot w_2 \\ &\geq 2 - w_1 - (k_{i_2, \max}(L_1) - 1) \cdot w_2 > 0.\end{aligned}$$

Then we obtain that $(E, \{l_{i, \bullet}\}_{i \in I})$ is \mathbf{w} -stable for the weights \mathbf{w} . □

By Proposition 35, Proposition 36, and Proposition 37, we obtain the following theorem:

Theorem 1. *A refined parabolic bundle $(E, \{l_{i, \bullet}\}_{i \in I})$ is tame and undecomposable if, and only if, it is stable for a convenient choice of weights \mathbf{w} .*

By Proposition 34, we have the following corollary:

Corollary 2. *Let $(E, \{l_i\}_{i \in I})$ be a parabolic bundle (Definition 6) and $(E, \{l_{i, \bullet}\}_{i \in I})$ be the corresponding refined parabolic bundle. The parabolic bundle $(E, \{l_i\}_{i \in I})$ is simple if, and only if, $(E, \{l_{i, \bullet}\}_{i \in I})$ is stable for a convenient choice of weights \mathbf{w}*

5. ELEMENTARY TRANSFORMATION OF REFINED PARABOLIC BUNDLES

For studying parabolic connections or parabolic bundles (where D is a reduced effective divisor), the elementary transformations play an important role. In this section, we will define elementary transformations for refined parabolic bundles and will show some properties of the elementary transformations.

5.1. Elementary transformations for Λ -connections. First we recall the elementary transformations for Λ -connections. Let (E, ∇) be a Λ -connection. For (E, ∇) we may define transformations as follows. Let $\mathbf{l} = \{l_i\}_{i \in I}$ be the corresponding parabolic structure of (E, ∇) . That is, l_i is a free submodule of $E|_{n_i[t_i]}$. We take an integer k where $1 \leq k \leq n_i$. Let $\varphi_k: E|_{n_i[t_i]} \rightarrow E|_{k[t_i]}$ be the natural morphism. For the free submodule $\varphi_k(l_i)$ of $E|_{k[t_i]}$, the vector bundle $E'_{\varphi_k(l_i)}$ is defined by the exact sequence of sheaves

$$0 \longrightarrow E'_{\varphi_k(l_i)} \longrightarrow E \longrightarrow E|_{k[t_i]}/\varphi_k(l_i) \longrightarrow 0.$$

The degree of $E'_{\varphi_k(l_i)}$ is $\deg(E) - k$. Let $\nabla_{\varphi_k(l_i)}$ be the induced connection by the morphism $E'_{\varphi_k(l_i)} \rightarrow E$. So we have a transformation $(E'_{\varphi_k(l_i)}, \nabla_{\varphi_k(l_i)})$. Let $\Lambda'_{i,k}$ be the formal data of $(E'_{\varphi_k(l_i)}, \nabla_{\varphi_k(l_i)})$. By this transformation, we have a map $\mathfrak{Con}_{\Lambda}(D) \rightarrow \mathfrak{Con}_{\Lambda'_{i,k}}(D)$.

5.2. Elementary transformations for parabolic bundles. Second we recall the elementary transformations for parabolic bundles. Let $(E, \{l_i\}_{i \in I})$ be a parabolic bundle. For $(E, \{l_i\}_{i \in I})$ we may define transformations as follows. For the free submodule l_i of $E|_{n_i[t_i]}$, the vector bundle E'_i is defined by the exact sequence of sheaves

$$0 \longrightarrow E'_i \longrightarrow E \longrightarrow E|_{n_i[t_i]}/l_i \longrightarrow 0.$$

The degree of E'_i is $\deg(E) - n_i$. We have a filtration of sheaves

$$E'_i \otimes \mathcal{O}(-n_i[t_i]) \xrightarrow{\subset} E \otimes \mathcal{O}(-n_i[t_i]) \xrightarrow{\subset} E'_i \xrightarrow{\subset} E.$$

We define a parabolic structure l'_i on E'_i by

$$l'_i = (E \otimes \mathcal{O}(-n_i[t_i])) / (E'_i \otimes \mathcal{O}(-n_i[t_i])).$$

So we have a transformation $(E'_i, \{l_j\}_{j \in I \setminus \{i\}} \cup \{l'_i\})$. We call this transformation the *elementary transformation* of a parabolic bundles $(E, \{l_i\}_{i \in I})$ at $n_i[t_i]$. We have the elementary transformations for Λ -connections and parabolic bundles:

$$(E, \nabla) \mapsto (E'_{\varphi_k(l_i)}, \nabla_{\varphi_k(l_i)}) \quad \text{and} \quad (E, \{l_i\}_{i \in I}) \mapsto (E'_i, \{l_j\}_{j \in I \setminus \{i\}} \cup \{l'_i\}),$$

respectively. On the other hand, we have a correspondence $(E, \nabla) \mapsto (E, \{l_i\}_{i \in I})$. If $k = n_i$, then these transformations are compatible with this correspondence.

5.3. Definition of elementary transformations for refined parabolic bundles. Now we consider the extension of the elementary transformation of parabolic bundles. That is, we will define the elementary transformation for refined parabolic bundles. For this purpose, we rephrase refined parabolic structures as filtrations of sheaves. By this rephrasing, we may define the elementary transformation for refined parabolic bundles simply. Let (E, \mathbf{l}) be a refined parabolic bundle, where $\mathbf{l} = \{l_{i, \bullet}\}_{i \in I}$. We take $i_0 \in I$ and consider the filtration $l_{i_0, \bullet} : E|_{n_{i_0}[t_{i_0}]} \supset l_{i_0, n_{i_0}} \supset l_{i_0, n_{i_0}-1} \supset \cdots \supset l_{i_0, 1} \supset 0$. We set

$$E_{i_0}^{(k)} := \ker(E \longrightarrow E|_{n_{i_0}[t_{i_0}]} / l_{i_0, k})$$

for $k = 0, 1, 2, \dots, n_{i_0}$. Then we have a filtration of sheaves

$$(5.1) \quad E \supset E_{i_0}^{(n_{i_0})} \supset E_{i_0}^{(n_{i_0}-1)} \supset \cdots \supset E_{i_0}^{(1)} \supset E_{i_0}^{(0)} = E(-n_{i_0}[t_{i_0}])$$

where $\text{length}(E/E_{i_0}^{(n_{i_0})}) = n_{i_0}$ and $\text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k-1)}) = 1$ for $k = 1, 2, \dots, n_{i_0}$. Conversely, if we set $l_{i_0, k} := E_{i_0}^{(k)}/E_{i_0}^{(0)}$ for $k = 0, 1, 2, \dots, n_{i_0}$, then we have a filtration

$$E/E_{i_0}^{(0)} = E|_{n_i[t_i]} \supset l_{i_0, n_{i_0}} \supset l_{i_0, n_{i_0}-1} \supset \cdots \supset l_{i_0, 1} \supset 0$$

of $\mathcal{O}_{n_{i_0}[t_{i_0}]}$ -modules where the length of $l_{i_0, k}$ is k .

Now we will transform the filtration (5.1) into a new filtration of sheaves. We set $E'_{i_0} = E_{i_0}^{(n_{i_0})}$. Since $\text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k-1)}) = 1$ for $k = 1, 2, \dots, n_{i_0}$, we may check that $E_{i_0}^{(k-1)} \supset E_{i_0}^{(k)}(-[t_{i_0}])$ for $k = 1, 2, \dots, n_{i_0}$. So we can define a filtration of sheaves as

$$\begin{aligned} E'_{i_0} &= E_{i_0}^{(n_{i_0})} \supset E_{i_0}^{(0)} \supset E_{i_0}^{(1)}(-[t_{i_0}]) \supset E_{i_0}^{(2)}(-2[t_{i_0}]) \supset \cdots \\ &\quad \cdots \supset E_{i_0}^{(n_{i_0}-1)}(-(n_{i_0}-1)[t_{i_0}]) \supset E_{i_0}^{(n_{i_0})}(-n_{i_0}[t_{i_0}]). \end{aligned}$$

This is a new filtration of sheaves. For $k = 0, 1, \dots, n_{i_0}$, we set

$$(E'_{i_0})^{(n_{i_0}-k)} := E^{(k)}(-k[t_{i_0}]).$$

We will check that $\text{length}(E'_{i_0}/(E'_{i_0})^{(n_{i_0})}) = n_{i_0}$ and $\text{length}((E'_{i_0})^{(k)}/(E'_{i_0})^{(k-1)}) = 1$ for $k = 1, 2, \dots, n_{i_0}$ as follows. For $k = 1, 2, \dots, n_{i_0}$, we have the following equalities:

$$\begin{aligned} &\text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k-1)}) + \text{length}((E'_{i_0})^{(n_{i_0}-k+1)}/(E'_{i_0})^{(n_{i_0}-k)}) \\ &= \text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k-1)}) + \text{length}(E_{i_0}^{(k-1)}(-(k-1)[t_{i_0}])/E_{i_0}^{(k)}(-k[t_{i_0}])) \\ &= \text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k)}(-[t_{i_0}])) = 2. \end{aligned}$$

Since $\text{length}(E_{i_0}^{(k)}/E_{i_0}^{(k-1)}) = 1$ for $k = 1, 2, \dots, n_{i_0}$, we have

$$\text{length}((E'_{i_0})^{(n_{i_0}-k+1)}/(E'_{i_0})^{(n_{i_0}-k)}) = 1$$

for $k = 1, 2, \dots, n_{i_0}$. Moreover we may check that $\text{length}(E'_{i_0}/(E'_{i_0})^{(n_{i_0})}) = n_{i_0}$. Finally, if we set $l'_{i_0, k} := (E'_{i_0})^{(k)}/(E'_{i_0})^{(0)}$ for $k = 0, 1, 2, \dots, n_{i_0}$, then we have a filtration

$$(5.2) \quad E|_{n_i[t_i]} \supset l'_{i_0, n_{i_0}} \supset l'_{i_0, n_{i_0}-1} \supset \cdots \supset l'_{i_0, 1} \supset 0$$

of $\mathcal{O}_{n_{i_0}[t_{i_0}]}$ -modules where the length of $l'_{i_0, k}$ is k .

Definition 38. We fix i_0 where $i_0 \in I$. We define the elementary transformation of $(E, \{l_{i,\bullet}\}_{i \in I})$ at $n_{i_0}[t_{i_0}]$ by

$$\text{elm}_{i_0}^-(E, \{l_{i,\bullet}\}_{i \in I}) = (E'_{i_0}, \{l_{i,\bullet}\}_{i \in I \setminus \{i_0\}} \cup \{l'_{i_0,\bullet}\}).$$

The degree of E'_{i_0} is $d - n_{i_0}$.

5.4. Some properties of elementary transformations for refined parabolic bundles. We take $i \in I$. Let $l_{i,\bullet}: l_{i,n_i} \supset \cdots \supset l_{i,1} \supset 0$ be a refined parabolic structure at $n_i[t_i]$ and $l'_{i,\bullet}$ be its transformation. For $l_{i,\bullet}$ and $l'_{i,\bullet}$, we can define standard tableaux T and T' , respectively (see Definition 18). We will consider the relation between these standard tableaux T and T' . Let $\lambda_i^{(k)}$ be the Young diagram which is the type of $l_{i,k}$. We define integers a_1^k and a_2^k so that $\lambda_i^{(k)} = (1^{a_1^k}, 2^{a_2^k})$ for $k = 1, 2, \dots, n_i$. We set

$$X_1(k) := a_1^{n_i} + a_2^{n_i} - a_1^k - a_2^k \quad \text{and} \quad X_2(k) := a_2^{n_i} - a_2^k.$$

Definition 39. We say a refined parabolic structure $l_{i,\bullet}$ is generic if

- $a_2^{n_i} = 0$, or
- $a_2^{n_i} \neq 0$ and we can take generators $\mathbf{v}_1, \mathbf{v}_2 \in l_{i,n_i}$ such that $\text{length}(\langle \mathbf{v}_1 \rangle) = a_1^{n_i} + a_2^{n_i}$, $\text{length}(\langle \mathbf{v}_2 \rangle) = a_2^{n_i}$, and $l_{i,\bullet}$ satisfies

$$l_{i,k} / \langle f_i^{X_1(k)} \mathbf{v}_1, f_i^{X_2(k)} \mathbf{v}_2 \rangle \neq 0$$

for any k with $a_2^{k+1} - a_2^k > 0$.

Now we consider examples of generic $l_{i,\bullet}$. Set $\lambda_i^{(n_i)} = (1^{n_i-2}, 2^1)$. So $a_1^{n_i} = n_i - 2$ and $a_2^{n_i} = 1$. Let T_k be the standard tableau defined in (3.4) for $k = 1, 2, \dots, n-1$. We fix k_0 ($1 \leq k_0 \leq n-1$). If the standard tableau of a refined parabolic bundle is T_{k_0} , then $a_2^{n-k_0+1} - a_2^{n-k_0} = 1 > 0$ and $a_2^{k+1} - a_2^k = 0$ when $k \neq n - k_0$. We take elements $\mathbf{v}_1, \mathbf{v}_2 \in E|_{n[t]}$ such that $\text{length}(\langle \mathbf{v}_1 \rangle) = n_i - 1$, and $\text{length}(\langle \mathbf{v}_2 \rangle) = 1$, and $l_{i,n_i} = \langle \mathbf{v}_1, \mathbf{v}_2 \rangle_{\mathcal{O}_{n[t]}}$. In particular, $f_i^{n_i-1} \mathbf{v}_1 = f_i \mathbf{v}_2 = 0$. We can describe a refined parabolic structure as follows:

$$\begin{aligned} l_{i,n_i} &= \langle \mathbf{v}_1, \mathbf{v}_2 \rangle \supset \langle f_i \mathbf{v}_1, \mathbf{v}_2 \rangle \supset \langle f_i^2 \mathbf{v}_1, \mathbf{v}_2 \rangle \supset \cdots \supset \langle f_i^{k_0-1} \mathbf{v}_1, \mathbf{v}_2 \rangle \\ &\supset \langle f_i^{k_0-1} \mathbf{v}_1 + \alpha \mathbf{v}_2 \rangle \supset \langle f_i^{k_0} \mathbf{v}_1 \rangle \supset \cdots \supset \langle f_i^{n_i-2} \mathbf{v}_1 \rangle, \end{aligned}$$

where $\alpha \in \mathbb{C}$. This refined parabolic structure is generic if, and only if, $\alpha \neq 0$.

To describe the standard tableau of the transformation $l'_{i,\bullet}$, first, we define integers $b_{n_i}, b_{n_i-1}, \dots, b_1$ by $b_{n_i} = a_1^{n_i}$ and $b_k = \min\{b_{k+1}, a_1^k\}$ for $k = n_i - 1, \dots, 2, 1$. Second, we modify $X_1(k)$ and $X_2(k)$ by using b_k as follows:

$$X_1^b(k) := X_1(k) + a_1^k - b_k \quad \text{and} \quad X_2^b(k) := X_2(k) - a_1^k + b_k.$$

We claim that $X_2^b(k) \geq 0$. Indeed, $X_2^b(k) = a_2^{n_i} - a_1^k - a_2^k + b_k$. When $a_1^k + a_2^k > a_2^{n_i}$, we have $\min\{a_1^{n-1}, a_1^{n-2}, \dots, a_1^k\} \geq a_1^k + a_2^k - a_2^{n_i}$ since we have the inclusions $l_{i,n_i} \supset l_{i,n_i-1} \supset l_{i,n_i-2} \supset \cdots \supset l_{i,k}$. Then we obtain $X_2^b(k) \geq 0$.

Proposition 40. Let $l'_{i,\bullet}$ be the transformation (5.2) of $l_{i,\bullet}$. Assume that $l_{i,\bullet}$ is generic. Then the type of l'_{i,n_i-k} is the following Young diagram:

$$\begin{cases} (1^{X_2^b(k)-X_1^b(k)}, 2^{X_1^b(k)}) & \text{when } X_2^b(k) - X_1^b(k) \geq 0 \\ (1^{X_1^b(k)-X_2^b(k)}, 2^{X_2^b(k)}) & \text{when } X_2^b(k) - X_1^b(k) < 0. \end{cases}$$

Proof. When $a_2^{n_i} = 0$, it is clear. So we consider the case where $a_2^{n_i} \neq 0$. We take generators $\mathbf{v}_1, \mathbf{v}_2$ of l_{i, n_i} as in the definition of the genericness of $l_{i, \bullet}$. We can take $F(f_i) \in \mathcal{O}_{n_i[t_i]}$ (where $F(0) \neq 0$) so that

$$(5.3) \quad l_{i, k} = \left\langle f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} F(f_i) \mathbf{v}_2, f_i^{X_2(k)} \mathbf{v}_2 \right\rangle.$$

Indeed, if $k = n_i$, then

$$\left\langle f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} F(f_i) \mathbf{v}_2, f_i^{X_2(k)} \mathbf{v}_2 \right\rangle = \langle \mathbf{v}_1, \mathbf{v}_2 \rangle$$

since $X_1(n_i) = X_2(n_i) = 0$ and $b_{n_i} = a_1^{n_i}$. We fix k ($1 \leq k \leq n_i$). We assume that $l_{i, k}$ has the description (5.3) for the fixed k . Since $X_2^b(k) \leq X_2(k)$ and $F(0) \neq 0$, we have an inequality

$$\text{length} \left(f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} F(f_i) \mathbf{v}_2 \right) \geq \text{length} \left(f_i^{X_2(k)} \mathbf{v}_2 \right).$$

First, we consider the case $a_1^k = a_1^{k-1} + 1$ and $a_2^k = a_2^{k-1}$ for the fixed k . In this case, the submodule $l_{i, k-1}$ has the following descriptions:

$$l_{i, k-1} = \left\langle f_i \cdot \left(f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} F(f_i) \mathbf{v}_2 \right), f_i^{X_2(k)} \mathbf{v}_2 \right\rangle.$$

If $b_k > b_{k-1}$, then $b_k = b_{k-1} + 1$ and $b_{k-1} = a_1^{k-1}$. So $X_2^b(k) = X_2^b(k-1) = X_2(k-1)$. Since $X_1(k) + 1 = X_1(k-1)$, $X_2(k) = X_2(k-1)$ and $X_2^b(k) + 1 > X_2(k-1)$, there exists

$$\begin{aligned} l_{i, k-1} &= \left\langle f_i^{X_1(k)+1} \mathbf{v}_1 + f_i^{X_2^b(k)+1} F(f_i) \mathbf{v}_2, f_i^{X_2(k)} \mathbf{v}_2 \right\rangle \\ &= \left\langle f_i^{X_1(k-1)} \mathbf{v}_1, f_i^{X_2(k-1)} \mathbf{v}_2 \right\rangle. \end{aligned}$$

Here remark that $X_2^b(k-1) = X_2(k-1)$. If $b_k = b_{k-1}$, then $X_2^b(k) + 1 = X_2^b(k-1)$. Since $X_2(k) = X_2(k-1)$, we have

$$\begin{aligned} l_{i, k-1} &= \left\langle f_i^{X_1(k)+1} \mathbf{v}_1 + f_i^{X_2^b(k)+1} F(f_i) \mathbf{v}_2, f_i^{X_2(k)} \mathbf{v}_2 \right\rangle \\ &= \left\langle f_i^{X_1(k-1)} \mathbf{v}_1 + f_i^{X_2^b(k-1)} F(f_i) \mathbf{v}_2, f_i^{X_2(k-1)} \mathbf{v}_2 \right\rangle. \end{aligned}$$

Second, we consider the case $a_1^k = a_1^{k-1} - 1$ and $a_2^k = a_2^{k-1} + 1$ for the fixed k . In this case, the submodule $l_{i, k-1}$ has the following descriptions:

$$l_{i, k-1} = \left\langle f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} \tilde{F}(f_i) \mathbf{v}_2, f_i^{X_2(k)+1} \mathbf{v}_2 \right\rangle,$$

since \mathbf{v}_1 and \mathbf{v}_2 satisfy the condition of the genericness of $l_{i, \bullet}$. Here remark that $X_2(k) \geq X_2^b(k)$. Since $a_1^k = a_1^{k-1} - 1$, $a_1^k \geq b_k$, and $b_{k-1} = \min\{b_k, a_1^{k-1}\}$, we have $b_k = b_{k-1}$. We may check that $X_1(k) = X_1(k-1)$, $X_2^b(k) = X_2^b(k-1)$ and $X_2(k) = X_2(k-1) - 1$. So

$$\begin{aligned} l_{i, k-1} &= \left\langle f_i^{X_1(k)} \mathbf{v}_1 + f_i^{X_2^b(k)} \tilde{F}(f_i) \mathbf{v}_2, f_i^{X_2(k)+1} \mathbf{v}_2 \right\rangle \\ &= \left\langle f_i^{X_1(k-1)} \mathbf{v}_1 + f_i^{X_2^b(k-1)} \tilde{F}(f_i) \mathbf{v}_2, f_i^{X_2(k-1)} \mathbf{v}_2 \right\rangle. \end{aligned}$$

Finally, we have the claim that we may take $F(f_i) \in \mathcal{O}_{n_i[t_i]}$ (where $F(0) \neq 0$) so that (5.3).

We take $\tilde{\mathbf{v}}_1, \tilde{\mathbf{v}}_2 \in E \otimes \mathcal{O}_{\mathbb{P}^1, t_i}$ such that $\tilde{\mathbf{v}}_1|_{n_i[t_i]} = \mathbf{v}_1$ and $\tilde{\mathbf{v}}_2|_{n_i[t_i]} = \mathbf{v}_2$. We define $\tilde{\mathbf{v}}'_1, \tilde{\mathbf{v}}'_2, \tilde{\mathbf{v}}''_1, \tilde{\mathbf{v}}''_2 \in E \otimes \mathcal{O}_{\mathbb{P}^1, t_i}$ as

$$\begin{aligned} \tilde{\mathbf{v}}'_1 &= f_i^{a_1^k + 2a_2^k} (f_i^{X_1(k)} \tilde{\mathbf{v}}_1 + f_i^{X_2^b(k)} F(f_i) \tilde{\mathbf{v}}_2), & \tilde{\mathbf{v}}'_2 &= f_i^{a_1^k + 2a_2^k} f_i^{X_2(k)} \tilde{\mathbf{v}}_2, \\ \tilde{\mathbf{v}}''_1 &= f_i^{a_1^{n_i} + 2a_2^{n_i}} \tilde{\mathbf{v}}_1, & \tilde{\mathbf{v}}''_2 &= f_i^{a_1^{n_i} + 2a_2^{n_i}} \tilde{\mathbf{v}}_2, \end{aligned}$$

respectively. Then we have $(E'_i)^{(k)} \otimes \mathcal{O}_{\mathbb{P}^1, t_i} \cong \langle \tilde{\mathbf{v}}'_1 \rangle \oplus \langle \tilde{\mathbf{v}}'_2 \rangle$ and $(E'_i)^{(n_i)} \otimes \mathcal{O}_{\mathbb{P}^1, t_i} \cong \langle \tilde{\mathbf{v}}''_1 \rangle \oplus \langle \tilde{\mathbf{v}}''_2 \rangle$. Now we compute the quotient $l'_{i,k} = (E'_i)^{(k)} / (E'_i)^{(n_i)}$. We may check the following equalities

$$\tilde{\mathbf{v}}'_1 = f_i^{-X_2(k)} \tilde{\mathbf{v}}''_1 + f_i^{-X_1^b(k)} F(f_i) \tilde{\mathbf{v}}''_2, \quad \tilde{\mathbf{v}}'_2 = f_i^{-X_1(k)} \tilde{\mathbf{v}}''_2,$$

and

$$f_i^{X_1^b(k) - X_1(k)} \tilde{\mathbf{v}}'_1 - F(f_i) \tilde{\mathbf{v}}'_2 = f_i^{-X_2^b(k)} \tilde{\mathbf{v}}''_1.$$

We set $\tilde{\mathbf{v}}'_3 := f_i^{X_1^b(k) - X_1(k)} \tilde{\mathbf{v}}'_1 - F(f_i) \tilde{\mathbf{v}}'_2$, which is an element of $l'_{i,k}$. Here remark that $X_1^b(k) - X_1(k) = a_1^k - b_k \geq 0$. We have

$$f_i^{X_1^b(k)} \tilde{\mathbf{v}}'_1 = f_i^{X_1(k)} \tilde{\mathbf{v}}'_2 = f_i^{X_2^b(k)} \tilde{\mathbf{v}}'_3 = 0$$

in $l'_{i,k}$. Here remark that $X_1^b(k) - X_2(k) = a_1^{n_i} - b_k \geq 0$. So, when $X_2^b(k) - X_1^b(k) \geq 0$, the type of $l'_{i,k}$ is $(1^{X_2^b(k) - X_1^b(k)}, 2^{X_1^b(k)})$. When $X_2^b(k) - X_1^b(k) < 0$ the type of $l'_{i,k}$ is $(1^{X_1^b(k) - X_2^b(k)}, 2^{X_2^b(k)})$. \square

If $a_2^{n_i} = 0$ or $a_2^{n_i} = 1$, then we have a simple description of the standard tableau of the transformation $l'_{i,\bullet}$. If $a_2^{n_i} = 0$, the standard tableau of $l_{i,\bullet}$ is unique. We denote by T_0 this standard tableau. Let T' be the standard tableau of the transformation $l'_{i,\bullet}$ of $l_{i,\bullet}$. Then we have $T' = T_0$ by the proposition above. That is, if $(E, \{l_{i,\bullet}\}_{i \in I})$ is an ordinary parabolic bundle, then $\text{elm}_{i_0}^-(E, \{l_{i,\bullet}\}_{i \in I})$ is also an ordinary parabolic bundle. When $a_2^{n_i} = 1$, description of the standard tableau of the transformation $l'_{i,\bullet}$ is as follows:

Corollary 41. *We assume that $\lambda^{(n)} = (1^{n-2}, 2^1)$. Let T_k be the standard tableau defined in (3.4) for $k = 1, 2, \dots, n-1$. Let $l_{i,\bullet}$ be a generic refined parabolic structure with the standard tableau T_k . Let $l'_{i,\bullet}$ be the transformation (5.2) of $l_{i,\bullet}$ and T' be the standard tableau of $l'_{i,\bullet}$. Then $T' = T_{n-k}$.*

Proof. By direct computation, we may check that

$$a_1^{k'} = \begin{cases} k' - 2 & \text{when } n - k + 1 \leq k' \leq n \\ k' & \text{when } 1 \leq k' \leq n - k \end{cases} \quad \text{and} \quad a_2^{k'} = \begin{cases} 1 & \text{when } n - k + 1 \leq k' \leq n \\ 0 & \text{when } 1 \leq k' \leq n - k \end{cases}$$

and

$$b_{k'} = \begin{cases} a_1^{k'} & \text{when } k' \neq n - k \\ a_1^{k'} - 1 & \text{when } k' = n - k. \end{cases}$$

Then we have the following equalities

$$X_1^b(k') = \begin{cases} n - k' & \text{when } n - k \leq k' \leq n - 1 \\ n - k' - 1 & \text{when } 1 \leq k' \leq n - k - 1 \end{cases} \quad \text{and} \quad X_2^b(k') = \begin{cases} 0 & \text{when } n - k \leq k' \leq n - 1 \\ 1 & \text{when } 1 \leq k' \leq n - k - 1. \end{cases}$$

By Proposition 40, we have the claim. \square

Now, we discuss a relation between the elementary transformations and the stability condition.

Proposition 42. *We take $i_0 \in I$. Let $(E'_{i_0}, \mathbf{l}'_{i_0}) = \text{elm}_{n_{i_0}[t_{i_0}]}^-(E, \{l_{i, \bullet}\}_{i \in I})$. Let L be a line subbundle of E and L' be a induced line subbundle of E'_{i_0} via $E'_{i_0} \subset E$. We set*

$$w'_{i,k} = \begin{cases} 1 - w_{i, n_{i_0} - k + 1} & \text{for } i = i_0 \text{ and } k = 1, \dots, n_{i_0} \\ w_{i,k} & \text{for } i \neq i_0 \text{ and } k = 1, \dots, n_i. \end{cases}$$

Let $\text{Stab}_{\mathbf{w}'}(L')$ be the stability index of L' with respect to $(E'_{i_0}, \mathbf{l}'_{i_0})$ and the weight $\mathbf{w}' = \{w'_{i,k}\}$. Then

$$\text{Stab}_{\mathbf{w}}(L) = \text{Stab}_{\mathbf{w}'}(L').$$

Proof. Since we have a short exact sequence

$$0 \longrightarrow E^{(n_{i_0})} \longrightarrow E \longrightarrow E/E^{(n_{i_0})} \longrightarrow 0$$

and $\text{length}(E/E^{(n_{i_0})}) = n_{i_0}$, we have $\deg(E) = \deg(E'_{i_0}) + n_{i_0}$. We have a short exact sequence

$$0 \longrightarrow L' \longrightarrow L \longrightarrow E|_L/E^{(n_{i_0})}|_L \longrightarrow 0.$$

Since

$$\begin{aligned} \text{length}(E|_L/E^{(n_{i_0})}|_L) &= \text{length}(E|_L/E(-n_{i_0}[t_{i_0}])|_L) - \sum_{k=1}^{n_{i_0}} \text{length}(E^{(k)}|_L/E^{(k-1)}|_L) \\ &= n_{i_0} - \sum_{k=1}^{n_{i_0}} \text{length}(E^{(k)}|_L/E^{(k-1)}|_L), \end{aligned}$$

we have an equality:

$$\deg(L) = \deg(L') + n_{i_0} - \sum_{k=1}^{n_{i_0}} \text{length}(E^{(k)}|_L/E^{(k-1)}|_L).$$

Moreover, we have the following equalities

$$\begin{aligned} &\text{length}(E^{(k)}|_L/E^{(k-1)}|_L) + \text{length}((E'_{i_0})^{(n_{i_0}-k+1)}|_{L'}/(E'_{i_0})^{(n_{i_0}-k)}|_{L'}) \\ &= \text{length}(E^{(k)}|_L/E^{(k-1)}|_L) + \text{length}(E^{(k-1)}(-k[t_{i_0}])|_L/E^{(k)}(-(k+1)[t_{i_0}])|_L) \\ &= \text{length}(E^{(k)}|_L/E^{(k)}(-[0])|_L) = 1. \end{aligned}$$

Then we obtain that

$$\begin{aligned}
 \text{Stab}_{\mathbf{w}}(L) &= \deg(E) - 2 \deg(L) + \sum_{k=1}^{n_{i_0}} w_{i_0,k} (1 - 2 \cdot \text{length}(E^{(k)}|_L/E^{(k-1)}|_L)) + \sum_{i \in I \setminus \{i_0\}} \sum_k \epsilon_{i,k}(L) w_{i,k} \\
 &= \deg(E'_{i_0}) + n_{i_0} - 2 \left(\deg(L') + n_{i_0} - \sum_{k=1}^{n_{i_0}} \text{length}(E^{(k)}|_L/E^{(k-1)}|_L) \right) \\
 &\quad + \sum_{k=1}^{n_{i_0}} w_{i_0,k} \left(2 \cdot \text{length}((E'_{i_0})^{(n_{i_0}-k+1)}|_{L'}/(E'_{i_0})^{(n_{i_0}-k)}|_{L'}) - 1 \right) + \sum_{i \in I \setminus \{i_0\}} \sum_k \epsilon_{i,k}(L) w_{i,k} \\
 &= \deg(E'_{i_0}) - 2 \deg(L') + \sum_{k'=1}^{n_{i_0}} (1 - 2 \cdot \text{length}((E'_{i_0})^{(k')}|_{L'}/(E'_{i_0})^{(k'-1)}|_{L'})) \\
 &\quad + \sum_{k'=1}^{n_{i_0}} (-w_{i_0,n_{i_0}-k'+1}) (1 - 2 \cdot \text{length}((E'_{i_0})^{(k')}|_{L'}/(E'_{i_0})^{(k'-1)}|_{L'})) + \sum_{i \in I \setminus \{i_0\}} \sum_k \epsilon_{i,k}(L) w_{i,k} \\
 &= \text{Stab}_{\mathbf{w}'}(L').
 \end{aligned}$$

□

Corollary 43. *If $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame and undecomposable, then $\text{elm}_{n_{i_0}[t_{i_0}]}^-(E, \{l_{i,\bullet}\}_{i \in I})$ is also tame and undecomposable.*

Proof. Since $(E, \{l_{i,\bullet}\}_{i \in I})$ is tame and undecomposable, $(E, \{l_{i,\bullet}\}_{i \in I})$ is stable for some weights \mathbf{w} (Theorem 1). By Proposition 42, the elementary transformation $\text{elm}_{n_{i_0}[t_{i_0}]}^-(E, \{l_{i,\bullet}\}_{i \in I})$ is stable for some weights \mathbf{w}' . By Theorem 1, $\text{elm}_{n_{i_0}[t_{i_0}]}^-(E, \{l_{i,\bullet}\}_{i \in I})$ is also tame and undecomposable. □

6. GEOMETRY OF MODULI SPACES AND WEAK DEL PEZZO SURFACES

Let $\overline{\mathfrak{Bun}}(D, 1)$ be the moduli space of tame and undecomposable refined parabolic bundles on (\mathbb{P}^1, D) whose degree are one. For a weight \mathbf{w} , let $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ be the moduli space of \mathbf{w} -semistable refined parabolic bundles on (\mathbb{P}^1, D) whose degrees are one. This moduli space is constructed as a projective scheme [27, Corollary 5.1]. The open subset of \mathbf{w} -stable parabolic bundles is smooth [28]. For generic weights \mathbf{w} , we have that \mathbf{w} -semistable = \mathbf{w} -stable. So $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ is a smooth projective scheme. If $n = 5$, then the dimension of $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ is two.

A free $\mathcal{O}_{n_i[t_i]}$ -submodule l_i of $E|_{n_i[t_i]}$ induces a one dimensional subspace l_i^{red} of $E|_{t_i}$.

Definition 44. *Assume that $E = \mathcal{O} \oplus \mathcal{O}(1)$. Let l_i be a free $\mathcal{O}_{n_i[t_i]}$ -submodule of $E|_{n_i[t_i]}$. We say l_i is in general position if $l_i^{\text{red}} \notin \mathcal{O}(1) \subset \mathbb{P}(E)$ for each for $i \in I$.*

We set $\tilde{I} = \{(i, k) \mid i \in I, k \in \{1, 2, \dots, n_i\}\}$.

Proposition 45. *We set $w_{i,k} = w$ (for any $(i, k) \in \tilde{I}$) with $\frac{1}{n} < w < \frac{1}{n-2}$. A refined parabolic bundle (E, \mathbf{l}) of degree 1 is \mathbf{w} -stable if, and only if, (E, \mathbf{l}) satisfies the following conditions:*

- $E \cong \mathcal{O} \oplus \mathcal{O}(1)$;
- (E, \mathbf{l}) is an undecomposable parabolic bundle (in particular, l_i is free for any $i \in I$);
- \mathbf{l} is in general position.

Proof. We assume that $E \cong \mathcal{O}(-d_0 + 1) \oplus \mathcal{O}(d_0)$ where $d_0 \geq 2$. We have the following inequality:

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(d_0)) \leq 1 - 2d_0 + nw < -3 + \frac{n}{n-2} < 0.$$

So this refined parabolic bundle is not stable. We assume that $E = \mathcal{O} \oplus \mathcal{O}(1)$ and there is a refined parabolic structure which intersects $\mathcal{O}(1)$. Then

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(1)) \leq -1 + (n-1)w - w = -1 + (n-2)w < 0.$$

So this refined parabolic bundle is not stable. This implies that \mathbf{l} is a (ordinary) parabolic structure and \mathbf{l} is in general position. We assume that $E = \mathcal{O} \oplus \mathcal{O}(1)$, the parabolic bundle is decomposable, and there is no parabolic structure which intersects $\mathcal{O}(1)$. Then there is a subbundle $\mathcal{O} \subset E$ such that any parabolic structures are contained in \mathcal{O} . Then

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}) \leq 1 - nw < 0.$$

Finally we have the following claim: If (E, \mathbf{l}) is \mathbf{w} -stable, then $E \cong \mathcal{O} \oplus \mathcal{O}(1)$, (E, \mathbf{l}) is an undecomposable parabolic bundle, and \mathbf{l} is in general position.

We assume that $(\mathcal{O} \oplus \mathcal{O}(1), \mathbf{l})$ is undecomposable and \mathbf{l} is in generic position. Since there is no refined parabolic structure which intersects $\mathcal{O}(1)$, we have

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(1)) = -1 + nw > 0.$$

Since $(\mathcal{O} \oplus \mathcal{O}(1), \mathbf{l})$ is undecomposable,

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}) \geq 1 + w(n-1) - w = 1 + w(n-2) > 0.$$

For any subbundle $\mathcal{O}(d_0)$ of $\mathcal{O} \oplus \mathcal{O}(1)$ where $d_0 \leq -1$, we have

$$\text{Stab}_{\mathbf{w}}(\mathcal{O}(d_0)) > 1 - 2d_0 - nw \geq 3 - nw > 3 - \frac{n}{n-2} > 0.$$

Then $(\mathcal{O} \oplus \mathcal{O}(1), \mathbf{l})$ is \mathbf{w} -stable. □

Proposition 46. *We set $w_{i,k} = w$ (for any $(i, k) \in \tilde{I}$) with $\frac{1}{n} < w < \frac{1}{n-2}$. We have an isomorphism $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \cong \mathbb{P}^{n-3}$.*

Proof. By Proposition 45, a \mathbf{w} -stable refined parabolic bundle (E, \mathbf{l}) of degree 1 is a nontrivial extension

$$0 \rightarrow (\mathcal{O}(1), \emptyset) \rightarrow (E, \mathbf{l}) \rightarrow (\mathcal{O}, \mathbf{l}) \rightarrow 0.$$

The obstruction to split the extension is measured by an element of

$$H^1(\mathbb{P}^1, \text{Hom}(\mathcal{O}(D), \mathcal{O}(1))) \cong H^1(\mathbb{P}^1, \mathcal{O}(1) \otimes \mathcal{O}(-D)) \cong H^0(\mathbb{P}^1, \mathcal{O}(-1) \otimes \Omega_{\mathbb{P}^1}^1(D))^*.$$

Any two nontrivial extensions define isomorphic parabolic bundles if, and only if, the corresponding cocycles are proportional. So the moduli space of extension is parametrized by $\mathbb{P}H^0(\mathbb{P}^1, \mathcal{O}(-1) \otimes \Omega_{\mathbb{P}^1}^1(D))^* \cong \mathbb{P}^{n-3}$. □

6.1. Moduli of stable refined parabolic bundles with democratic weights for $n = 5$ (Proof of Theorem B). We assume that $n = 5$. Let D be one of the effective divisors in the following list:

$$D_{2111} := 2[t_1] + [t_2] + [t_3] + [t_4], \quad D_{221} := 2[t_1] + 2[t_2] + [t_3], \quad D_{311} := 3[t_1] + [t_2] + [t_3],$$

$$D_{32} := 3[t_1] + 2[t_2], \quad D_{41} := 4[t_1] + [t_2], \quad D_5 := 5[t_1].$$

Our purpose here is to describe moduli spaces of stable refined parabolic bundles with democratic weights $\mathbf{w} = (w_1, \dots, w_\nu)$ for $n = \deg(D) = 5$. Here the democratic weights means that

$$\mathbf{w}_i = (w_{i,n_i}, \dots, w_{i,2}, w_{i,1}) = (w, \dots, w, w)$$

for $0 < w < 1$. If $\frac{1}{5} < w < \frac{1}{3}$, then this moduli space is \mathbb{P}^2 (see Proposition 46). We would like to describe the moduli spaces for another democratic weight w . If w is moved and w crosses walls, then some “special bundles” (which are stable for $\frac{1}{5} < w < \frac{1}{3}$) become unstable and some “special bundles” (which are unstable for $\frac{1}{5} < w < \frac{1}{3}$) become stable. Here, the “special bundles” are defined in the appendix below. To describe the moduli spaces, we consider such a wall-crossing.

Definition 47. We define special bundles and the types of special bundles as in the appendix below. There are 6 types: type A, type B, ..., type F. We say a tame and indecomposable refined parabolic bundle is generic if this refined parabolic bundle is not a special bundle.

We may check that the following claims:

- If a tame and indecomposable refined parabolic bundle is generic, then this refined parabolic bundle is stable for any w ($\frac{1}{5} < w < 1$);
- If $\frac{1}{5} < w < \frac{1}{3}$, then special bundles with type A, type B, and type C are stable. On the other hand, special bundles with type D and type F are unstable;
- If $\frac{1}{3} < w < \frac{3}{5}$, then special bundles with type C become unstable. Instead of it, special bundles with type D become stable;
- If $\frac{3}{5} < w < 1$, then special bundles with type A become unstable. Instead of it, special bundles with type F become stable;
- Special bundles with type E are unstable for any democratic weights \mathbf{w} .

These can be summarized as Table 1.

TABLE 1. Table of stability of special bundles

	type A	type B	type C	type D	type E	type F
$0 < w < \frac{1}{5}$	unstable	unstable	unstable	unstable	unstable	unstable
$\frac{1}{5} < w < \frac{1}{3}$	stable	stable	stable	unstable	unstable	unstable
$\frac{1}{3} < w < \frac{3}{5}$	stable	stable	unstable	stable	unstable	unstable
$\frac{3}{5} < w < 1$	unstable	stable	unstable	stable	unstable	stable

When $\frac{1}{5} < w < \frac{1}{3}$, $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \cong \mathbb{P}^2$. So there are loci on \mathbb{P}^2 corresponding to special bundles with type A, type B, and type C. The locus corresponding special bundles with type A is a conic on \mathbb{P}^2 . The loci corresponding special bundles with type B are lines on \mathbb{P}^2 . The loci corresponding special bundles with type C are intersections the conic and the lines. We may define an embedding

$$(6.1) \quad \Pi: \mathbb{P}^1 \longrightarrow \overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \cong \mathbb{P}^2$$

such that the image is the conic corresponding to special bundles with type A and the images of the points t_1, t_2, \dots, t_ν are points corresponding to special bundles with type C. We may describe the configuration of the loci of special bundles of type A, type B, and type C as follows.

- When $D = D_{2111}$, the configuration of the loci of special bundles is as follows: $P_{[t_i]} = \Pi(t_i)$ ($i = 1, 2, 3, 4$), the line $L_{2[t_1]}$ is tangent to the conic C at $P_{[t_1]}$, and the line $L_{[t_i],[t_j]}$ is intersect to the conic C at $P_{[t_i]}$ and $P_{[t_j]}$ for $i, j \in \{1, 2, 3, 4\}$ with $i \neq j$.
- When $D = D_{221}$, the configuration of the loci of special bundles is as follows: $P_{[t_i]} = \Pi(t_i)$ ($i = 1, 2, 3$), the line $L_{2[t_{i'}]}$ is tangent to the conic C at $P_{[t_{i'}]}$ ($i' = 1, 2$), and the line $L_{[t_i],[t_j]}$ is intersect to the conic C at $P_{[t_i]}$ and $P_{[t_j]}$ for $i, j \in \{1, 2, 3\}$ with $i \neq j$.
- When $D = D_{311}$, the configuration of the loci of special bundles is as follows: $P_{[t_i]} = \Pi(t_i)$ ($i = 1, 2, 3$), the line $L_{2[t_1]}$ is tangent to the conic C at $P_{[t_1]}$, and the line $L_{[t_i],[t_j]}$ is intersect to the conic C at $P_{[t_i]}$ and $P_{[t_j]}$ for $i, j \in \{1, 2, 3\}$ with $i \neq j$.
- When $D = D_{32}$, the configuration of the loci of special bundles is as follows: $P_{[t_i]} = \Pi(t_i)$ ($i = 1, 2$), the line $L_{2[t_i]}$ is tangent to the conic C at $P_{[t_i]}$ ($i = 1, 2$), and the line $L_{[t_1],[t_2]}$ is intersect to the conic C at $P_{[t_1]}$ and $P_{[t_2]}$.
- When $D = D_{41}$, the configuration of the loci of special bundles is as follows: $P_{[t_i]} = \Pi(t_i)$ ($i = 1, 2$), the line $L_{2[t_1]}$ is tangent to the conic C at $P_{[t_1]}$, and the line $L_{[t_1],[t_2]}$ is intersect to the conic C at $P_{[t_1]}$ and $P_{[t_2]}$.
- When $D = D_5$, the configuration of the loci of special bundles is as follows: $P_{[t_1]} = \Pi(t_1)$, the line $L_{2[t_1]}$ is tangent to the conic C at $P_{[t_1]}$.

Now, we will describe the moduli space $\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1)$ with $\frac{1}{3} < w < \frac{3}{5}$. Let \mathbf{w}' be democratic weights with $\frac{1}{5} < w' < \frac{1}{3}$. We identify $\overline{\mathfrak{Bun}}_{\mathbf{w}'}(D, 1)$ with \mathbb{P}^2 . We consider the sequence of blowing ups

$$X_5(D) \xrightarrow{\pi_5} X_4(D) \xrightarrow{\pi_4} X_3(D) \xrightarrow{\pi_3} X_2(D) \xrightarrow{\pi_2} X_1(D) \xrightarrow{\pi_1} \mathbb{P}^2.$$

of $\overline{\mathfrak{Bun}}_{\mathbf{w}'}(D, 1)$ at points which have the following position:

- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}, P_{[t_2]}, P_{[t_3]},$ and $P_{[t_4]}$ when $D = D_{2111}$;
- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}, P_{[t_2]}^{(0)} \succ_C P_{[t_2]},$ and $P_{[t_3]}$ when $D = D_{221}$;
- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}^{(1)} \succ_C P_{[t_1]}, P_{[t_2]},$ and $P_{[t_3]}$ when $D = D_{311}$;
- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}^{(1)} \succ_C P_{[t_1]},$ and $P_{[t_2]}^{(0)} \succ_C P_{[t_2]}$ when $D = D_{32}$;
- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}^{(1)} \succ_C P_{[t_1]}^{(2)} \succ_C P_{[t_1]},$ and $P_{[t_2]}$ when $D = D_{41}$;
- $P_{[t_1]}^{(0)} \succ_C P_{[t_1]}^{(1)} \succ_C P_{[t_1]}^{(2)} \succ_C P_{[t_1]}^{(3)} \succ_C P_{[t_1]}$ when $D = D_5$.

Here $P' \succ_C P$ means P' is the intersection of the exceptional divisor of the blowing up at P and the strict transformation of the conic C . Here P is a point on C or a strict transformation of C . We denote by $\text{wdP}_X^{(n)}$ the weak del Pezzo surface of degree n whose configuration of (-2) -curves is X . We have that

$$\begin{aligned} X_5(D_{2111}) &= \text{wdP}_{A_1}^{(4)}, & X_5(D_{221}) &= \text{wdP}_{2A_1}^{(4)}, & X_5(D_{311}) &= \text{wdP}_{A_2}^{(4)}, \\ X_5(D_{32}) &= \text{wdP}_{A_2+A_1}^{(4)}, & X_5(D_{41}) &= \text{wdP}_{A_3}^{(4)}, & X_5(D_5) &= \text{wdP}_{A_4}^{(4)}, \end{aligned}$$

and

$$\overline{\mathfrak{Bun}}_{\mathbf{w}}(D, 1) \cong X_5(D), \quad \frac{1}{3} < w < \frac{3}{5}.$$

Let $E_{[t_i]}^{(k)}$ ($k = 0, 1, 2, \dots, n_i - 1$) be the locus of special bundles with type D on the moduli space $\overline{\mathfrak{Bun}}_w(D, 1)$. By using Corollary 26, $E_{[t_i]}^{(k)}$ ($k = 1, 2, \dots, n_i - 1$) compose a chain of $(n_i - 1)$ projective lines. We may check that $E_{[t_i]}^{(k)}$ is the exceptional divisor of the blowing up at $P_{[t_i]}^{(k)}$ ($k = 0, 1, 2, \dots, n_i - 1$). Here we set $P_{[t_i]}^{(n_i-1)} = P_{[t_i]}$.

Now, we will describe the moduli space $\overline{\mathfrak{Bun}}_w(D, 1)$ with $\frac{3}{5} < w < 1$. We may check that the strict transformation of the conic C on $X_5(D)$ is a (-1) -curve. We contract this (-1) -curve on $X_5(D)$:

$$\pi'_5: X_5(D) \longrightarrow X'_5(D).$$

Then we have weak del Pezzo surfaces of degree 5:

$$\begin{aligned} X'_5(D_{2111}) &= \text{wdP}_{A_1}^{(5)}, & X'_5(D_{221}) &= \text{wdP}_{2A_1}^{(5)}, & X'_5(D_{311}) &= \text{wdP}_{A_2}^{(5)}, \\ X'_5(D_{32}) &= \text{wdP}_{A_2+A_1}^{(5)}, & X'_5(D_{41}) &= \text{wdP}_{A_3}^{(5)}, & X'_5(D_5) &= \text{wdP}_{A_4}^{(5)}. \end{aligned}$$

We have that

$$\overline{\mathfrak{Bun}}_w(D, 1) \cong X'_5(D), \quad \frac{3}{5} < w < 1.$$

Let P_C be the locus of special bundles with type F on the moduli space $\overline{\mathfrak{Bun}}_w(D, 1)$. We may check that $P_C = \pi'_5(C)$. These can be summarized as Table 2.

TABLE 2. Table of moduli spaces $\overline{\mathfrak{Bun}}_w(D, 1)$

	$D = D_{2111}$	$D = D_{221}$	$D = D_{311}$	$D = D_{32}$	$D = D_{41}$	$D = D_5$
$0 < w < \frac{1}{5}$	\emptyset	\emptyset	\emptyset	\emptyset	\emptyset	\emptyset
$\frac{1}{5} < w < \frac{1}{3}$	\mathbb{P}^2	\mathbb{P}^2	\mathbb{P}^2	\mathbb{P}^2	\mathbb{P}^2	\mathbb{P}^2
$\frac{1}{3} < w < \frac{3}{5}$	$\text{wdP}_{A_1}^{(4)}$	$\text{wdP}_{2A_1}^{(4)}$	$\text{wdP}_{A_2}^{(4)}$	$\text{wdP}_{A_1+A_2}^{(4)}$	$\text{wdP}_{A_3}^{(4)}$	$\text{wdP}_{A_4}^{(4)}$
$\frac{3}{5} < w < 1$	$\text{wdP}_{A_1}^{(5)}$	$\text{wdP}_{2A_1}^{(5)}$	$\text{wdP}_{A_2}^{(5)}$	$\text{wdP}_{A_1+A_2}^{(5)}$	$\text{wdP}_{A_3}^{(5)}$	$\text{wdP}_{A_4}^{(5)}$

Now we check the assertion (ii) in Theorem B. Let $\Pi_{-2}(D)$ be the effective divisors consisting of all (-2) -curves on $\overline{\mathfrak{Bun}}_w(D, 1)$ with $\frac{1}{3} < w < \frac{3}{5}$. We have that

$$\Pi_{-2}(D) = \sum_{\{i \in I | n_i > 1\}} (E_{[t_i]}^{(1)} + E_{[t_i]}^{(2)} + \dots + E_{[t_i]}^{(n_i-1)}).$$

By the list in the appendix (Section 7), we may check the following claim: If $(E, \{l_{i,\bullet}\}_i)$ be a refined parabolic bundle corresponding to a point on $\Pi_{-2}(D)$, then there exists i ($1 \leq i \leq \nu$) such that l_{i,n_i} is not free, that is, $l_{i,\bullet}$ is not a parabolic structure for this i .

Next we check the assertion (iii) in Theorem B. Let $(E, \{l_{i,\bullet}\}_i)$ be a refined parabolic bundle corresponding to a point on $\overline{\mathfrak{Bun}}_w(D, 1)$ with $\frac{1}{5} < w < \frac{1}{3}$. By Proposition 45, $(E, \{l_{i,\bullet}\}_i)$ is an ordinary parabolic bundle. By Lemma 12 (or Proposition 34), we have that $(E, \{l_{i,n_i}\}_i)$ is a simple parabolic bundle. Let $\Pi_{-2}(D)'$ be the effective divisor on $\overline{\mathfrak{Bun}}_w(D, 1)$ with $\frac{3}{5} < w < 1$ induced by $\Pi_{-2}(D)$. Let $(E, \{l_{i,\bullet}\}_i)$ be a refined parabolic bundle corresponding to a point on $\overline{\mathfrak{Bun}}_w(D, 1) \setminus \Pi_{-2}(D)$ with $\frac{1}{3} < w < \frac{3}{5}$, or $\overline{\mathfrak{Bun}}_w(D, 1) \setminus \Pi_{-2}(D)$ with $\frac{3}{5} < w < 1$. By the argument of the effect on the moduli space of varying the weight and the list in the appendix (Section 7), we may check that $(E, \{l_{i,\bullet}\}_i)$ is an ordinary parabolic bundle. Moreover, by Lemma 12 (or Proposition 34), we have that $(E, \{l_{i,n_i}\}_i)$ is a simple parabolic bundle. Finally, we obtain the statements in Theorem B.

6.2. Automorphisms of the moduli spaces for the democratic weight with $w = \frac{1}{2}$. Now we introduce some automorphisms of the moduli space $\overline{\mathfrak{Bun}}_w(D, 1)$ for the democratic weight with $w = \frac{1}{2}$. By Proposition 42, we have the following claim: If a refined parabolic bundle $(E, \mathbf{1})$ is stable for democratic weights for w ($0 < w < 1$), then the elementary transformation of $(E, \mathbf{1})$ at t_{i_0} is stable for the weight $\mathbf{w}' = (\mathbf{w}'_i)_{i \in I}$, where

$$\mathbf{w}'_i = \begin{cases} (1 - w, \dots, 1 - w, 1 - w) & \text{when } i = i_0 \\ (w, \dots, w, w) & \text{when } i \neq i_0. \end{cases}$$

So we have an isomorphism between moduli spaces:

$$\text{elm}_{n_{i_0}[t_{i_0}]}^- : \overline{\mathfrak{Bun}}_w(D, 1) \longrightarrow \overline{\mathfrak{Bun}}_{\mathbf{w}'}(D, 1 - n_{i_0}).$$

Note that $\mathbf{w}'_i = \mathbf{w}_i$ when $w = \frac{1}{2}$. If $w = \frac{1}{2}$ and n_{i_0} is even, we set

$$\text{elm}_{n_{i_0}[t_{i_0}]} := \mathcal{O}((n_{i_0}/2) \cdot t_{i_0}) \otimes \text{elm}_{n_{i_0}[t_{i_0}]}^-.$$

Then $\text{elm}_{n_{i_0}[t_{i_0}]}$ induces an automorphism of $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D, 1)$. By Corollary 41, we have the claim that the image of (-2) -curves under the automorphism $\mathcal{O}((n_{i_0}/2) \cdot t_{i_0}) \otimes \text{elm}_{n_{i_0}[t_{i_0}]}^-$ (where $n_{i_0} \geq 2$ and n_{i_0} is even) is the following:

$$\mathcal{O}((n_{i_0}/2) \cdot t_{i_0}) \otimes \text{elm}_{n_{i_0}[t_{i_0}]}^- : E_{[t_{i_0}]}^{(k)} \longrightarrow E_{[t_{i_0}]}^{(n_{i_0}-k)}$$

for $k = 1, 2, \dots, n_{i_0} - 1$.

We take $i_1 \in I \setminus \{i_0\}$ so that $n_{i_0} + n_{i_1}$ is even. We set


$$\text{elm}_{n_{i_0}[t_{i_0}] + n_{i_1}[t_{i_1}]} := \mathcal{O}\left(\frac{n_{i_0} + n_{i_1}}{2}\right) \otimes \left(\text{elm}_{n_{i_0}[t_{i_0}]}^- \circ \text{elm}_{n_{i_1}[t_{i_1}]}^-\right),$$

which is an automorphism of $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D, 1)$. Since the automorphisms $\text{elm}_{n_{i_0}[t_{i_0}]}$ and $\text{elm}_{n_{i_1}[t_{i_1}]}$ are commutative, the automorphism $\text{elm}_{n_{i_0}[t_{i_0}] + n_{i_1}[t_{i_1}]}$ is independent of the order of $n_{i_0}[t_{i_0}]$ and $n_{i_1}[t_{i_1}]$.

6.3. Geometry of the weak del Pezzo surfaces from the modular point of view. Now we impose that $n = 5$ and $w_{i,1} = w_{i,2} = \dots = w_{i,n_i} = \frac{1}{2}$ for any $i \in I$. Then the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D, 1)$ is a weak del Pezzo surface $\text{wdP}_X^{(4)}$. Here X means the configuration of (-2) -curves on the weak del Pezzo surface. The purposes of this section is to recover the geometry of the weak del Pezzo surfaces of degree 4 from the modular point of view. First we will describe the graph of negative curves on the surface $\text{wdP}_X^{(4)}$ for each D . Second we will reconstruct all automorphisms of the surface $\text{wdP}_X^{(4)}$ via elementary transformations of refined parabolic bundles and automorphisms of the base curve with D .

Now we set $D = D_{2111}$. The left-hand-side of Figure 1 is the projective plane \mathbb{P}^2 with the locus of special bundles. Here the projective plane \mathbb{P}^2 is identified with the moduli space $\overline{\mathfrak{Bun}}_w(D_{2111}, 1)$ with $\frac{1}{5} < w < \frac{1}{3}$. The right-hand-side of Figure 1 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{A_1}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{2111}, 1)$. Remark that some lines are repeated in order to see all intersections. Below, the curve C is identified with the base curve of the parabolic bundle for simplicity by (6.1). Note that the one thick line is (-2) -curves.

In the previous section, we had seen that some elementary transformations of refined parabolic bundles induce automorphisms on the moduli space $\overline{\mathfrak{Bun}}_w(D_{2111}, 1) \cong \text{wdP}_{A_1}^{(4)}$. Now we reconstruct


 FIGURE 1. \mathbb{P}^2 and $\text{wdP}_{A_1}^{(4)}$

all automorphisms of the weak del Pezzo surface $\text{wdP}_{A_1}^{(4)}$ via these elementary transformations. Let $\text{Aut}(C, D_{2111})$ be the automorphism of C sending D_{2111} to itself. Remark that any automorphisms in $\text{Aut}(C, D_{2111})$ fix the point t_1 . If the points t_1, t_2, t_3 , and t_4 have a generic cross-ratio, then the non-trivial automorphisms in $\text{Aut}(C, [t_1] + [t_2] + [t_3] + [t_4])$ act on the 4 points by double-transposition: That is,

$$(t_1 \leftrightarrow t_2, t_3 \leftrightarrow t_4), \quad (t_1 \leftrightarrow t_3, t_2 \leftrightarrow t_4), \quad \text{and} \quad (t_1 \leftrightarrow t_4, t_2 \leftrightarrow t_3).$$

So $\text{Aut}(C, D_{2111}) = \langle \text{id} \rangle$.

Proposition 48. *If $\text{Aut}(C, D_{2111}) = \langle \text{id} \rangle$, then the automorphism group of the weak del Pezzo surface is $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/2$, generated by elementary transformations:*

$$\text{Aut} \left(\text{wdP}_{A_1}^{(4)} \right) = \langle \text{elm}_{[t_2]+[t_3]}, \text{elm}_{[t_3]+[t_4]}, \text{elm}_{2[t_1]} \rangle.$$

Proof. These three elementary transformations act as follows on the negative curves:

- all of them stabilize the unique (-2) -curve $E_{[t_1]}^{(1)}$ (and so does any automorphism), and therefore permute the 4 intersecting (-1) -curves $E_{[t_1]}^{(0)}, L_{[t_1], [t_2]}, L_{[t_1], [t_3]}, L_{[t_1], [t_4]}$;
- the action on these 4 curves is by double-transposition, since the crossratio of their intersection point with the (-2) -curve $E_{[t_1]}^{(1)}$ should be preserved and is generic (coinciding with the cross-ratio of $(C, [t_1] + [t_2] + [t_3] + [t_4])$);
- $\text{elm}_{[t_2]+[t_3]}$ permutes $L_{[t_1], [t_2]}$ with $L_{[t_1], [t_3]}$;
- $\text{elm}_{[t_3]+[t_4]}$ permutes $L_{[t_1], [t_3]}$ with $L_{[t_1], [t_4]}$;
- $\text{elm}_{2[t_1]}$ stabilizes all 4 curves $E_{[t_1]}^{(0)}, L_{[t_1], [t_2]}, L_{[t_1], [t_3]}, L_{[t_1], [t_4]}$, but permutes C and $L_{2[t_1]}$.

One easily check these properties by studying the action of elementary transformations on parabolic structure and special line bundles. One also deduce that the 3 elementary transformations (which commute) indeed generate a order 8 group of automorphisms isomorphic to $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/2$.

It remains to check that any automorphism ϕ belongs to that 8 order group. One can assume, after composition by an element of the 4 order group $\langle \text{elm}_{[t_2]+[t_3]}, \text{elm}_{[t_3]+[t_4]} \rangle$, that ϕ stabilizes all 4

curves $E_{[t_1]}^{(0)}$, $L_{[t_1],[t_2]}$, $L_{[t_1],[t_3]}$, $L_{[t_1],[t_4]}$. Since $E_{[t_1]}^{(0)}$ intersects only 3 negative curves, namely C and $L_{2[t_1]}$ and the (-2) -curve $E_{[t_1]}^{(1)}$, it has to permute the two first ones. By composing with $\text{elm}_{2[t_1]}$, we can assume that ϕ also stabilizes each of the two curves C and $L_{2[t_1]}$. Then, the intersection combinatorics shows that ϕ eventually stabilizes all negative curves. Therefore, we can blow-down back to \mathbb{P}^2 and get a linear automorphism $\bar{\phi}$ which fixes the 4 points supporting the divisor on the conic. There exists an one-to-one correspondence between linear automorphisms on \mathbb{P}^2 which preserve the conic and automorphisms on the conic. Therefore $\bar{\phi}$ is the identity, ϕ as well, obviously belonging to the 8 order group. \square

If D has special configuration, then we might have non trivial automorphism $\varphi \in \text{Aut}(C, D_{2111})$. For the non trivial automorphism φ , we can pull-back a refined parabolic bundle by φ . Since we impose that the weights are democratic, the stability condition of this pull-back is the same as the stability condition of the refined parabolic bundle. So, in that case, the automorphism φ induces an extra automorphism for the weak del Pezzo surface.

Now we set $D = D_{221}$. The left-hand-side of Figure 2 is the projective plane \mathbb{P}^2 with the locus of special bundles. The right-hand-side of Figure 2 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{2A_1}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{221}, 1)$. Note that C and $L_{[t_3]}$ intersect each other, although we don't see on the picture.




FIGURE 2. \mathbb{P}^2 and $\text{wdP}_{2A_1}^{(4)}$

We take the automorphism $\varphi \in \text{Aut}(C, D_{221})$ permuting t_1 and t_2 , fixing t_3 . Let φ^* be the automorphism of $\text{wdP}_{2A_1}^{(4)}$ induced by the pull-backs of refined parabolic bundles by φ .

Proposition 49. *The automorphism group of the weak del Pezzo surface is $\mathbb{Z}/2 \times (\mathbb{Z}/2 \times \mathbb{Z}/2)$, generated by:*

$$\text{Aut}\left(\text{wdP}_{2A_1}^{(4)}\right) = \langle \varphi^*, \text{elm}_{2[t_1]}, \text{elm}_{2[t_2]} \rangle.$$

Proof. It is similar to the previous one. One can check:

- any automorphism permutes the two (-2) -curves $E_{[t_1]}^{(1)}$ and $E_{[t_2]}^{(1)}$, and therefore stabilize the unique (-1) -curve intersecting them, namely $L_{[t_1],[t_2]}$;
- φ permutes non trivially $E_{[t_1]}^{(1)}$ and $E_{[t_2]}^{(1)}$;

- $\text{elm}_{2[t_1]}$ stabilizes $E_{[t_1]}^{(1)}$, $E_{[t_1]}^{(0)}$, $L_{[t_1],[t_3]}$, $E_{[t_2]}^{(1)}$ but permutes $E_{[t_2]}^{(0)}$ and $L_{[t_2],[t_3]}$ non trivially;
- in a similar way, $\text{elm}_{2[t_2]}$ stabilizes $E_{[t_1]}^{(1)}$, $E_{[t_2]}^{(1)}$, $E_{[t_2]}^{(0)}$ and $L_{[t_2],[t_3]}$, but permutes $E_{[t_1]}^{(0)}$ and $L_{[t_1],[t_3]}$ non trivially.

Given an automorphism ϕ , one can assume, after composition by an element of the 8 order group of the statement, that it stabilizes all 7 curves $E_{[t_j]}^{(i)}$, $L_{[t_i],[t_j]}$; but combinatorics imply that ϕ stabilizes all negative curves: its blow-down $\bar{\phi}$ on \mathbb{P}^2 has to preserve the conic and fix support of D_{221} . It is the identity. \square

Now we set $D = D_{311}$. The left-hand-side of Figure 3 is the projective plane \mathbb{P}^2 with the locus of special bundles. The right-hand-side of Figure 3 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{A_2}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{311}, 1)$. Note that the two thick lines are (-2) -curves.




FIGURE 3. \mathbb{P}^2 and $\text{wdP}_{A_2}^{(4)}$

We take the automorphism $\varphi \in \text{Aut}(C, D_{311})$ permuting t_2 and t_3 , fixing t_1 . Let φ^* be the automorphism of $\text{wdP}_{A_2}^{(4)}$ induced by the pull-backs of refined parabolic bundles by φ .

Proposition 50. *The automorphism group of the weak del Pezzo surface is $\mathbb{Z}/2 \times (\mathbb{Z}/2 \times \mathbb{Z}/2)$, generated by:*

$$\text{Aut}\left(\text{wdP}_{A_2}^{(4)}\right) = \langle \varphi^*, \text{elm}_{3[t_1]+[t_2]}, \text{elm}_{[t_2]+[t_3]} \rangle.$$

Proof. It is similar to the previous ones. One can check:

- any automorphism permutes the two (-2) -curves $E_{[t_1]}^{(2)}$ and $E_{[t_1]}^{(1)}$;
- $\text{elm}_{3[t_1]+[t_2]}$ permutes non trivially $E_{[t_1]}^{(2)}$ and $E_{[t_1]}^{(1)}$;
- φ stabilizes $E_{[t_1]}^{(2)}$, $E_{[t_1]}^{(1)}$, $E_{[t_1]}^{(0)}$ and $L_{2[t_1]}$, but permutes non trivially the two lines $L_{[t_1],[t_2]}$ and $L_{[t_1],[t_3]}$;
- $\text{elm}_{[t_2]+[t_3]}$ stabilizes $E_{[t_1]}^{(2)}$ and $E_{[t_1]}^{(1)}$, but permutes non trivially the two lines $E_{[t_1]}^{(0)}$ and $L_{2[t_1]}$.

Given an automorphism ϕ , one can assume, after composition by an element of the 8 order group of the statement, that it stabilizes all 6 curves $E_{[t_1]}^{(i)}$, $L_{[t_1],[t_j]}$ and $L_{2[t_1]}$; but combinatorics imply that

ϕ stabilizes all negative curves: its blow-down $\bar{\phi}$ on \mathbb{P}^2 has to preserve the conic and fix support of D_{311} . It is the identity. \square

Now we set $D = D_{32}$. The left-hand-side of Figure 4 is the projective plane \mathbb{P}^2 with the locus of special bundles. The right-hand-side of Figure 4 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{A_2+A_1}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{32}, 1)$. Note that the three thick lines are (-2) -curves.




FIGURE 4. \mathbb{P}^2 and $\text{wdP}_{A_2+A_1}^{(4)}$

Let $\text{Aut}(C, D_{32})$ be the group of automorphisms fixing t_1 and t_2 . We have an isomorphism $\text{Aut}(C, D_{32}) \cong \mathbb{C}^*$.

Proposition 51. *The automorphism group of the weak del Pezzo surface is $\mathbb{Z}/2 \times \mathbb{C}^*$, generated by:*

$$\text{Aut}\left(\text{wdP}_{A_1+A_2}^{(4)}\right) = \langle \text{elm}_{2[t_1]}, \text{Aut}(C, D_{32}) \rangle.$$

Proof. It is similar to the previous ones. One can check:

- any automorphism stabilizes $E_{[t_1]}^{(i)}$, $i = 1, 2$, $E_{[t_2]}^{(i)}$, $i = 0, 1$, and $L_{[t_1],[t_2]}$ by intersection combinatorics between (-2) and (-1) -curves;
- $\text{elm}_{2[t_2]}$ permutes $L_{2[t_1]}$ with $E_{[t_1]}^{(0)}$, and $L_{2[t_2]}$ with C .

Given an automorphism ϕ , one can assume, after composition by $\text{elm}_{2[t_2]}$ that is stabilizes all negative curves and blow-down as a linear automorphism of \mathbb{P}^2 stabilizing the conic C and fixing the two points $P_{[t_1]}$ and $P_{[t_2]}$. It is determined by its action on the conic; there is an element $\varphi \in \text{Aut}(C, D_{32})$ inducing the same action on the conic: ϕ coincides with the natural action of φ on refined parabolic bundles. So $\text{Aut}\left(\text{wdP}_{A_1+A_2}^{(4)}\right)$ is generated by $\text{elm}_{2[t_1]}$ and $\text{Aut}(C, D_{32})$.

Since $\varphi \in \text{Aut}(C, D_{32})$ fixes t_1 and t_2 , we may check that

$$\text{elm}_{2[t_1]}(\varphi^* E, \varphi^* \mathbf{1}) = \varphi^* (\text{elm}_{2[t_1]}(E, \mathbf{1}))$$

for any $(E, \mathbf{1}) \in \overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{32}, 1)$. That is, the automorphisms φ^* and $\text{elm}_{2[t_1]}$ commute. So the automorphism group of the weak del Pezzo surface is isomorphic to $\mathbb{Z}/2 \times \mathbb{C}^*$. \square

Now we set $D = D_{41}$. The left-hand-side of Figure 5 is the projective plane \mathbb{P}^2 with the locus of special bundles. The right-hand-side of Figure 5 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{A_3}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_{41}, 1)$. Note that the three thick lines are (-2) -curves.




FIGURE 5. \mathbb{P}^2 and $\text{wdP}_{A_3}^{(4)}$

Let $\text{Aut}(C, D_{41})$ be the group of automorphisms fixing t_1 and t_2 .

Proposition 52. *The automorphism group of the weak del Pezzo surface is $\mathbb{Z}/2 \times \mathbb{C}^*$, generated by:*

$$\text{Aut}\left(\text{wdP}_{A_3}^{(4)}\right) = \langle \text{elm}_{4[t_1]}, \text{Aut}(C, D_{41}) \rangle.$$

Proof. One can check:

- any automorphism stabilizes $E_{[t_1]}^{(2)}$ and $L_{2[t_1]}$;
- $\text{elm}_{4[t_1]}$ permutes $E_{[t_1]}^{(3)}$ with $E_{[t_1]}^{(1)}$.

Given an automorphism ϕ , one can assume, after composition by $\text{elm}_{4[t_1]}$ that is stabilizes all negative curves, and therefore blow-down as a linear automorphism of \mathbb{P}^2 stabilizing the conic C and fixing the two points $P_{[t_1]}$ and $P_{[t_2]}$. So $\text{Aut}\left(\text{wdP}_{A_3}^{(4)}\right)$ is generated by $\text{elm}_{4[t_1]}$ and $\text{Aut}(C, D_{41})$. Since $\varphi \in \text{Aut}(C, D_{41})$ fixes t_1 and t_2 , the automorphisms $\text{elm}_{4[t_1]}$ and φ^* commute. \square


Now we set $D = D_5$. The left-hand-side of Figure 6 is the projective plane \mathbb{P}^2 with the locus of special bundles. The right-hand-side of Figure 6 is the graph of negative curves on the weak del Pezzo surface obtained by blowing-up of the left-hand-side. This weak del Pezzo surface $\text{wdP}_{A_4}^{(4)}$ is just the moduli space $\overline{\mathfrak{Bun}}_{\frac{1}{2}}(D_5, 1)$. Note that the three four lines are (-2) -curves.

Let $\text{Aut}(C, D_5)$ be the group of automorphisms fixing t_1 .

Proposition 53. *The automorphism group of the weak del Pezzo surface is:*

$$\text{Aut}\left(\text{wdP}_{A_4}^{(4)}\right) = \text{Aut}(C, D_5).$$

Proof. By intersection combinatorics, any automorphism ϕ must stabilize all negative curves, and therefore blow-down as a linear automorphism of \mathbb{P}^2 stabilizing the conic C and fixing the point $P_{[t_1]}$. \square

FIGURE 6. \mathbb{P}^2 and $\text{wdP}_{A_4}^{(4)}$

7. APPENDIX: LIST OF SPECIAL BUNDLES

In this appendix, we will give a list of special bundles for $n = 5$. First, we give a list of standard tableaux of refined parabolic structures, which appear in the cases where $n = 5$. If the multiplicity of a point is 2, then the standard tableau of a refined parabolic structure at this point has two possibility:

$$T_I^{(2)} : (1, 1) \supset (1) \quad T_{II}^{(2)} : (2) \supset (1).$$

If the multiplicity of a point is 3, then the standard tableau of a refined parabolic structure at this point has three possibility:

$$T_I^{(3)} : (1, 1, 1) \supset T_I^{(2)} \quad T_{II}^{(3)} : (1, 2) \supset T_{II}^{(2)} \quad T_{III}^{(3)} : (1, 2) \supset T_I^{(2)}.$$

If the multiplicity of a point is 4, then the standard tableau of a refined parabolic structure at this point has six possibility:

$$\begin{aligned} T_I^{(4)} : (1, 1, 1, 1) \supset T_I^{(3)} & \quad T_{II}^{(4)} : (1, 1, 2) \supset T_{II}^{(3)} & \quad T_{III}^{(4)} : (1, 1, 2) \supset T_{III}^{(3)} \\ T_{IV}^{(4)} : (1, 1, 2) \supset T_I^{(3)} & \quad T_V^{(4)} : (2, 2) \supset T_{II}^{(3)} & \quad T_{VI}^{(4)} : (2, 2) \supset T_{III}^{(3)}. \end{aligned}$$

If the multiplicity of a point is 5, then the standard tableau of a refined parabolic structure at this point has ten possibility:

$$\begin{aligned} T_I^{(5)} : (1, 1, 1, 1, 1) \supset T_I^{(4)} & \quad T_{II}^{(5)} : (1, 1, 1, 2) \supset T_{II}^{(4)} & \quad T_{III}^{(5)} : (1, 1, 1, 2) \supset T_{III}^{(4)} \\ T_{IV}^{(5)} : (1, 1, 1, 2) \supset T_{IV}^{(4)} & \quad T_V^{(5)} : (1, 1, 1, 2) \supset T_I^{(4)} & \quad T_{VI}^{(5)} : (1, 2, 2) \supset T_V^{(4)} \\ T_{VII}^{(5)} : (1, 2, 2) \supset T_{VI}^{(4)} & \quad T_{VIII}^{(5)} : (1, 2, 2) \supset T_{II}^{(4)} & \quad T_{IX}^{(5)} : (1, 2, 2) \supset T_{III}^{(4)} \\ T_X^{(5)} : (1, 2, 2) \supset T_{IV}^{(4)}. & & \end{aligned}$$

For giving a list of special bundles for $n = 5$, now, we discuss some condition for refined parabolic structures. For a subbundle $L \subset E$, we define an integer $m_{t_i, L}$ by

$$m_{t_i, L} = \sum_{k=1}^{n_i} \text{length}((l_{i,k} \cap L|_{n_i[t_i]}) / (l_{i,k-1} \cap L|_{n_i[t_i]})).$$

We assume that $(E, \mathbf{1})$ is tame and undecomposable and $\deg(E) = 1$. In particular, $(E, \mathbf{1})$ is admissible. So we have the inequality

$$\sum_{i \in I} m_{t_i, L} \leq 4 - 2 \deg(L)$$

for any line subbundles L such that $\deg(E) \leq 2 \deg(L)$. So we have the following facts:

- $(E, \mathbf{1})$ with $E \cong \mathcal{O}(-k) \oplus \mathcal{O}(1+k)$ (where $k \geq 2$) disappears;
- if $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $L = \mathcal{O}(1)$, then $\sum_{i \in I} m_{t_i, \mathcal{O}(1)} \leq 2$;
- if $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $L = \mathcal{O}(2)$, then $m_{t_1, \mathcal{O}(2)} = \dots = m_{t_n, \mathcal{O}(2)} = 0$.

If l_{i, n_i} is not free, then $m_{t_i, L} \geq 1$ for any line subbundles L such that $\deg(E) \leq 2 \deg(L)$. So if $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$, then l_{i, n_i} are free for any i .

Now we consider the cases where $E = \mathcal{O} \oplus \mathcal{O}(1)$. We impose $m_{t_i, \mathcal{O}(1)} \leq 2$ for $i \in I$. We set $\epsilon_{i, k} := \epsilon_{i, k}(\mathcal{O}(1))$ for $i \in I$ and $k \in \{1, 2, \dots, n_i\}$. When we will check the tameness of refined parabolic bundles, we will calculate $N_i(\mathcal{O}(1))$, which is defined in (4.2). Now the tameness means that

$$2 \leq \sum_{i \in I_{\mathcal{O}(1)}^+} N_i(\mathcal{O}(1)).$$

To calculate $N_i(\mathcal{O}(1))$, we consider the tuple $(\epsilon_{i, n_i}, \dots, \epsilon_{i, 1})$. The tuple $(\epsilon_{i, n_i}, \dots, \epsilon_{i, 1})$ is generically determined by the pair $(T_{l_{i, \bullet}}, m_{t_i, \mathcal{O}(1)})$. Here $T_{l_{i, \bullet}}$ is the standard tableau of the refined parabolic structure $l_{i, \bullet}$ at t_i . Now we will describe the correspondence between $(\epsilon_{i, n_i}, \dots, \epsilon_{i, 1})$ and $(T_{l_{i, \bullet}}, m_{t_i, \mathcal{O}(1)})$ in Table 3, Table 4, Table 5, and Table 6.

TABLE 3. The correspondence with $n_i = 2$

$(T_{l_{i, \bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i, 2}, \epsilon_{i, 1})$	$N_i(\mathcal{O}(1))$
$(T_I^{(2)}, 1)$	$(+, -)$	0
$(T_{II}^{(2)}, 1)$	$(-, +)$	1
$(T_{III}^{(2)}, 2)$	$(-, -)$	-2

TABLE 4. The correspondence with $n_i = 3$

$(T_{l_{i, \bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i, 3}, \epsilon_{i, 2}, \epsilon_{i, 1})$	$N_i(\mathcal{O}(1))$	$(T_{l_{i, \bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i, 3}, \epsilon_{i, 2}, \epsilon_{i, 1})$	$N_i(\mathcal{O}(1))$
$(T_I^{(3)}, 1)$	$(+, +, -)$	1	$(T_I^{(3)}, 2)$	$(+, -, -)$	-1
$(T_{II}^{(3)}, 1)$	$(+, -, +)$	1	$(T_{II}^{(3)}, 2)$	$(-, -, +)$	1
$(T_{III}^{(3)}, 1)$	$(-, +, +)$	2	$(T_{III}^{(3)}, 2)$	$(-, +, -)$	0

TABLE 5. The correspondence with $n_i = 4$

$(T_{l_{i,\bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i,4}, \epsilon_{i,3}, \epsilon_{i,2}, \epsilon_{i,1})$	$N_i(\mathcal{O}(1))$	$(T_{l_{i,\bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i,4}, \epsilon_{i,3}, \epsilon_{i,2}, \epsilon_{i,1})$	$N_i(\mathcal{O}(1))$
$(T_I^{(4)}, 1)$	$(+, +, +, -)$	2	$(T_I^{(4)}, 2)$	$(+, +, -, -)$	0
$(T_{II}^{(4)}, 1)$	$(+, +, -, +)$	2	$(T_{II}^{(4)}, 2)$	$(+, -, -, +)$	1
$(T_{III}^{(4)}, 1)$	$(+, -, +, +)$	2	$(T_{III}^{(4)}, 2)$	$(+, -, +, -)$	0
$(T_{IV}^{(4)}, 1)$	$(-, +, +, +)$	3	$(T_{IV}^{(4)}, 2)$	$(-, +, +, -)$	1
$(T_V^{(4)}, 2)$	$(-, +, -, +)$	1	$(T_{VI}^{(4)}, 2)$	$(-, -, +, +)$	2

TABLE 6. The correspondence with $n_i = 5$

$(T_{l_{i,\bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i,5}, \epsilon_{i,4}, \epsilon_{i,3}, \epsilon_{i,2}, \epsilon_{i,1})$	$N_i(\mathcal{O}(1))$	$(T_{l_{i,\bullet}}, m_{t_i, \mathcal{O}(1)})$	$(\epsilon_{i,5}, \epsilon_{i,4}, \epsilon_{i,3}, \epsilon_{i,2}, \epsilon_{i,1})$	$N_i(\mathcal{O}(1))$
$(T_I^{(5)}, 1)$	$(+, +, +, +, -)$	3	$(T_I^{(5)}, 2)$	$(+, +, +, -, -)$	1
$(T_{II}^{(5)}, 1)$	$(+, +, +, -, +)$	3	$(T_{II}^{(5)}, 2)$	$(+, +, -, -, +)$	1
$(T_{III}^{(5)}, 1)$	$(+, +, -, +, +)$	3	$(T_{III}^{(5)}, 2)$	$(+, +, -, +, -)$	1
$(T_{IV}^{(5)}, 1)$	$(+, -, +, +, +)$	3	$(T_{IV}^{(5)}, 2)$	$(+, -, +, +, -)$	1
$(T_V^{(5)}, 1)$	$(-, +, +, +, +)$	4	$(T_V^{(5)}, 2)$	$(-, +, +, +, -)$	2
$(T_{VI}^{(5)}, 2)$	$(+, -, +, -, +)$	1	$(T_{VII}^{(5)}, 2)$	$(+, -, -, +, +)$	2
$(T_{VIII}^{(5)}, 2)$	$(-, +, +, -, +)$	2	$(T_{IX}^{(5)}, 2)$	$(-, +, -, +, +)$	2
$(T_X^{(5)}, 2)$	$(-, -, +, +, +)$	3			

7.1. **Case $D = D_{2111}$.** Now we describe special refined parabolic bundles which are indecomposable and tame when $D = D_{2111}$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = \cdots = m_{t_4, \mathcal{O}(1)} = 0$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(-1)} = 2, m_{t_2, \mathcal{O}(-1)} = m_{t_3, \mathcal{O}(-1)} = m_{t_4, \mathcal{O}(-1)} = 1$

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = \cdots = m_{t_4, \mathcal{O}(1)} = 0$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{[t_i], [t_j]}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = 2, m_{t_k, \mathcal{O}} = 1$ where $k \in \{2, 3, 4\} \setminus \{i, j\}$.
$L_{[t_1], [t_i]}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = m_{t_j, \mathcal{O}} = m_{t_k, \mathcal{O}} = 1$ where $\{j, k\} = \{2, 3, 4\} \setminus \{i\}$.
$L_{2[t_1]}$	$T_I^{(2)}$	$m_{t_2, \mathcal{O}} = m_{t_3, \mathcal{O}} = m_{t_4, \mathcal{O}} = 1$

for $i, j \in \{2, 3, 4\}$ with $i \neq j$.

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}, \quad P_{[t_i]} := C \cap \left(\bigcap_{j \in \{2, 3, 4\} \setminus \{i\}} L_{[t_i], [t_j]} \right) \quad i \in \{2, 3, 4\}.$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_i]}^{(0)}$	$T_I^{(2)}$	$m_{t_i, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 1$

for $i, j \in \{1, 2, 3, 4\}$.

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{[t_i], [t_j]}^{(0)}$	$T_I^{(2)}$	$m_{t_i, \mathcal{O}(1)} = m_{t_j, \mathcal{O}(1)} = 1$.
$Q_{2[t_1]}^{(0)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{[t_1], [t_{i'}]}^{(1)}$	$T_{II}^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_{i'}, \mathcal{O}(1)} = 1$.

for $i, j \in \{1, 2, 3, 4\}$ with $i \neq j$ and for $i' \in \{2, 3, 4\}$.

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{l_1, \bullet} : (1, 1) \supset (1)$.

7.2. **Case $D = D_{221}$.** Now we describe special refined parabolic bundles which are indecomposable and tame when $D = D_{221}$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = m_{t_3, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(-1)} = m_{t_2, \mathcal{O}(-1)} = 2, m_{t_3, \mathcal{O}(-1)} = 1$.

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = m_{t_3, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{[t_1], [t_2]}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = m_{t_2, \mathcal{O}} = m_{t_3, \mathcal{O}} = 1$.
$L_{[t_i], [t_3]}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_{i'}, \mathcal{O}} = 2, m_{t_3, \mathcal{O}} = 1$ where $i' \in \{1, 2\} \setminus \{i\}$.
$L_{2[t_i]}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_{i'}, \mathcal{O}} = 2, m_{t_3, \mathcal{O}} = 1$ where $i' \in \{1, 2\} \setminus \{i\}$.

for $i \in \{1, 2\}$.

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}, \quad P_{[t_2]} := C \cap L_{2[t_2]}, \quad P_{[t_3]} := C \cap L_{[t_1], [t_3]} \cap L_{[t_2], [t_3]}.$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_i]}^{(0)}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_i, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(2)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_2]}^{(1)}$	$T_I^{(2)}$	$T_{II}^{(2)}$	$m_{t_2, \mathcal{O}(1)} = 1$

for $i \in \{1, 2, 3\}$.

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_{1,\bullet}}$	$T_{l_{2,\bullet}}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{[t_i],[t_3]}^{(0)}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_i,\mathcal{O}(1)} = m_{t_3,\mathcal{O}(1)} = 1.$
$Q_{[t_{j'}],[t_2]}^{(1)}$	$T_I^{(2)}$	$T_{II}^{(2)}$	$m_{t_{j'},\mathcal{O}(1)} = m_{t_2,\mathcal{O}(1)} = 1.$
$Q_{[t_1],[t_{j'}]}^{(2)}$	$T_{II}^{(2)}$	$T_I^{(2)}$	$m_{t_1,\mathcal{O}(1)} = m_{t_{j'},\mathcal{O}(1)} = 1.$
$Q_{[t_1],[t_2]}^{(3)}$	$T_{II}^{(2)}$	$T_{II}^{(2)}$	$m_{t_1,\mathcal{O}(1)} = m_{t_2,\mathcal{O}(1)} = 1$
$Q_{2[t_i]}^{(0)}$	$T_I^{(2)}$	$T_I^{(2)}$	$m_{t_i,\mathcal{O}(1)} = 2.$

for $i \in \{1, 2\}$, and $i' \in \{1, 3\}$, $j' \in \{2, 3\}$.

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{l_{1,\bullet}} : (1, 1) \supset (1)$.

7.3. **Case $D = D_{311}$.** Now we describe special refined parabolic bundles which are indecomposable and tame when $D = D_{311}$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1,\mathcal{O}(1)} = m_{t_2,\mathcal{O}(1)} = m_{t_3,\mathcal{O}(1)} = 0$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(3)}$	$m_{t_1,\mathcal{O}(-1)} = 3, m_{t_2,\mathcal{O}(-1)} = m_{t_3,\mathcal{O}(-1)} = 1$

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1,\mathcal{O}(1)} = m_{t_2,\mathcal{O}(1)} = m_{t_3,\mathcal{O}(1)} = 0$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{[t_2],[t_3]}$	$T_I^{(3)}$	$m_{t_1,\mathcal{O}} = 3.$
$L_{[t_1],[t_i]}$	$T_I^{(3)}$	$m_{t_1,\mathcal{O}} = 2, m_{t_{i'},\mathcal{O}} = 1$ where $i' \in \{2, 3\} \setminus \{i\}$.
$L_{2[t_1]}$	$T_I^{(3)}$	$m_{t_1,\mathcal{O}} = m_{t_2,\mathcal{O}} = m_{t_3,\mathcal{O}} = 1$

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}, \quad P_{[t_2]} := C \cap L_{[t_1],[t_2]} \cap L_{[t_2],[t_3]}, \quad P_{[t_3]} := C \cap L_{[t_1],[t_3]} \cap L_{[t_2],[t_3]}.$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_i]}^{(0)}$	$T_I^{(3)}$	$m_{t_i,\mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = 1$
$E_{[t_1]}^{(2)}$	$T_{III}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = 1$

for $i \in \{1, 2, 3\}$.

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_{1,\bullet}}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{[t_i],[t_j]}^{(0)}$	$T_I^{(3)}$	$m_{t_i,\mathcal{O}(1)} = m_{t_j,\mathcal{O}(1)} = 1.$
$Q_{[t_1],[t_{j'}]}^{(0)}$	$T_{II}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = m_{t_{j'},\mathcal{O}(1)} = 1.$
$Q_{[t_1],[t_{j'}]}^{(0)}$	$T_{III}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = m_{t_{j'},\mathcal{O}(1)} = 1.$
$Q_{2[t_1]}^{(0)}$	$T_I^{(3)}$	$m_{t_1,\mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(1)}$	$T_{II}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(2)}$	$T_{III}^{(3)}$	$m_{t_1,\mathcal{O}(1)} = 2$

for $i, j \in \{1, 2, 3\}$ with $i \neq j$ and $j' \in \{2, 3\}$.

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{l_{1,\bullet}} : (1, 1, 1) \supset (1, 1) \supset (1)$.

7.4. **Case $D = D_{32}$.** Now we describe special refined parabolic bundles which are indecomposable and tame when $D = D_{32}$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(-1)} = 3, m_{t_2, \mathcal{O}(-1)} = 2$

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{2[t_2]}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = 3$.
$L_{[t_1], [t_2]}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = 2, m_{t_2, \mathcal{O}} = 1$.
$L_{2[t_1]}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}} = 1, m_{t_2, \mathcal{O}} = 2$.

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}, \quad P_{[t_2]} := C \cap L_{2[t_2]}.$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_i]}^{(0)}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_i, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_2]}^{(1)}$	$T_I^{(3)}$	$T_{II}^{(2)}$	$m_{t_2, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(2)}$	$T_{III}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 1$

for $i \in \{1, 2\}$.

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{[t_1], [t_2]}^{(0)}$	$T_{II}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$
$Q_{[t_1], [t_2]}^{(1)}$	$T_{III}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$
$Q_{[t_1], [t_2]}^{(2)}$	$T_I^{(3)}$	$T_{II}^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$
$Q_{[t_1], [t_2]}^{(3)}$	$T_{II}^{(3)}$	$T_{II}^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$
$Q_{[t_1], [t_2]}^{(4)}$	$T_{III}^{(3)}$	$T_{II}^{(2)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$

and

	$T_{l_1, \bullet}$	$T_{l_2, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{2[t_1]}^{(0)}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(1)}$	$T_{II}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(2)}$	$T_{III}^{(3)}$	$T_I^{(2)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_2]}^{(0)}$	$T_I^{(3)}$	$T_I^{(2)}$	$m_{t_2, \mathcal{O}(1)} = 2$
$Q_{2[t_2]}^{(1)}$	$T_I^{(3)}$	$T_{II}^{(2)}$	$m_{t_2, \mathcal{O}(1)} = 2$

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{l_1, \bullet} : (1, 1, 1) \supset (1, 1) \supset (1)$.

7.5. **Case $D = D_{41}$.** Now we describe special refined parabolic bundles which are indecomposable and tame when $D = D_{41}$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(4)}$	$m_{t_1, \mathcal{O}(-1)} = 4, m_{t_2, \mathcal{O}(-1)} = 1$

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{[t_1], [t_2]}$	$T_I^{(4)}$	$m_{t_1, \mathcal{O}} = 3$.
$L_{2[t_1]}$	$T_I^{(4)}$	$m_{t_1, \mathcal{O}} = 2, m_{t_2, \mathcal{O}} = 1$.

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}, \quad P_{[t_2]} := (C \cap L_{[t_1], [t_2]}) \setminus P_{[t_1]}.$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_1]}^{(0)}$	$T_I^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(2)}$	$T_{III}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(3)}$	$T_{IV}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 1$

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{[t_1], [t_2]}^{(0)}$	$T_I^{(4)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$.
$Q_{[t_1], [t_2]}^{(1)}$	$T_{II}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$.
$Q_{[t_1], [t_2]}^{(2)}$	$T_{III}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$.
$Q_{[t_1], [t_2]}^{(3)}$	$T_{IV}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = m_{t_2, \mathcal{O}(1)} = 1$.

and

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{2[t_1]}^{(0)}$	$T_{II}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(1)}$	$T_{IV}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(2)}$	$T_V^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(3)}$	$T_{VI}^{(4)}$	$m_{t_1, \mathcal{O}(1)} = 2$

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{l_1, \bullet} : (1, 1, 1) \supset (1, 1) \supset (1)$.

7.6. **Case $D = D_5$.** Now we describe special refined parabolic bundles which are undecomposable and tame when $D = D_5$.

[Type A]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O}(-1) \subset E$ as below:
C	$T_I^{(5)}$	$m_{t_1, \mathcal{O}(-1)} = 5$

[Type B]. $E = \mathcal{O} \oplus \mathcal{O}(1)$ and $m_{t_1, \mathcal{O}(1)} = 0$.

	$T_{l_1, \bullet}$	there exists a subbundle $\mathcal{O} \subset E$ as below:
$L_{2[t_1]}$	$T_I^{(5)}$	$m_{t_1, \mathcal{O}} = 3$.

[Type C].

$$P_{[t_1]} := C \cap L_{2[t_1]}$$

[Type D]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{I,\bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$E_{[t_1]}^{(0)}$	$T_I^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 1$
$E_{[t_1]}^{(1)}$	$T_{II}^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 1$
\vdots	\vdots	\vdots
$E_{[t_1]}^{(4)}$	$T_V^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 1$

[Type E]. $E = \mathcal{O} \oplus \mathcal{O}(1)$.

	$T_{I,\bullet}$	there exists a subbundle $\mathcal{O}(1) \subset E$ as below:
$Q_{2[t_1]}^{(0)}$	$T_V^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(1)}$	$T_{VII}^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(2)}$	$T_{VIII}^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(3)}$	$T_{IX}^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 2$
$Q_{2[t_1]}^{(4)}$	$T_X^{(5)}$	$m_{t_1, \mathcal{O}(1)} = 2$

[Type F]. $E = \mathcal{O}(-1) \oplus \mathcal{O}(2)$ and $T_{I,\bullet} : (1, 1, 1) \supset (1, 1) \supset (1)$.

REFERENCES

- [1] C. Araujo, T. Fassarella, I. Kaur, A. Massarenti, *On automorphisms of moduli spaces of parabolic vector bundles*. Int. Math. Res. Not. IMRN 2021, no. **3**, 2261–2283.
- [2] D. Arinkin, S. Lysenko. *Isomorphisms between moduli spaces of $SL(2)$ -bundles with connections on $\mathbb{P}^1 \setminus \{x_1, \dots, x_4\}$* . Math. Res. Lett., **4** (2-3):181–190, 1997.
- [3] D. Arinkin, S. Lysenko, *On the moduli of $SL(2)$ -bundles with connections on $\mathbb{P}^1 \setminus \{x_1, \dots, x_4\}$* , Internat. Math. Res. Notices (1997), no. **19**, 983–999.
- [4] D. Arinkin, R. Fedorov, *An example of the Langlands correspondance for irregular rank two connections on \mathbb{P}^1* . Advances in Math. **230** (2012) 1078–1123.
- [5] S. Bauer, *Parabolic bundles, elliptic surfaces and $SU(2)$ -representation spaces of genus zero Fuchsian groups*. Math. Ann. **290** (1991), no. 3, 509–526.
- [6] U. N. Bhosle. *Parabolic vector bundles on curves*. Ark. Mat., **27** (1): 15–22, 1989.
- [7] U. N. Bhosle. *Generalised parabolic bundles and applications to torsionfree sheaves on nodal curves*. Ark. Mat. **30** (1992), no. 2, 187–215.
- [8] U. N. Bhosle. *Generalized parabolic bundles and applications. II*. Proc. Indian Acad. Sci. Math. Sci. **106** (1996), no. 4, 403–420.
- [9] U. N. Bhosle, I. Biswas, *Moduli space of parabolic vector bundles on a curve*. Beitr. Algebra Geom. **53** (2012), no. 2, 437–449.
- [10] I. Biswas, Y. I. Holla, C. Kumar. *On moduli spaces of parabolic vector bundles of rank 2 over $\mathbb{C}\mathbb{P}^1$* . Michigan Math. J., **59** (2): 467–475, 2010.
- [11] H. U. Boden, Y. Hu. *Variations of moduli of parabolic bundles*. Math. Ann., **301**(3): 539–559, 1995.
- [12] I. Dolgachev, *Classical algebraic geometry. A modern view*. Cambridge University Press, Cambridge, 2012. xii+639 pp. ISBN: 978-1-107-01765-8
- [13] R. Donagi, T. Pantev *Parabolic Hecke eigensheaves* arXiv:1910.02357.
- [14] H.-B. Moon, S.-B. Yoo, *Birational geometry of the moduli space of rank 2 parabolic vector bundles on a rational curve*. Int. Math. Res. Not. IMRN 2016, no. **3**, 827–859.
- [15] M.-A. Inaba, *Moduli of parabolic connections on curves and the Riemann-Hilbert correspondence*. J. Algebraic Geom. **22** (2013), no. 3, 407–480.

- [16] M.-A. Inaba, K. Iwasaki, M.-H. Saito, *Moduli of stable parabolic connections, Riemann-Hilbert correspondence and geometry of Painlevé equation of type VI. I.* Publ. Res. Inst. Math. Sci. **42** (2006), no. 4, 987–1089.
- [17] M.-A. Inaba, K. Iwasaki, M.-H. Saito, *Moduli of stable parabolic connections, Riemann-Hilbert correspondence and geometry of Painlevé equation of type VI. II.* Moduli spaces and arithmetic geometry, 387–432, Adv. Stud. Pure Math., **45**, Math. Soc. Japan, Tokyo, 2006.
- [18] M.-A. Inaba, M.-H. Saito, *Moduli of unramified irregular singular parabolic connections on a smooth projective curve.* Kyoto J. Math. **53** (2013) 433–482.
- [19] H. Kawakami, A. Nakamura, H. Sakai, *Toward a classification of four-dimensional Painlevé-type equations.* Algebraic and geometric aspects of integrable systems and random matrices, 143–161, Contemp. Math., **593**, Amer. Math. Soc., Providence, RI, 2013.
- [20] H. Kimura, *The degeneration of the two-dimensional Garnier system and the polynomial Hamiltonian structure,* Ann. Mat. Pura Appl. (4) **155** (1989), 25–74.
- [21] F. Loray, M.-H. Saito, *Lagrangian fibrations in duality on moduli spaces of rank 2 logarithmic connections over the projective line.* Internat. Math. Res. Notices (2015), no. 4, 995–1043.
- [22] F. Loray, M.-H. Saito, C. Simpson. *Foliations on the moduli space of rank two connections on the projective line minus four points.* In Geometric and differential Galois theories, volume **27** of Sémin. Congr., pages 115–168. Soc. Math. France, Paris, 2012.
- [23] V. B. Mehta, C. S. Seshadri. *Moduli of vector bundles on curves with parabolic structures.* Math. Ann., **248** (3): 205–239, 1980.
- [24] K. Miyazaki *On compactifications of moduli of unramified irregular singular connections and Okamoto-Painlevé pairs.* Doctoral thesis, Kobe university (2013).
- [25] S. Mukai, *Finite generation of the nagata invariant rings in A-D-E cases,* RIMS Preprint, 1502, 2005.
- [26] M.-H. Saito, T. Takebe, H. Terajima, *Deformation of Okamoto-Painlevé pairs and Painlevé equations,* J. Algebraic Geom. **11** (2002), no. 2, 311–362. **220** (2001), 165–229.
- [27] K. Yokogawa, *Compactification of moduli of parabolic sheaves and moduli of parabolic Higgs sheaves.* Journal of Mathematics of Kyoto University **33**, no. 2 (1993): 451–504.
- [28] K. Yokogawa, *Infinitesimal deformation of parabolic Higgs sheaves.* Internat. J. Math. **6** (1995), no. 1, 125–148.

CENTER FOR MATHEMATICAL AND DATA SCIENCES, KOBE UNIV., 1-1 ROKKODAI-CHO, NADA-KU, KOBE, 657-8501, JAPAN

Email address: akomyo@math.kobe-u.ac.jp

UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE

Email address: frank.loray@univ-rennes1.fr

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KOBE UNIVERSITY, 1-1 ROKKODAI-CHO, NADA-KU, KOBE, 657-8501, JAPAN

Email address: mhsaito@math.kobe-u.ac.jp